

25 YEARS ANNIVERSARY
SOICT

HA NOI UNIVERSITY OF SCIENCE AND TECHNOLOGY
SCHOOL OF INFORMATION AND COMMUNICATION TECHNOLOGY



HA NOI UNIVERSITY OF SCIENCE AND TECHNOLOGY
SCHOOL OF INFORMATION AND COMMUNICATION TECHNOLOGY

FUNDAMENTALS OF OPTIMIZATION

Linear Programming

CONTENT

- Linear Programs
- Geometric approach
- Simplex method

Linear Programs

Standard form

$$f(x) = c_1x_1 + c_2x_2 + \dots + c_nx_n \rightarrow \max$$

$$a_{1,1}x_1 + a_{1,2}x_2 + \dots + a_{1,n}x_n \leq b_1$$

$$a_{2,1}x_1 + a_{2,2}x_2 + \dots + a_{2,n}x_n \leq b_2$$

...

$$a_{m,1}x_1 + a_{m,2}x_2 + \dots + a_{m,n}x_n \leq b_m$$

$$x_1, x_2, \dots, x_n \in R, x_1, x_2, \dots, x_n \geq 0$$

Linear Programs

- Convert general linear program forms into standard form
 - $f(x) \rightarrow \min \Leftrightarrow -f(x) \rightarrow \max$
 - $g(x) \geq b \Leftrightarrow -g(x) \leq -b$
 - $A = B \Leftrightarrow (A \leq B) \text{ and } (A \geq B)$
 - A variable $x_j \in \mathbb{R}$ can be represented by $x_j = x_j^+ - x_j^-$ where $x_j^+, x_j^- \geq 0$

Linear Programs

- Example: Convert general linear program forms into standard form

$$f(x_1, x_2) = 3x_1 + 2x_2 \rightarrow \min$$

$$2x_1 + x_2 \leq 7$$

$$x_1 + 2x_2 = 8$$

$$x_1 - x_2 \geq 2$$

$$x_1, x_2 \in \mathbb{R}, x_2 \geq 0$$

Linear Programs

- Example: Convert general linear program forms into standard form
 - Represent: $x_1 = x_1^+ - x_1^-$

$$f(x_1^+, x_1^-, x_2) = -3 x_1^+ + 3x_1^- - 2x_2 \rightarrow \max$$

$$2 x_1^+ - 2x_1^- + x_2 \leq 7$$

$$x_1^+ - x_1^- + 2x_2 \leq 8$$

$$-x_1^+ + x_1^- - 2x_2 \leq -8$$

$$-x_1^+ + x_1^- + x_2 \leq 2$$

$$x_1^+, x_1^-, x_2 \in \mathbb{R}, x_1^+, x_1^-, x_2 \geq 0$$

Geometric approach

- Constraints (inequalities) form a feasible region
- Optimal points will be one of the corners of the feasible region

Geometric approach

$$\begin{aligned} f(x_1, x_2) = \quad & 3x_1 + 2x_2 \rightarrow \max \\ & 2x_1 + x_2 \leq 7 \\ & x_1 + 2x_2 \leq 8 \\ & x_1 - x_2 \leq 2 \\ & x_1, x_2 \geq 0 \end{aligned}$$


Geometric approach

$$2x_1 + x_2 \leq 7$$

$$x_1 + 2x_2 \leq 8$$

$$x_1 - x_2 \leq 2$$

$$x_1, x_2 \geq 0$$




Geometric approach

$$2x_1 + x_2 \leq 7$$

$$x_1 + 2x_2 \leq 8$$

$$x_1 - x_2 \leq 2$$

$$x_1, x_2 \geq 0$$



Geometric approach

- Special cases
 - Problem does not have optimal solutions
 - Problem does not have feasible solutions

$$f(x_1, x_2) = 3x_1 + 2x_2 \rightarrow \max$$

$$-2x_1 - x_2 \leq -7$$

$$x_1 - x_2 \leq 2$$

$$x_1, x_2 \in \mathbb{R}, x_1, x_2 \geq 0$$

$$f(x_1, x_2) = 3x_1 + 2x_2 \rightarrow \max$$

$$2x_1 + x_2 \leq 7$$

$$-4x_1 - 2x_2 \leq -16$$

$$x_1, x_2 \in \mathbb{R}, x_1, x_2 \geq 0$$

Example

- A company must decide to make a plan to produce 2 products P1, P2.
 - The revenue received when selling 1 unit of P1 and P2 are respectively 5\$ and 7\$
 - The manufacturing cost when producing P1 and P2 are respectively 5\$ and 3\$
 - The storage cost in warehouses for 1 unit of P1 and P2 are respectively 2\$ and 3\$
- Compute the production plan so that
 - Total manufacturing cost is less than or equal to 200\$
 - Total storage cost is less than or equal to 150\$
 - Total revenue is maximal

Simplex method

$$f(x_1, x_2, \dots, x_n) = c_1x_1 + c_2x_2 + \dots + c_nx_n \rightarrow \max$$

$$a_{1,1}x_1 + a_{1,2}x_2 + \dots + a_{1,n}x_n \leq b_1$$

$$a_{2,1}x_1 + a_{2,2}x_2 + \dots + a_{2,n}x_n \leq b_2$$

. . .

$$a_{m,1}x_1 + a_{m,2}x_2 + \dots + a_{m,n}x_n \leq b_m$$

$$x_1, x_2, \dots, x_n \geq 0$$

Simplex method

Add slack variables $s_1, s_2, \dots, s_m \geq 0$

$$f(x_1, x_2, \dots, x_n, s_1, s_2, \dots, s_m) = c_1x_1 + c_2x_2 + \dots + c_nx_n \rightarrow \max$$

$$a_{1,1}x_1 + a_{1,2}x_2 + \dots + a_{1,n}x_n + s_1 = b_1$$

$$a_{2,1}x_1 + a_{2,2}x_2 + \dots + a_{2,n}x_n + 0 + s_2 + \dots = b_2$$

. . .

$$a_{m,1}x_1 + a_{m,2}x_2 + \dots + a_{m,n}x_n + 0 + \dots + s_m = b_m$$

$$x_1, x_2, \dots, x_n, s_1, s_2, \dots, s_m \geq 0$$

Simplex method

Consider the linear program under augmented form

$$f(x) = cx \rightarrow \max$$

$$Ax = b$$

$$x \geq 0$$

$$x = \begin{pmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{pmatrix}$$

$$c = (c_1, c_2, \dots, c_n)$$

$$A = \begin{pmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} \\ a_{2,1} & a_{2,2} & \dots & a_{2,n} \\ \dots & \dots & \dots & \dots \\ a_{m,1} & a_{m,2} & \dots & a_{m,n} \end{pmatrix}$$

$$b = \begin{pmatrix} b_1 \\ b_2 \\ \dots \\ b_m \end{pmatrix}$$

Denote $J = (1, 2, \dots, n)$, $I = (1, 2, \dots, m)$ - vector of indices

Basic feasible solutions

- Denote
 - $A(j)$ the j^{th} column of matrix A
 - If $J = (j_1, j_2, \dots, j_k)$ – vector of indices (we also use J as a set of indices), then $x_J = (x_{j_1}, \dots, x_{j_k})$, $A(J) = (A(j_1), \dots, A(j_k))$ submatrix of A composed by selecting columns $A(j_1), \dots, A(j_k)$
- Suppose $\text{rank}(A) = m$
- $J_B = (j_1, j_2, \dots, j_m)$ such that columns $A(j_1), \dots, A(j_m)$ are linear independent, J_N – vector of remaining indices of J (out of J_B)
- $B = A(J_B)$ - basic matrix
- $N = A(J_N)$
- $x = (x_B, x_N)$, $x_B = x(J_B) = (x_{j_1}, \dots, x_{j_m})$ such that $Bx = b$, $x_B = B^{-1} b$, $x_N = 0$

Basic feasible solutions

Example

$$f(x) = cx \rightarrow \max$$

$$Ax = b$$

$$x \geq 0$$

$$x = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix}$$

$$c = (3, 2, 0, 0, 0)$$

$$A = \begin{pmatrix} 2 & 1 & 1 & 0 & 0 \\ 1 & 2 & 0 & 1 & 0 \\ 1 & -1 & 0 & 0 & 1 \end{pmatrix}$$

$$b = \begin{pmatrix} 7 \\ 8 \\ 2 \end{pmatrix}$$

$J = (1,2,3,4,5)$ – vector of variable indices, $I = (1,2,3)$ – vector of constraint indices

$$J_B = (3,4,5), J_N = (1,2),$$

$$x_B = \begin{pmatrix} x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 7 \\ 8 \\ 2 \end{pmatrix} \quad x_N = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad N = \begin{pmatrix} 2 & 1 \\ 1 & 2 \\ 1 & -1 \end{pmatrix}$$

Simplex method

- Consider another feasible solution $x' = x + \Delta x$
 $\rightarrow Ax' = Ax \rightarrow A\Delta x = 0 \rightarrow B\Delta x_B + N\Delta x_N = 0 \rightarrow \Delta x_B = -B^{-1}N\Delta x_N$
 $\rightarrow \Delta f = c\Delta x = c_B\Delta x_B + c_N\Delta x_N = -(c_B B^{-1}N - c_N)\Delta x_N = -\Delta_N\Delta x_N =$
 $-\sum_{j \in J_N} \Delta_j \Delta x_j$, where $u = c_B B^{-1}$, $\Delta_N = uN - c_N$
- $\Delta x_j = x_j \geq 0, \forall j \in J_N$
- If $\Delta_j \geq 0, \forall j \in J_N$ then $\Delta f \leq 0$: x is a maximizer

Simplex method

- If there exists index $p \in J_N$ such that $\Delta_p < 0$, build another feasible solution $x' = x + \Delta x$ as follows
 - $\Delta x_p = \theta \geq 0$
 - $\Delta x_j = 0, \forall j \in J_N \setminus \{p\}$
 - $\Delta x_B = -B^{-1}N\Delta x_N = -\theta B^{-1}A(p), x'_B = x_B - \theta B^{-1}A(p),$
 - $\Delta f = -\theta \Delta_p$
- with $\theta > 0$ and small, we have $\Delta f > 0$: x is not an optimal solution because x' is better than x .
- If $B^{-1}A(p) \leq 0$, then $x'_B = x_B - \theta B^{-1}A(p) \geq 0, \forall \theta > 0$. It means that $f \rightarrow +\infty$ when $\theta \rightarrow +\infty$ (cannot find maximizer)

Simplex method

- Denote $B^{-1}A(p) = (x_{j_1,p}, x_{j_2,p}, \dots, x_{j_m,p})^\top$ and

$$\theta_i = \begin{cases} \frac{x_i}{x_{i,p}} & \text{if } x_{i,p} > 0 \\ +\infty, & \text{otherwise} \end{cases} \quad \forall i \in J_B$$

→ select $\theta = \theta_q = \min\{\theta_i \mid i \in J_B\}$ (we have $\theta < +\infty$, otherwise the objective function is unbounded)

Simplex method

```
 $J_B = (j_1, j_2, \dots, j_m)$  s.t.  $B = A(J_B)$  is a basic  
 $J_N$  – vector of remaining indices of  $J$  (out of  $J_B$ )  
While stop condition not reach{  
   $u = c_B B^{-1}$ ,  $\Delta_N = uN - c_N$ ,  $x_N = 0$ ,  $x_B = B^{-1}b$   
  if  $\Delta_N \geq 0$  then {  
    print('found optimal solution!'), break  
  } else {  
     $p = \arg\min\{\Delta_j \mid j \in J_N\}$   
     $Y = B^{-1}A(p)$   
    if  $Y \leq 0$  then {  
      print('objective is unbounded'), Return null  
    } else {  
       $q = \arg\min\{x_i/Y_i \mid i \in J_B \text{ s.t. } Y_i > 0\}$   
      remove  $q$  from  $J_B$  and add  $p$  to  $J_B$   
      remove  $p$  from  $J_N$  and add  $q$  to  $J_N$   
       $B = A(J_B)$ ,  $N = A(J_N)$   
    }  
  }  
}  
Return  $(x_B, x_N)$  where  $x_B = B^{-1}b$ ,  $x_N = 0$ 
```

Simplex method

Example

$$f(x_1, x_2) = 3x_1 + 2x_2 \rightarrow \max$$

$$2x_1 + x_2 \leq 7$$

$$x_1 + 2x_2 \leq 8$$

$$x_1 - x_2 \leq 2$$

$$x_1, x_2 \geq 0$$



Add 3 slack variables x_3, x_4, x_5

$$f(x_1, x_2) = 3x_1 + 2x_2 \rightarrow \max$$

$$2x_1 + x_2 + x_3 = 7$$

$$x_1 + 2x_2 + x_4 = 8$$

$$x_1 - x_2 + x_5 = 2$$

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

Simplex method

Init: $J_B = (3,4,5)$, $J_N = (1,2)$

$$x_B = \begin{pmatrix} x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 7 \\ 8 \\ 2 \end{pmatrix} \quad x_N = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$B = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad N = \begin{pmatrix} 2 & 1 \\ 1 & 2 \\ 1 & -1 \end{pmatrix}$$

$$f(x) = 0$$

Simplex method

Step 1: $\Delta_N = (-3 \quad -2) \rightarrow \text{select } p = 1$

$Y = \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} \rightarrow \text{select } q = 5, J_B = (1,3,4), J_N = (2,5)$

$$x_B = \begin{pmatrix} x_1 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \\ 6 \end{pmatrix} \quad x_N = \begin{pmatrix} x_2 \\ x_5 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$f(x) = 6$$

$$B = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix} \quad N = \begin{pmatrix} 1 & 0 \\ 2 & 0 \\ -1 & 1 \end{pmatrix}$$

$$B^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & -2 \\ 0 & 1 & -1 \end{pmatrix}$$

Simplex method

Step 2: $\Delta_N = (-5 \ 3) \rightarrow \text{select } p = 2$

$Y = \begin{bmatrix} -1 \\ 3 \\ 3 \end{bmatrix} \rightarrow \text{select } q = 3, J_B = (1,2,4), J_N = (3,5)$

$$x_B = \begin{bmatrix} x_1 \\ x_2 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 3 \end{bmatrix} \quad x_N = \begin{bmatrix} x_3 \\ x_5 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$f(x) = 11$$

$$B = \begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 1 & -1 & 0 \end{bmatrix} \quad N = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$$

$$B^{-1} = \begin{bmatrix} 1/3 & 0 & 1/3 \\ 1/3 & 0 & -2/3 \\ -1 & 1 & 1 \end{bmatrix}$$

Simplex method

Step 3: $\Delta_N = (5/3 \quad -1/3) \rightarrow \text{select } p = 5$

$Y = \begin{pmatrix} 1/3 \\ -2/3 \\ 1 \end{pmatrix} \rightarrow \text{select } q = 4, J_B = (1,2,5), J_N = (3,4)$

$$x_B = \begin{pmatrix} x_1 \\ x_2 \\ x_5 \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \\ 3 \end{pmatrix} \quad x_N = \begin{pmatrix} x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$f(x) = 12$$

$$B = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 1 & -1 & 1 \end{pmatrix} \quad N = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$B^{-1} = \begin{pmatrix} 2/3 & -1/3 & 0 \\ -1/3 & 2/3 & 0 \\ -1 & 1 & 1 \end{pmatrix}$$

Simplex method

Step 3: $\Delta_N = (4/3 \quad 1/3) > 0 \rightarrow \text{STOP, found optimal solution}$

$$x_B = \begin{pmatrix} x_1 \\ x_2 \\ x_5 \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \\ 3 \end{pmatrix} \quad x_N = \begin{pmatrix} x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$f(x) = 12$$

Simplex method: Tabular form

c_B	B	x_B	c_1	...	c_p	...	c_n	θ
			A_1	...	A_p	...	A_n	
c_{j_1}	A_{j_1}	x_{j_1}	$x_{j_1,1}$		$x_{j_1,p}$		$x_{j_1,n}$	θ_{j_1}
...
c_q	A_{i_0}	x_q	$x_{q,1}$		$x_{q,p}$		$x_{q,n}$	θ_q
...
c_{j_m}	A_{j_m}	x_{j_m}	$x_{j_m,1}$		$x_{j_m,p}$		$x_{j_m,n}$	θ_{j_m}
Δ			Δ_1	...	Δ_p	...	Δ_n	

Simplex method: Tabular form

c_B	B	x_B	c_1	...	c_p	...	c_n	θ
			A_1	...	A_p	...	A_n	
c_{j_1}	A_{j_1}	x_{j_1}	$x_{j_1,1}$		$x_{j_1,p}$		$x_{j_1,n}$	θ_{j_1}
...
c_q	A_{i_0}	x_q	$x_{q,1}$		$x_{q,p}$		$x_{q,n}$	θ_q
...
c_{j_m}	A_{j_m}	x_{j_m}	$x_{j_m,1}$		$x_{j_m,p}$		$x_{j_m,n}$	θ_{j_m}
Δ			Δ_1	...	Δ_p	...	Δ_n	

Simplex method: Tabular form

- Update schema
 - Update $x_{i,j}$: $x'_{i,j} = x_{i,j} - (x_{q,j} * x_{i,p}) / x_{q,p}, \forall i \in J_B \setminus \{q\}, \forall j \in J \setminus \{p\}$
 - $\Delta'_j = \Delta_j - (x_{q,j} * \Delta_p) / x_{q,p}, \forall j \in J \setminus \{p\}$
 - $J_B = J_B \setminus \{q\} \cup \{p\}$
 - $x'_q = x_q / x_{q,p}$
 - On row q : $x'_{q,j} = x_{q,j} / x_{q,p}, \forall j \in J$
 - On column p : $x'_{q,p} = 1, x'_{i,p} = 0, \forall i \in J_B \setminus \{q\}$
 - $\Delta_p = 0$

Simplex method: Tabular form

Example

$$f(x_1, x_2, x_3, x_4, x_5) = 3x_1 + 2x_2 \rightarrow \max$$

$$2x_1 + x_2 + x_3 = 7$$

$$x_1 + 2x_2 + x_4 = 8$$

$$x_1 - x_2 + x_5 = 2$$

$$x_1, x_2, x_3, x_4, x_5 \geq 0$$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	7	2	1	1	0	0	
0	A_4	8	1	2	0	1	0	
0	A_5	2	1	-1	0	0	1	
Δ			-3	-2	0	0	0	

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	7	2	1	1	0	0	
0	A_4	8	1	2	0	1	0	
0	A_5	2	1	-1	0	0	1	
Δ			-3	-2	0	0	0	

Select $p = 1$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	7	2	1	1	0	0	7/2
0	A_4	8	1	2	0	1	0	8
0	A_5	2	1	-1	0	0	1	2
Δ			-3	-2	0	0	0	

Select $p = 1$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	7	2	1	1	0	0	7/2
0	A_4	8	1	2	0	1	0	8
0	A_5	2	1	-1	0	0	1	2
Δ			-3	-2	0	0	0	

Select $p = 1$

Select $q = 5$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	3	2	3	1	0	-2	
0	A_4	6	1	3	0	1	-1	
0	A_5	2	1	-1	0	0	1	
Δ			-3	-5	0	0	3	

Update Δ , x_B , $x_{i,j}$ except row A_5 , and except columns 1

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	3	0	3	1	0	-2	
0	A_4	6	0	3	0	1	-1	
3	A_1	2	1	-1	0	0	1	
Δ			0	-5	0	0	3	

Replace A_5 by A_1 in the B , Update row corresponding to A_1 , and column 1

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	3	0	3	1	0	-2	
0	A_4	6	0	3	0	1	-1	
3	A_1	2	1	-1	0	0	1	
Δ			0	-5	0	0	3	

Select $p = 2$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	3	0	3	1	0	-2	1
0	A_4	6	0	3	0	1	-1	2
3	A_1	2	1	-1	0	0	1	$+\infty$
Δ			0	-5	0	0	3	

Select $p = 2$

Select $q = 3$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
0	A_3	3	0	3	1	0	-2	
0	A_4	3	0	3	-1	1	1	
3	A_1	3	1	-1	1/3	0	1/3	
Δ			0	-5	5/3	0	-1/3	

Update Δ , x_B , $x_{i,j}$ except row A_3 , and except columns 2

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
2	A_2	1	0	1	1/3	0	-2/3	
0	A_4	3	0	0	-1	1	1	
3	A_1	3	1	0	1/3	0	1/3	
Δ			0	0	5/3	0	-1/3	

Replace A_3 by A_2 in B , update x_B , $x_{i,j}$ on row A_3 , and columns 2

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
2	A_2	1	0	1	1/3	0	-2/3	$+\infty$
0	A_4	3	0	0	-1	1	1	3
3	A_1	3	1	0	1/3	0	1/3	9
Δ			0	0	5/3	0	-1/3	

Select $p = 5$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
2	A_2	1	0	1	1/3	0	-2/3	$+\infty$
0	A_4	3	0	0	-1	1	1	3
3	A_1	3	1	0	1/3	0	1/3	9
Δ			0	0	5/3	0	-1/3	

Select $p = 5$

Select $q = 4$

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
2	A_2	3	0	1	-1	2/3	-2/3	
0	A_4	3	0	0	-1	1	1	
3	A_1	2	1	0	2/3	-1/3	1/3	
Δ			0	0	4/3	1/3	-1/3	

Simplex method: Tabular form

c_B	B	x_B	3	2	0	0	0	θ
			A_1	A_2	A_3	A_4	A_5	
2	A_2	3	0	1	-1/3	2/3	0	
0	A_5	3	0	0	-1	1	1	
3	A_1	2	1	0	2/3	-1/3	0	
Δ			0	0	4/3	1/3	0	

STOP, found optimal solution!!!!



25 YEARS ANNIVERSARY
SOICT

VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG
SCHOOL OF INFORMATION AND COMMUNICATION TECHNOLOGY

**Thank you
for your
attentions!**

 soict.hust.edu.vn/  fb.com/groups/soict

