

GENERAL PHYSICS PH1110

Nam Le, Dr.

School of Engineering Physics

Hanoi University of Science and Technology

2. THERMODYNAMICS

2.2 FIRST LAW OF THERMODYNAMICS

- 1 STATEMENT OF THE FIRST LAW
- 2 WORK AND HEAT IN EQUILIBRIUM PROCESSES
- 3 EQUILIBRIUM PROCESSES OF IDEAL GASES

1. Statement of the first law

1. Statement of the first law

♣ The law was stated by **Rudolf Clausius** in 1850.

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

$$\Delta U = Q + W$$

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

$$\Delta U = Q + W$$

or

$$dU = \delta Q + \delta W$$

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

$$\Delta U = Q + W$$

or

$$dU = \delta Q + \delta W$$

- The change, $\Delta U = U_{\text{final}} - U_{\text{initial}}$,

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

$$\Delta U = Q + W$$

or

$$dU = \delta Q + \delta W$$

- The change, $\Delta U = U_{\text{final}} - U_{\text{initial}}$, is the same as the increase in internal energy.

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

$$\Delta U = Q + W$$

or

$$dU = \delta Q + \delta W$$

- The change, $\Delta U = U_{\text{final}} - U_{\text{initial}}$, is the same as the increase in internal energy.
- If Q' is the heat transferred from the system

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

$$\Delta U = Q + W$$

or

$$dU = \delta Q + \delta W$$

- The change, $\Delta U = U_{\text{final}} - U_{\text{initial}}$, is the same as the increase in internal energy.
- If Q' is the heat transferred from the system and W' is the work done by the system:

1. Statement of the first law

- ♣ The law was stated by **Rudolf Clausius** in 1850.

The change in internal energy of a system is equal to the heat added to the system plus the work done on the system.

$$\Delta U = Q + W$$

or

$$dU = \delta Q + \delta W$$

- The change, $\Delta U = U_{\text{final}} - U_{\text{initial}}$, is the same as the increase in internal energy.
- If Q' is the heat transferred from the system and W' is the work done by the system:

$$\Delta U + Q' + W' = 0$$

2. THERMODYNAMICS

2.2 FIRST LAW OF THERMODYNAMICS

- 1 STATEMENT OF THE FIRST LAW
- 2 WORK AND HEAT IN EQUILIBRIUM PROCESSES
- 3 EQUILIBRIUM PROCESSES OF IDEAL GASES

2. Work and heat in equilibrium processes

2. Work and heat in equilibrium processes

♣ Equilibrium states

2. Work and heat in equilibrium processes

♣ **Equilibrium states**

- For an ideal gas,

2. Work and heat in equilibrium processes

♣ Equilibrium states

- For an ideal gas, the variables N , p , V , and T are related by the EOS.

2. Work and heat in equilibrium processes

♣ Equilibrium states

- For an ideal gas, the variables N , p , V , and T are related by the EOS. If N is fixed, only two of them are independent.

2. Work and heat in equilibrium processes

♣ Equilibrium states

- For an ideal gas, the variables N , p , V , and T are related by the EOS. If N is fixed, only two of them are independent.
- In case p and V are chosen as independent variables,

2. Work and heat in equilibrium processes

♣ Equilibrium states

- For an ideal gas, the variables N , p , V , and T are related by the EOS. If N is fixed, only two of them are independent.
- In case p and V are chosen as independent variables, each state is represented by a point in the **PV diagram**.

2. Work and heat in equilibrium processes

♣ Equilibrium states

- For an ideal gas, the variables N , p , V , and T are related by the EOS. If N is fixed, only two of them are independent.
- In case p and V are chosen as independent variables, each state is represented by a point in the **PV diagram**. However, we can only represent equilibrium states.

2. Work and heat in equilibrium processes

♣ Equilibrium states

- For an ideal gas, the variables N , p , V , and T are related by the EOS. If N is fixed, only two of them are independent.
- In case p and V are chosen as independent variables, each state is represented by a point in the **PV diagram**. However, we can only represent equilibrium states.

Each variable has a single value in an equilibrium state, and its change between any two equilibrium states is single-valued.

2. Work and heat in equilibrium processes

♣ Equilibrium processes

2. Work and heat in equilibrium processes

♣ **Equilibrium processes**

An equilibrium process is a series of equilibrium states.

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.
- In practice, a quasi-static process (a sufficiently slow process)

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.
- In practice, a quasi-static process (a sufficiently slow process) can be considered as an equilibrium process.

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.
- In practice, a quasi-static process (a sufficiently slow process) can be considered as an equilibrium process.
- Work done in an equilibrium process (quasi-static work):

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.
- In practice, a quasi-static process (a sufficiently slow process) can be considered as an equilibrium process.
- Work done in an equilibrium process (quasi-static work):

$$dW = -pdV$$

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.
- In practice, a quasi-static process (a sufficiently slow process) can be considered as an equilibrium process.
- Work done in an equilibrium process (quasi-static work):

$$\boxed{dW = -pdV} \quad \rightarrow \quad W = - \int_{12} p dV$$

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.
- In practice, a quasi-static process (a sufficiently slow process) can be considered as an equilibrium process.
- Work done in an equilibrium process (quasi-static work):

$$\boxed{dW = -pdV} \quad \rightarrow \quad W = - \int_{12} p dV \quad (\text{path dependent})$$

2. Work and heat in equilibrium processes

♣ Equilibrium processes

An equilibrium process is a series of equilibrium states.

- Each equilibrium process is represented by a curve segment in the PV diagram.
- In practice, a quasi-static process (a sufficiently slow process) can be considered as an equilibrium process.
- Work done in an equilibrium process (quasi-static work):

$$\boxed{dW = -pdV} \quad \rightarrow \quad W = - \int_{12} p dV \quad (\text{path dependent})$$

This work is equal to the area under the pV curve.

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV$$

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{\overline{1a2b1}} p dV = \text{area limited by the path}$$

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = m c dT = \frac{m}{\mu} C dT$$

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = mcdT = \frac{m}{\mu} C dT \quad \rightarrow \quad Q = \int_{12} \frac{m}{\mu} C dT$$

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = m c dT = \frac{m}{\mu} C dT \quad \rightarrow \quad Q = \int_{12} \frac{m}{\mu} C dT$$

where c is the **specific heat capacity**

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = m c dT = \frac{m}{\mu} C dT \quad \rightarrow \quad Q = \int_{12} \frac{m}{\mu} C dT$$

where c is the **specific heat capacity** and C is the **molar heat capacity**.

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = mcdT = \frac{m}{\mu}CdT \quad \rightarrow \quad Q = \int_{12} \frac{m}{\mu}CdT$$

where c is the **specific heat capacity** and C is the **molar heat capacity**. The heat Q is path dependent.

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{1a2b1} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = mcdT = \frac{m}{\mu}CdT \quad \rightarrow \quad Q = \int_{12} \frac{m}{\mu}CdT$$

where c is the **specific heat capacity** and C is the **molar heat capacity**. The heat Q is path dependent.

- Definition of molar heat capacity:

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{\overline{1a2b1}} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = m c dT = \frac{m}{\mu} C dT \quad \rightarrow \quad Q = \int_{\widehat{12}} \frac{m}{\mu} C dT$$

where c is the **specific heat capacity** and C is the **molar heat capacity**. The heat Q is path dependent.

- Definition of molar heat capacity:

$$C \equiv \frac{\mu}{m} \frac{\delta Q}{dT}$$

2. Work and heat in equilibrium processes

- Work done in an equilibrium cycle (closed process):

$$W = - \int_{\overline{1a2b1}} p dV = \text{area limited by the path}$$

- Heat added to the system can be written in the form

$$\delta Q = m c dT = \frac{m}{\mu} C dT \quad \rightarrow \quad Q = \int_{\widehat{12}} \frac{m}{\mu} C dT$$

where c is the **specific heat capacity** and C is the **molar heat capacity**. The heat Q is path dependent.

- Definition of molar heat capacity:

$$C \equiv \frac{\mu}{m} \frac{\delta Q}{dT} \quad \rightarrow \quad \boxed{C = \frac{\mu}{m} \frac{\partial Q}{\partial T}}$$

2. THERMODYNAMICS

2.2 FIRST LAW OF THERMODYNAMICS

- 1 STATEMENT OF THE FIRST LAW
- 2 WORK AND HEAT IN EQUILIBRIUM PROCESSES
- 3 EQUILIBRIUM PROCESSES OF IDEAL GASES

3. Equilibrium processes of ideal gases

3. Equilibrium processes of ideal gases

♣ Isochoric process

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

• Equation:

$$\frac{p}{T} = \text{constant}$$

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:

$$\frac{p}{T} = \text{constant}$$

- Work:

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:

$$\frac{p}{T} = \text{constant}$$

- Work: $W = \int dW$

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\boxed{\frac{p}{T} = \text{constant}}$$

- Work:
$$W = \int dW = - \int p dV$$

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\boxed{\frac{p}{T} = \text{constant}}$$

- Work:
$$W = \int dW = - \int p dV \rightarrow \boxed{W = 0}$$

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\boxed{\frac{p}{T} = \text{constant}}$$

- Work:
$$W = \int dW = - \int p dV \rightarrow \boxed{W = 0}$$

- Heat:

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\boxed{\frac{p}{T} = \text{constant}}$$

- Work:
$$W = \int dW = - \int p dV \rightarrow \boxed{W = 0}$$

- Heat:
$$Q = \Delta U$$

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\boxed{\frac{p}{T} = \text{constant}}$$

- Work:
$$W = \int dW = - \int p dV \quad \rightarrow \quad \boxed{W = 0}$$

- Heat:
$$Q = \Delta U \quad \rightarrow \quad \boxed{Q = \frac{m}{\mu} \frac{i}{2} R \Delta T}$$

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\frac{p}{T} = \text{constant}$$
- Work:
$$W = \int dW = - \int p dV \quad \rightarrow \quad W = 0$$
- Heat:
$$Q = \Delta U \quad \rightarrow \quad Q = \frac{m}{\mu} \frac{i}{2} R \Delta T$$
- Molar heat capacity at constant volume:

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\boxed{\frac{p}{T} = \text{constant}}$$

- Work:
$$W = \int dW = - \int p dV \quad \rightarrow \quad \boxed{W = 0}$$

- Heat:
$$Q = \Delta U \quad \rightarrow \quad \boxed{Q = \frac{m}{\mu} \frac{i}{2} R \Delta T}$$

- Molar heat capacity at constant volume:

$$C_V = \frac{\mu}{m} \left(\frac{\partial Q}{\partial T} \right)_V$$

3. Equilibrium processes of ideal gases

♣ **Isochoric process** (constant-volume process, $V = \text{constant}$)

- Equation:
$$\frac{p}{T} = \text{constant}$$

- Work:
$$W = \int dW = - \int p dV \rightarrow W = 0$$

- Heat:
$$Q = \Delta U \rightarrow Q = \frac{m}{\mu} \frac{i}{2} R \Delta T$$

- Molar heat capacity at constant volume:

$$C_V = \frac{\mu}{m} \left(\frac{\partial Q}{\partial T} \right)_V \rightarrow C_V = \frac{i}{2} R$$

3. Equilibrium processes of ideal gases

♣ Isobaric process

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

3. Equilibrium processes of ideal gases

- ♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)
 - Equation:

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work:

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV$

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V$

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V \rightarrow W = -\frac{m}{\mu}R\Delta T$

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V \rightarrow W = -\frac{m}{\mu} R \Delta T$

- Heat:

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V \rightarrow W = -\frac{m}{\mu}R\Delta T$

- Heat: $Q = \Delta U - W$

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V \rightarrow W = -\frac{m}{\mu}R\Delta T$

- Heat: $Q = \Delta U - W \rightarrow Q = \frac{m}{\mu} \left(\frac{i}{2} + 1 \right) R\Delta T$

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V \rightarrow W = -\frac{m}{\mu}R\Delta T$

- Heat: $Q = \Delta U - W \rightarrow Q = \frac{m}{\mu} \left(\frac{i}{2} + 1 \right) R\Delta T$

- Molar heat capacity at constant pressure:

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V \rightarrow W = -\frac{m}{\mu}R\Delta T$

- Heat: $Q = \Delta U - W \rightarrow Q = \frac{m}{\mu} \left(\frac{i}{2} + 1 \right) R\Delta T$

- Molar heat capacity at constant pressure:

$$C_p = \frac{\mu}{m} \left(\frac{\partial Q}{\partial T} \right)_p$$

3. Equilibrium processes of ideal gases

♣ **Isobaric process** (constant-pressure process, $p = \text{constant}$)

- Equation:

$$\frac{V}{T} = \text{constant}$$

- Work: $W = -p \int dV = -p\Delta V \rightarrow W = -\frac{m}{\mu}R\Delta T$

- Heat: $Q = \Delta U - W \rightarrow Q = \frac{m}{\mu} \left(\frac{i}{2} + 1 \right) R\Delta T$

- Molar heat capacity at constant pressure:

$$C_p = \frac{\mu}{m} \left(\frac{\partial Q}{\partial T} \right)_p \rightarrow C_p = \frac{i+2}{2} R$$

3. Equilibrium processes of ideal gases

♣ Heat capacity ratio

3. Equilibrium processes of ideal gases

♣ **Heat capacity ratio** (adiabatic index)

3. Equilibrium processes of ideal gases

♣ **Heat capacity ratio** (adiabatic index)

- Definition:

3. Equilibrium processes of ideal gases

♣ **Heat capacity ratio** (adiabatic index)

- Definition:

$$\gamma = \frac{C_p}{C_v}$$

3. Equilibrium processes of ideal gases

♣ **Heat capacity ratio** (adiabatic index)

- Definition:

$$\gamma = \frac{C_p}{C_v}$$

- For ideal gases:

3. Equilibrium processes of ideal gases

♣ **Heat capacity ratio** (adiabatic index)

- Definition:

$$\gamma = \frac{C_p}{C_v}$$

- For ideal gases:

$$C_v = \frac{i}{2}R$$

$$C_p = \frac{i+2}{2}R$$

3. Equilibrium processes of ideal gases

♣ Heat capacity ratio (adiabatic index)

- Definition:

$$\gamma = \frac{C_p}{C_v}$$

- For ideal gases:

$$C_v = \frac{i}{2}R$$

$$C_p = \frac{i+2}{2}R$$

→

$$\gamma = \frac{i+2}{i}$$

3. Equilibrium processes of ideal gases

♣ **Heat capacity ratio** (adiabatic index)

- Definition:

$$\gamma = \frac{C_p}{C_v}$$

- For ideal gases:

$$C_v = \frac{i}{2}R$$

$$C_p = \frac{i+2}{2}R$$

→

$$\gamma = \frac{i+2}{i}$$

$$i = \frac{2}{\gamma - 1}$$

3. Equilibrium processes of ideal gases

♣ Heat capacity ratio (adiabatic index)

- Definition:

$$\gamma = \frac{C_p}{C_v}$$

- For ideal gases:

$$C_v = \frac{i}{2}R$$

$$C_p = \frac{i+2}{2}R$$

→

$$\gamma = \frac{i+2}{i}$$

$$i = \frac{2}{\gamma - 1}$$

→

$$C_v = \frac{R}{\gamma - 1}$$

3. Equilibrium processes of ideal gases

♣ Heat capacity ratio (adiabatic index)

- Definition:

$$\gamma = \frac{C_p}{C_v}$$

- For ideal gases:

$$C_v = \frac{i}{2}R$$

$$C_p = \frac{i+2}{2}R$$

→

$$\gamma = \frac{i+2}{i}$$

$$i = \frac{2}{\gamma - 1}$$

→

$$C_v = \frac{R}{\gamma - 1}$$

$$C_p = \frac{\gamma R}{\gamma - 1}$$

3. Equilibrium processes of ideal gases

♣ Isothermal process

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation:

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work:

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work: $W = - \int_{\widehat{12}} p dV$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work:
$$W = - \int_{\widehat{12}} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work: $W = - \int_{12} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$

$$W = \frac{m}{\mu} RT \ln \frac{V_1}{V_2}$$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work: $W = - \int_{V_1}^{V_2} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$

$$W = \frac{m}{\mu} RT \ln \frac{V_1}{V_2} = \frac{m}{\mu} RT \ln \frac{p_2}{p_1}$$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work: $W = - \int_{V_1}^{V_2} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$

$$W = \frac{m}{\mu} RT \ln \frac{V_1}{V_2} = \frac{m}{\mu} RT \ln \frac{p_2}{p_1}$$

- Heat:

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work: $W = - \int_{V_1}^{V_2} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$

$$W = \frac{m}{\mu} RT \ln \frac{V_1}{V_2} = \frac{m}{\mu} RT \ln \frac{p_2}{p_1}$$

- Heat: $\Delta T = 0$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work: $W = - \int_{V_1}^{V_2} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$

$$W = \frac{m}{\mu} RT \ln \frac{V_1}{V_2} = \frac{m}{\mu} RT \ln \frac{p_2}{p_1}$$

- Heat: $\Delta T = 0 \rightarrow \Delta U = 0$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

• Equation: $pV = \text{constant}$

• Work: $W = - \int_{V_1}^{V_2} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$

$$W = \frac{m}{\mu} RT \ln \frac{V_1}{V_2} = \frac{m}{\mu} RT \ln \frac{p_2}{p_1}$$

• Heat: $\Delta T = 0 \rightarrow \Delta U = 0 \rightarrow Q = -A.$

3. Equilibrium processes of ideal gases

♣ **Isothermal process** (constant-temperature, $T = \text{constant}$)

- Equation: $pV = \text{constant}$

- Work: $W = - \int_{V_1}^{V_2} p dV = - \frac{m}{\mu} RT \int_{V_1}^{V_2} \frac{dV}{V}$

$$W = \frac{m}{\mu} RT \ln \frac{V_1}{V_2} = \frac{m}{\mu} RT \ln \frac{p_2}{p_1}$$

- Heat: $\Delta T = 0 \rightarrow \Delta U = 0 \rightarrow Q = -A.$

$$Q = \frac{m}{\mu} RT \ln \frac{V_2}{V_1} = \frac{m}{\mu} RT \ln \frac{p_1}{p_2}$$

3. Equilibrium processes of ideal gases

♣ Adiabatic process

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

3. Equilibrium processes of ideal gases

- ♣ **Adiabatic process** (no heat transferred, $Q = 0$)
 - Equation:

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

- Equation:

$$\delta Q = 0$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

● Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT = \frac{1}{\gamma - 1} (dpV + p dV)$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT = \frac{1}{\gamma - 1} (dpV + p dV)$$

and $\delta W = -p dV$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT = \frac{1}{\gamma - 1} (dpV + p dV)$$

$$\underline{\text{and}} \quad \delta W = -p dV \quad \rightarrow \quad \frac{1}{\gamma - 1} (dpV + p dV) = -p dV$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT = \frac{1}{\gamma - 1} (dpV + p dV)$$

and $\delta W = -p dV \quad \rightarrow \quad \frac{1}{\gamma - 1} (dpV + p dV) = -p dV$

$$\frac{dp}{p} + \gamma \frac{dV}{V} = 0$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT = \frac{1}{\gamma - 1} (dpV + p dV)$$

and $\delta W = -p dV \quad \rightarrow \quad \frac{1}{\gamma - 1} (dpV + p dV) = -p dV$

$$\frac{dp}{p} + \gamma \frac{dV}{V} = 0 \quad \rightarrow \quad \boxed{pV^\gamma = \text{constant}}$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT = \frac{1}{\gamma - 1} (dpV + p dV)$$

and $\delta W = -p dV \quad \rightarrow \quad \frac{1}{\gamma - 1} (dpV + p dV) = -p dV$

$$\frac{dp}{p} + \gamma \frac{dV}{V} = 0 \quad \rightarrow \quad \boxed{pV^\gamma = \text{constant}}$$

• The PV diagram of an adiabatic process is similar to that of an isothermal process,

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Equation:

$$\delta Q = 0 \quad \rightarrow \quad \delta W = dU = \frac{m}{\mu} \frac{i}{2} R dT = \frac{1}{\gamma - 1} (dpV + p dV)$$

and $\delta W = -p dV \quad \rightarrow \quad \frac{1}{\gamma - 1} (dpV + p dV) = -p dV$

$$\frac{dp}{p} + \gamma \frac{dV}{V} = 0 \quad \rightarrow \quad \boxed{pV^\gamma = \text{constant}}$$

• The PV diagram of an adiabatic process is similar to that of an isothermal process, but the magnitude of its gradient is greater.

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

- Work:

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

- Work:

$$W = - \int_{\widehat{12}} p dV$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma}$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma + 1}$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

● Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma+1}$$

$$W = \frac{p_2 V_2 - p_1 V_1}{\gamma - 1}$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

● Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma + 1}$$

$$W = \frac{p_2 V_2 - p_1 V_1}{\gamma - 1}$$

In terms of change in temperature:

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

● Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma + 1}$$

$$W = \frac{p_2 V_2 - p_1 V_1}{\gamma - 1}$$

In terms of change in temperature:

$$\frac{pV}{\gamma - 1}$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

● Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma + 1}$$

$$W = \frac{p_2 V_2 - p_1 V_1}{\gamma - 1}$$

In terms of change in temperature:

$$\frac{pV}{\gamma - 1} = pV \frac{C_V}{R}$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

● Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma+1}$$

$$W = \frac{p_2 V_2 - p_1 V_1}{\gamma - 1}$$

In terms of change in temperature:

$$\frac{pV}{\gamma - 1} = pV \frac{C_V}{R} = C_V \frac{m}{\mu} T$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

• Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma+1}$$

$$W = \frac{p_2 V_2 - p_1 V_1}{\gamma - 1}$$

In terms of change in temperature:

$$\frac{pV}{\gamma - 1} = pV \frac{C_V}{R} = C_V \frac{m}{\mu} T$$

$$W = \frac{m}{\mu} C_V (T_2 - T_1)$$

3. Equilibrium processes of ideal gases

♣ **Adiabatic process** (no heat transferred, $Q = 0$)

● Work:

$$W = - \int_{\widehat{12}} p dV = -p_1 V_1^\gamma \int_{V_1}^{V_2} \frac{dV}{V^\gamma} = -p_1 V_1^\gamma \frac{V_2^{-\gamma+1} - V_1^{-\gamma+1}}{-\gamma+1}$$

$$W = \frac{p_2 V_2 - p_1 V_1}{\gamma - 1}$$

In terms of change in temperature:

$$\frac{pV}{\gamma - 1} = pV \frac{C_V}{R} = C_V \frac{m}{\mu} T$$

$$W = \frac{m}{\mu} C_V (T_2 - T_1) = \frac{m}{\mu} C_V \Delta T$$