

# Groundwater flow modeling: uncertain boundary conditions and their impact on the forecast

---

Emanuel Huber<sup>1</sup>, Jef Caers<sup>1</sup>, Peter Huggenberger<sup>2</sup>, Thomas Vienken<sup>3</sup>,  
Manuel Kreck<sup>3</sup>, Peter Dietrich<sup>3,4</sup>

<sup>1</sup> Department of Geological Sciences, Stanford University

<sup>2</sup> Applied and Environmental Geology, University of Basel

<sup>3</sup> UFZ-Helmholtz Centre for Environmental Research, Leipzig

<sup>4</sup> Eberhard Karls University of Tübingen

## Subsurface flow

## An interplay between

- subsurface heterogeneity
- boundary conditions (inflow/outflow, pressure)





## Objectives


- Uncertainty model for boundary conditions
- Uncertainty quantification of forecast




# Objectives

- Uncertainty model for boundary conditions
- Uncertainty quantification of forecast

## Synthetic case study





**Prediction** = bacteria concentration for the next 10 days



# Objectives

- Uncertainty model for boundary conditions
- Uncertainty quantification of forecast

## Synthetic case study





# Uncertain boundary conditions

## River water–groundwater interaction

- river bed conductivity  $C_{\text{riv}}$   $\rightarrow$  spatially constant, uniform prior.
- river heads ( $h_{\text{riv}}$ )  $\rightarrow$  measured (assumed to be known everywhere)

# Uncertain boundary conditions

## River water–groundwater interaction

- river bed conductivity  $C_{\text{riv}}$   $\rightarrow$  spatially constant, uniform prior.
- river heads ( $h_{\text{riv}}$ )  $\rightarrow$  measured (assumed to be known everywhere)

## Specified heads

Interpolation: space-time Gaussian process conditioned to measured groundwater heads

- Matérn (space) and Gaussian (time) covariance functions
- (unknown) linear mean function
- no-flow boundary conditions (specified derivatives)

# Uncertain boundary conditions

## River water–groundwater interaction

- river bed conductivity  $C_{riv}$   $\rightarrow$  spatially constant, uniform prior.
- river heads ( $h_{riv}$ )  $\rightarrow$  measured (assumed to be known everywhere)

## Specified heads

Interpolation: space-time Gaussian process conditioned to measured groundwater heads


- Matérn (space) and Gaussian (time) covariance functions
- (unknown) linear mean function
- no-flow boundary conditions (specified derivatives)

But river and groundwater heads  
are unknown for the next 10 days!

# Uncertain future boundary conditions

## Convolution (\*) model

Strong relationship between precipitation and river heads




river heads



# Uncertain future boundary conditions

## Convolution (\*) model


Strong relationship between precipitation and river heads




# Uncertain future boundary conditions

## Convolution (\*) model

Strong relationship between precipitation and river heads




Uncertain weather forecast → random sampling




# Uncertain future boundary conditions

## Convolution (\*) model

Strong relationship between precipitation and river heads




Uncertain weather forecast  $\rightarrow$  random sampling




# Uncertain future boundary conditions

## Convolution (\*) model

Strong relationship between precipitation and river heads



Uncertain weather forecast  $\rightarrow$  random sampling






# Uncertain future boundary conditions

## Convolution (\*) model

Strong relationship between precipitation and river heads




Uncertain weather forecast  $\rightarrow$  random sampling




# Uncertain future boundary conditions

## Convolution (\*) model

Strong relationship between precipitation and river heads



Uncertain weather forecast  $\rightarrow$  random sampling



☞ Same approach for the groundwater heads

# Spatial uncertainty

## Subsurface heterogeneity uncertainty

Hydraulic conductivity

→ Gaussian random field (Matérn covariance function)

Porosity, specific storage, specific yield

→ spatially constant, uniform prior

# Impact of uncertain boundary conditions

## Distance-based general sensitivity analysis

1. Sample 1000 realisations and compute the model response
2. Cluster the realisations based on the model response
3. For each parameter: compare the parameter distribution in each cluster with the global parameter distribution
4. If the distance between both distributions is significant, then the parameter is influential (on the model response)

# Impact of uncertain boundary conditions

## Distance-based general sensitivity analysis

1. Sample 1000 realisations and compute the model response
2. Cluster the realisations based on the model response
3. For each parameter:  
compare the parameter distribution in each cluster with the global parameter distribution
4. If the distance between both distributions is significant, then the parameter is influential (on the model response)

# Impact of uncertain boundary conditions

## Distance-based general sensitivity analysis

1. Sample 1000 realisations and compute the model response
2. Cluster the realisations based on the model response
3. For each parameter: compare the parameter distribution in each cluster with the global parameter distribution
4. If the distance between both distributions is significant, then the parameter is influential (on the model response)

# Impact of uncertain boundary conditions


## Distance-based general sensitivity analysis

1. Sample 1000 realisations and compute the model response
2. Cluster the realisations based on the model response
3. For each parameter: compare the parameter distribution in each cluster with the global parameter distribution
4. If the distance between both distributions is significant, then the parameter is influential (on the model response)

# Impact of uncertain boundary conditions

## Distance-based general sensitivity analysis

1. Sample 1000 realisations and compute the model response
2. Cluster the realisations based on the model response
3. For each parameter: compare the parameter distribution in each cluster with the global parameter distribution
4. If the distance between both distributions is significant, then the parameter is influential (on the model response)







# Statistical prediction

Data (1000 realisations)

groundwater heads (day 1 - day 31)

7 wells  $\times$  31 time steps




# Statistical prediction

Data (1000 realisations)



groundwater heads (day 1 - day 31)

7 wells  $\times$  31 time steps



tracer concentration (day 1 - day 5)

7 wells  $\times$  500 time steps





# Statistical prediction

## Forecast (1000 realisations)

bacteria concentration (day 1 - day 31)

1 wells  $\times$  10 time steps



# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform  
**data**

**prediction**

# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform

**data**




basis function expansion

Time-series  $d(t)$  approximated by a linear combination of B-splines  $\Phi_j(t)$   
$$d(t) \approx \sum_j c_j \Phi_j(t)$$

**prediction**

# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform




functional version of PCA  
summations change into integrations

**prediction**

# Statistical prediction – Method


Dimension reduction + CCA + linear regression + backtransform



**prediction**

# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform





**prediction**




# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform



# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform



find two bases **A** and **B** in which the correlation matrix between the variables is diagonal and the correlations on the diagonal are maximized.

$$\mathbf{U} = \mathbf{XA}$$

$$\mathbf{V} = \mathbf{YB}$$

with



$(\mathbf{U}, \mathbf{V})$  = canonical variables

$\mathbf{X}$  = reduced data variable

$\mathbf{Y}$  = reduced forecast variable



# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform





# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform



# Statistical prediction – Method

Dimension reduction + CCA + linear regression + backtransform




# Statistical prediction – Results

## Canonical correlation space

tracer data (99.9% of variance)

data ( $1000 \times 64$ ), forecast ( $1000 \times 7$ )







# Statistical prediction – Results

## Canonical correlation space





tracer data (99.9% of variance)

data ( $1000 \times 64$ ), forecast ( $1000 \times 7$ )



head data (99.9% of variance)


data ( $1000 \times 16$ ), forecast ( $1000 \times 7$ )




# Statistical prediction – Results

## Prediction

tracer data



head data




More accurate results with head data (larger log predictive density)




# Statistical prediction – Results

## Decision

tracer data



head data



👉 Turn off the drinking water extraction well!

# Conclusion

- model of uncertain boundary conditions
- distance-based general sensitivity analysis
  - importance of boundary conditions (specified heads)
- statistical prediction
  - circumvent classical inversion
  - relevance of data for prediction

# Conclusion

- model of uncertain boundary conditions
- distance-based general sensitivity analysis
  - importance of boundary conditions (specified heads)
- statistical prediction
  - circumvent classical inversion
  - relevance of data for prediction

## Further research

- model for riverbed conductivity (spatial and temporal)
- statistical prediction for designing monitoring network
- use statistical prediction for resampling