

Perfect state transfer in graphs with an involution

C.M. van Bommel

May 16, 2021

1 Preliminaries

Lemma 1. [4] Let G be a (weighted) graph with an involution σ , which respects loops and edge weights. Then the characteristic polynomial of the (weighted) adjacency matrix A of G factors into two factors Π_+ and Π_- which are, respectively, the characteristic polynomials of $A_+ := \begin{bmatrix} A' + A_\sigma & A_\delta \\ 2A_\delta^T & A_S \end{bmatrix}$ and $A_- := A' - A_\sigma$.

Furthermore, there is an eigenbasis for A consisting of vectors that take the form $\begin{bmatrix} a & a & b \end{bmatrix}^T$ and $\begin{bmatrix} c & -c & 0 \end{bmatrix}^T$, where $\begin{bmatrix} a & b \end{bmatrix}^T$ is an eigenvector for A_+ , and c an eigenvector for A_- .

Theorem 2 (Weyl's Inequality). If A and B are Hermitian, then $\lambda_{i+j-1}(A+B) \leq \lambda_i(A) + \lambda_j(B)$, and equivalently, $\lambda_{i+j-n}(A+B) \geq \lambda_i(A) + \lambda_j(B)$.

Lemma 3. [3] Let u, v be vertices of G , and H the Hamiltonian. Then perfect state transfer from u to v occurs at some time if and only if the following two conditions are satisfied:

1. Every eigenvector x of H satisfies either $x(u) = x(v)$ or $x(u) = -x(v)$.
2. If $\{\lambda_i\}$ are the eigenvalues with $x(u) = x(v)$, and $\{\mu_j\}$ are the eigenvalues for the eigenvectors with $x(u) = -x(v)$, and $x(u)$ and $x(v)$ are non-zero, then there exists some time T such that

$$e^{iT\lambda_1} = e^{iT\lambda_i} = -e^{iT\mu_j}$$

for all i, j .

Corollary 4. [3] Using the notation of Lemma 3, if perfect state transfer occurs from u to v , then

$$\frac{\lambda_i - \lambda_j}{\lambda_k - \lambda_\ell} \in \mathbb{Q}$$

and

$$\frac{\lambda_i - \mu_j}{\lambda_k - \lambda_\ell} = \frac{\text{odd}}{\text{even}}$$

for all i, j, k, ℓ .

Theorem 5. [2] Suppose X is a graph with at least two vertices. Then X is periodic at a if and only if either:

1. The eigenvalues in the eigenvalue support of a are integers.
2. There is a square-free integer Δ , the eigenvalues in the eigenvalue support of a are quadratic integers in $\mathbb{Q}(\sqrt{\Delta})$, and the difference of any two eigenvalues in the eigenvalue support of a is an integer multiple of $\sqrt{\Delta}$.

Theorem 6. [1] There is no perfect state transfer on the double star graph $S_{k,\ell}$.

2 Result

Theorem 7. *Let X be a graph, v a vertex of X , and let Y be the graph formed from two copies of X by adding an edge between the copies of v . If $|V(X)| \geq 2$, then Y does not admit perfect state transfer between the copies of v .*

Proof. Suppose Y admits perfect state transfer between the copies of v . By construction, Y has an involution σ mapping each vertex to the corresponding vertex in the other copy of X . Hence, by Lemma 1, the eigenvalues of $A(Y)$ are the eigenvalues of $A(X) + e_v e_v^T$ and $A(X) - e_v e_v^T$.

Let $\{x_i\}$ be an eigenbasis for $A(X) + e_v e_v^T$ such that the number of eigenvectors for which $x_i(v) = 0$ is maximized. Then every such eigenvector is an eigenvector of $A(X) - e_v e_v^T$. We can extend this set of eigenvectors to an eigenbasis $\{y_j\}$ of $A(X) - e_v e_v^T$ such that the number of eigenvectors for which $y_j(v) = 0$ is maximized. If there is an eigenvector $y_j \notin \{x_i\}$ such that $y_j(v) = 0$, then y_j is an eigenvector of $A(X) + e_v e_v^T$, contradicting maximality. Hence, $A(X) + e_v e_v^T$ and $A(X) - e_v e_v^T$ have the same number of eigenvalues in the eigenvalue support of v , and the remaining eigenvalues (with multiplicity) are both eigenvalues of $A(X) + e_v e_v^T$ and $A(X) - e_v e_v^T$.

Let $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ be the eigenvalues of $A(X) + e_v e_v^T$ in the support of v and $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$ be the eigenvalues of $A(X) - e_v e_v^T$ in the support of v . It follows from Weyl's Inequality (Theorem 2) that $\lambda_i \geq \mu_i$ for $1 \leq i \leq n$. Moreover, it follows from Corollary 4 that $\lambda_i \neq \mu_i$. Finally, it follows from Theorem 5 that $\lambda_i \geq \mu_i + 1$. Therefore, if $\{\theta_k\}$ is the multiset of remaining eigenvalues of $A(X) \pm e_v e_v^T$ not in the support of v , we have

$$1 = \text{tr}(A(X) + e_v e_v^T) = \sum_{i=1}^n \lambda_i + \sum_k \theta_k \geq \sum_{i=1}^n (\mu_i + 1) + \sum_k \theta_k = \text{tr}(A(X) - e_v e_v^T) + n = n - 1,$$

whence $n \leq 2$.

Suppose $n = 2$, then the inequality holds with equality, and we have $\lambda_i = \mu_i + 1$. As the covering radius of v is at most 1, v is a universal vertex. Now, there exists an eigenbasis of $A(X)$ such that $|V(X)| - 2$ of the vectors x are such that $x(v) = 0$. Then each such eigenvector x is an eigenvector of $X \setminus v$ and $x \cdot \mathbf{1} = 0$. Thus, the remaining eigenvector of $X \setminus v$ is $\mathbf{1}$, so $X \setminus v$ is regular; we assume of degree k .

It follows that if $m = |V(X \setminus v)|$, then λ_1, λ_2 are eigenvalues of the quotient matrix

$$\begin{bmatrix} k & \sqrt{m} \\ \sqrt{m} & 1 \end{bmatrix}$$

and μ_1, μ_2 are eigenvalues of the quotient matrix

$$\begin{bmatrix} k & \sqrt{m} \\ \sqrt{m} & -1 \end{bmatrix}.$$

Hence, we have

$$\begin{aligned} \mu_1 + \mu_2 &= k - 1 \\ \mu_1 \mu_2 &= -k - m \\ \lambda_1 \lambda_2 &= k - m \\ (\mu_1 + 1)(\mu_2 + 1) &= k - m \\ \mu_1 \mu_2 + \mu_1 + \mu_2 + 1 &= k - m \\ -k - m + k - 1 + 1 &= k - m \\ 0 &= k, \end{aligned}$$

so $X \setminus v = \overline{K}_m$ and therefore Y is a double star, which contradicts Theorem 6.

Therefore, $n = 1$, so $X = K_1$, which completes the proof. \square

References

- [1] X. Fan, C. Godsil. Pretty good state transfer on double stars. *Linear Algebra and its Applications*, 438: 2346–2358, 2013.
- [2] C. Godsil. Periodic Graphs. *Electronic Journal of Combinatorics*, 18(1): P23, 2011.
- [3] M. Kempton, G. Lippner, S.-T. Yau. Perfect state transfer on graphs with a potential. *Quantum Information and Computation*, 17 (3 & 4):0303–0327, 2017.
- [4] M. Kempton, G. Lippner, and S.-T. Yau. Pretty good state transfer in graphs with an involution. arXiv:1702.07000, 2017.