

lab10: Structural Bioinformatics

Ebony Michelle Argaez (PID: A59026556)

Uniprot has 251600,768 proteins

```
round(183201/251600768*100, 2)
```

```
[1] 0.07
```

```
string <-c("10", "100", 1, "1,000")  
as.numeric(string) + 1
```

Warning: NAs introduced by coercion

```
[1] 11 101 2 NA
```

```
string<- gsub(",", "", string)  
  
##my function  
remove_commas <- function(df) {  
  df[] <-lapply(df, function(x) gsub(",", "", x))  
  return(df)  
}  
  
remove_commas(string)
```

```
[[1]]  
[1] "10"
```

```
[[2]]  
[1] "100"
```

```
[[3]]  
[1] "1"
```

```
[[4]]  
[1] "1000"
```

```
#Barry's function  
rm_comma <-function(x){  
  as.numeric(gsub(",", "", x))  
}
```

1:Introduction to the RCSB Protein Data Bank (PDB)

PDB statistics

```
stats<-(read.csv("Data Export Summary.csv", row.names = 1))  
  
pdbstats<-apply(stats, 2, rm_comma)  
  
rownames(pdbstats) <-rownames(stats)  
pdbstats
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	158844	11759	12296	197	73	32
Protein/Oligosaccharide	9260	2054	34	8	1	0
Protein/NA	8307	3667	284	7	0	0
Nucleic acid (only)	2730	113	1467	13	3	1
Other	164	9	32	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	183201					
Protein/Oligosaccharide	11357					
Protein/NA	12265					
Nucleic acid (only)	4327					
Other	205					
Oligosaccharide (only)	22					

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

93.16%

```
totals<-apply(pdbstats, 2, sum)

round(totals/totals["Total"]*100,2)
```

X.ray	EM	NMR	Multiple.methods
84.83	8.33	6.68	0.11
Neutron	Other	Total	
0.04	0.02	100.00	


```
84.83+8.33
```

[1] 93.16

Skipping Q2-3 >Q2: What proportion of structures in the PDB are protein?

Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?

2. Visualizing the HIV-1 protease structure



Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

The resolution is too low. It's 2Å.

Q5: There is a critical “conserved” water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have

308

Q6: Generate and save a figure clearly showing the two distinct chains of HIV-protease along with the ligand. You might also consider showing the catalytic residues ASP 25 in each chain and the critical water (we recommend “Ball & Stick” for these side-chains). Add this figure to your Quarto document.

adding a nicer picture, colored by secondary structure with catalytic site ASP 25 shown in each chain along with MK1 drug.



Figure 1: Lovely image

Using the bio3d Package

```
library(bio3d)

pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

Call: read.pdb(file = "1hsg")

```
Total Models#: 1
  Total Atoms#: 1686,  XYZs#: 5058  Chains#: 2  (values: A B)

Protein Atoms#: 1514  (residues/Calpha atoms#: 198)
Nucleic acid Atoms#: 0  (residues/phosphate atoms#: 0)

Non-protein/nucleic Atoms#: 172  (residues: 128)
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

Protein sequence:

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

Q7: How many amino acid residues are there in this pdb object?

198

Q8: Name one of the two non-protein residues?

HOH(127), MK(1)

Q9: How many protein chains are in this structure?

2

```
attributes(pdb)
```

```
$names
```

```
[1] "atom" "xyz" "seqres" "helix" "sheet" "calpha" "remark" "call"
```

```
$class
```

```
[1] "pdb" "sse"
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

```
head(pdb$atom$resid)
```

```
[1] "PRO" "PRO" "PRO" "PRO" "PRO" "PRO"
```

```
aa321(pdb$atom$resid[pdb$calpha])
```

```
[1] "P" "Q" "I" "T" "L" "W" "Q" "R" "P" "L" "V" "T" "I" "K" "I" "G" "G" "Q"
[19] "L" "K" "E" "A" "L" "L" "D" "T" "G" "A" "D" "D" "T" "V" "L" "E" "E" "M"
[37] "S" "L" "P" "G" "R" "W" "K" "P" "K" "M" "I" "G" "G" "I" "G" "G" "F" "I"
[55] "K" "V" "R" "Q" "Y" "D" "Q" "I" "L" "I" "E" "I" "C" "G" "H" "K" "A" "I"
[73] "G" "T" "V" "L" "V" "G" "P" "T" "P" "V" "N" "I" "I" "G" "R" "N" "L" "L"
[91] "T" "Q" "I" "G" "C" "T" "L" "N" "F" "P" "Q" "I" "T" "L" "W" "Q" "R" "P"
```

```
[109] "L" "V" "T" "I" "K" "I" "G" "G" "Q" "L" "K" "E" "A" "L" "L" "D" "T" "G"
[127] "A" "D" "D" "T" "V" "L" "E" "E" "M" "S" "L" "P" "G" "R" "W" "K" "P" "K"
[145] "M" "I" "G" "G" "I" "G" "G" "F" "I" "K" "V" "R" "Q" "Y" "D" "Q" "I" "L"
[163] "I" "E" "I" "C" "G" "H" "K" "A" "I" "G" "T" "V" "L" "V" "G" "P" "T" "P"
[181] "V" "N" "I" "I" "G" "R" "N" "L" "L" "T" "Q" "I" "G" "C" "T" "L" "N" "F"
```

Predicting functional motions of a single structure

Run a Normal Mode analysis (NMA) - a bioinformatics method to predict functional motions

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file


PDB has ALT records, taking A only, rm.alt=TRUE

```
modes<-nma(adk)
```

Building Hessian... Done in 0.015 seconds.

Diagonalizing Hessian... Done in 0.272 seconds.

```
plot(modes)
```

```
mktrj(modes, pdb=adk, file="modes.pdb")
```