

# 1 Phosphorus flows mapping and economic analysis for its recovery in 2 the province of Ontario, Canada

3 Université Laval team<sup>a,\*</sup>, McGill University team<sup>b</sup>, University of Waterloo team<sup>c</sup>

4 <sup>a</sup>*Université Laval*  
5 <sup>b</sup>*McGill University*  
6 <sup>c</sup>*University of Waterloo*

---

## 7 Abstract

Phosphorus is a key non-renewable element for food production and other economic activities. Phosphorus is used intensively in the food production and consumption system, however, its sustained emissions over time lead to nutrient pollution and eutrophication of ecosystems. In this work, we map the phosphorus flows through Ontario's economic sectors, and we identify the phosphorus recovery opportunities and recycling potential. These mainly target flows associated with food production and processing, including wastewater and food waste. Up to 86% of phosphorus imports for food production could be covered by recycled phosphorus, with an average recovery cost of 36 EUR/kg. This is lower than the estimated economic losses derived from the release of phosphorus into the environment, but significantly higher than fossil-based phosphorus products. Additionally, there exist a wide variation on phosphorus recovery costs from different streams, which suggest the need of exploring cooperative approaches for effective phosphorus recovery at regional scale.

---

## 8 1. Introduction

9 Phosphorus is an essential for production of food which has been intensively used for crop and  
10 livestock production since the development of synthetic fertilizers in the XIX and XX centuries  
11 ([Samreen and Kausar, 2019](#)). The combination of synthetic fertilizers with other modern intensive  
12 agricultural techniques have increased the productivity of agriculture and farming industries ([Pin-  
13 gali, 2012](#)). However, the intensive use of fertilizers in agriculture has resulted in the over-application  
14 of phosphorus ([Reid and Schneider, 2019](#)), while the run of intensive livestock production facilities,

---

\*Corresponding author  
Email address: [edgar.martin-hernandez.1@ulaval.ca](mailto:edgar.martin-hernandez.1@ulaval.ca) (Université Laval team)

also known as concentrated animal feeding operations (CAFOs) ([U.S. Department of Agriculture, 2011](#)), result in important difficulties in the management of the large amounts of manure produced, which is often spread in lands in the vicinity of CAFOs, which also leads to the accumulation of phosphorus in soil. Soil acts as a phosphorus reservoir ([Ehlert et al., 2003](#)), building-up a legacy P that can be used for future crops, but also can be transported to waterbodies by erosion and runoff leading to the eutrophication of aquatic ecosystems.

The dual dimension of the anthropogenic use of phosphorus, i.e., its key role in the food production system and the negative environmental impacts associated with the phosphorus used in intensive agricultural techniques, as it has been stated by the United Nations Environment Assembly in the resolution UNEP/EA.5/Res.2 ([United Nations Environment Programme, 2022](#)). An additional factor to be considered for addressing the phosphorus challenge is the non-renewable nature of phosphorus, since the phosphorus consumed is not replenished by natural means at human time scale, and there is currently no known synthetic substitute for this material ([Cordell et al., 2009](#)). Since the global phosphorus reserves are concentrated in a few number of regions, the supply of phosphorus from a limited number of global supply chains lacks resiliency and it has been proven that it can be globally disrupted by regional events and conflicts, resulting geopolitical tensions ([Food and Agriculture Organization of the United Nations, 2022](#)). As a consequence, the recovery and recycling of phosphorus is not just a desirable but also a necessary approach to assure a sustainable, reliable, and sovereign food production system.

Although the main uses of phosphorus are in the agri-food sector, phosphorus is also involved in other industrial activities, including steel, chemical, and forestry industries. Henceforth, phosphorus is a key material for many aspects of human development. As a result, mapping the phosphorus flows involved in human activities to detect opportunities for recovery and recycling is essential for, in a second stage, assess amount of phosphorus that is viable to recover, the economical costs involved, and the enhancement in terms of resiliency of the regional food production system, savings from the reduction of phosphorus imports, and the mitigation of phosphorus pollution on the region implementing strategies for phosphorus recovery and recycling. The quantification of phosphorus flows has been addressed in previous works in the literature for certain sectors such as the agri-

43 food sector (Boh and Clark, 2020; Zhou et al., 2021; Nesme et al., 2018). Additionally, studies  
44 on the global phosphorus flows have also been performed (Villalba et al., 2008; Chen and Graedel,  
45 2016), although these studies tend to have a low flow resolution since these are aggregated by major  
46 sectors. Additionally, the works quantifying phosphorus often include qualitative recommendations  
47 to improve the phosphorus use efficiency and recycling (Van Dijk et al., 2016; Senthilkumar et al.,  
48 2012), but often they do not include quantitative assessments on the amount of phosphorus which  
49 recovery is feasible along with the costs involved. Conversely, those works focused on estimating  
50 the recoverable phosphorus and the associated recovery cost target specific flows, lacking a holistic  
51 perspective of the phosphorus flows in the various human activities (Martín-Hernández et al., 2021;  
52 Sampat et al., 2018).

53 In this work, we intend to perform a holistic approach to phosphorus management, recovery,  
54 and recycling using in the Canadian province of Ontario. In a first stage, we proceed to map the  
55 phosphorus flows involved in the economical sectors of Ontario, i.e., the agricultural, industrial,  
56 and urban sectors. This data is used in a second stage to identify the flows in which phospho-  
57 rus recovery is feasible, estimating the amount of phosphorus that could be recovered within the  
58 province considering different phosphorus recovery technologies with technology readiness levels  
59 equal or above 6, as well as the costs associated with phosphorus recovery. Finally, we discuss the  
60 implications that would be derived from implementing active phosphorus recovery and recycling  
61 approaches regarding phosphorus supply and use in Ontario.

## 62 2. Methods

### 63 2.1. Spatial and temporal boundaries

64 Phosphorus flows have been mapped through a material flow analysis (MFA) (Brunner and  
65 Rechberger, 2016) conducted within the political boundaries of the Canadian province of Ontario  
66 using data reported for the year 2019.

## 67 2.2. Estimation of phosphorus flows

68 The estimation of phosphorus flows in Ontario’s economic sectors is based on the use open  
69 data sources, often from governmental institutions, complemented with information from scientific  
70 articles when needed. In the next sections we describe the general procedure followed to estimate  
71 the phosphorus flows of each sector. For a comprehensive description of the procedure followed for  
72 estimating each particular phosphorus flow, we refers the reader to the methodology described in  
73 [Pollution Probe \(2022\)](#).

### 74 2.2.1. Agricultural sector

75 Phosphorus flows in the agricultural sector are estimated based on production data of livestock  
76 and crop products, as well as data on fertilizer application.

77 Phosphorus in livestock feeding and manure is estimated based on the number and type of  
78 animals reported for Ontario in the Census of Agriculture ([Statistics Canada – Statistique Canada, 2021a,e,c,d,b](#)), multiplied by the phosphorus feeding requirements and concentration of phosphorus  
79 in manure ([Statistics Netherlands, 2012](#); [Brown, Christine, 2013](#); [Van Staden et al., 2021](#)). We  
80 assumed that the number of animals reported is throughout the year (i.e., the animals culled are  
81 replaced by new ones). However, in the case of broilers and turkeys, the number of animals reported  
82 by the livestock census have been reduced by a factor of 0.68 (broilers) and 0.80 (turkeys), since  
83 these animals have life cycles of 43 and 80 days respectively, meaning barns are empty for 20 days  
84 between cycles. 305 ([Yang et al., 2007](#)).

86 Phosphorus contained in meat and slaughterhouse waste is based on the number of animals  
87 slaughtered reported by both federally and provincially licensed meat plants ([Agriculture and Agri-  
88 Food Canada, 2021c,b](#)) multiplied by the concentration of phosphorus in carcasses ([Agriculture  
89 and Agri-Food Canada, 2021d](#); [Hayse and Marion, 1973](#); [Brake et al., 1995](#); [Statistics Netherlands, 2012](#)).

91 Phosphorus flows associated with the production of milk and eggs is based on provincial produc-  
92 tion data ([Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2020b,a](#)),  
93 multiplying these products by their average phosphorus concentration ([Health Canada, 2008](#); [Cham-](#)

94 [bers et al., 2017](#)).

95 Phosphorus in fertilizer applied to open fields in Ontario is estimated based on the amount  
96 of fertilizer products traded to Ontario’s agricultural markets containing phosphorus ([Statistics  
97 Canada – Statistique Canada, 2022](#)). Regarding manure, we assume that all of the manure generated  
98 by livestock is applied in crop fields ([van Bochove et al., 2010](#)).

99 The uptake of phosphorus by crops is determined based on the area used in each census division  
100 ([Opendatasoft, 2019](#)) to grow each type of crops by census division ([Agriculture and Agri-Food  
101 Canada, 2022a,b,c](#)) multiplied by the specific yield and phosphorus content for each crop type  
102 ([United States Department of Agriculture, 2009](#)). The phosphorus uptake by crops is divided  
103 according to whether it uptake in the grain, fruit or vegetable, or straw and stover components of  
104 each type of crop. This is necessary to determine the amount of phosphorus that flows within food  
105 or feed (i.e., grains, fruits and vegetables) while straw and stover remain in the field after harvesting  
106 as crop residues.

107 A fraction of the phosphorus applied to crop fields as manure of synthetic fertilizer is lost  
108 through erosion, runoff, and drainage. This transportation of phosphorus depends on a range  
109 of factor, including the amount of phosphorus applied; soil composition, texture, and slope; and  
110 precipitation, resulting in a complex and data-intensive process for estimating the phosphorus  
111 transported out of the crop fields. As an approximation, we have estimated the phosphorus losses  
112 by using export coefficients determined for crop fields in Ontario corrected to account for both  
113 surface and subsurface runoffs for synthetic fertilizers (1.267 kg/ha/year ), and liquid and solid  
114 manure (2.548 kg/ha/year and 1.717 kg/ha/year respectively) ([Zhang et al., 2015](#); [Wang et al.,  
115 2018](#); [Tan and Zhang, 2011](#)).

116 A fraction of the P supplied to crop fields is not taken up by the plants and remains in soil,  
117 resulting in the accumulation of P over time as a result of synthetic fertilizer and manure over  
118 over sustained periods of time, often applying phosphorus in greater quantities than crops require  
119 to ensure satisfactory yields ([Reid et al., 2019](#)). This buildup is often referred to as “legacy P”,  
120 and it is estimated as the balance between phosphorus inflows to crop fields (application of manure  
121 and synthetic fertilizers) and outflows (crop food and feed products, crop residues, and phosphorus

122 losses).

123       Regarding greenhouse crops, the data available was limited, resulting in an estimation of phos-  
124 phorus applied as synthetic fertilizers based on the sum of phosphorus uptake by greenhouse crops  
125 (i.e., tomatoes, peppers, and cucumbers)119 and the phosphorus releases from greenhouse irriga-  
126 tion systems (greenhouse nutrient feedwater systems ([Ontario Ministry of Agriculture and Food](#)  
127 [and the Ministry of Rural Affairs, 2021](#)) systems. The phosphorus uptake by greenhouse crops is  
128 determined by multiplying the production of greenhouse crops ([Ontario Ministry of Agriculture and](#)  
129 [Food and the Ministry of Rural Affairs, 2022](#)) by the phosphorus content of each vegetable type  
130 ([United States Department of Agriculture, 2009](#)). The phosphorus releases from the GNF systems  
131 was estimated based on the average concentration of phosphorus in GNF outlet streams of Ontario  
132 (33.6 mg/L) ([Ontario Ministry of the Environment, Conservation and Parks, 2012](#)) and the total  
133 water discharges from GNF systems, assuming that water discharges from GNF systems is equiv-  
134 alent to 25% of the total water applied in greenhouses, which corresponding with the worst-case  
135 scenario of no water recirculation in the GNF system ([Ontario Ministry of Agriculture and Food](#)  
136 [and the Ministry of Rural Affairs, 2021](#)). The average water consumption in greenhouses in Ontario  
137 was assumed to be 1,000 L/m<sup>2</sup>/year ([Ontario Ministry of Agriculture and Food and the Ministry](#)  
138 [of Rural Affairs, 2011](#)). We have also estimated the phosphorus releases from the seasonal workers  
139 live in households in the vicinity of the greenhouses that may use septic systems, considering that  
140 the seasonal labour force in Ontario greenhouses is estimated to be 6,699 workers ([Government](#)  
141 [of Canada, 2022](#)), and an average phosphorus load rate of f 0.0156 kg P/person/week from septic  
142 systems ([Oldfield et al., 2020](#)).

143       Food imports and exports (other than livestock) are estimated scaling each type of food traded  
144 in Canada ([Statistics Canada - Statistique Canada, 2022e](#)) with the population of Ontario ([Statistics](#)  
145 [Canada - Statistique Canada, 2022c](#)). The phosphorus contained in each type of imported and  
146 exported food is estimated multiplying the amount of ech type of traded food by its phosphorus  
147 content ([Health Canada, 202](#)).

### 148 2.2.2. Industrial sector

149 Phosphorus flows through imports, production, exports and waste for the steel, forestry, and  
150 food and beverage, industries of Ontario were mapped. The steel industry is the first non-food sector  
151 in terms of phosphorus use. The main phosphorus inflows of steel manufacturing are associated  
152 with the use of iron ore, coal, and coke, while the main outflow of phosphorus is within slag, which  
153 remove most of the impurities from steel, including phosphorus. It must be noted that, although  
154 some minor amounts of phosphorus can be desired in steel for making anti-corrosion surface  
155 coatings, it is largely considered an impurity in the steel manufacturing process. Phosphorus in  
156 these flows is estimated multiplying their average phosphorus content (0.06% P in iron ore, 0.05%  
157 P for coal, 0.4% P in slag, and 0.01% in steel) ([Yokoyama et al., 2007](#)) by the steel production  
158 capacity of the facilities located in Ontario ([Cheminfo Services Inc., 2019](#); [Algoma Steel Inc., 2022](#);  
159 [Stelco Inc., 2022](#); [Pollution Probe, 2022](#)) and the imports and exports of these materials ([World](#)  
160 [Integrated Trade Solution, 2022](#); [Statistics Canada - Statistique Canada, 2022a](#)).

161 Phosphorus flows in Ontario's forestry industry includes wood harvesting, wood products man-  
162 ufacturing, as well as the production of pulp and paper. The estimation of these phosphorus flows  
163 are the result of multiplying the production data of wood, wood products, pulp and paper, and  
164 their retrospectives imports, exports, and waste streams ([Canadian Forest Service, 2020](#); [Statistics](#)  
165 [Canada - Statistique Canada, 2022a](#)), by the average phosphorus content, which is assumed to be  
166 0.01% for wood ([Sardans and Peñuelas, 2013](#)) and 0.005% for pulp and paper products.

167 Phosphorus in aquaculture are mainly due to supply of feed as part of fish feed the grow of  
168 trouts, part of which is uptake by fishes, while the rest of phosphorus is released into aquatic  
169 ecosystems since aquaculture effluents are directly discharged to the environment ([Ontario Ministry](#)  
170 [of the Environment, Conservation and Parks, 2019](#)). The amount phosphorus uptakes by fishes is  
171 calculated multiplying the fish production ([Statistics Canada - Statistique Canada, 2021](#)), by their  
172 phosphorus content ([Health Canada, 202](#)), while the phosphorus content in the aquaculture waste  
173 effluents of Ontario is estimated to be 10 kg of phosphorus per ton of fish produced ([Bureau et al.,](#)  
174 [2003](#)). The sum of phosphorus uptakes by fishes and phosphorus in aquaculture waste effluents  
175 result in the phosphorus supplied to aquaculture as fish feed.

176 Regarding other industrial activities which could involve the use of phosphorus, the local pro-  
177 duction of phosphorus is assumed to be negligible since phosphorus is not mined or refined in  
178 Ontario, and the synthetic phosphorus fertilizer imports are accounted in the agricultural section.  
179 The general chemical facilities located in Ontario report 350 t/year of phosphorus as waste ??, in  
180 addition of imports and exports of chemical products. However, there exist a significant fraction of  
181 phosphorus used in the industrial sector that cannot be tracked due to the lack of data.

### 182 2.2.3. Urban sector

183 In this section we include the phosphorus inflows and outflows through wastewater treatment  
184 plants (WWTPs), septic systems, and food and organic waste management facilities (landfills,  
185 composting sites, and anaerobic digestion facilities).

186 Phosphorus flows through WWTPs is estimated combining data from the National Pollutant  
187 Release Inventory (NPRI) ([Environment and Climate Change Canada, 2021a](#)), a public database  
188 of releases, disposals and transfers of pollutants, including industrial facilities, and data from the  
189 Wastewater Systems Effluent Regulations (WSER) database ([Environment and Climate Change  
190 Canada, 2021b](#)). Since the NPRI only contains data of those facilities that meet certain regulatory  
191 requirements, the information of this database must be complemented with the data from the  
192 WSER database, which includes information of Canadian WWTPs at the federal, provincial, and  
193 municipal level. The estimations on phosphorus flows through WWTPs are validated using the  
194 Municipal Treated Wastewater Effluent (MTWE) database ([Ontario Ministry of the Environment,  
195 Conservation and Parks, 2021](#)), which collects annual data on water quality data and effluent levels  
196 for WWTPs in Ontario. We note that this data set only provides information about phosphorus  
197 releases from municipal WWTPs, but it does not collect phosphorus disposals and transfers. This  
198 methodology is shown in Figure 1






Figure 1: Procedure for estimating phosphorus flows through wastewater treatment plants.

However, there exist households that are not connected to any sewer systems. These households are equipped with septic systems to perform a rough treatment of the wastewater produced prior to its release into the environment, which typically consist into a septic tank that separates solid matter from the wastewater, and a drainfield where the effluent is discharged. The estimation of phosphorus releases from septic systems is based on the fraction of households equipped with these systems, estimated on 13% (Statistics Canada - Statistique Canada, 2015), which are inhabited by an average of 2.58 individuals (Statistics Canada - Statistique Canada, 2017), and the average phosphorus load rate from septic systems, which is estimated on 0.81 kg of phosphorus per person per year for the Lake Erie Basin in Ontario by Oldfield et al. (2020).

Phosphorus flows in the form of food and organic waste are based on applying food loss factors for the steps associated with food processing (FAO, 2011), considering the food production and import values estimated in Section 2.2.1.

### 2.3. Phosphorus recovery techniques

There currently exist different processes for phosphorus recovery from different sources which technical viability has been proven or is at advanced development stage, i.e., systems with technologies readiness level (TRL) ([National Aeronautics and Space Administration, 2022](#)) of 6 or above (commercial or pilot plant stage). Since the flows from different processes have different properties, the techniques for phosphorus recovery vary between sectors and flows and, therefore, their recovery efficiencies, costs, and products obtained. Table 1 shows a summary of the specifications of the phosphorus recovery technologies for different flows, including literature references where comprehensive descriptions of each system and its specifications can be found. We noted that the phosphorus recovery processes currently available exceed the systems included in this work, nonetheless the processes considered in this study are a selection of the main techniques for phosphorus recovery, although different processes may have been developed on the foundations of the same technique, e.g., the multiple processes based on struvite precipitation.

Phosphorus recovery costs include operating and annualized capital costs. Capital costs are annualized through the application of an annual capital charge ratio ( $ACCR$ ) as defined by [Towler and Sinnott \(2013\)](#), assuming a typical interest rate of 5% and a plant lifetime of 20 years. Dynamic phosphorus recovery costs in function of the processing capacity have been considered in order to capture the economies of scale for those technologies for which sufficient data are available.

In general terms, the process for phosphorus recovery and recycling can be classified into those employed for the treatment of liquid streams, which are based on the formation of precipitated through the direct processing of the liquid effluent, and those processes employed for the treatment of solid fractions, which usually require the pretreatment of the waste through an incineration stage.

Phosphorus in manure represent an important flow within the agricultural sector. The techniques used for phosphorus recovery from cattle and swine manure differ for the liquid and solid fractions. Struvite precipitation is the dominant technology for phosphorus recovery from liquid manure, existing different processes for struvite production based on the type of reactors used with similar recovery efficiencies but different treatment capacities, and thus different recovery costs ([Martín-Hernández et al., 2021](#)). Additionally, there exist modular processes based on physical

Table 1: Phosphorus recovery techniques considered in the study.  $F$  denotes the phosphorus recovered as kg  $P_{\text{recovered}}/\text{year}$ , while  $[x]$  represent the ceiling function applied to  $x$ . The definition of annual capital charge ratio ( $ACCR$ ) can be found in the Supplementary Material, Section 1.1. Refs: [1]: [Martín-Hernández et al. \(2021\)](#), [2]: [Jupp et al. \(2021\)](#), [3]: [Egle et al. \(2016\)](#), [4]: [Schoumans et al. \(2010\)](#), [5]: [Sögi et al. \(2008\)](#), [6]: [AMPC \(2018\)](#), [7]: [Zagklis et al. \(2020\)](#), [8]: [Fernández-Delgado et al. \(2022\)](#), [9]: [Ohtake and Tsumeda \(2019\)](#), [10]: [Sharma and Chandel \(2021\)](#)

Sector	Inflow	Pretreatment	Pretreatment cost (EUR/kg $P_{\text{recovered}}$ )	Technology	Type	P recovery potential (% related to inflow)	P recovery cost (EUR/kg $P_{\text{recovered}}$ )	TRL	Ref tech
Agriculture	Cattle and swine manure, liquid phase (30% of total manure P)	Solid-liquid separation (screw press)	See [1]	Multiform	Struvite	60	$25.7 + 1.10 \cdot 10^{-6} \cdot [1.19 \cdot 10^{-4} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Crystallator	Struvite/ Calcium phosphate	60	$3.53 + (2.30 \cdot 10^6 + 0.71 \cdot [3.32 \cdot 10^{-5} \cdot F]) [3.32 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Ostara Pearl 500	Struvite	60	$12.57 + 2.30 \cdot 10^6 \cdot [7.02 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Ostara Pearl 2K	Struvite	60	$12.57 + 3.10 \cdot 10^6 \cdot [1.83 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Ostara Pearl 10K	Struvite	60	$12.57 + 10.00 \cdot 10^6 \cdot [3.65 \cdot 10^{-6} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[1]
	Poultry litter	Solid-liquid separation (screw press)	See [1]	Nurelys	Struvite	60	$10.37 + 1.38 \cdot 10^6 \cdot [2.24 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	MAPHEX	Solid	90	$184.67 + 0.30 \cdot 10^6 \cdot [2.47 \cdot 10^{-4} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	6	[1]
		Incineration	8.9	EcoPhos	Phosphoric acid	82	4.5	6	[2,3,4]
		Incineration	8.9	AshDec depollution	Calcium phosphate	86	1.8	6	[2,3,4]
		Incineration	8.9	AshDec Rheumia	Calcium phosphate	86	1.9	6	[2,3,4]
	Slaughterhouse waste, liquid phase (14% of total slaughterhouse P)	Incineration	8.9	PASCH	Calcium phosphate	79	4.7	6	[2,3,4]
		Incineration	8.9	LEACHPHOS	Calcium phosphate	78	5.1	9	[2,3,4]
		Incineration	8.9	RecoPhos	Mineral	87	2.5	9	[2,3,4]
		Incineration	8.9	Thermophos	P4	81	2.7	9	[2,3,4]
		Incineration	-	Quick wash	Solid precipitate	70	4.4	3	[5]
Urban & industrial	WWTPs (liquid phase)	-	-	Multiform	Struvite	84	$22.6 + 1.10 \cdot 10^6 \cdot [1.05 \cdot 10^{-4} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[6]
		-	-	Ostara Pearl 500	Struvite	58	$15.60 + 2.30 \cdot 10^6 \cdot [8.70 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[6]
		-	-	Ostara Pearl 2K	Struvite	58	$15.60 + 3.10 \cdot 10^6 \cdot [2.26 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[6]
		-	-	Ostara Pearl 10K	Struvite	58	$15.60 + 10.00 \cdot 10^6 \cdot [4.53 \cdot 10^{-6} \cdot F] \cdot ACCR \cdot \frac{1}{P}$	9	[6]
		Incineration	14.6	EcoPhos	Phosphoric acid	82	4.5	6	[2,3,7]
	Slaughterhouse waste, solid phase (86% of total slaughterhouse P)	Incineration	14.6	AshDec depollution	Calcium phosphate	86	1.8	6	[2,3,7]
		Incineration	14.6	AshDec Rheumia	Calcium phosphate	86	1.9	6	[2,3,7]
		Incineration	14.6	PASCH	Calcium phosphate	79	4.7	6	[2,3,7]
		Incineration	14.6	LEACHPHOS	Calcium phosphate	78	5.1	9	[2,3,7]
		Incineration	14.6	RecoPhos	Mineral	87	2.5	9	[2,3,7]
	WWTPs (sewage sludge, 60-90% of P)	Incineration	14.6	Thermophos	P4	81	2.7	9	[2,3,7]
		-	-	Crystallator	Struvite/ Calcium phosphate	38	$305,920 \cdot \left(\frac{F}{34,506}\right)^{0.99} \cdot \frac{1}{P}$	9	[3]
		-	-	Ostara Pearl	Struvite	20	$130,856 \cdot \left(\frac{F}{53,140}\right)^{0.86} \cdot \frac{1}{P}$	9	[3]
		-	-	P-RoC	Calcium phosphate	27	$75,970 \cdot \left(\frac{F}{17,789}\right)^{0.78} \cdot \frac{1}{P}$	6	[3]
		-	-	REM-NUT	Struvite	47	$977,933 \cdot \left(\frac{F}{30,875}\right)^{0.94} \cdot \frac{1}{P}$	6	[3]
Urban & industrial	WWTPs (sewage sludge, 60-90% of P)	-	-	AirPrex	Struvite	15	$74,195 \cdot \left(\frac{F}{9,532}\right)^{0.38} \cdot \frac{1}{P}$	9	[3]
		-	-	PRISA	Struvite	18	$186,923 \cdot \left(\frac{F}{11,326}\right)^{0.43} \cdot \frac{1}{P}$	6	[3]
		-	-	Stutgart process	Struvite	40	$581,730 \cdot \left(\frac{F}{20,280}\right)^{0.89} \cdot \frac{1}{P}$	9	[3]
		-	-	Gifhorn process	Struvite	40	$400,384 \cdot \left(\frac{F}{26,280}\right)^{0.82} \cdot \frac{1}{P}$	9	[3]
		-	-	PHOXNAN	Struvite	51	$891,667 \cdot \left(\frac{F}{30,507}\right)^{0.84} \cdot \frac{1}{P}$	6	[3]
	WWTPs (sewage sludge ash, SSA, 60-90% of P)	-	-	Aqua Reci	Calcium phosphate	61	$939,605 \cdot \left(\frac{F}{30,077}\right)^{0.82} \cdot \frac{1}{P}$	6	[3]
		-	-	MEPREX	P rich slag	68	$1,154,473 \cdot \left(\frac{F}{44,676}\right)^{0.61} \cdot \frac{1}{P}$	6	[3]
		Incineration	8	EcoPhos	Phosphoric acid	82	4.5	6	[3]
		Incineration	8	AshDec depollution	Calcium phosphate	86	1.8	6	[3]
		Incineration	8	AshDec Rheumia	Calcium phosphate	86	1.9	6	[3]
	Organic municipal & food waste	Incineration	8	PASCH	Calcium phosphate	79	4.7	6	[3]
		Incineration	8	LEACHPHOS	Calcium phosphate	78	5.1	9	[3]
		Incineration	8	RecoPhos	Mineral	87	2.5	9	[3]
		Incineration	8	Thermophos	P4	81	2.7	9	[3]
		Incineration	6.43	AshDec Rheumia	Calcium phosphate	86	1.9	6	[3,9,10]
	Organic municipal & food waste	Incineration	6.43	PASCH	Calcium phosphate	79	4.7	6	[3,9,10]
		Incineration	6.43	LEACHPHOS	Calcium phosphate	78	5.1	9	[3,9,10]
		Incineration	6.43	RecoPhos	Mineral	87	2.5	9	[3,9,10]
		Incineration	6.43	Thermophos	P4	81	2.7	9	[3,9,10]

separations oriented to small-scale intensive livestock facilities (Church et al., 2018). The recovery of phosphorus from the solid fraction of manure involves the incineration of the waste, and the further processing of the ashes, recovering phosphorus precipitates or phosphoric acid (Jupp et al., 2021; Egle et al., 2016). Phosphorus recovery from poultry litter is based on acid extraction and further precipitation (Szögi et al., 2008).

Slaughterhouse waste is flow from food industry which can be targeted for phosphorus recovery. It should be noted that slaughterhouse is comprised by a liquid (slaughterhouse wastewater) and a solid fraction (animal carcass waste) and, therefore, the phosphorus recovery systems for each flow will differ. Similarly to the case of phosphorus recovery from manure, phosphorus recovery from slaughterhouse wastewater is performed through struvite precipitation (AMPC, 2018), while the animal carcass waste is incinerated and phosphorus is recovered from ashes in form of calcium carbonate or phosphoric acid (Jupp et al., 2021).

Municipal wastewater contains significant amounts of phosphorus that can be recovered. It must be noted that phosphorus outflows from WWTPs are divided into phosphorus contained in the treated water and phosphorus contained in sludge. Phosphorus contained in water can be recovered through the formation of precipitates such as struvite, while phosphorus contained in sludge can be recovered either through the direct processing of sludge producing precipitates, or from sludge ashes after an incineration stage, obtaining different products such as phosphoric acid or calcium phosphate.

### 3. Results and discussion

#### 3.1. Phosphorus flows in Ontario

Figure 2 summarizes the phosphorus flows in the province of Ontario. It can be observed that phosphorus flows in anthropogenic activities can be divided into 3 independent networks, i.e., the flow of phosphorus involved the production and processing of food (including the treatment of wastewater), the flow of phosphorus used in the steel and chemical industries, and the phosphorus involved in the forestry industry.




Figure 2: Phosphorus flows in the province of Ontario (kt/year). The streams within red rectangles denote phosphorus inflows into the province, while those streams within blue rectangles denote phosphorus outflows out of the province.

265 The main inflows food production and processing network are those associated with the supply  
 266 of synthetic fertilizers and livestock feed, accounting for 55.6 and 21.8 kt/year respectively. Other  
 267 phosphorus imports in Ontario are made in the form of food products (4.7 kt/year) and aquacul-  
 268 ture feed imports (0.07 kt/year). Crop residues and manure are the main phosphorus waste effluent  
 269 from the agricultural sector, accounting for 32.5 and 30.6 kt/year. However, it must be noted the  
 270 different properties of these materials. While crop residues can be left in the cropfields, transfer-  
 271 ring part of the phosphorus taken by crops back to soil and acting as soil amendment materials  
 272 due to their carbon content, manure produced in intensive livestock facilities is a point source of  
 273 phosphorus releases highly spatially concentrated, resulting in the accumulation of phosphorus in  
 274 the vicinity of these facilities. As a consequence, the production of manure has the potential of  
 275 being environmentally harmful and requires of adequate management strategies. The food pro-  
 276 cessing industry involves the largest flows within the province, which can be classified plant and  
 277 animal-based product, and slaughter house waste, resulting in phosphorus flows of 43.2, 10.3, and  
 278 3.8 kt/year respectively. A significant fraction of end-flows are waste flows in the form of landfill  
 279 (37.5 kt/year), composting (2.1 kt/year), and anaerobic digestion (1.4 kt/year) disposals, as well  
 280 as wastewater (7.8 kt/year). Phosphorus exports out of Ontario are performed through the exports  
 281 of food products and livestock, accounting for 10.8 and 1.8 kt/year respectively.

282 Chemical industry involve significant phosphorus flows, importing 22.2 kt/year into the province,  
 283 while the exports represent 14.0 kt/year. This sector produces 0.35 kt/year of phosphorus that is  
 284 classified as waste. However, an important fraction of phosphorus used by this sector (11.7 kt/ton)  
 285 cannot be tracked and. therefore, it is unknown what fraction of this phosphorus can result as  
 286 waste. Steel production imports 8.3 kt/year of phosphorus, of which 7.0 kt/year ends as steel wast  
 287 or slag, while the phosphorus flows in steel materials are 1.4 kt/year.

288 Phosphorus imports from the forestry industry are 0.38 kt/year, while 2.3 kton/year of phospho-  
 289 rus is taken from wood harvested in Ontario. This sector releases 0.4 kton/year of phosphorus in  
 290 the form of wood and pulp waste, while 0.81 kt/year of phosphorus is exported out of the province  
 291 as wood, wood products or pulp and paper.

### 292 3.2. Potential of phosphorus recovery in Ontario

293 The potential for phosphorus recovery in the province of Ontario through the deployment of  
 294 different processes for the recovery of phosphorus from different flows is assess in this section. As  
 295 shown in Section 2.3, different processes can be employed for the recovery of phosphorus from  
 296 the same stream. However, each system is design for operating under certain conditions and they  
 297 have different processing capacities. As a result, phosphorus recovery efficiency and cost will differ  
 298 between technologies for the treatment of the same flow. In order to explore this variability be-  
 299 tween phosphorus recovery systems, all the systems described in Table 1 are evaluated. The results  
 300 obtained in terms of phosphorus recovered and recovery cost for each technology and flow are col-  
 301 lected in the Supplementary Material. Two scenarios are selected for deeper analysis, the minimum  
 302 cost scenario that selects the most economical technology, and the maximum recovery scenario,  
 303 comprised by the phosphorus recovery system which deployment result in the largest phosphorus  
 304 recovery.




Figure 3: Phosphorus recovered and phosphorus recovery cost in the province of Ontario by flow. Note that the scale of bottom x-axis is different for left and right axes.




Figure 4: Average phosphorus recovered and average phosphorus recovery cost per kilogram of phosphorus recovered in the province of Ontario by flow.

### 3.2.1. Agricultural sector

Manure is an agricultural flow from which effective phosphorus recovery might be achieved since it can be collected from the intensive livestock facilities and further treated (Schoumans et al., 2010). The inventory of cattle, swine, poultry, and other livestock for Ontario in 2019 is 1,376,984, 506,768, 148,508, and 135,725 animal units respectively, resulting in the release of 13,274, 8,569, 6,457, and 2,283 metric tonnes of phosphorus per year through manure respectively. Phosphorus recovery from manure is highly influenced by the economies of scale and, therefore, by the scale of the CAFOs (Martín-Hernández et al., 2021). Since no data on the size distribution of CAFOs in Ontario is available, the average sizes of livestock facilities reported by Statistics Canada - Statistique Canada (2022b) for the year 2019 are considered, resulting in average sizes for cattle, swine, poultry and other livestock (primarily sheep and lambs) facilities of 98, 168, 15 and 16 animal units respectively. The number of cattle, swine, poultry, and other livestock CAFOs obtained is 14,051, 3,022, 10,069, and 8,636 respectively, which is in alignment with the number of livestock facilities reported by Statistics Canada - Statistique Canada (2022d).



319 Phosphorus is divided between the liquid and solid fractions of manure, being the solid fraction  
 320 the one containing the largest amount of phosphorus, and thus the fraction from which larger  
 321 quantities amounts of phosphorus can be recovered with lower costs, as observed in Figure 3.  
 322 However, it must be noted that phosphorus recovery from solid manure involved more complex  
 323 processes that include the incineration of the waste, which in turn makes the process more energy  
 324 intensive and may result in environmentally harmful emissions of gases. Cattle manure contains  
 325 the largest amount of phosphorus as a consequence of being the largest manure flow, followed by  
 326 swine and poultry manure. However, the comparison of the average phosphorus recovery costs  
 327 per kilogram of phosphorus recovered shows that phosphorus recovery from swine liquid manure is  
 328 lowest, as shown in Figure 4. This is due to the size of the swine intensive facilities, which in average  
 329 are comprised by a larger number of animal units than cattle intensive facilities. This reveals the  
 330 important role of the economies of scale in phosphorus recovery. Moreover, the small size of the  
 331 CAFOs in Ontario result in high phosphorus recovery costs, whose values range between 200 and  
 332 700 EUR/kg P recovered. These costs are significantly higher than the phosphorus recovery costs  
 333 reported by [Martín-Hernández et al. \(2022\)](#) for the comparatively larger CAFOs of the U.S. states  
 334 in the Great Lakes area, which average sizes range from 630 and 2,600 animal units, resulting  
 335 in phosphorus recovery costs between 13 and 73 USD/kilogram of phosphorus recovered. The  
 336 phosphorus recovery efficiency is similar for all livestock types since all the process selected is the  
 337 modular physical separation system due to the small scale of the livestock facilities in Ontario.  
 338 For the case of solid manure it can be observed that all livestock types show a similar average  
 339 phosphorus recovery cost as a result of the lack of data to estimate the effect of the economies of  
 340 scale of these processes.

### 341 *3.2.2. Industrial and urban sector*

342 Industrial and urban sectors are grouped since some flows belong to both sectors, particularly  
 343 those related with wastewater, and the organic fraction of industrial and municipal solid waste,  
 344 including food waste.

345 Slaughterhouse waste is an industrial flow from which phosphorus can be effectively recovered

(AMPC, 2018). Data on individual capacities for the slaughterhouses in Ontario is not available for estimating the effects of the economies of scale on the cost of phosphorus recovery and, therefore, average slaughterhouse capacities are considered, which values are 104,017, 802,186, and  $14.4 \cdot 10^6$  cattle, hog, and poultry heads slaughtered/(facility  $\cdot$  year) respectively (Agriculture and Agri-Food Canada, 2021a; INAC Services, 2014). Considering the inventory of slaughtered animals reported by Agriculture and Agri-Food Canada (2021c,b), 7, 6, and 17 cattle, hog, and poultry slaughtering facilities are obtained, with associated phosphorus flows of 317.4, 103.5, and 53.2 metric tonnes/(facility  $\cdot$  year) respectively. Phosphorus flows from sheep and rabbit slaughtered are considered negligible due to the low number of animals slaughtered. Figure 3 shows that most of phosphorus can be recovered from the solid fraction of waste due to its larger phosphorus content. The variations between the minimum cost and maximum recovery scenarios are not significant for the solid slaughterhouse waste flow, however, for the liquid fraction the phosphorus recovery the difference between these two scenarios increase by a factor of 2.3, while the total recovery cost in the maximum recovery scenario increases by a factor of 3.3 times larger, as shown in Figure 4, showing that the increase of phosphorus recovery efficiency results in a non-linear increase in the phosphorus recovery cost. The numerical results are collected in the Supplementary Material

Wastewater is a large flow containing significant amounts of phosphorus. Wastewater is collected and directed to wastewater treatment plants (WWTPs). These facilities produces a liquid water effluent with adequate environmental parameters for its being releas into the environment, and a sludge flow from the primary and second treatments. Phosphorus can be recovered from both flows, although most of the phosphorus is contained in the sludge fraction. The distribution of phosphorus between treated water and sludge considered is 14.1% - 85.9% respectively (Pollution Probe, 2022), based on the data reproted by NPRI and WSER databases REFs, which is in alignment with the distribution values reported by (Egle et al., 2016). The capacity of the wastewater treatment plants installed in Ontario, together with their phosphorus flows, have been considered to determine the effect of the economies of scale in the cost of phosphorus recovery. Figure 3 shows that the potential for phosphorus recovery from sludge is greater than from the WWTPs liquid fraction, as mentioned before. Little variation is observed between the minum cost and maximum recovery scenarios for the

recovery of phosphorus from sledge, which implies that there exist a certain degree of homogeneity in the current technologies for phosphorus recovery from sludge, which can be appreciated in Figure 4. It must be noted that the phosphorus recovery systems from sludge ashes reveal to be more effective, including the incineration cost, than the direct recovery of phosphorus from sludge due to the higher recovery efficiency of the former ones. Conversely, the phosphorus recovery from the liquid wastewater fraction show a larger variability between both scenarios considered. The phosphorus recovered in the maximum recovery scenario is 1.7 times larger than in the minimum cost scenarios. However, this increase in the phosphorus recovery efficiency result in the increase of the recovery cost by a factor of 9.6, showing that similarly to the case of the liquid fraction of slaughterhouse waste, the achievement of greater recovery efficiencies through the use of more effective technologies result in an exponential increase of recovery costs.

Figure 3 shows that the food and organic waste represent the largest potential for phosphorus recovery. However, it must be noted that this flow includes all those streams comprised by solid organic wastes other than slaughterhouse waste, including food processing industry waste, household food waste, and other municipal organic waste. Similarly to other solid fractions, the minimum cost and maximum recovery scenario shows a narrow variability regarding phosphorus recovered and recovery costs.

Finally, there are some processes under development for the recovery of phosphorus from steel production industry, particularly from steelmaking slag. These processes are based on selective leaching and further chemical precipitation to obtain solid phosphates (Du et al., 2022, 2019) or magnetic-aid calcium phosphate precipitation (Yokoyama et al., 2007). Although these are promising processes that can result in an effective recovery of phosphorus which can be further recycled, they are at early development stages and, thus, they have not been considered in this assessment.

### 3.3. *Economic implications of phosphorus recovery in Ontario*

In order to compare the costs derived from the recovery of anthropogenic phosphorus, Figure 5 shows the average phosphorus recovery cost in Ontario along with the long-term social and envi-

ronmental economical losses derived from uncontrolled releases of phosphorus into the environment estimated by [Sampat et al. \(2021\)](#). and the temporal evolution of prices for different phosphorus commodities reported by the World Bank’s Commodity Markets report ([The World Bank, 2022](#)), i.e., phosphate rock, diammonium phosphate (DAP), and triple superphosphate (TSP) together. The costs are referred per mass unit of phosphorus, assuming that the phosphorus content of commercial phosphorus rock ranges from 28 to 34% in mass basis ([FAO & IAEA, 2004](#); [Kaiser and Pagliari, 2018](#)), while phosphorus content of DAP and TSP is 20% and 19% respectively. The prices have been adjusted to the year 2020 through the Consumer Price Index (CPI) elaborated by the U.S. Census Bureau ([U.S. Census Bureau, 2021](#)).

It can be observed that the average cost of phosphorus recovery in Ontario, valued around 36 EUR/kg of phosphorus recovered, is significantly lower than the economic losses derived from the release of phosphorus into the environment, estimated at 74.5 EUR/kg P in the context of state of Wisconsin, U.S. This provides an economical support to the recovery of phosphorus, since high long-term costs derived from the social and environmental damages caused by phosphorus releases into the environment could be avoided through the recovery of phosphorus from waste streams. Nevertheless, it can be observed that the cost of phosphorus obtained from recovery processes is more costly than phosphorus in commercial fertilizers obtained from mining, which reduces the economic incentives for the recovery and uses of phosphorus, and further support in form of environmental and/or agricultural regulations, phosphorus recovery incentives, or sustainability arguments are needed in order to promote the recovery and recycling of phosphorus.

Additionally, [Figure 5](#) shows that average cost of phosphorus recovery within the province is around 36 EUR/kg of phosphorus recovered, however, in [Figure 4](#) we can observe that phosphorus recovery costs can reach significantly higher values, particularly for the recovery of phosphorus from manure due to the lack of economies of scale in Ontario’s CAFOs. Therefore, regional cooperative strategies can be implemented to develop a coordinated action against nutrient pollution, distributing the benefits and costs of phosphorus recovery, and prioritizing the total utility rather than the utility of each stakeholder, similarly to the cooperative approaches proposed for minimizing the cost of reducing greenhouse gases emissions from electricity generation ([Galán-Martín et al., 2018](#)).




Figure 5: Average phosphorus recovery cost in Ontario, economical losses derived from the releases of phosphorus into the environment (Sampat et al., 2021), and temporal evolution of prices for different phosphorus commodities. Note that the scale of y-axis is different for top and bottom axes.

#### 3.4. Potential for phosphorus recycling in Ontario

Phosphorus recovered can be further recycled within the food production systems, developing a circular economy around the use of phosphorus. Phosphorus recovery and recycling would result in curbing the depletion of phosphorus rock reserves and reduce the dependency on phosphorus supply from other regions. Considering these factors, there exist some governmental initiatives that, through the creation of different forums and platforms, aim to promote the recovery and recycling of phosphorus (IISD, 2018; Pollution Probe, 2022). In addition, the European Union is setting specific targets to reduce the use of non-renewable materials in fertilizer production (European Commission, 2018) and to promote the use of waste-based fertilizers (European Commission, 2022), encouraging the effective recovery and recycling of phosphorus, and they could serve as a guideline to support phosphorus recovery in other regions.

The comparison between the phosphorus imported for the production of food within Ontario, i.e., in the form of fertilizer or animal feed, and the phosphorus that could be recovered, as shown in Figure 6, is a useful metric to determine the effectiveness of phosphorus recovery, and the potential reduction of phosphorus supplies from external sources. This assessment is limited to the food

444 production sector depicted in Figure 2 since the recovery of phosphorus is performed in different  
 445 flows belonging to this sector.




Figure 6: Phosphorus inputs for food production and phosphorus recovered per sector. S1 refers to the scenario minimizing phosphorus recovery cost, while S2 refers to the scenario maximizing phosphorus recovery.

446 Figure 6 shows that up to 88% of the phosphorus used for food production, excluding imports of  
 447 food that have been produce elsewhere, could be recovered and recycled in the best case scenarios  
 448 where no phosphorus losses occur between the recovery and the reuse stages. An effective recycling  
 449 of phosphorus implies the redistribution of this material from the locations where it is released  
 450 to the CAFOs where it is used for livestock raise, or to phosphorus-deficient cropfields, which in  
 451 turn involves the transportation of phosphorus products as the last stage before the use final use of  
 452 the recovered phosphorus. It must be considered that the transportation of phosphorus products  
 453 result in additional costs that can hinder the recycling of phosphorus, as well as environmental  
 454 impacts related with the current use of fossil fuels-based transportation vehicles. In this context,

coordinated-market approaches have been proposed to optimize the spatial distribution of phosphorus from P-surplus to P-deficient locations, minimizing the cost and environmental footprint of the transportation of phosphorus products (Sampat et al., 2019).

#### 4. Conclusions

Phosphorus is a key non-renewable element for many economic sectors, and particularly for the production of food. The current linear economy scheme deplete phosphorus reserves, resulting in supply dependencies from regions holding phosphorus rock reserves, and it is the sources of nutrient pollution, eutrophication, and other environmental concerns relates with the end-of-life release of phosphorus into the environment. As a consequence, the recovery and recycling of phosphorus is not only a desirable but a necessary approach for the development of phosphorus sustainable systems. For achieving this goal, the mapping of phosphorus across the different economic sectors is the first stage to identify the main streams for phosphorus recovery. This information allows the estimation of the potential for the recycling of phosphorus in a region.

For the case of Ontario, the best case scenario result in a phosphorus recycling potential up to 86% over the phosphorus imported in the province for food production (i.e., excluding the imports of livestock and food produced in other regions). An average phosphorus recovery cost is estimated, although it shows a large variation among different flows. Phosphorus recovery costs is particularly large for phosphorus recovery from manure due to the small scale of the livestock facilities in Ontario. Conversely, phosphorus recovery costs estimated for other regions with larger livestock facilities such as the U.S. Great Lakes area result in significantly lower values, showing the important role of the economizes of scale for phosphorus recovery. Nevertheless, considering the region studied as a whole, the average phosphorus recovery cost estimated is around 36 EUR/kg P recovered, which is lower than the economic losses of phosphorus releases into the environment estimated at 74.5 EUR/kg of phosphorus.

The wide difference of costs for the recovery of phosphorus from different flows suggest the need develop regional cooperative mechanisms to distribute the phosphorus recovery costs. Moreover, further research in the effective distribution of the phosphorus recovered from regions with phos-

phorus surplus to phosphorus-deficient locations is needed, although there exist some research on the development of coordinated markets for phosphorus recovery and recycling.

## 5. Acknowledgments

Please add your acknowledgements

## References

- Agriculture and Agri-Food Canada, 2021a. Distribution of slaughtering activity. <https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/red-meat-and-livestock-market-information/slaughter-and-carcass-weights/distribution-slaughtering-activity>. [Online; accessed 22-August-2022].
- Agriculture and Agri-Food Canada, 2021b. Poultry slaughter reports. <https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/poultry-and-egg-market-information/slaughter>. [Online; accessed 16-December-2021].
- Agriculture and Agri-Food Canada, 2021c. Red meat and livestock slaughter and carcass weights. <https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/red-meat-and-livestock-market-information/slaughter-and-carcass-weights>. [Online; accessed 15-December-2021].
- Agriculture and Agri-Food Canada, 2021d. Red meat conversion factors. <https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/red-meat-and-livestock-market-information/slaughter-and-carcass-weights/conversion-factors>. [Online; accessed 16-December-2021].
- Agriculture and Agri-Food Canada, 2022a. Field crops and hay, Census of Agriculture, 2011 and 2016. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210041601>. [Online; accessed 16-December-2021].



505 Agriculture and Agri-Food Canada, 2022b. Field vegetables,, Census of Agriculture, 2011 and 2016.  
 506 <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210041801>. [Online; accessed  
 507 16-December-2021].

508 Agriculture and Agri-Food Canada, 2022c. Greenhouse products and mushrooms,, 2011 and 2016.  
 509 <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042001>. [Online; accessed  
 510 16-December-2021].

511 Algoma Steel Inc., 2022. Corporate Profile. [https://algoma.com/about-algoma/  
 512 corporate-profile/](https://algoma.com/about-algoma/corporate-profile/). [Online; accessed 13-July-2022].

513 AMPC, 2018. Struvite or Traditional Chemical Phosphorus Precipitation – What Option Rocks?  
 514 [https://www.ampc.com.au/uploads/cgblog/id408/2018-1026\\_-\\_Final\\_Report.pdf](https://www.ampc.com.au/uploads/cgblog/id408/2018-1026_-_Final_Report.pdf). [On-  
 515 line; accessed 20-March-2019].

516 van Bochove, E., Thériault, G., Denault, J.T., 2010. Indicator of risk of water contamination by  
 517 phosphorus (IROWC P): a handbook for presenting the IROWC P algorithms. Technical Report.  
 518 Agriculture and Agri-Food Canada.

519 Boh, M.Y., Clark, O.G., 2020. Nitrogen and phosphorus flows in Ontario’s food systems. Resources,  
 520 Conservation and Recycling 154, 104639.

521 Brake, J., Havenstein, G., Ferket, P., Rives, D., Giesbrecht, F., 1995. Relationship of sex, strain,  
 522 and body weight to carcass yield and offal production in turkeys. Poultry science 74, 161–168.

523 Brown, Christine, 2013. Available Nutrients and Value for Manure From Various Livestock Types.  
 524 Technical Report. Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs.

525 Brunner, P.H., Rechberger, H., 2016. Handbook of material flow analysis: For environmental,  
 526 resource, and waste engineers. CRC press.

527 Bureau, D.P., Gunther, S.J., Cho, C.Y., 2003. Chemical composition and preliminary theoretical  
 528 estimates of waste outputs of rainbow trout reared in commercial cage culture operations in  
 529 Ontario. North American Journal of Aquaculture 65, 33–38.

- 530 Canadian Forest Service, 2020. Statistical data. <https://cfs.nrcan.gc.ca/statsprofile/>. [On-  
531 line; accessed 13-July-2022].
- 532 Chambers, J.R., Zaheer, K., Akhtar, H., Abdel-Aal, E.S.M., 2017. Chicken eggs, in: Egg innovations  
533 and strategies for improvements. Elsevier, pp. 1–9.
- 534 Cheminfo Services Inc., 2019. Economic Assessment of the Inte-  
535 grated Steel Industry. [https://www.canadiansteel.ca/files/resources/  
536 Final-Report-Economic-Assessment-of-the-Integrated-Steel-Industry.pdf](https://www.canadiansteel.ca/files/resources/Final-Report-Economic-Assessment-of-the-Integrated-Steel-Industry.pdf). [Online;  
537 accessed 13-July-2022].
- 538 Chen, M., Graedel, T., 2016. A half-century of global phosphorus flows, stocks, production, con-  
539 sumption, recycling, and environmental impacts. *Global Environmental Change* 36, 139–152.
- 540 Church, C.D., Hristov, A.N., Kleinman, P.J., Fishel, S.K., Reiner, M.R., Bryant, R.B., 2018. Ver-  
541 satility of the MANure PHosphorus EXtraction (MAPHEX) System in Removing Phosphorus,  
542 Odor, Microbes, and Alkalinity from Dairy Manures: A Four-Farm Case Study. *Applied En-  
543 gineering in Agriculture* 34, 567–572. URL: [http://elibrary.asabe.org/abstract.asp?AID=  
544 48976&t=3&dabs=Y&redir=&redirType=](http://elibrary.asabe.org/abstract.asp?AID=48976&t=3&dabs=Y&redir=&redirType=), doi:10.13031/aea.12632.
- 545 Cordell, D., Drangert, J.O., White, S., 2009. The story of phosphorus: global food security and  
546 food for thought. *Global environmental change* 19, 292–305.
- 547 Du, C.M., Gao, X., Ueda, S., Kitamura, S.Y., 2019. Separation and recovery of phosphorus from  
548 steelmaking slag via a selective leaching–chemical precipitation process. *Hydrometallurgy* 189,  
549 105109.
- 550 Du, C.m., Gao, X., Ueda, S., Kitamura, S.y., 2022. Recovery of high-quality phosphate from  
551 steelmaking slag by a hydrometallurgical process. *Science of The Total Environment* 819, 153125.
- 552 Egle, L., Rechberger, H., Krampe, J., Zessner, M., 2016. Phosphorus recovery from mu-  
553 nicipal wastewater: An integrated comparative technological, environmental and economic  
554 assessment of P recovery technologies. *Science of The Total Environment* 571, 522–542.

555 URL: <https://linkinghub.elsevier.com/retrieve/pii/S0048969716314656>, doi:10.1016/  
556 [j.scitotenv.2016.07.019](https://linkinghub.elsevier.com/retrieve/pii/S0048969716314656).

557 Ehlert, P., Morel, C., Fotyma, M., Destain, J.P., 2003. Potential role of phosphate buffering capacity  
558 of soils in fertilizer management strategies fitted to environmental goals. *Journal of plant nutrition*  
559 *and soil science* 166, 409–415.

560 Environment and Climate Change Canada, 2021a. About the Na-  
561 tional Pollutant Release Inventory. [https://www.canada.ca/en/](https://www.canada.ca/en/environment-climate-change/services/national-pollutant-release-inventory/about-national-pollutant-release-inventory.html)  
562 [environment-climate-change/services/national-pollutant-release-inventory/](https://www.canada.ca/en/environment-climate-change/services/national-pollutant-release-inventory/about-national-pollutant-release-inventory.html)  
563 [about-national-pollutant-release-inventory.html](https://www.canada.ca/en/environment-climate-change/services/national-pollutant-release-inventory/about-national-pollutant-release-inventory.html). [Online; accessed 29-December-  
564 2021].

565 Environment and Climate Change Canada, 2021b. Wastewater Systems Effluent Regulations.  
566 <https://open.canada.ca/data/en/dataset/9e11e114-ef0d-4814-8d93-24af23716489>.  
567 [Online; accessed 29-December-2021].

568 European Commission, 2018. Circular Economy: Agreement on Commission proposal to boost the  
569 use of organic and waste-based fertilisers. [https://ec.europa.eu/commission/presscorner/](https://ec.europa.eu/commission/presscorner/detail/en/IP_18_6161)  
570 [detail/en/IP\\_18\\_6161](https://ec.europa.eu/commission/presscorner/detail/en/IP_18_6161). [Online; accessed 25-August-2022].

571 European Commission, 2022. Consolidated text: Regulation (EU) 2019/1009 of the European Parlia-  
572 ment and of the Council of 5 June 2019 laying down rules on the making available on the market  
573 of EU fertilising products and amending Regulations (EC) No 1069/2009 and (EC) No 1107/2009  
574 and repealing Regulation (EC) No 2003/2003 (Text with EEA relevance)Text with EEA rele-  
575 vance. <https://eur-lex.europa.eu/eli/reg/2019/1009>. [Online; accessed 25-August-2022].

576 FAO, 2011. Global food losses and food waste - Extent, causes and prevention. [https://www.fao.](https://www.fao.org/3/i2697e/i2697e.pdf)  
577 [org/3/i2697e/i2697e.pdf](https://www.fao.org/3/i2697e/i2697e.pdf). [Online; accessed 15-July-2022].

578 FAO & IAEA, 2004. Use of Phosphate Rocks for Sustainable Agriculture. [https://www.fao.org/](https://www.fao.org/3/y5053e/y5053e00.htm)  
579 [3/y5053e/y5053e00.htm](https://www.fao.org/3/y5053e/y5053e00.htm). [Online; accessed 23-August-2022].

580 Fernández-Delgado, M., del Amo-Mateos, E., García-Cubero, M.T., Coca, M., Lucas, S., 2022.  
 581 Phosphorus recovery from organic waste for its agronomic valorization: technical and economic  
 582 evaluation. *Journal of Chemical Technology & Biotechnology* 97, 167–178.

583 Food and Agriculture Organization of the United Nations, 2022. The importance of Ukraine and  
 584 the Russian Federation for global agricultural markets and the risks associated with the war in  
 585 Ukraine.

586 Galán-Martín, A., Pozo, C., Azapagic, A., Grossmann, I.E., Mac Dowell, N., Guillén-Gosálbez,  
 587 G., 2018. Time for global action: an optimised cooperative approach towards effective climate  
 588 change mitigation. *Energy & Environmental Science* 11, 572–581.

589 Government of Canada, 2022. Greenhouse Worker in Ontario - Job prospects. [https:](https://www.jobbank.gc.ca/marketreport/outlook-occupation/23339/ON)  
 590 [//www.jobbank.gc.ca/marketreport/outlook-occupation/23339/ON](https://www.jobbank.gc.ca/marketreport/outlook-occupation/23339/ON). [Online; accessed 22-  
 591 December-2021].

592 Hayse, P.L., Marion, W.W., 1973. Eviscerated yield, component parts, and meat, skin and bone  
 593 ratios in the chicken broiler. *Poultry Science* 52, 718–722.

594 Health Canada, 2008. Nutrient Value of Some Common Foods. [https://www.canada.ca/content/](https://www.canada.ca/content/dam/hc-sc/migration/hc-sc/fn-an/alt_formats/pdf/nutrition/fiche-nutri-data/nvscf-vnqau-eng.pdf)  
 595 [dam/hc-sc/migration/hc-sc/fn-an/alt\\_formats/pdf/nutrition/fiche-nutri-data/](https://www.canada.ca/content/dam/hc-sc/migration/hc-sc/fn-an/alt_formats/pdf/nutrition/fiche-nutri-data/nvscf-vnqau-eng.pdf)  
 596 [nvscf-vnqau-eng.pdf](https://www.canada.ca/content/dam/hc-sc/migration/hc-sc/fn-an/alt_formats/pdf/nutrition/fiche-nutri-data/nvscf-vnqau-eng.pdf). [Online; accessed 14-December-2021].

597 Health Canada, 202. Canadian Nutrient File. [https://food-nutrition.canada.ca/cnf-fce/](https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp)  
 598 [index-eng.jsp](https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp). [Online; accessed 15-July-2022].

599 IISD, 2018. Nutrient Recovery and Reuse in Canada: Foundations for a national framework. [https:](https://www.iisd.org/sites/default/files/meterial/nutrient-recovery-reuse-canada.pdf)  
 600 [//www.iisd.org/sites/default/files/meterial/nutrient-recovery-reuse-canada.pdf](https://www.iisd.org/sites/default/files/meterial/nutrient-recovery-reuse-canada.pdf).  
 601 [Online; accessed 25-August-2022].

602 INAC Services, 2014. Poultry processing in Canada. [http://inacservices.com/](http://inacservices.com/poultry-processing-in-canada/)  
 603 [poultry-processing-in-canada/](http://inacservices.com/poultry-processing-in-canada/). [Online; accessed 22-August-2022].

- 604 Jupp, A.R., Beijer, S., Narain, G.C., Schipper, W., Slootweg, J.C., 2021. Phosphorus recovery and  
605 recycling-closing the loop. *Chemical Society Reviews* 50, 87–101.
- 606 Kaiser, D.E., Pagliari, P., 2018. Understanding phosphorus fertilizers. [https://extension.](https://extension.umn.edu/phosphorus-and-potassium/understanding-phosphorus-fertilizers)  
607 [umn.edu/phosphorus-and-potassium/understanding-phosphorus-fertilizers](https://extension.umn.edu/phosphorus-and-potassium/understanding-phosphorus-fertilizers). [Online; ac-  
608 cessed 23-August-2022].
- 609 Martín-Hernández, E., Hu, Y., Zavala, V.M., Martín, M., Ruiz-Mercado, G.J., 2022. Analysis of  
610 incentive policies for phosphorus recovery at livestock facilities in the great lakes area. *Resources,*  
611 *Conservation and Recycling* 177, 105973.
- 612 Martín-Hernández, E., Martín, M., Ruiz-Mercado, G.J., 2021. A geospatial environmental and  
613 techno-economic framework for sustainable phosphorus management at livestock facilities. *Re-*  
614 *sources, Conservation and Recycling* 175, 105843.
- 615 National Aeronautics and Space Administration, 2022. Technology Readiness Level Definitions.  
616 [https://www.nasa.gov/pdf/458490main\\_TRL\\_Definitions.pdf](https://www.nasa.gov/pdf/458490main_TRL_Definitions.pdf). [Online; accessed 15-August-  
617 2022].
- 618 Nesme, T., Metson, G.S., Bennett, E.M., 2018. Global phosphorus flows through agricultural trade.  
619 *Global Environmental Change* 50, 133–141.
- 620 Ohtake, H., Tsuneda, S., 2019. Phosphorus recovery and recycling. Springer.
- 621 Oldfield, L., Rakhimbekova, S., Roy, J.W., Robinson, C.E., 2020. Estimation of phosphorus loads  
622 from septic systems to tributaries in the canadian lake erie basin. *Journal of Great Lakes Research*  
623 46, 1559–1569.
- 624 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2011. Growing Green-  
625 house Vegetables in Ontario.
- 626 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2020a. Ontario egg  
627 production. <https://data.ontario.ca/dataset/ontario-egg-production>. [Online; accessed  
628 14-December-2021].

629 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2020b.  
630 Ontario milk shipments by county. [https://data.ontario.ca/en/dataset/](https://data.ontario.ca/en/dataset/ontario-milk-shipments-by-county)  
631 [ontario-milk-shipments-by-county](https://data.ontario.ca/en/dataset/ontario-milk-shipments-by-county). [Online; accessed 14-December-2021].

632 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2021. Greenhouse  
633 Nutrient Feedwater Regulation. [http://www.omafra.gov.on.ca/english/nm/regs/gnfpro/](http://www.omafra.gov.on.ca/english/nm/regs/gnfpro/gnfreg.htm)  
634 [gnfreg.htm](http://www.omafra.gov.on.ca/english/nm/regs/gnfpro/gnfreg.htm). [Online; accessed 23-December-2021].

635 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2022. Horticultural  
636 Crops. <http://www.omafra.gov.on.ca/english/stats/hort/index.html>. [Online; accessed  
637 18-December-2021].

638 Ontario Ministry of the Environment, Conservation and Parks, 2012. Greenhouse  
639 Wastewater Monitoring Project (2010 and 2011). [https://www.ontario.ca/page/](https://www.ontario.ca/page/greenhouse-wastewater-monitoring-project-2010-and-2011)  
640 [greenhouse-wastewater-monitoring-project-2010-and-2011](https://www.ontario.ca/page/greenhouse-wastewater-monitoring-project-2010-and-2011). [Online; accessed 22-  
641 December-2021].

642 Ontario Ministry of the Environment, Conservation and Parks, 2019. Provincial policy objectives for  
643 managing effects of cage aquaculture operations on the quality of water and sediment in Ontario's  
644 waters. <https://ero.ontario.ca/notice/012-7186>. [Online; accessed 15-July-2022].

645 Ontario Ministry of the Environment, Conservation and Parks, 2021. Municipal Treated Wastewater  
646 Effluent. <https://data.ontario.ca/dataset/municipal-treated-wastewater-effluent>.  
647 [Online; accessed 29-December-2021].

648 Opendatasoft, 2019. Census divisions - Canada. [https://public.opendatasoft.com/](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario)  
649 [explore/dataset/georef-canada-census-division/table/?disjunctive.prov\\_name\\_](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario)  
650 [en&disjunctive.cd\\_name\\_en&sort=year&refine.prov\\_name\\_en=Ontario](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario). [Online; accessed  
651 29-November-2021].

652 Pingali, P.L., 2012. Green revolution: impacts, limits, and the path ahead. Proceedings of the  
653 National Academy of Sciences 109, 12302–12308.

654 Pollution Probe, 2022. Mapping Phosphorus Flows in the Ontario Economy. Exploring Nutrient  
655 Recovery and Reuse Opportunities in a Provincial Context. [https://www.pollutionprobe.org/  
656 mapping-phosphorus-flows-in-the-ontario-economy/](https://www.pollutionprobe.org/mapping-phosphorus-flows-in-the-ontario-economy/). [Online; accessed 13-July-2022].

657 Reid, K., Schneider, K., Joosse, P., 2019. Addressing imbalances in phosphorus accumulation in  
658 canadian agricultural soils. *Journal of environmental quality* 48, 1156–1166.

659 Reid, K., Schneider, K.D., 2019. Phosphorus accumulation in canadian agricultural soils over 30  
660 yr. *Canadian Journal of Soil Science* 99, 520–532.

661 Sampat, A.M., Hicks, A., Ruiz-Mercado, G.J., Zavala, V.M., 2021. Valuing economic impact  
662 reductions of nutrient pollution from livestock waste. *Resources, Conservation and Recycling*  
663 164, 105199.

664 Sampat, A.M., Hu, Y., Sharara, M., Aguirre-Villegas, H., Ruiz-Mercado, G., Larson, R.A., Zavala,  
665 V.M., 2019. Coordinated management of organic waste and derived products. *Computers &  
666 chemical engineering* 128, 352–363.

667 Sampat, A.M., Martin-Hernandez, E., Martín, M., Zavala, V.M., 2018. Technologies and logistics  
668 for phosphorus recovery from livestock waste. *Clean Technologies and Environmental Policy* 20,  
669 1563–1579.

670 Samreen, S., Kausar, S., 2019. Phosphorus fertilizer: The original and commercial sources, in:  
671 Zhang, T. (Ed.), *Phosphorus*. IntechOpen. chapter 6.

672 Sardans, J., Peñuelas, J., 2013. Tree growth changes with climate and forest type are associated  
673 with relative allocation of nutrients, especially phosphorus, to leaves and wood. *Global Ecology  
674 and Biogeography* 22, 494–507.

675 Schoumans, O., Rulkens, W., Oenema, O., Ehlert, P., 2010. Phosphorus recovery from animal  
676 manure: Technical opportunities and agro-economical perspectives. Technical Report. Alterra,  
677 Wageningen UR.

678 Senthilkumar, K., Nesme, T., Mollier, A., Pellerin, S., 2012. Conceptual design and quantification  
679 of phosphorus flows and balances at the country scale: The case of France. *Global Biogeochemical*  
680 *Cycles* 26.

681 Sharma, B.K., Chandel, M.K., 2021. Life cycle cost analysis of municipal solid waste management  
682 scenarios for mumbai, india. *Waste Management* 124, 293–302.

683 Statistics Canada – Statistique Canada, 2021a. Cattle and calves on census day. [https:](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042401)  
684 [//www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042401](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042401). [Online; accessed 29-  
685 November-2019].

686 Statistics Canada – Statistique Canada, 2021b. Other livestock on census day. [https://www150.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042701)  
687 [statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042701](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042701). [Online; accessed 29-November-  
688 2019].

689 Statistics Canada – Statistique Canada, 2021c. Pigs on census day. [https://www150.statcan.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042601)  
690 [gc.ca/t1/tbl1/en/tv.action?pid=3210042601](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042601). [Online; accessed 29-November-2019].

691 Statistics Canada – Statistique Canada, 2021d. Poultry inventory on census day. [https:](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042801)  
692 [//www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042801](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042801). [Online; accessed 29-  
693 November-2019].

694 Statistics Canada – Statistique Canada, 2021e. Sheep and lambs on census day. [https:](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042501)  
695 [//www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042501](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042501). [Online; accessed 29-  
696 November-2019].

697 Statistics Canada – Statistique Canada, 2022. Fertilizer shipments to Canadian agriculture markets,  
698 by nutrient content and fertilizer year, cumulative data (x 1,000). [https://www150.statcan.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210003901)  
699 [gc.ca/t1/tbl1/en/tv.action?pid=3210003901](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210003901). [Online; accessed 09-February-2022].

700 Statistics Canada - Statistique Canada, 2015. Sewer and septic system connections. House-  
701 holds and the Environment. [https://www150.statcan.gc.ca/n1/pub/11-526-x/2013001/](https://www150.statcan.gc.ca/n1/pub/11-526-x/2013001/t059-eng.htm)  
702 [t059-eng.htm](https://www150.statcan.gc.ca/n1/pub/11-526-x/2013001/t059-eng.htm). [Online; accessed 15-July-2022].



703 Statistics Canada - Statistique Canada, 2017. Estimated number of households and average house-  
704 hold size by domain, Canada. User guide for the survey of household spending. <https://www150.statcan.gc.ca/n1/pub/62f0026m/2017002/app-ann-g-eng.htm>. [Online; accessed  
705 15-July-2022].  
706

707 Statistics Canada - Statistique Canada, 2021. Aquaculture, production and value. [https://www150.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210010701)  
708 [statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210010701](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210010701). [Online; accessed 15-July-2022].

709 Statistics Canada - Statistique Canada, 2022a. Interprovincial and international trade flows,  
710 basic prices, summary level. [https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1210008801)  
711 [1210008801](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1210008801). [Online; accessed 13-July-2022].

712 Statistics Canada - Statistique Canada, 2022b. Livestock estimates. [https://www150.statcan.gc.](https://www150.statcan.gc.ca/n1/daily-quotidien/220228/dq220228d-cansim-eng.htm)  
713 [ca/n1/daily-quotidien/220228/dq220228d-cansim-eng.htm](https://www150.statcan.gc.ca/n1/daily-quotidien/220228/dq220228d-cansim-eng.htm). [Online; accessed 15-August-  
714 2022].

715 Statistics Canada - Statistique Canada, 2022c. Population estimates, quarterly. [https://www150.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901)  
716 [statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901). [Online; accessed 15-July-2022].

717 Statistics Canada - Statistique Canada, 2022d. Selected livestock and poultry, Census of Agriculture  
718 historical data. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210015501>.  
719 [Online; accessed 18-August-2022].

720 Statistics Canada - Statistique Canada, 2022e. Trade Data Online. [https://www.ic.gc.ca/app/](https://www.ic.gc.ca/app/scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng)  
721 [scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng](https://www.ic.gc.ca/app/scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng). [Online; accessed 15-July-2022].

722 Statistics Netherlands, 2012. Standardised calculation methods for animal manure and nutrients.  
723 Standard data 1990–2008. Technical Report. Statistics Netherlands.

724 Stelco Inc., 2022. Our Facilities. <https://www.stelco.com/about-us/our-facilities>. [Online;  
725 accessed 13-July-2022].

726 Szögi, A., Vanotti, M., Hunt, P., 2008. Phosphorus recovery from poultry litter. Transactions of  
727 the ASABE 51, 1727–1734.

728 Tan, C., Zhang, T., 2011. Surface runoff and sub-surface drainage phosphorus losses under regular  
 729 free drainage and controlled drainage with sub-irrigation systems in southern ontario. Canadian  
 730 Journal of Soil Science 91, 349–359.

731 The World Bank, 2022. Annual Prices - Commodity Markets. [https://www.worldbank.org/en/  
 732 research/commodity-markets](https://www.worldbank.org/en/research/commodity-markets). [Online; accessed 23-August-2022].

733 Towler, G., Sinnott, R., 2013. Chemical engineering design: principles, practice and economics of  
 734 plant and process design. Butterworth-Heinemann.

735 United Nations Environment Programme, 2022. Resolution adopted by the United Nations Envi-  
 736 ronment Assembly on 2 March 2022, UNEP/EA.5 Res.2, on Sustainable nitrogen management.

737 United States Department of Agriculture, 2009. Agricultural Wate Management Field Handbook.  
 738 Technical Report. United States Department of Agriculture (USDA).

739 U.S. Census Bureu, 2021. Current versus Constant (or Real) Dollars. [https://www.census.  
 740 gov/topics/income-poverty/income/guidance/current-vs-constant-dollars.html](https://www.census.gov/topics/income-poverty/income/guidance/current-vs-constant-dollars.html). [On-  
 741 line; accessed 23-August-2022].

742 U.S. Department of Agriculture, 2011. Animal Feeding Operations (AFO) and Concentrated  
 743 Animal Feeding Operations (CAFO). [https://www.nrcs.usda.gov/wps/portal/nrcs/main/  
 744 national/plantsanimals/livestock/afo/](https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/plantsanimals/livestock/afo/). [Online; accessed 10-August-2020].

745 Van Dijk, K.C., Lesschen, J.P., Oenema, O., 2016. Phosphorus flows and balances of the European  
 746 Union Member States. Science of the Total Environment 542, 1078–1093.

747 Van Staden, T., Van Meter, K., Saurette, D., Basu, N., Parsons, C., Van Cappellen, P., 2021.  
 748 Anthropogenic phosphorus mass balance in ontario counties and watersheds. 10.20383/101.0208.

749 Villalba, G., Liu, Y., Schroder, H., Ayres, R.U., 2008. Global phosphorus flows in the industrial  
 750 economy from a production perspective. Journal of Industrial ecology 12, 557–569.

751 Wang, Y., Zhang, T., Tan, C., Qi, Z., Welacky, T., 2018. Solid cattle manure less prone to  
 752 phosphorus loss in tile drainage water. Journal of environmental quality 47, 318–325.

- 753 World Integrated Trade Solution, 2022. Data on Export, Import, Tariff. [https://wits.worldbank.](https://wits.worldbank.org/)  
754 [org/](https://wits.worldbank.org/). [Online; accessed 13-July-2022].
- 755 Yang, J., De Jong, R., Drury, C., Huffman, E., Kirkwood, V., Yang, X., 2007. Development of a  
756 canadian agricultural nitrogen budget (canb v2. 0) model and the evaluation of various policy  
757 scenarios. *Canadian Journal of Soil Science* 87, 153–165.
- 758 Yokoyama, K., Kubo, H., Mori, K., Okada, H., Takeuchi, S., Nagasaka, T., 2007. Separation and  
759 recovery of phosphorus from steelmaking slags with the aid of a strong magnetic field. *ISIJ*  
760 *international* 47, 1541–1548.
- 761 Zagklis, D., Konstantinidou, E., Zafiri, C., Kornaros, M., 2020. Assessing the economic viability of  
762 an animal byproduct rendering plant: Case study of a slaughterhouse in greece. *Sustainability*  
763 12, 5870.
- 764 Zhang, T., Tan, C., Zheng, Z., Drury, C., 2015. Tile drainage phosphorus loss with long-term  
765 consistent cropping systems and fertilization. *Journal of environmental quality* 44, 503–511.
- 766 Zhou, J., Jiao, X., Ma, L., de Vries, W., Zhang, F., Shen, J., 2021. Model-based analysis of  
767 phosphorus flows in the food chain at county level in china and options for reducing the losses  
768 towards green development. *Environmental Pollution* 288, 117768.