

# Phosphorus in Ontario's economic sectors: mapping flows and assessing recovery and recycling potential

Université Laval team<sup>a,\*</sup>, McGill University team<sup>b</sup>, University of Waterloo team<sup>c</sup>

<sup>a</sup> *Université Laval*

<sup>b</sup> *McGill University*

<sup>c</sup> *University of Waterloo*

---

## Abstract

The dual dimension of the anthropogenic use of phosphorus, i.e., its key role in the food production system and the negative environmental impacts associated with the phosphorus used in intensive agricultural techniques, has been stated by the United Nations Environment Assembly. In addition, phosphorus is a non-renewable material which reserves are concentrated in a few number of regions, making global supply chains vulnerable to regional events and conflicts. As a consequence, the recovery and recycling of phosphorus is not just a desirable but also a necessary approach to assure a sustainable, reliable, and sovereign food production system. In this work we map the phosphorus flows through the economic sectors of the Canadian province of Ontario, and phosphorus recovery and recycling opportunities are identified. These mainly belong to the agricultural sector, including manure (30.5 kt/year) and slaughterhouse waste (3.7 kt/year), although significant amounts of P are also found in food and organic waste, including municipal wastewater (6.4 kt/year). Different scenarios are studied to determine the amount of phosphorus that could be recovered within the province considering according with the technology readiness level of different phosphorus recovery processes, as well as the costs associated with phosphorus recovery [Add some more numbers here](#). Finally, we discuss the implications that would be derived from implementing active phosphorus recovery and recycling approaches regarding phosphorus supply and use in Ontario.

---

\*Corresponding author

Email address: [edgar.martin-hernandez.1@ulaval.ca](mailto:edgar.martin-hernandez.1@ulaval.ca) (Université Laval team)

## 8 1. Introduction

9 Phosphorus is an essential for production of food which has been intensively used for crop and  
10 livestock production since the development of synthetic fertilizers in the XIX and XX centuries  
11 ([Samreen and Kausar, 2019](#)). The combination of synthetic fertilizers with other modern inten-  
12 sive agricultural techniques have increased the productivity of agriculture and farming industries  
13 ([Pingali, 2012](#)). However, the intensive use of fertilizers in agriculture has resulted in the over-  
14 application of phosphorus in many regions worldwide REF, while the run of intensive livestock  
15 production facilities, also known as concentrated animal feeding operations (CAFOs) ([U.S. Depart-  
16 ment of Agriculture, 2011](#)), result in important difficulties in the management of the large amounts  
17 of manure produced, which is often spread in lands in the vicinity of CAFOs, which also leads to  
18 the accumulation of phosphorus in soil. Soil acts as a phosphorus reservoir ([Ehlert et al., 2003](#)),  
19 building-up a legacy P that can be used for future crops, but also can be transported to waterbodies  
20 by erosion and runoff leading to the eutrophication of aquatic ecosystems.

21 The dual dimension of the anthropogenic use of phosphorus, i.e., its key role in the food pro-  
22 duction system and the negative environmental impacts associated with the phosphorus used in  
23 intensive agricultural techniques, has been stated by the United Nations Environment Assembly in  
24 the resolution UNEP/EA.5/Res.2 ([United Nations Environment Programme, 2022](#)). An additional  
25 factor to be considered for addressing the phosphorus challenge is the non-renewable nature of  
26 phosphorus, since the phosphorus consumed is not replenished by natural means at human time  
27 scale, and there is currently no known synthetic substitute for this material ([Cordell et al., 2009](#)).  
28 Since the global phosphorus reserves are concentrated in a few number of regions, the supply of  
29 phosphorus from a limited number of global supply chains lacks resiliency and it has been proven  
30 that it can be globally disrupted by regional events and conflicts, resulting geopolitical tensions  
31 ([Food and Agriculture Organization of the United Nations, 2022](#)). As a consequence, the recov-  
32 ery and recycling of phosphorus is not just a desirable but also a necessary approach to assure a  
33 sustainable, reliable, and sovereign food production system.

34 Although the main uses of phosphorus are in the agri-food sector, phosphorus is also involved in  
35 other industrial activities, including steel, chemical, and forestry industries. Henceforth, phosphorus

is a key material for many aspects of human development. As a result, mapping the phosphorus flows involved in human activities to detect opportunities for recovery and recycling is essential for, in a second stage, assess amount of phosphorus that is viable to recover, the economical costs involved, and the enhancement in terms of resiliency of the regional food production system, savings from the reduction of phosphorus imports, and the mitigation of phosphorus pollution on the region implementing strategies for phosphorus recovery and recycling. The quantification of phosphorus flows has been addressed in previous works in the literature for certain sectors such as the agri-food sector (Boh and Clark, 2020; Zhou et al., 2021; Nesme et al., 2018). Additionally, studies on the global phosphorus flows have also been performed (Villalba et al., 2008; Chen and Graedel, 2016), although these studies tend to have a low flow resolution since these are aggregated by major sectors. Additionally, the works quantifying phosphorus often include qualitative recommendations to improve the phosphorus use efficiency and recycling (Van Dijk et al., 2016; Senthilkumar et al., 2012), but often they do not include quantitative assessments on the amount of phosphorus which recovery is feasible along with the costs involved. Conversely, those works focused on estimating the recoverable phosphorus and the associated recovery cost target specific flows, lacking a holistic perspective of the phosphorus flows in the various human activities (Martín-Hernández et al., 2021; Sampat et al., 2018).

In this work, we intend to perform a holistic approach to the opportunities for phosphorus recovery and recycling in the Canadian province of Ontario. In a first stage, we proceed to map the phosphorus flows involved in the economical sectors of Ontario, i.e., the agri-food, industrial, and urban sectors. This data is used in a second stage to identify the flows in which phosphorus recovery is feasible, determining the amount of phosphorus that could be recovered within the province considering different scenarios regarding the technology readiness level of different phosphorus recovery processes, as well as the costs associated with phosphorus recovery. Finally, we discuss the implications that would be derived from implementing active phosphorus recovery and recycling approaches regarding phosphorus supply and use in Ontario.

## 2. Methods

### 2.1. Spatial resolution

Phosphorus flows have been mapped within the Canadian province of Ontario, and thus the political borders of Ontario has been considered as the boundaries for the substance flow analysis performed. In those cases where the data was available, the distribution of phosphorus flows within Ontario has also been studied at Census Division level ([Statistics Canada – Statistique Canada, 2017](#)). The database collecting the IDs of Ontario Census Divisions, their names, and geospatial information is taken from [Opendatasoft \(2019\)](#).

ADD MAP WITH CENSUS DIVISIONS????

### 2.2. Temporal resolution

The study has being performed for year 2019 since the most of data required is available for this year. In addition, the temporal evolution of the largest phosphorus flows, i.e., agricultural and wastewater phosphorus flows, has been studied for a period of 13 years from 2007 to 2019.

### 2.3. Estimation of phosphorus flows

The estimation of phosphorus flows within the Ontario’s agricultural sectors is based on open data sources, often from governmental institutions, complemented with information from scientific articles when needed. The particular procedure followed for each flow depends on the information publicly available. In the next sections we depict the main lines of the estimating procedure for each sector, while we refer the reader to [Pollution Probe \(2022\)](#) for a more comprehensive description of the procedure followed for estimating each phosphorus flow.

#### 2.3.1. Agricultural sector

Phosphorus flows in the agricultural sector are estimated based on production data of livestock and crop products, as well as data on fertilizer application.

For those production data were not available, a number of different methods were used to estimate the P flow based on approaches established in the literature. For example, P inflows associated

with synthetic fertilizers could be directly estimated based on application data reported in the Fertilizer Shipments Survey (FSS).<sup>37</sup> Conversely, P flows associated with manure were determined indirectly by accounting for the magnitude from which the flow of P could be derived. In this case,

Phosphorus in livestock imports and exports is estimated from livestock trading data REF, multiplying the number of animals by the concentration of phosphorus in the different types of livestock REF,

Phosphorus in livestock feeding and manure is estimated based on the number and type of animals reported for Ontario at Census Division level in the Census of Agriculture REF!, multiplied by the phosphorus feeding requirements REF, and concentration of phosphorus in manure REF. The Census of Agriculture is published by Statistics Canada every five years (i.e., 2001, 2006, 2011, and 2016) for cattle<sup>52</sup> REF, sheep<sup>53</sup> REF, swine<sup>54</sup> REF, poultry<sup>55</sup> REF, and other livestock<sup>56</sup> REF, with the exception of rabbits, where data is not available prior to 2009. The number of animals for the years in between census reporting have been estimated using a linear interpolation. We assumed that the number of animals reported is throughout the year (i.e., the animals culled are replaced by new ones). However, in the case of broilers and turkeys, the number of animals reported by the livestock census have been reduced by a factor of 0.68 (broilers) and 0.80 (turkeys), since these animals have life cycles of 43 and 80 days respectively, meaning barns are empty for 20 days between cycles. <sup>305</sup> REF

Phosphorus contained in meat and slaughterhouse waste is based on the number of animals slaughtered reported by both federally and provincially licensed meat plants.<sup>59, 60</sup> REF multiplied by the concentration of phosphorus in carcasses REF.

Phosphorus flows associated with the production of milk and eggs is based on provincial production data, multiplying these products by their average phosphorus concentration <sup>57, 58</sup> REF.

Phosphorus in fertilizer applied to open fields in Ontario is estimated based on the amount of fertilizer products traded to Ontario's agricultural markets containing P <sup>100</sup> REF. The distribution of phosphorus fertilizers among the Census Division of the province is based on the fraction of fertilized area of each census division, i.e., dividing the reported area of land fertilized for each census division by the total fertilized area of land in Ontario, removing the areas that correspond

115 with greenhouse crops<sup>101, 102 103 REF</sup>. Regarding manure, we assume that all of the manure  
116 generated by livestock is applied in crop fields <sup>50 REF</sup>.

117 The uptake of phosphorus by crops is determined based on the area used in each Census Division  
118 to grow each type of crops by census division<sup>104, 105, 106</sup> and its yield<sup>107, 108</sup> multiplied by the  
119 specific P content for each crop type.<sup>109, 110</sup> The phosphorus uptake by crops is divided according  
120 to whether it uptake in the grain, fruit or vegetable, or straw and stover components of each type  
121 of crop. This is necessary to determine the amount of phosphorus that flows within food or feed  
122 (i.e., grains, fruits and vegetables) while straw and stover remain in the field after harvesting as crop  
123 residues.

124 A fraction of the phosphorus applied to crop fields as manure of synthetic fertilizer is lost  
125 through erosion, runoff, and drainage. This transportation of phosphorus depends on a range  
126 of factor, including the amount of phosphorus applied; soil composition, texture, and slope; and  
127 precipitation, resulting in a complex and data-intensive process for estimating the phosphorus  
128 transported out of the crop fields. As an approximation, we have estimated the phosphorus losses  
129 by using export coefficients determined for crop fields in Ontario <sup>112 REF 113 REF</sup> corrected to  
130 account for both surface and subsurface runoffs for synthetic fertilizers (1.267 kg/ha/year ), and  
131 liquid and solid manure (2.548 kg/ha/year and 1.717 kg/ha/year respectively) <sup>113 REF</sup> ([Pollution  
132 Probe, 2022](#)).

133 A fraction of the P supplied to crop fields is not taken up by the plants and remains in soil,  
134 resulting in the accumulation of P over time as a result of synthetic fertilizer and manure over  
135 over sustained periods of time, often applying phosphorus in greater quantities than crops require  
136 to ensure satisfactory yields <sup>132 REF</sup>. This buildup is often referred to as “legacy P”, and it is  
137 estimated as the balance between phosphorus inflows to crop fields (application of manure and  
138 synthetic fertilizers) and outflows (crop food and feed products, crop residues, and phosphorus  
139 losses).

140 Regarding greenhouse crops, the data available was limited, resulting in an estimation of phos-  
141 phorus applied as synthetic fertilizers based on the sum of phosphorus uptake by greenhouse crops  
142 (i.e., tomatoes, peppers, and cucumbers)<sup>119</sup> and the phosphorus releases from greenhouse irrigation

systems (greenhouse nutrient feedwater systems (GNF) REF ONTARIO) systems. The phosphorus uptake by greenhouse crops is determined by multiplying the production of greenhouse crops REF by the phosphorus content of each vegetable type 121, 122 REF. The phosphorus releases from the GNF systems was estimated based on the average concentration of phosphorus in GNF outlet streams of Ontario (33.6 mg/L) 123 REF and the total water discharges from GNF systems REF, assuming that water discharges from GNF systems is equivalent to 25% of the total water applied in greenhouses, which corresponding with the worst-case scenario of no water recirculation in the GNF system. The average water consumption in greenhouses in Ontario was assumed to be 1,000 L/m<sup>2</sup>/year 125 REF. We have also estimated the phosphorus releases from the seasonal workers live in households in the vicinity of the greenhouses that may use septic systems, considering that the seasonal labour force in Ontario greenhouses is estimated to be 6,699 workers REF, and an average phosphorus load rate of 0.0156 kg P/person/week from septic systems REF.

### 2.3.2. Industrial sector

Phosphorus flows through imports, production, exports and waste for the steel, forestry, and food and beverage, industries of Ontario were mapped. The steel industry is the first non-food sector in terms of phosphorus use. The main phosphorus inflows of steel manufacturing are associated with the use of iron ore, coal, and coke, while the main outflow of phosphorus is within slag, which remove most of the impurities from steel, including phosphorus. It must be noted that, although some minor amounts of phosphorus can be desired in steel for making anti-corrosion surface coatings, it is largely considered an impurity in the steel manufacturing process. Phosphorus in these flows is estimated multiplying their average phosphorus content (0.06% P in iron ore, 0.05% P for coal, 0.4% P in slag, and 0.01% in steel) 176 REF by the steel production capacity of the facilities located in Ontario ([Cheminfo Services Inc., 2019](#); [Algoma Steel Inc., 2022](#); [Stelco Inc., 2022](#); [Pollution Probe, 2022](#)) and the imports and exports of these materials ([World Integrated Trade Solution, 2022](#); [Statistics Canada - Statistique Canada, 2022](#)).

Phosphorus flows in Ontario's forestry industry includes wood harvesting, wood products man-

170 ufacturing, as well as the production of pulp and paper. The estimation of these phosphorus flows  
171 are the result of multiplying the production data of wood, wood products, pulp and paper, and  
172 their retrospectives imports, exports, and waste streams ([Canadian Forest Service, 2020](#); [Statistics  
173 Canada - Statisque Canada, 2022](#)), by the average phosphorus content, which is assumed to be  
174 0.01% for wood 181 REF and 0.005% for pulp and paper products REF.

175 Regarding other industrial activities which could involve the use of phosphorus, the local pro-  
176 duction of phosphorus is assumed to be negligible since phosphorus is not mined or refined in  
177 Ontario, and the synthetic phosphorus fertilizer imports are accounted in the agricultural section.  
178 The general chemical facilities located in Ontario report 350 t/year of phosphorus as waste REF,  
179 in addition of imports and exports of chemical products REF. However, there exist a significant  
180 fraction of untracked phosphorus used in the industrial sector.

### 181 *2.3.3. Urban sector*

## 182 **3. Results and discussion**

### 183 *3.1. Phosphorus flows in Ontario*

184 *Showing an overview of the P flows in the province. The use of figures summarizing all the flows*  
185 *of the province in the shape of Sankey or network flow figures could be so great*






Figure 1: Phosphorus flows in the province of Ontario (kt/year). The streams within red rectangles denote phosphorus inflows into the province, while those streams within blue rectangles denote phosphorus outflows out of the province.

186 Figure 1 summarizes the phosphorus flows in the province of Ontario. It can be observed that  
187 the flow of of phosphorus through the anthropogenic activities are divided into 3 independent  
188 networks, i.e., the flow of phosphorus involved the production and processing of food (including the  
189 treatment of wastewater), the flow of phosphorus used in the steel and chemical industries, and the  
190 phosphorus involved in the forestry industry.

191 The production of animal food products exhibits a lower phosphorus use efficiency than the  
192 production of plant base products, similarly to the use efficiency of other resources such as water  
193 CITE HERE, CALCULAR ENTRA VS SALE!

### 194 3.2. Phosphorus recovery techniques (*This section could be Supplementary Material*)

195 Brief overview of potential P recovery techniques for each sector

#### 196 3.2.1. Agricultural sector

#### 197 3.2.2. Industrial sector

#### 198 3.2.3. Urban sector

### 199 3.3. Potential of phosphorus recovery in Ontario

200 Assessment of different scenarios of P recovery in Ontario, P imports that would be saved,  
201 reduction of P dependency of the province, etc (all implications related with mass-balances)

### 202 3.4. Economic implications of phosphorus recovery in Ontario

203 Economic costs or saving derived from the recovery of P in the province and all implications  
204 related with economy

### 205 3.5. Implications on food sovereignty of phosphorus recovery in Ontario

206 Implications on food production self-sufficiency derived from the (partial) recycling of P. Discus-  
207 sion on the improvement of the food production system resiliency against disruptions of the global  
208 supply supply chains (e.g., current context derived from the COVID-19 pandemia and the war in  
209 Ukraine)

210 3.6. *Gaps of knowledge*

## 211 4. Conclusions

## 212 5. Acknowledgments

213 Pollution Probe

214 ECCC

## 215 References

216 Algoma Steel Inc., 2022. Corporate Profile. [https://algoma.com/about-algoma/  
217 corporate-profile/](https://algoma.com/about-algoma/corporate-profile/). [Online; accessed 13-July-2022].

218 Boh, M.Y., Clark, O.G., 2020. Nitrogen and phosphorus flows in Ontario’s food systems. *Resources,  
219 Conservation and Recycling* 154, 104639.

220 Canadian Forest Service, 2020. Statistical data. <https://cfs.nrcan.gc.ca/statsprofile/>. [On-  
221 line; accessed 13-July-2022].

222 Cheminfo Services Inc., 2019. Economic Assessment of the Inte-  
223 grated Steel Industry. [https://www.canadiansteel.ca/files/resources/  
224 Final-Report-Economic-Assessment-of-the-Integrated-Steel-Industry.pdf](https://www.canadiansteel.ca/files/resources/Final-Report-Economic-Assessment-of-the-Integrated-Steel-Industry.pdf). [Online;  
225 accessed 13-July-2022].

226 Chen, M., Graedel, T., 2016. A half-century of global phosphorus flows, stocks, production, con-  
227 sumption, recycling, and environmental impacts. *Global Environmental Change* 36, 139–152.

228 Cordell, D., Drangert, J.O., White, S., 2009. The story of phosphorus: global food security and  
229 food for thought. *Global environmental change* 19, 292–305.

230 Ehlert, P., Morel, C., Fotyma, M., Destain, J.P., 2003. Potential role of phosphate buffering capacity  
231 of soils in fertilizer management strategies fitted to environmental goals. *Journal of plant nutrition  
232 and soil science* 166, 409–415.

233 Food and Agriculture Organization of the United Nations, 2022. The importance of Ukraine and  
 234 the Russian Federation for global agricultural markets and the risks associated with the war in  
 235 Ukraine.

236 Martín-Hernández, E., Martín, M., Ruiz-Mercado, G.J., 2021. A geospatial environmental and  
 237 techno-economic framework for sustainable phosphorus management at livestock facilities. *Re-*  
 238 *sources, Conservation and Recycling* 175, 105843.

239 Nesme, T., Metson, G.S., Bennett, E.M., 2018. Global phosphorus flows through agricultural trade.  
 240 *Global Environmental Change* 50, 133–141.

241 Opendatasoft, 2019. Census divisions - Canada. [https://public.opendatasoft.com/](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario)  
 242 [explore/dataset/georef-canada-census-division/table/?disjunctive.prov\\_name\\_](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario)  
 243 [en&disjunctive.cd\\_name\\_en&sort=year&refine.prov\\_name\\_en=Ontario](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario). [Online; accessed  
 244 29-November-2021].

245 Pingali, P.L., 2012. Green revolution: impacts, limits, and the path ahead. *Proceedings of the*  
 246 *National Academy of Sciences* 109, 12302–12308.

247 Pollution Probe, 2022. Mapping Phosphorus Flows in the Ontario Economy. Exploring Nutrient  
 248 Recovery and Reuse Opportunities in a Provincial Context. [https://www.pollutionprobe.org/](https://www.pollutionprobe.org/mapping-phosphorus-flows-in-the-ontario-economy/)  
 249 [mapping-phosphorus-flows-in-the-ontario-economy/](https://www.pollutionprobe.org/mapping-phosphorus-flows-in-the-ontario-economy/). [Online; accessed 13-July-2022].

250 Sampat, A.M., Martin-Hernandez, E., Martín, M., Zavala, V.M., 2018. Technologies and logistics  
 251 for phosphorus recovery from livestock waste. *Clean Technologies and Environmental Policy* 20,  
 252 1563–1579.

253 Samreen, S., Kausar, S., 2019. Phosphorus fertilizer: The original and commercial sources, in:  
 254 Zhang, T. (Ed.), *Phosphorus*. IntechOpen. chapter 6.

255 Senthilkumar, K., Nesme, T., Mollier, A., Pellerin, S., 2012. Conceptual design and quantification  
 256 of phosphorus flows and balances at the country scale: The case of France. *Global Biogeochemical*  
 257 *Cycles* 26.

258 Statistics Canada – Statistique Canada, 2017. Census division (CD). [https://www150.](https://www150.statcan.gc.ca/n1/pub/92-195-x/2016001/geo/cd-dr/cd-dr-eng.htm)  
259 [statcan.gc.ca/n1/pub/92-195-x/2016001/geo/cd-dr/cd-dr-eng.htm](https://www150.statcan.gc.ca/n1/pub/92-195-x/2016001/geo/cd-dr/cd-dr-eng.htm). [Online; accessed 23-  
260 November-2021].

261 Statistics Canada - Statistique Canada, 2022. Interprovincial and international trade flows,  
262 basic prices, summary level. [https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1210008801)  
263 [1210008801](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1210008801). [Online; accessed 13-July-2022].

264 Stelco Inc., 2022. Our Facilities. <https://www.stelco.com/about-us/our-facilities>. [Online;  
265 accessed 13-July-2022].

266 United Nations Environment Programme, 2022. Resolution adopted by the United Nations Envi-  
267 ronment Assembly on 2 March 2022, UNEP/EA.5 Res.2, on Sustainable nitrogen management.

268 U.S. Department of Agriculture, 2011. Animal Feeding Operations (AFO) and Concentrated  
269 Animal Feeding Operations (CAFO). [https://www.nrcs.usda.gov/wps/portal/nrcs/main/](https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/plantsanimals/livestock/afo/)  
270 [national/plantsanimals/livestock/afo/](https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/plantsanimals/livestock/afo/). [Online; accessed 10-August-2020].

271 Van Dijk, K.C., Lesschen, J.P., Oenema, O., 2016. Phosphorus flows and balances of the European  
272 Union Member States. *Science of the Total Environment* 542, 1078–1093.

273 Villalba, G., Liu, Y., Schroder, H., Ayres, R.U., 2008. Global phosphorus flows in the industrial  
274 economy from a production perspective. *Journal of Industrial ecology* 12, 557–569.

275 World Integrated Trade Solution, 2022. Data on Export, Import, Tariff. [https://wits.worldbank.](https://wits.worldbank.org/)  
276 [org/](https://wits.worldbank.org/). [Online; accessed 13-July-2022].

277 Zhou, J., Jiao, X., Ma, L., de Vries, W., Zhang, F., Shen, J., 2021. Model-based analysis of  
278 phosphorus flows in the food chain at county level in china and options for reducing the losses  
279 towards green development. *Environmental Pollution* 288, 117768.