

Mapping of phosphorus flows and analysis of the potential for recovery and reuse in Ontario, Canada

Edgar Martín-Hernández^{a,*}, Jorge A. Garcia Hernandez^d, Samantha Gangapersad^c, Tian Zhao^c, Roy Brouwer^{d,e}, Sidney Omelon^c, Céline Vaneeckhaute^{a,b}

^a*BioEngine - Research Team on Green Process Engineering and Biorefineries, Chemical Engineering Department, Université Laval, 1065 Ave. de la Médecine, Québec, QC, G1V 0A6, Canada*

^b*CentrEau, Centre de recherche sur l'eau, Université Laval, 1065 Avenue de la Médecine, Québec, QC, G1V 0A6, Canada*

^c*Department of Mining and Materials Engineering, McGill University, Montréal, Canada*

^d*Department of Economics, University of Waterloo, 200 University Avenue West, Waterloo, ON, N2L 3G1, Canada*

^e*The Water Institute, University of Waterloo, 200 University Avenue West, Waterloo, ON, N2L 3G1, Canada*

Abstract

Phosphorus is a key non-renewable element used in multiple economic activities, and notably for food production. However, sustained releases over time have led to nutrient pollution and eutrophication of ecosystems. In this work, we map the phosphorus flows through Ontario's economic sectors, and we identify the potential for phosphorus recovery and recycling. These mainly target flows associated with food production and processing, including wastewater and food waste. Up to 86% of phosphorus imports for food production could be covered by recycled phosphorus, with an average recovery cost of 36 EUR/kg. This is lower than the estimated economic losses derived from the release of phosphorus into the environment, but significantly higher than fossil-based commercial phosphorus products. Additionally, there exist a wide variation on phosphorus recovery costs from different streams, which suggest the need of exploring cooperative approaches for effective phosphorus recovery at regional scale.

Keywords: Phosphorus recovery, Circular economy, Nutrient pollution, Eutrophication, Food sovereignty

1. Introduction

Phosphorus is an essential element for the production of food. It has been intensively used for crop and livestock production since the development of synthetic fertilizers and feed supplements

*Corresponding author
Email address: edmah2@ulaval.ca (Edgar Martín-Hernández)

19 in the XIX and XX centuries ([Samreen and Kausar, 2019](#)). The combination of synthetic fertilizers
20 with other modern intensive agricultural techniques have increased the productivity of agriculture
21 and farming industries ([Pingali, 2012](#)). However, the intensive use of fertilizers in agriculture has
22 resulted in the over-application of phosphorus ([Reid and Schneider, 2019](#)), while the run of intensive
23 livestock operations, result in important difficulties in the management of the large amounts of
24 manure produced. This is often spread on lands in the vicinity of the livestock operations, which
25 in turn leads to the accumulation of phosphorus in the soil. Although soil acts as a phosphorus
26 reservoir ([Ehlert et al., 2003](#)), building-up a legacy P that can be used for future crops, it can
27 also be transported to waterbodies by erosion and runoff, resulting in the eutrophication of aquatic
28 ecosystems.

29 The dual dimension of the anthropogenic use of phosphorus, i.e., its key role in the food pro-
30 duction system and the negative environmental impacts associated with the phosphorus used in
31 intensive agricultural techniques, has been stated by the United Nations Environment Assembly in
32 the resolution UNEP/EA.5/Res.2 ([United Nations Environment Programme, 2022](#)). An additional
33 factor to be considered for addressing the phosphorus challenge is the non-renewable nature of
34 phosphorus, since the phosphorus consumed is not replenished by natural means at human time
35 scale, and there is currently no known synthetic substitute for this material ([Cordell et al., 2009](#)).
36 Since phosphorus reserves are concentrated in a few number of regions, the supply of phosphorus
37 from a limited number of suppliers lacks resilience and it has been proven that it can be globally
38 disrupted by regional events and conflicts ([Food and Agriculture Organization of the United Na-
39 tions, 2022](#)). As a consequence, the recovery and recycling of phosphorus is not just a desirable but
40 also a necessary approach to assure sustainable, reliable, and sovereign food production systems.

41 Although the main uses of phosphorus are in the agri-food sector, phosphorus is also involved
42 in other industrial activities, including steel, chemical, and forestry industries. Henceforth, phos-
43 phorus is a key material for many aspects of human development. As a result, the mapping of
44 phosphorus flows involved in human activities to detect opportunities for recovery and recycling
45 is essential to determine the fraction of phosphorus which recovery is viable, promoting a circular
46 economy that enhances the sustainability of food production systems in terms of resiliency, savings

47 from the reduction of phosphorus imports, and mitigation of phosphorus pollution. on the regions
48 implementing strategies for phosphorus recovery and recycling. The detailed quantification of phos-
49 phorus flows has been addressed in the literature for certain sectors, particularly for the agri-food
50 sector (Boh and Clark, 2020; Zhou et al., 2021; Nesme et al., 2018). In addition, phosphorus flows
51 have also been studied at global (Villalba et al., 2008; Chen and Graedel, 2016) and national scales
52 (Van Dijk et al., 2016; Klinglmair et al., 2015), although these studies tend to aggregate the flows
53 by major sectors, resulting in a lower flow resolution.

54 The works quantifying phosphorus often include qualitative recommendations to improve the
55 phosphorus use efficiency and recycling (Van Dijk et al., 2016; Senthilkumar et al., 2012), but they
56 do not include quantitative assessments on the amount of phosphorus that is feasible to recover along
57 with the costs involved. Conversely, those works focused on estimating the recoverable phosphorus
58 and the associated recovery costs target specific activities such as livestock production (Martín-
59 Hernández et al., 2021; Sampat et al., 2018) and wastewater treatment (Egle et al., 2016; Nättorp
60 et al., 2017). However, a holistic approach mapping the phosphorus flows and identifying the key
61 streams for phosphorus recovery and reuse is a crucial stage to promote the debate about global
62 and regional circular nutrient economies and redistribution systems (Kahiluoto et al., 2021), and to
63 design important aspects of future phosphorus recycling strategies such as the design of coordinated
64 markets (Sampat et al., 2019) and incentive policies (Martín-Hernández et al., 2022).

65 In this work, we intend to perform a holistic approach to phosphorus management, recovery,
66 and recycling through the study of the Canadian province of Ontario. In a first stage, we proceed
67 to map the phosphorus flows involved in the economical sectors of Ontario, i.e., the agricultural,
68 industrial, and urban sectors. These data is used in a second stage to identify the flows in which
69 phosphorus recovery is feasible, estimating the amount of phosphorus that could be recovered within
70 the province. Different phosphorus recovery technologies with technology readiness levels equal or
71 above 6 are evaluated, as well as the phosphorus recovery costs of each one of them. Finally, we
72 discuss the implications that would be derived from implementing active phosphorus recovery and
73 recycling approaches regarding phosphorus supply and use in Ontario.

2. Methods

2.1. Spatial and temporal boundaries

Phosphorus flows have been mapped through a material flow analysis (MFA) (Brunner and Rechberger, 2016) conducted within the political boundaries of the Canadian province of Ontario using data reported for the year 2019.

2.2. Estimation of phosphorus flows

The estimation of phosphorus flows in Ontario’s economic sectors is based on the use open data sources, often from governmental institutions, complemented with information from scientific articles when needed. In the next sections, we describe the general procedure followed to estimate the phosphorus flows of each sector. For a comprehensive description of the procedure followed for estimating each particular phosphorus flow, we refer the reader to the methodology described in Pollution Probe (2022).

2.2.1. Agriculture and aquaculture sector

Phosphorus flows in the agricultural sector are estimated based on production data of livestock and crop products, as well as on fertilizer application data.

Phosphorus in livestock feeding and manure is estimated based on the number and type of animals reported for Ontario in the Census of Agriculture, including cattle (Statistics Canada – Statistique Canada, 2021a), swine (Statistics Canada – Statistique Canada, 2021e), poultry (Statistics Canada – Statistique Canada, 2021f), and other livestock (Statistics Canada – Statistique Canada, 2021g,d), multiplied by the phosphorus feeding requirements and concentration of phosphorus in manure (Statistics Netherlands, 2012; Brown, Christine, 2013; Van Staden et al., 2021). We assumed that the number of animals reported is throughout the year (i.e., the animals culled are replaced by new ones). However, in the case of broilers and turkeys, the number of animals reported by the livestock census have been reduced by a factor of 0.68 (broilers) and 0.80 (turkeys), since these animals have life cycles of 43 and 80 days respectively, meaning barns are empty for 20 days between cycles (Yang et al., 2007).

100 Phosphorus flows through the imports and exports of animals is estimated using data on ani-
101 mal imports and exports ([Statistics Canada – Statistique Canada, 2021b,c,h](#)) multiplied by their
102 phosphorus to live weight ratios ([Statistics Netherlands, 2012](#)).

103 Phosphorus contained in meat and slaughterhouse waste is based on the number of animals
104 slaughtered reported by both federally and provincially licensed meat plants ([Agriculture and Agri-
105 Food Canada, 2021c,b](#)) multiplied by the concentration of phosphorus in carcasses ([Agriculture
106 and Agri-Food Canada, 2021d](#); [Hayse and Marion, 1973](#); [Brake et al., 1995](#); [Statistics Netherlands,
107 2012](#)).

108 Phosphorus flows associated with the production of milk and eggs is based on provincial produc-
109 tion data ([Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2020a,b](#)),
110 multiplying these products by their average phosphorus concentration ([Health Canada, 2008](#); [Cham-
111 bers et al., 2017](#)).

112 Phosphorus applied to open fields as synthetic fertilizer is estimated based on the amount
113 of fertilizer products traded to Ontario’s agricultural markets containing phosphorus ([Statistics
114 Canada – Statistique Canada, 2022](#)). Regarding manure, we assume that all of the manure generated
115 by livestock is applied in crop fields ([van Bochove et al., 2010](#)).

116 The uptake of phosphorus by crops is determined based on the area used in each census division
117 ([Opendatasoft, 2019](#)) to grow each type of crops by census division ([Agriculture and Agri-Food
118 Canada, 2022a,b,c](#)) multiplied by the specific yield and phosphorus content for each crop type
119 ([United States Department of Agriculture, 2009](#)). The phosphorus uptake by crops is divided
120 according to whether it uptake in the grain, fruit or vegetable, or straw and stover components of
121 each type of crop. This is necessary to determine the amount of phosphorus that flows within food
122 or feed (i.e., grains, fruits and vegetables), while straw and stover remain in the field after harvesting
123 as crop residues.

124 A fraction of the phosphorus applied to crop fields as manure of synthetic fertilizer is lost through
125 erosion, runoff, and drainage. The magnitude of this flow depends on a range of factors, including
126 the amount of phosphorus applied; soil composition, texture, and slope; and precipitations, resulting
127 in a complex and data-intensive process for estimating the phosphorus transported out of the crop

fields. As an approximation, we have estimated the phosphorus losses by using export coefficients determined for crop fields in Ontario corrected to account for both surface and subsurface runoffs for synthetic fertilizers (1.267 kg/ha/year), and liquid and solid manure (2.548 kg/ha/year and 1.717 kg/ha/year respectively) (Zhang et al., 2015; Wang et al., 2018; Tan and Zhang, 2011). In addition, a fraction of the P supplied to crop fields is not taken up by the plants and remains in soil, resulting in the accumulation of P over time as a result of synthetic fertilizer and manure over sustained periods of time, often applying phosphorus in greater quantities than crops require to ensure satisfactory yields (Reid et al., 2019). This buildup is often referred to as “legacy P”, and it is estimated as the balance between phosphorus inflows to crop fields (application of manure and synthetic fertilizers) and outflows (crop food and feed products, crop residues, and phosphorus losses by erosion and runoff).

Regarding greenhouse crops, the data available was limited, resulting in an estimation of phosphorus applied as synthetic fertilizers based on the sum of phosphorus uptake by greenhouse crops and phosphorus releases from greenhouse irrigation systems also known as greenhouse nutrient feed-water (GNF) systems (Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2021). The phosphorus uptake by greenhouse crops is determined by multiplying the production of greenhouse crops (Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2022) by the phosphorus content of each vegetable type (United States Department of Agriculture, 2009). The phosphorus releases from the GNF systems was estimated based on the average concentration of phosphorus in GNF outlet streams for Ontario, 33.6 mg/L (Ontario Ministry of the Environment, Conservation and Parks, 2012), and the total water discharges from GNF systems, assuming that the water discharges are equivalent to 25% of the total water applied in greenhouses, which corresponds with the worst-case scenario of no water recirculation in the GNF systems (Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2021). The average water consumption in greenhouses in Ontario was assumed to be 1,000 L/m²/year (Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2011). We have also estimated the phosphorus releases from the seasonal workers live in households in the vicinity of the greenhouses that may use septic systems, considering that the seasonal labour force in Ontario greenhouses is

156 estimated to be 6,699 workers ([Government of Canada, 2022](#)), and an average phosphorus load rate
157 of f 0.0156 kg P/person/week from septic systems ([Oldfield et al., 2020](#)).

158 Phosphorus in aquaculture are mainly due to supply of feed as part of fish feed the grow of
159 trouts, part of which is uptake by fishes, while the rest of phosphorus is released into aquatic
160 ecosystems since aquaculture effluents are directly discharged to the environment ([Ontario Ministry
161 of the Environment, Conservation and Parks, 2019](#)). The amount phosphorus uptakes by fishes is
162 calculated multiplying the fish production ([Statistics Canada - Statistique Canada, 2021](#)), by their
163 phosphorus content ([Health Canada, 202](#)), while the phosphorus content in the aquaculture waste
164 effluents of Ontario is estimated to be 10 kg of phosphorus per ton of fish produced ([Bureau et al.,
165 2003](#)). The sum of phosphorus uptakes by fishes and phosphorus in aquaculture waste effluents
166 result in the phosphorus supplied to aquaculture as fish feed.

167 *2.2.2. Industrial sector*

168 Phosphorus flows through imports, production, exports and waste for the food, steel, and
169 forestry industries of Ontario were mapped.

170 Processed food imports and exports are estimated scaling each type of food traded in Canada
171 ([Statistics Canada - Statistique Canada, 2022e](#)) with the population of Ontario ([Statistics Canada -
172 Statistique Canada, 2022c](#)). The phosphorus contained in each type of imported and exported food
173 is estimated multiplying the amount of ech type of traded food by its phosphorus content ([Health
174 Canada, 202](#)). Phosphorus flows in the form of food and organic waste are based on applying food
175 loss factors for the steps associated with food processing, from the production of food raw materials
176 to consumption ([FAO, 2011](#)), considering the food production and import values estimated in
177 Section 2.2.1.

178 The steel industry is the first non-food sector in terms of phosphorus use. The main phosphorus
179 inflows of steel manufacturing are associated with the use of iron ore, coal, and coke, while the
180 main outflow of phosphorus is within slag, which remove most of the impurities from steel, includ-
181 ing phosphorus. It must be noted that, although some minor amounts of phophosphorus can be
182 desired in steel for making anti-corrosion surface coatings, it is largely considered an impurity in

183 the steel manufacturing process. Phosphorus in these flows is estimated multiplying their average
184 phosphorus content (0.06% P in iron ore, 0.05% P for coal, 0.4% P in slag, and 0.01% in steel)
185 ([Yokoyama et al., 2007](#)) by the steel production capacity of the facilities located in Ontario ([Chem-](#)
186 [info Services Inc., 2019](#); [Algoma Steel Inc., 2022](#); [Stelco Inc., 2022](#); [Pollution Probe, 2022](#)) and the
187 imports and exports of these materials ([World Integrated Trade Solution, 2022](#); [Statistics Canada](#)
188 [- Statistique Canada, 2022a](#)).

189 Phosphorus flows in Ontario’s forestry industry includes wood harvesting, wood products man-
190 ufacturing, as well as the production of pulp and paper. The estimation of these phosphorus flows
191 are the result of multiplying the production data of wood, wood products, pulp and paper, and
192 their retrospectives imports, exports, and waste streams ([Canadian Forest Service, 2020](#); [Statistics](#)
193 [Canada - Statistique Canada, 2022a](#)), by the average phosphorus content, which is assumed to be
194 0.01% for wood ([Sardans and Peñuelas, 2013](#)) and 0.005% for pulp and paper products.

195 The general chemical facilities located in Ontario report 350 t/year of phosphorus as waste
196 ([Pollution Probe, 2022](#)), in addition of imports and exports of chemical products. However, there
197 exist a significant fraction of phosphorus used in the industrial sector that cannot be tracked due to
198 the lack of data. Regarding other industrial activities which could involve the use of phosphorus,
199 the local production of phosphorus is assumed to be negligible since phosphorus is not mined or
200 refined in Ontario, and the synthetic phosphorus fertilizer imports are accounted in the agricultural
201 section.

202 *2.2.3. Urban sector*

203 In this section we include the phosphorus inflows and outflows through wastewater treatment
204 plants (WWTPs), septic systems, and food and organic waste management facilities (landfills,
205 composting sites, and anaerobic digestion facilities).

206 Phosphorus flows through WWTPs is estimated combining data from the National Pollutant
207 Release Inventory (NPRI) ([Environment and Climate Change Canada, 2021a](#)), a public database
208 of releases, disposals and transfers of pollutants , and data from the Wastewater Systems Effluent
209 Regulations (WSER) database ([Environment and Climate Change Canada, 2021b](#)). Since the NPRI

only contains data of those facilities that meet certain regulatory requirements, the information of this database must be complemented with the data from the WSER database, which includes information of Canadian WWTPs at federal, provincial, and municipal levels. The estimations on phosphorus flows through WWTPs are validated using the Municipal Treated Wastewater Effluent (MTWE) database (Ontario Ministry of the Environment, Conservation and Parks, 2021), which collects annual data on water quality data and effluent levels for WWTPs in Ontario. We note that this data set only provides information about phosphorus releases from municipal WWTPs, but it does not collect phosphorus flows through sludge disposals. This methodology is shown in Figure 1.




Figure 1: Procedure for estimating phosphorus flows through wastewater treatment plants.

There exist households that are not connected to any sewer systems, but they are equipped with septic systems to perform a rough treatment of the wastewater produced prior to its release into the environment. This typically consist into a septic tank that separates solid matter from the wastewater, and a drainfield where the effluent is discharged. The estimation of phosphorus releases from septic systems is based on the fraction of households equipped with these systems, estimated

on 13% (Statistics Canada - Statistique Canada, 2015), which are inhabited by an average of 2.58 individuals (Statistics Canada - Statistique Canada, 2017). The average phosphorus load rate from septic systems assumed is 0.81 kg of phosphorus per person per year (Oldfield et al., 2020).

2.3. Phosphorus recovery techniques

There exist different processes for phosphorus recovery from different sources which technical viability has been proven or is at advanced development stage, i.e., systems with technologies readiness level (TRL) (National Aeronautics and Space Administration, 2022) of 6 or above (commercial or pilot plant stage). Since the flows from different processes have different properties, the techniques for phosphorus recovery vary between sectors and flows and, therefore, their recovery efficiencies, costs, and products obtained are different. Table A.1, which is included in the Appendix, shows a summary of the specifications of the phosphorus recovery technologies for different flows, including literature references where comprehensive descriptions of each system and its specifications can be found. We note that the phosphorus recovery processes currently available exceed the systems included in this work, nonetheless, the processes considered in this study are a selection of the main techniques for phosphorus recovery. However, different processes may have been developed on the foundations of the same technique, e.g., the multiple processes are based on struvite precipitation.

Phosphorus recovery costs include operating and annualized capital costs. Capital costs are annualized through the application of an annual capital charge ratio ($ACCR$) as defined by Towler and Sinnott (2013), assuming a typical interest rate of 5% and a plant lifetime of 20 years. Dynamic phosphorus recovery costs in function of the processing capacity have been considered in order to capture the economies of scale for those technologies for which sufficient data are available.

In general terms, the processes for phosphorus recovery and recycling can be classified into those employed for the treatment of liquid streams, which are based on the formation of precipitates through the direct treatment of the liquid effluent, and those processes employed for the treatment of solid fractions, which usually require the pretreatment of the waste, e.g., through an incineration stage.

Phosphorus in manure represent an important flow within the agricultural sector. The tech-

251 niques used for phosphorus recovery from cattle and swine manure differ for the liquid and solid
252 fractions. Struvite precipitation is the dominant technology for phosphorus recovery from liquid
253 manure, existing different processes for struvite production based on the type of reactors used with
254 similar recovery efficiencies but different treatment capacities, and thus different recovery costs
255 ([Martín-Hernández et al., 2021](#)). Additionally, there exist modular processes based on physical
256 separations oriented to small-scale intensive livestock facilities ([Church et al., 2018](#)). The recovery
257 of phosphorus from the solid fraction of manure involves the incineration of the waste, and the
258 further processing of the ashes, recovering phosphorus precipitates or phosphoric acid ([Jupp et al.,](#)
259 [2021](#); [Egle et al., 2016](#)). Phosphorus recovery from poultry litter is based on acid extraction and
260 further precipitation ([Szögi et al., 2008](#)).

261 Slaughterhouse waste is a flow from the food processing industry which can be targeted for
262 phosphorus recovery. It should be noted that slaughterhouse is comprised by a liquid (slaughter-
263 house wastewater) and a solid fraction (animal carcass waste). Similarly to phosphorus recovery
264 from liquid manure, phosphorus recovery from slaughterhouse wastewater is performed through
265 through struvite precipitation ([AMPC, 2018](#)), while the animal animal carcass waste is incinerated
266 and phosphorus is recovered from ashes in form of calcium carbonate or phosphoric acid ([Jupp](#)
267 [et al., 2021](#)).

268 Municipal wastewater contains significant amounts of phosphorus that can be recovered. It
269 must be noted that phosphorus outflows from WWTPs are divided into phosphorus contained
270 in the treated water and phosphorus contained in sludge. Phosphorus contained in water can
271 be recovered through the formation of precipitates such as struvite or calcium phosphate, while
272 phosphorus contained in sludge can be recovered either through the direct processing of sludge
273 producing precipitates, or from sludge ashes after an incineration stage, obtaining different products
274 such as phosphoric acid or calcium phosphate.

275 3. Results and discussion

276 3.1. Phosphorus flows in Ontario

277 Figure 2 summarizes the phosphorus flows in the province of Ontario. It can be observed
278 that phosphorus flows in anthropogenic activities can be divided into 3 independent networks, i.e.,
279 the flow of phosphorus involved the production and processing of food (including the treatment
280 of organic solid waste and wastewater), the flow of phosphorus used in the steel and chemical
281 industries, and the phosphorus involved in the forestry industry.

282 The main inflows of the food production and processing network are those associated with the
283 supply of synthetic fertilizers and livestock feed, accounting for 55.6 and 21.8 kt/year respectively.
284 Other phosphorus imports in Ontario are made in the form of food products (4.7 kt/year) and
285 aquaculture feed (0.07 kt/year). Crop residues and manure are the main phosphorus waste ef-
286 fluent from the agricultural sector, accounting for 32.5 and 30.6 kt/year respectively. While crop
287 residues are usually left in the cropfields, transferring part of the phosphorus taken by crops back
288 to soil and acting as soil amendment materials, manure produced in intensive livestock operations
289 is a spatially concentrated point source of phosphorus releases, resulting in the accumulation of
290 phosphorus in the vicinity of these facilities. As a consequence, the production of manure can re-
291 sult in negative environmental impacts and requires of adequate management strategies. The food
292 processing industry involves the largest flows within the province, which can be classified as plant
293 and animal-based food products, and slaughterhouse waste, resulting in phosphorus flows of 43.2,
294 10.3, and 3.8 kt/year respectively. A significant fraction of end-flows are waste flows in the form of
295 landfill (37.5 kt/year), composting (2.1 kt/year), and anaerobic digestion (1.4 kt/year) disposals,
296 as well as wastewater (7.8 kt/year). Phosphorus exports out of Ontario are performed through the
297 exports of food products and livestock, accounting for 10.8 and 1.8 kt/year respectively.




Figure 2: Phosphorus flows in the province of Ontario (kt/year). The streams within red rectangles denote phosphorus inflows into the province, while those streams within blue rectangles denote phosphorus outflows out of the province.

298 Chemical industry involve significant phosphorus flows, importing 22.2 kt/year into the province,
299 while the exports represent 14.0 kt/year. This sector produces 0.35 kt/year of phosphorus that is
300 classified as waste. However, an important fraction of phosphorus used by this sector (11.7 kt/ton)
301 cannot be tracked and, therefore, it is unknown what is the real amount of phosphorus disposed
302 as waste. Steel production imports 8.3 kt/year of phosphorus, of which 7.0 kt/year ends as steel
303 waste or slag, while the phosphorus outflows through steel materials are 1.4 kt/year.

304 Phosphorus imports from the forestry industry are 0.38 kt/year, while 2.3 kton/year of phos-
305 phorus is taken from local wood harvested within Ontario. This sector releases 0.4 kton/year of
306 phosphorus in the form of wood and pulp waste, while 0.81 kt/year of phosphorus is exported out
307 of the province as wood, manufactured wood products, pulp, and paper.

308 *3.2. Potential of phosphorus recovery in Ontario*

309 The potential for phosphorus recovery in the province of Ontario is assessed in this section.
310 As shown in Section 2.3, different processes can be employed for the recovery of phosphorus from
311 the same stream. However, each system is designed for operating under certain conditions and
312 they have different processing capacities. As a result, phosphorus recovery efficiency and cost may
313 vary between technologies for the treatment of the same flow. In order to explore this variability
314 between phosphorus recovery systems, all the systems described in Table A.1 are evaluated. The
315 results obtained in terms of phosphorus recovered and recovery cost for each technology and flow
316 are collected in the Supplementary Material. Two scenarios are selected for deeper analysis, the
317 minimum cost scenario, in which the most economical technology is selected, and the maximum
318 recovery scenario, in which the phosphorus recovery system selected is the one recovery the largest
319 amount of phosphorus.




Figure 3: Phosphorus recovered and phosphorus recovery cost in the province of Ontario by flow. Note that the scale of bottom x-axes is different.




Figure 4: Average phosphorus recovered and average phosphorus recovery cost per kilogram of phosphorus recovered in the province of Ontario by flow.

3.2.1. Agricultural sector

Manure is an agricultural flow from which effective phosphorus recovery might be achieved since it can be collected from the intensive livestock operations and be further treated (Schoumans et al., 2010). The inventory of cattle, swine, poultry, and other livestock for Ontario in 2019 is 1,376,984, 506,768, 148,508, and 135,725 animal units respectively, resulting in the release of 13,274, 8,569, 6,457, and 2,283 metric tonnes of phosphorus per year through manure respectively. An animal unit is defined as an animal equivalent of 1,000 pounds (453.6 kg) live weight (U.S. Department of Agriculture, 2011).

Phosphorus recovery from manure is highly influenced by the economies of scale and, therefore, by the scale of the livestock operations (Martín-Hernández et al., 2021). Since no data on the size distribution of livestock operations in Ontario is available, the average sizes of livestock facilities reported by Statistics Canada - Statistique Canada (2022b) for the year 2019 are considered, resulting in average sizes for cattle, swine, poultry and other livestock (primarily sheep and lambs) facilities of 98, 168, 15 and 16 animal units respectively. The number of cattle, swine, poultry, and other livestock operations obtained is 14,051, 3,022, 10,069, and 8,636 respectively, which is in alignment with the number of livestock facilities reported by Statistics Canada - Statistique Canada (2022d).

Phosphorus is divided between the liquid and solid fractions of manure, being the solid fraction the one containing the largest amount of phosphorus, and thus the fraction from which larger quantities amounts of phosphorus can be recovered with lower costs, as it can be observed in Figure 3. However, it must be noted that phosphorus recovery from solid manure involved more complex processes, since they include the incineration of the waste, which in turn makes the process more energy intensive and may result in environmentally harmful emissions of gases. Cattle manure contains the largest amount of phosphorus as a consequence of being the largest manure flow, followed by swine and poultry manure. However, the comparison of the average phosphorus recovery costs per kilogram of phosphorus recovered shows that phosphorus recovery from swine liquid manure is more cost-effective, as shown in Figure 4. This is due to the size of the swine intensive facilities, which in average are comprised by a larger number of animal units than cattle intensive facilities. This reveals the important role of the economies of scale in phosphorus recovery. However, in

348 general terms the small size of the livestock operations in Ontario results in high phosphorus
 349 recovery costs, whose values range between 200 and 700 EUR/kg P recovered. These costs are
 350 significantly higher than the phosphorus recovery costs reported by [Martín-Hernández et al. \(2022\)](#)
 351 for the comparatively larger livestock operations of the U.S. states in the Great Lakes area, whose
 352 average sizes range from 630 and 2,600 animal units, resulting in phosphorus recovery costs between
 353 13 and 73 USD/kilogram of phosphorus recovered. The phosphorus recovery efficiency is similar for
 354 all livestock types since the processes selected are the modular physical separation system MAPHEX
 355 due to the small scale of the livestock facilities. It was not possible to evaluate the effect of the
 356 economies of scale for the processes intended to phosphorus recovery from the solid fraction of
 357 manure due to the lack of techno-economic data available for them.

358 *3.2.2. Industrial and urban sector*

359 Industrial and urban sectors are grouped since some flows belong to both sectors, particularly
 360 those related with wastewater and the organic fraction of industrial and municipal solid waste,
 361 including food waste.

362 Slaughterhouse waste is an industrial flow from which phosphorus can be effectively recovered
 363 ([AMPC, 2018](#)). Data on individual capacities for the slaughterhouses in Ontario is not available
 364 to estimate the effect of the economies of scale on the cost of phosphorus recovery. Therefore,
 365 average slaughterhouse capacities are considered, which values are 104,017, 802,186, and $14.4 \cdot$
 366 10^6 cattle, hog, and poultry heads slaughtered/(facility \cdot year) respectively ([Agriculture and Agri-](#)
 367 [Food Canada, 2021a; INAC Services, 2014](#)). Considering the inventory of slaughtered animals
 368 reported by [Agriculture and Agri-Food Canada \(2021c,b\)](#), 7, 6, and 17 cattle, hog, and poultry
 369 slaughtering facilities are obtained, with associated phosphorus flows of 317.4, 103.5, and 53.2
 370 metric tonnes/(facility \cdot year) respectively. Phosphorus flows from slaughtered sheep and rabbit are
 371 considered negligible due to the low number of animals slaughtered. Figure 3 shows that most of
 372 phosphorus can be recovered from the solid fraction of waste due to its larger phosphorus content.
 373 The variations between the minimum cost and maximum recovery scenarios are not significant for
 374 the solid slaughterhouse waste flow. However, for slaughterhouse wastewater phosphorus recovery

for the maximum recovery scenario increase by a factor of 2.3 over the minimum cost scenario, while the total recovery cost increases by a factor of 3.3, as it can be observed in Figure 4. Therefore, the increase of phosphorus recovery efficiency results in a non-linear increase in the phosphorus recovery cost. The numerical results are collected in the Supplementary Material.

Wastewater is a large flow containing significant amounts of phosphorus. Wastewater is collected and directed to wastewater treatment plants (WWTPs). These facilities produce a liquid water effluent with adequate environmental parameters for being released into the environment, and a sludge flow from the primary and second treatments. Phosphorus can be recovered from both flows, although most of the phosphorus is contained in the sludge fraction. The distribution of phosphorus assumed for treated liquid water and sludge is 14.1% - 85.9% respectively, based on the data reported by NPRI and WSER databases ([Environment and Climate Change Canada, 2021a,b](#)), which is in alignment with the distribution values reported by [Egle et al. \(2016\)](#). The capacity of the wastewater treatment plants installed in Ontario, together with their phosphorus flows, have been analyzed to determine the effect of the economies of scale in the cost of phosphorus recovery. Figure 3 shows that the potential for phosphorus recovery from sludge is greater than from the liquid fraction, as mentioned before. Little variation is observed between the minimum cost and maximum recovery scenarios for the recovery of phosphorus from sludge, which implies that there exist a certain degree of homogeneity in the current technologies for phosphorus recovery from sludge. This can also be appreciated in Figure 4. It must be noted that the phosphorus recovery systems from sludge ashes reveal to be more effective, including the incineration cost, than the direct recovery of phosphorus from sludge due to the higher recovery efficiency of the former ones. Conversely, the phosphorus recovery from the liquid wastewater fraction shows a larger variability between both scenarios considered. The phosphorus recovered in the maximum recovery scenario is 1.7 times larger than in the minimum cost scenarios. However, this increase in the phosphorus recovery efficiency results in the increase of the recovery cost by a factor of 9.6, showing that similarly to the case of the liquid fraction of slaughterhouse waste, the achievement of greater recovery efficiencies through the use of more effective technologies result in an exponential increase of recovery costs.

Figure 3 shows that the food and organic waste represent the largest potential for phosphorus recovery. However, it must be noted that this flow includes all those streams comprised by solid organic wastes other than slaughterhouse waste, including food processing waste, household food waste, and other municipal organic waste. Similarly to other solid fractions, the minimum cost and maximum recovery scenarios show a narrow variability regarding phosphorus recovered and recovery costs.

Finally, there are some processes under development for the recovery of phosphorus from steel production industry, particularly from steelmaking slag. These processes are based on selective leaching and further chemical precipitation to obtain solid phosphates (Du et al., 2022, 2019) or magnetic-aid calcium phosphate precipitation (Yokoyama et al., 2007). Although these are promising processes that can result in an effective recovery of phosphorus which can be further recycled, they are at early development stages and, thus, they have not been considered in this assessment.

3.3. Economic implications of phosphorus recovery in Ontario

In order to compare the costs derived from the recovery of anthropogenic phosphorus, Figure 5 shows the average phosphorus recovery cost in Ontario along with the long-term social and environmental economical losses derived from uncontrolled releases of phosphorus into the environment estimated by Sampat et al. (2021), and the temporal evolution of prices for different phosphorus commodities reported by the World Bank's Commodity Markets report (The World Bank, 2022), i.e., phosphate rock, diammonium phosphate (DAP), and triple superphosphate (TSP) together. The costs are normalized per mass unit of phosphorus, assuming that the phosphorus content of commercial phosphorus rock ranges from 28 to 34% in mass basis (FAO & IAEA, 2004; Kaiser and Pagliari, 2018), while phosphorus content of DAP and TSP is 20% and 19% respectively. The prices have been adjusted to the year 2020 through the Consumer Price Index (CPI) elaborated by the U.S. Census Bureau (U.S. Census Bureau, 2021).

It can be observed that the average cost of phosphorus recovery in Ontario, valued around 36 EUR/kg of phosphorus recovered, is significantly lower than the economic losses derived from

the release of phosphorus into the environment, estimated at 74.5 EUR/kg P in the context of state of Wisconsin, U.S. This provides an economical support to the recovery of phosphorus, since high long-term costs derived from the social and environmental damages caused by phosphorus releases into the environment could be avoided through the recovery of phosphorus from waste streams. Nevertheless, it can be observed that the cost of phosphorus obtained from recovery processes is more costly than phosphorus in commercial fertilizers obtained from mining, which reduces the economic incentives for the recovery and reuse of phosphorus. As a consequence, further support in form of environmental and/or agricultural regulations, phosphorus recovery incentives, or sustainability arguments are needed in order to promote the recovery and recycling of phosphorus.




Figure 5: Average phosphorus recovery cost in Ontario, economical losses derived from the releases of phosphorus into the environment (Sampat et al., 2021), and temporal evolution of prices for different phosphorus commodities. Note that the scale of y-axis is different for top and bottom axes.

Additionally, Figure 5 shows that average cost of phosphorus recovery within the province is around 36 EUR/kg of phosphorus recovered, however, in Figure 4 we can observe that phosphorus recovery costs can reach significantly higher values, particularly for the recovery of phosphorus from manure due to the lack of economies of scale of Ontario's livestock operations. Therefore, regional cooperative strategies could be implemented to develop a coordinated action against nutrient pollution, distributing the benefits and costs of phosphorus recovery, and prioritizing the

total utility rather than the utility of each stakeholder, similarly to the cooperative approaches proposed for minimizing the cost of reducing greenhouse gases emissions from electricity generation (Galán-Martín et al., 2018).

3.4. Potential for phosphorus recycling in Ontario

Phosphorus recovered can be further recycled within the food production system, developing a circular economy around the use of phosphorus. Phosphorus recovery and recycling would result in curbing the depletion of phosphorus rock reserves and the reduction of the dependency on phosphorus supply from other regions. Considering these factors, there exist some governmental initiatives that, through the creation of different forums and platforms, aim to promote the recovery and recycling of phosphorus (IISD, 2018; Pollution Probe, 2022). In addition, the European Union is setting specific targets to reduce the use of non-renewable materials in fertilizer production (European Comission, 2018) and to promote the use of waste-based fertilizers (European Comission, 2022), encouraging the effective recovery and recycling of phosphorus, and they could serve as a guideline to support phosphorus recovery in other regions.

The comparison between the phosphorus imported for the production of food within Ontario, i.e., in the form of fertilizer or animal feed, and the phosphorus that could be recovered, as it is shown in Figure 6, is a useful metric to determine the effectiveness of phosphorus recovery, and the potential reduction of phosphorus supplies from external sources. This assessment is limited to the food production sector depicted in Figure 2 since the recovery of phosphorus is performed in different flows belonging to this sector.




Figure 6: Phosphorus inputs for food production and phosphorus recovered per sector. S1 refers to the scenario minimizing phosphorus recovery cost, while S2 refers to the scenario maximizing phosphorus recovery.

Figure 6 shows that up to 88% of the phosphorus used for food production, excluding imports of food that have been produce elsewhere, could be recovered and recycled in the best case scenarios where no phosphorus losses occur between the recovery and the reuse stages. An effective recycling of phosphorus implies the redistribution of this material from the locations where it is released to the livestock operations where it is used for livestock raise, or to phosphorus-deficient cropfields, which in turn involves the transportation of phosphorus products as the last stage before the use final use of the recovered phosphorus. It must be considered that the transportation of phosphorus products result in additional costs that can hinder the recycling of phosphorus, as well as environmental impacts related with the current use of fossil fuels-based transportation vehicles. In this context, coordinated-market approaches have been proposed to optimize the spatial distribution of phosphorus from regions with surplus of phosphorus to P-deficient locations, minimizing the cost

476 and environmental footprint of the transportation of phosphorus products ([Sampat et al., 2019](#)).

477 **4. Conclusions**

478 Phosphorus is a key non-renewable element for many economic sectors, and particularly for the
479 production of food. The current linear economy scheme deplete phosphorus reserves, resulting in
480 supply dependencies from regions holding phosphorus rock reserves, and it is the sources of nutrient
481 pollution, eutrophication, and other environmental concerns relates with the end-of-life release of
482 phosphorus into the environment. As a consequence, the recovery and recycling of phosphorus
483 is not only a desirable but a necessary approach for the development of phosphorus sustainable
484 systems. For achieving this goal, the mapping of phosphorus across the different economic sectors
485 is the first stage to identify the main streams for phosphorus recovery. This information allows the
486 estimation of the potential for the recycling of phosphorus in a region.

487 For the case of Ontario, the best case scenario result in a phosphorus recycling potential up
488 to 86% over the phosphorus imported in the province for food production (i.e., excluding the
489 imports of livestock and food produced in other regions). An average phosphorus recovery cost is
490 estimated, although it shows a large variation among different flows. Phosphorus recovery costs
491 is particularly large for phosphorus recovery from manure due to the small scale of the livestock
492 facilities in Ontario. Conversely, phosphorus recovery costs estimated for other regions with larger
493 livestock facilities such as the U.S. Great Lakes area result in significantly lower values, showing the
494 important role of the economizes of scale for phosphorus recovery. Nevertheless, considering the
495 region studied as a whole, the average phosphorus recovery cost estimated is around 36 EUR/kg
496 P recovered, which is lower than the economic losses of phosphorus releases into the environment
497 estimated at 74.5 EUR/kg of phosphorus.

498 The wide difference of costs for the recovery of phosphorus from different flows suggest the need
499 develop regional cooperative mechanisms to distribute the phosphorus recovery costs. Moreover,
500 further research in the effective distribution of the phosphorus recovered from regions with phos-
501 phorus surplus to phosphorus-deficient locations is needed, although there exist some research on
502 the development of coordinated markets for phosphorus recovery and recycling.

503 **5. Acknowledgments**

504 Please add your acknowledgements

Appendix A. Phosphorus recovery technologies techno-economic data

Table A.1: Phosphorus recovery technologies considered in the study. F denotes the phosphorus recovered as kg $P_{\text{recovered}}/\text{year}$, while $[x]$ represent the ceiling function applied to x . The definition of annual capital charge ratio ($ACCR$) can be found in the Supplementary Material, Section 1.1. Refs: [1]: Martín-Hernández et al. (2021), [2]: Jupp et al. (2021), [3]: Egle et al. (2010), [5]: Szögi et al. (2008), [6]: AMPC (2018), [7]: Zagklis et al. (2020), [8]: Fernández-Delgado et al. (2022), [9]: Ohtake and Tsuneda (2019), [10]: Sharma and Chandel (2021)

Sector	Inflow	Pretreatment	Pretreatment cost (EUR/kg $P_{\text{recovered}}$)	Technology	Type	P recovery potential (% related to inflow)	P recovery cost (EUR/kg $P_{\text{recovered}}$)	TRL	Ref (tech)
Agriculture	Cattle and swine manure, liquid phase (30% of total manure P)	Solid-liquid separation (screw press)	See [1]	Multiform	Struvite	60	$25.7 + 1.10 \cdot 10^6 \cdot [1.19 \cdot 10^{-4} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Crystallator	Calcium phosphate	60	$3.53 + (2.30 \cdot 10^6 + 0.71 \cdot [3.32 \cdot 10^{-5} \cdot F]) \cdot ACCR \cdot \frac{1}{F}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Ostara Pearl 500	Struvite	60	$12.57 + 2.30 \cdot 10^6 \cdot [7.02 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Ostara Pearl 2K	Struvite	60	$12.57 + 3.10 \cdot 10^6 \cdot [1.83 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Ostara Pearl 10K	Struvite	60	$12.57 + 10.00 \cdot 10^6 \cdot [3.05 \cdot 10^{-6} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	Nurelys	Struvite	60	$10.37 + 1.38 \cdot 10^6 \cdot [2.24 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[1]
		Solid-liquid separation (screw press)	See [1]	MAPHEX	Solid	90	$184.67 + 0.30 \cdot 10^6 \cdot [2.47 \cdot 10^{-4} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	6	[1]
	Poultry litter	Incineration	8.9	EcoPhos	Phosphoric acid	82	4.5	6	[2,3,4]
		Incineration	8.9	AshDec depollution	Calcium phosphate	86	1.8	6	[2,3,4]
		Incineration	8.9	AshDec Rheumia	Calcium phosphate	86	1.9	6	[2,3,4]
		Incineration	8.9	PASCH	Calcium phosphate	79	4.7	6	[2,3,4]
		Incineration	8.9	LEACHPHOS	Calcium phosphate	78	5.1	9	[2,3,4]
		Incineration	8.9	RecoPhos	Mineral	87	2.5	9	[2,3,4]
		Incineration	8.9	Thermophos	P4	81	2.7	9	[2,3,4]
		-	-	Quick wash	Solid precipitate	70	4.4	3	[5]
	Slaughterhouse waste, liquid phase (14% of total slaughterhouse P)	-	-	Multiform	Struvite	84	$22.6 + 1.10 \cdot 10^6 \cdot [1.05 \cdot 10^{-4} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[6]
		-	-	Ostara Pearl 500	Struvite	58	$15.60 + 2.30 \cdot 10^6 \cdot [8.70 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[6]
		-	-	Ostara Pearl 2K	Struvite	58	$15.60 + 3.10 \cdot 10^6 \cdot [2.26 \cdot 10^{-5} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[6]
		-	-	Ostara Pearl 10K	Struvite	58	$15.60 + 10.00 \cdot 10^6 \cdot [4.53 \cdot 10^{-6} \cdot F] \cdot ACCR \cdot \frac{1}{F}$	9	[6]
Urban & industrial	Slaughterhouse waste, solid phase (86% of total slaughterhouse P)	Incineration	14.6	EcoPhos	Phosphoric acid	82	4.5	6	[2,3,7]
		Incineration	14.6	AshDec depollution	Calcium phosphate	86	1.8	6	[2,3,7]
		Incineration	14.6	AshDec Rheumia	Calcium phosphate	86	1.9	6	[2,3,7]
		Incineration	14.6	PASCH	Calcium phosphate	79	4.7	6	[2,3,7]
		Incineration	14.6	LEACHPHOS	Calcium phosphate	78	5.1	9	[2,3,7]
		Incineration	14.6	RecoPhos	Mineral	87	2.5	9	[2,3,7]
		Incineration	14.6	Thermophos	P4	81	2.7	9	[2,3,7]
	WWTPs (liquid phase)	-	-	Crystallator	Struvite/Calcium phosphate	38	$305,920 \cdot \left(\frac{F}{34,966}\right) \cdot \frac{1}{F}$	9	[3]
		-	-	Ostara Pearl	Struvite	20	$130,856 \cdot \left(\frac{F}{13,110}\right) \cdot \frac{1}{F}$	9	[3]
		-	-	P-RoC	Calcium phosphate	27	$75,970 \cdot \left(\frac{F}{17,759}\right) \cdot \frac{1}{F}$	6	[3]
		-	-	REM-NUT	Struvite	47	$977,933 \cdot \left(\frac{F}{30,579}\right) \cdot \frac{1}{F}$	6	[3]
	WWTPs (sewage sludge, 60-90% of P)	-	-	AirPrex	Struvite	15	$74,195 \cdot \left(\frac{F}{6,587}\right) \cdot \frac{1}{F}$	9	[3]
		-	-	PRISA	Struvite	18	$186,923 \cdot \left(\frac{F}{11,876}\right) \cdot \frac{1}{F}$	6	[3]
		-	-	Stuttgart process	Struvite	40	$581,730 \cdot \left(\frac{F}{36,756}\right) \cdot \frac{1}{F}$	9	[3]
		-	-	Giffhorn process	Struvite	40	$400,384 \cdot \left(\frac{F}{36,756}\right) \cdot \frac{1}{F}$	9	[3]
Urban & industrial	WWTPs (sewage sludge ash, SSA, 60-90% of P)	-	-	PHOXNAN	Struvite	51	$891,667 \cdot \left(\frac{F}{35,597}\right) \cdot \frac{1}{F}$	6	[3]
		-	-	Aqua Reci	Calcium phosphate	61	$939,605 \cdot \left(\frac{F}{40,077}\right) \cdot \frac{1}{F}$	6	[3]
		-	-	MEPREX	P rich slag	68	$1,154,473 \cdot \left(\frac{F}{44,676}\right) \cdot \frac{1}{F}$	6	[3]
		Incineration	8	EcoPhos	Phosphoric acid	82	4.5	6	[3]
		Incineration	8	AshDec depollution	Calcium phosphate	86	1.8	6	[3]
		Incineration	8	AshDec Rheumia	Calcium phosphate	86	1.9	6	[3]
		Incineration	8	PASCH	Calcium phosphate	79	4.7	6	[3]
		Incineration	8	LEACHPHOS	Calcium phosphate	78	5.1	9	[3]
		Incineration	8	RecoPhos	Mineral	87	2.5	9	[3]
		Incineration	8	Thermophos	P4	81	2.7	9	[3]
Urban & industrial	Organic municipal & food waste	Incineration	6.43	AshDec Rheumia	Calcium phosphate	86	1.9	6	[3,9,10]
		Incineration	6.43	PASCH	Calcium phosphate	79	4.7	6	[3,9,10]
		Incineration	6.43	LEACHPHOS	Calcium phosphate	78	5.1	9	[3,9,10]
		Incineration	6.43	RecoPhos	Mineral	87	2.5	9	[3,9,10]
		Incineration	6.43	Thermophos	P4	81	2.7	9	[3,9,10]

505 References

- 506 Agriculture and Agri-Food Canada, 2021a. Distribution of slaughtering activity.
507 [https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/
508 red-meat-and-livestock-market-information/slaughter-and-carcass-weights/
509 distribution-slaughtering-activity](https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/red-meat-and-livestock-market-information/slaughter-and-carcass-weights/distribution-slaughtering-activity). [Online; accessed 22-August-2022].
- 510 Agriculture and Agri-Food Canada, 2021b. Poultry slaughter reports. [https:
511 //agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/
512 poultry-and-egg-market-information/slaughter](https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/poultry-and-egg-market-information/slaughter). [Online; accessed 16-December-2021].
- 513 Agriculture and Agri-Food Canada, 2021c. Red meat and livestock slaughter and carcass weights.
514 [https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/
515 red-meat-and-livestock-market-information/slaughter-and-carcass-weights](https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/red-meat-and-livestock-market-information/slaughter-and-carcass-weights). [Online;
516 accessed 15-December-2021].
- 517 Agriculture and Agri-Food Canada, 2021d. Red meat conversion factors. [https:
518 //agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/
519 red-meat-and-livestock-market-information/slaughter-and-carcass-weights/
520 conversion-factors](https://agriculture.canada.ca/en/canadas-agriculture-sectors/animal-industry/red-meat-and-livestock-market-information/slaughter-and-carcass-weights/conversion-factors). [Online; accessed 16-December-2021].
- 521 Agriculture and Agri-Food Canada, 2022a. Field crops and hay, Census of Agriculture, 2011 and
522 2016. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210041601>. [Online; ac-
523 cessed 16-December-2021].
- 524 Agriculture and Agri-Food Canada, 2022b. Field vegetables,, Census of Agriculture, 2011 and 2016.
525 <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210041801>. [Online; accessed
526 16-December-2021].
- 527 Agriculture and Agri-Food Canada, 2022c. Greenhouse products and mushrooms,, 2011 and 2016.
528 <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042001>. [Online; accessed
529 16-December-2021].

530 Algoma Steel Inc., 2022. Corporate Profile. [https://algoma.com/about-algoma/](https://algoma.com/about-algoma/corporate-profile/)
 531 [corporate-profile/](https://algoma.com/about-algoma/corporate-profile/). [Online; accessed 13-July-2022].

532 AMPC, 2018. Struvite or Traditional Chemical Phosphorus Precipitation – What Option Rocks?
 533 https://www.ampc.com.au/uploads/cgblog/id408/2018-1026_-_Final_Report.pdf. [On-
 534 line; accessed 20-March-2019].

535 van Bochove, E., Thériault, G., Denault, J.T., 2010. Indicator of risk of water contamination by
 536 phosphorus (IROWC P): a handbook for presenting the IROWC P algorithms. Technical Report.
 537 Agriculture and Agri-Food Canada.

538 Boh, M.Y., Clark, O.G., 2020. Nitrogen and phosphorus flows in Ontario’s food systems. Resources,
 539 Conservation and Recycling 154, 104639.

540 Brake, J., Havenstein, G., Ferket, P., Rives, D., Giesbrecht, F., 1995. Relationship of sex, strain,
 541 and body weight to carcass yield and offal production in turkeys. Poultry science 74, 161–168.

542 Brown, Christine, 2013. Available Nutrients and Value for Manure From Various Livestock Types.
 543 Technical Report. Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs.

544 Brunner, P.H., Rechberger, H., 2016. Handbook of material flow analysis: For environmental,
 545 resource, and waste engineers. CRC press.

546 Bureau, D.P., Gunther, S.J., Cho, C.Y., 2003. Chemical composition and preliminary theoretical
 547 estimates of waste outputs of rainbow trout reared in commercial cage culture operations in
 548 Ontario. North American Journal of Aquaculture 65, 33–38.

549 Canadian Forest Service, 2020. Statistical data. <https://cfs.nrcan.gc.ca/statsprofile/>. [On-
 550 line; accessed 13-July-2022].

551 Chambers, J.R., Zaheer, K., Akhtar, H., Abdel-Aal, E.S.M., 2017. Chicken eggs, in: Egg innovations
 552 and strategies for improvements. Elsevier, pp. 1–9.

553 Cheminfo Services Inc., 2019. Economic Assessment of the Inte-
 554 grated Steel Industry. <https://www.canadiansteel.ca/files/resources/>

555 [Final-Report-Economic-Assessment-of-the-Integrated-Steel-Industry.pdf](#). [Online;
556 accessed 13-July-2022].

557 Chen, M., Graedel, T., 2016. A half-century of global phosphorus flows, stocks, production, con-
558 sumption, recycling, and environmental impacts. *Global Environmental Change* 36, 139–152.

559 Church, C.D., Hristov, A.N., Kleinman, P.J., Fishel, S.K., Reiner, M.R., Bryant, R.B., 2018. Ver-
560 satility of the MANure PHosphorus EXtraction (MAPHEX) System in Removing Phosphorus,
561 Odor, Microbes, and Alkalinity from Dairy Manures: A Four-Farm Case Study. *Applied En-
562 gineering in Agriculture* 34, 567–572. URL: [http://elibrary.asabe.org/abstract.asp?AID=](http://elibrary.asabe.org/abstract.asp?AID=48976&t=3&dabs=Y&redir=&redirType=)
563 [48976&t=3&dabs=Y&redir=&redirType=](http://elibrary.asabe.org/abstract.asp?AID=48976&t=3&dabs=Y&redir=&redirType=), doi:10.13031/aea.12632.

564 Cordell, D., Drangert, J.O., White, S., 2009. The story of phosphorus: global food security and
565 food for thought. *Global environmental change* 19, 292–305.

566 Du, C.M., Gao, X., Ueda, S., Kitamura, S.Y., 2019. Separation and recovery of phosphorus from
567 steelmaking slag via a selective leaching–chemical precipitation process. *Hydrometallurgy* 189,
568 105109.

569 Du, C.m., Gao, X., Ueda, S., Kitamura, S.y., 2022. Recovery of high-quality phosphate from
570 steelmaking slag by a hydrometallurgical process. *Science of The Total Environment* 819, 153125.

571 Egle, L., Rechberger, H., Krampe, J., Zessner, M., 2016. Phosphorus recovery from mu-
572 nicipal wastewater: An integrated comparative technological, environmental and economic
573 assessment of P recovery technologies. *Science of The Total Environment* 571, 522–542.
574 URL: <https://linkinghub.elsevier.com/retrieve/pii/S0048969716314656>, doi:10.1016/
575 [j.scitotenv.2016.07.019](https://linkinghub.elsevier.com/retrieve/pii/S0048969716314656).

576 Ehlert, P., Morel, C., Fotyma, M., Destain, J.P., 2003. Potential role of phosphate buffering capacity
577 of soils in fertilizer management strategies fitted to environmental goals. *Journal of plant nutrition*
578 and soil science 166, 409–415.

579 Environment and Climate Change Canada, 2021a. About the Na-
580 tional Pollutant Release Inventory. [https://www.canada.ca/en/](https://www.canada.ca/en/environment-climate-change/services/national-pollutant-release-inventory/about-national-pollutant-release-inventory.html)
581 [environment-climate-change/services/national-pollutant-release-inventory/](https://www.canada.ca/en/environment-climate-change/services/national-pollutant-release-inventory/about-national-pollutant-release-inventory.html)
582 [about-national-pollutant-release-inventory.html](https://www.canada.ca/en/environment-climate-change/services/national-pollutant-release-inventory/about-national-pollutant-release-inventory.html). [Online; accessed 29-December-
583 2021].

584 Environment and Climate Change Canada, 2021b. Wastewater Systems Effluent Regulations.
585 <https://open.canada.ca/data/en/dataset/9e11e114-ef0d-4814-8d93-24af23716489>.
586 [Online; accessed 29-December-2021].

587 European Commission, 2018. Circular Economy: Agreement on Commission proposal to boost the
588 use of organic and waste-based fertilisers. [https://ec.europa.eu/commission/presscorner/](https://ec.europa.eu/commission/presscorner/detail/en/IP_18_6161)
589 [detail/en/IP_18_6161](https://ec.europa.eu/commission/presscorner/detail/en/IP_18_6161). [Online; accessed 25-August-2022].

590 European Commission, 2022. Consolidated text: Regulation (EU) 2019/1009 of the European Parlia-
591 ment and of the Council of 5 June 2019 laying down rules on the making available on the market
592 of EU fertilising products and amending Regulations (EC) No 1069/2009 and (EC) No 1107/2009
593 and repealing Regulation (EC) No 2003/2003 (Text with EEA relevance)Text with EEA rele-
594 vance. <https://eur-lex.europa.eu/eli/reg/2019/1009>. [Online; accessed 25-August-2022].

595 FAO, 2011. Global food losses and food waste - Extent, causes and prevention. [https://www.fao.](https://www.fao.org/3/i2697e/i2697e.pdf)
596 [org/3/i2697e/i2697e.pdf](https://www.fao.org/3/i2697e/i2697e.pdf). [Online; accessed 15-July-2022].

597 FAO & IAEA, 2004. Use of Phosphate Rocks for Sustainable Agriculture. [https://www.fao.org/](https://www.fao.org/3/y5053e/y5053e00.htm)
598 [3/y5053e/y5053e00.htm](https://www.fao.org/3/y5053e/y5053e00.htm). [Online; accessed 23-August-2022].

599 Fernández-Delgado, M., del Amo-Mateos, E., García-Cubero, M.T., Coca, M., Lucas, S., 2022.
600 Phosphorus recovery from organic waste for its agronomic valorization: technical and economic
601 evaluation. *Journal of Chemical Technology & Biotechnology* 97, 167–178.

602 Food and Agriculture Organization of the United Nations, 2022. The importance of Ukraine and
603 the Russian Federation for global agricultural markets and the risks associated with the war in
604 Ukraine.

Galán-Martín, A., Pozo, C., Azapagic, A., Grossmann, I.E., Mac Dowell, N., Guillén-Gosálbez, G., 2018. Time for global action: an optimised cooperative approach towards effective climate change mitigation. *Energy & Environmental Science* 11, 572–581.

Government of Canada, 2022. Greenhouse Worker in Ontario - Job prospects. <https://www.jobbank.gc.ca/marketreport/outlook-occupation/23339/ON>. [Online; accessed 22-December-2021].

Hayse, P.L., Marion, W.W., 1973. Eviscerated yield, component parts, and meat, skin and bone ratios in the chicken broiler. *Poultry Science* 52, 718–722.

Health Canada, 2008. Nutrient Value of Some Common Foods. https://www.canada.ca/content/dam/hc-sc/migration/hc-sc/fn-an/alt_formats/pdf/nutrition/fiche-nutri-data/nvscf-vnqau-eng.pdf. [Online; accessed 14-December-2021].

Health Canada, 202. Canadian Nutrient File. <https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp>. [Online; accessed 15-July-2022].

IISD, 2018. Nutrient Recovery and Reuse in Canada: Foundations for a national framework. <https://www.iisd.org/sites/default/files/meterial/nutrient-recovery-reuse-canada.pdf>. [Online; accessed 25-August-2022].

INAC Services, 2014. Poultry processing in Canada. <http://inacservices.com/poultry-processing-in-canada/>. [Online; accessed 22-August-2022].

Jupp, A.R., Beijer, S., Narain, G.C., Schipper, W., Slootweg, J.C., 2021. Phosphorus recovery and recycling—closing the loop. *Chemical Society Reviews* 50, 87–101.

Kahiluoto, H., Pickett, K.E., Steffen, W., 2021. Global nutrient equity for people and the planet. *Nature Food* 2, 857–861.

Kaiser, D.E., Pagliari, P., 2018. Understanding phosphorus fertilizers. <https://extension.umn.edu/phosphorus-and-potassium/understanding-phosphorus-fertilizers>. [Online; accessed 23-August-2022].

- 630 Klinglmair, M., Lemming, C., Jensen, L.S., Rechberger, H., Astrup, T.F., Scheutz, C., 2015. Phos-
 631 phorus in denmark: national and regional anthropogenic flows. *Resources, Conservation and*
 632 *Recycling* 105, 311–324.
- 633 Martín-Hernández, E., Hu, Y., Zavala, V.M., Martín, M., Ruiz-Mercado, G.J., 2022. Analysis of
 634 incentive policies for phosphorus recovery at livestock facilities in the great lakes area. *Resources,*
 635 *Conservation and Recycling* 177, 105973.
- 636 Martín-Hernández, E., Martín, M., Ruiz-Mercado, G.J., 2021. A geospatial environmental and
 637 techno-economic framework for sustainable phosphorus management at livestock facilities. *Re-*
 638 *sources, Conservation and Recycling* 175, 105843.
- 639 National Aeronautics and Space Administration, 2022. Technology Readiness Level Definitions.
 640 https://www.nasa.gov/pdf/458490main_TRL_Definitions.pdf. [Online; accessed 15-August-
 641 2022].
- 642 Nättorp, A., Remmen, K., Remy, C., 2017. Cost assessment of different routes for phosphorus recov-
 643 ery from wastewater using data from pilot and production plants. *Water Science and Technology*
 644 76, 413–424.
- 645 Nesme, T., Metson, G.S., Bennett, E.M., 2018. Global phosphorus flows through agricultural trade.
 646 *Global Environmental Change* 50, 133–141.
- 647 Ohtake, H., Tsuneda, S., 2019. Phosphorus recovery and recycling. Springer.
- 648 Oldfield, L., Rakhimbekova, S., Roy, J.W., Robinson, C.E., 2020. Estimation of phosphorus loads
 649 from septic systems to tributaries in the canadian lake erie basin. *Journal of Great Lakes Research*
 650 46, 1559–1569.
- 651 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2011. Growing Green-
 652 house Vegetables in Ontario.
- 653 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2020a. Ontario egg

654 production. <https://data.ontario.ca/dataset/ontario-egg-production>. [Online; accessed
655 14-December-2021].

656 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2020b.
657 Ontario milk shipments by county. [https://data.ontario.ca/en/dataset/
658 ontario-milk-shipments-by-county](https://data.ontario.ca/en/dataset/ontario-milk-shipments-by-county). [Online; accessed 14-December-2021].

659 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2021. Greenhouse
660 Nutrient Feedwater Regulation. [http://www.omafra.gov.on.ca/english/nm/regs/gnfpro/
661 gnfnreg.htm](http://www.omafra.gov.on.ca/english/nm/regs/gnfpro/gnfreg.htm). [Online; accessed 23-December-2021].

662 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs, 2022. Horticultural
663 Crops. <http://www.omafra.gov.on.ca/english/stats/hort/index.html>. [Online; accessed
664 18-December-2021].

665 Ontario Ministry of the Environment, Conservation and Parks, 2012. Greenhouse
666 Wastewater Monitoring Project (2010 and 2011). [https://www.ontario.ca/page/
667 greenhouse-wastewater-monitoring-project-2010-and-2011](https://www.ontario.ca/page/greenhouse-wastewater-monitoring-project-2010-and-2011). [Online; accessed 22-
668 December-2021].

669 Ontario Ministry of the Environment, Conservation and Parks, 2019. Provincial policy objectives for
670 managing effects of cage aquaculture operations on the quality of water and sediment in Ontario's
671 waters. <https://ero.ontario.ca/notice/012-7186>. [Online; accessed 15-July-2022].

672 Ontario Ministry of the Environment, Conservation and Parks, 2021. Municipal Treated Wastewater
673 Effluent. <https://data.ontario.ca/dataset/municipal-treated-wastewater-effluent>.
674 [Online; accessed 29-December-2021].

675 Opendatasoft, 2019. Census divisions - Canada. [https://public.opendatasoft.com/
676 explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_
677 en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario). [Online; accessed
678 29-November-2021].

- 679 Pingali, P.L., 2012. Green revolution: impacts, limits, and the path ahead. Proceedings of the
680 National Academy of Sciences 109, 12302–12308.
- 681 Pollution Probe, 2022. Mapping Phosphorus Flows in the Ontario Economy. Exploring Nutrient
682 Recovery and Reuse Opportunities in a Provincial Context. [https://www.pollutionprobe.org/
683 mapping-phosphorus-flows-in-the-ontario-economy/](https://www.pollutionprobe.org/mapping-phosphorus-flows-in-the-ontario-economy/). [Online; accessed 13-July-2022].
- 684 Reid, K., Schneider, K., Joosse, P., 2019. Addressing imbalances in phosphorus accumulation in
685 canadian agricultural soils. Journal of environmental quality 48, 1156–1166.
- 686 Reid, K., Schneider, K.D., 2019. Phosphorus accumulation in canadian agricultural soils over 30
687 yr. Canadian Journal of Soil Science 99, 520–532.
- 688 Sampat, A.M., Hicks, A., Ruiz-Mercado, G.J., Zavala, V.M., 2021. Valuing economic impact
689 reductions of nutrient pollution from livestock waste. Resources, Conservation and Recycling
690 164, 105199.
- 691 Sampat, A.M., Hu, Y., Sharara, M., Aguirre-Villegas, H., Ruiz-Mercado, G., Larson, R.A., Zavala,
692 V.M., 2019. Coordinated management of organic waste and derived products. Computers &
693 chemical engineering 128, 352–363.
- 694 Sampat, A.M., Martin-Hernandez, E., Martín, M., Zavala, V.M., 2018. Technologies and logistics
695 for phosphorus recovery from livestock waste. Clean Technologies and Environmental Policy 20,
696 1563–1579.
- 697 Samreen, S., Kausar, S., 2019. Phosphorus fertilizer: The original and commercial sources, in:
698 Zhang, T. (Ed.), Phosphorus. IntechOpen. chapter 6.
- 699 Sardans, J., Peñuelas, J., 2013. Tree growth changes with climate and forest type are associated
700 with relative allocation of nutrients, especially phosphorus, to leaves and wood. Global Ecology
701 and Biogeography 22, 494–507.
- 702 Schoumans, O., Rulkens, W., Oenema, O., Ehlert, P., 2010. Phosphorus recovery from animal

703 manure: Technical opportunities and agro-economical perspectives. Technical Report. Alterra,
704 Wageningen UR.

705 Senthilkumar, K., Nesme, T., Mollier, A., Pellerin, S., 2012. Conceptual design and quantification
706 of phosphorus flows and balances at the country scale: The case of France. *Global Biogeochemical*
707 *Cycles* 26.

708 Sharma, B.K., Chandel, M.K., 2021. Life cycle cost analysis of municipal solid waste management
709 scenarios for mumbai, india. *Waste Management* 124, 293–302.

710 Statistics Canada – Statistique Canada, 2021a. Cattle and calves on census day. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042401>. [Online; accessed 29-
711 November-2019].
712

713 Statistics Canada – Statistique Canada, 2021b. Cattle statistics, supply and disposition of cat-
714 tle. <https://open.canada.ca/data/en/dataset/e8785f41-2d3c-4dcc-a1ce-e70dac519b66>.
715 [Online; accessed 29-December-2021].

716 Statistics Canada – Statistique Canada, 2021c. Hogs statistics, supply and dis-
717 position of hogs, semi-annual. [https://open.canada.ca/data/en/dataset/](https://open.canada.ca/data/en/dataset/e3c88421-4c7e-4594-9b73-e04a958061e4)
718 [e3c88421-4c7e-4594-9b73-e04a958061e4](https://open.canada.ca/data/en/dataset/e3c88421-4c7e-4594-9b73-e04a958061e4). [Online; accessed 29-December-2021].

719 Statistics Canada – Statistique Canada, 2021d. Other livestock on census day. [https://www150.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042701)
720 [statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042701](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042701). [Online; accessed 29-November-
721 2019].

722 Statistics Canada – Statistique Canada, 2021e. Pigs on census day. [https://www150.statcan.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042601)
723 [gc.ca/t1/tbl1/en/tv.action?pid=3210042601](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042601). [Online; accessed 29-November-2019].

724 Statistics Canada – Statistique Canada, 2021f. Poultry inventory on census day. [https://www150.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042801)
725 [statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042801](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042801). [Online; accessed 29-
726 November-2019].

727 Statistics Canada – Statistique Canada, 2021g. Sheep and lambs on census day. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210042501>. [Online; accessed 29-
728 November-2019].

730 Statistics Canada – Statistique Canada, 2021h. Sheep statistics, supply and
731 disposition of sheep and lambs. <https://open.canada.ca/data/en/dataset/12d4e931-c6b8-4df3-bf5b-0a25524fc6c1>. [Online; accessed 29-December-2021].
732

733 Statistics Canada – Statistique Canada, 2022. Fertilizer shipments to Canadian agriculture markets,
734 by nutrient content and fertilizer year, cumulative data (x 1,000). <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210003901>. [Online; accessed 09-February-2022].
735

736 Statistics Canada - Statistique Canada, 2015. Sewer and septic system connections. House-
737 holds and the Environment. <https://www150.statcan.gc.ca/n1/pub/11-526-x/2013001/t059-eng.htm>. [Online; accessed 15-July-2022].
738

739 Statistics Canada - Statistique Canada, 2017. Estimated number of households and average house-
740 hold size by domain, Canada. User guide for the survey of household spending. <https://www150.statcan.gc.ca/n1/pub/62f0026m/2017002/app-ann-g-eng.htm>. [Online; accessed
741 15-July-2022].
742

743 Statistics Canada - Statistique Canada, 2021. Aquaculture, production and value. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210010701>. [Online; accessed 15-July-2022].
744

745 Statistics Canada - Statistique Canada, 2022a. Interprovincial and international trade flows,
746 basic prices, summary level. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1210008801>. [Online; accessed 13-July-2022].
747

748 Statistics Canada - Statistique Canada, 2022b. Livestock estimates. <https://www150.statcan.gc.ca/n1/daily-quotidien/220228/dq220228d-cansim-eng.htm>. [Online; accessed 15-August-
749 2022].
750

751 Statistics Canada - Statistique Canada, 2022c. Population estimates, quarterly. [https://www150.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901)
752 [statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901). [Online; accessed 15-July-2022].

753 Statistics Canada - Statistique Canada, 2022d. Selected livestock and poultry, Census of Agriculture
754 historical data. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210015501>.
755 [Online; accessed 18-August-2022].

756 Statistics Canada - Statistique Canada, 2022e. Trade Data Online. [https://www.ic.gc.ca//app/](https://www.ic.gc.ca//app/scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng)
757 [scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng](https://www.ic.gc.ca//app/scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng). [Online; accessed 15-July-2022].

758 Statistics Netherlands, 2012. Standardised calculation methods for animal manure and nutrients.
759 Standard data 1990–2008. Technical Report. Statistics Netherlands.

760 Stelco Inc., 2022. Our Facilities. <https://www.stelco.com/about-us/our-facilities>. [Online;
761 accessed 13-July-2022].

762 Szögi, A., Vanotti, M., Hunt, P., 2008. Phosphorus recovery from poultry litter. Transactions of
763 the ASABE 51, 1727–1734.

764 Tan, C., Zhang, T., 2011. Surface runoff and sub-surface drainage phosphorus losses under regular
765 free drainage and controlled drainage with sub-irrigation systems in southern ontario. Canadian
766 Journal of Soil Science 91, 349–359.

767 The World Bank, 2022. Annual Prices - Commodity Markets. [https://www.worldbank.org/en/](https://www.worldbank.org/en/research/commodity-markets)
768 [research/commodity-markets](https://www.worldbank.org/en/research/commodity-markets). [Online; accessed 23-August-2022].

769 Towler, G., Sinnott, R., 2013. Chemical engineering design: principles, practice and economics of
770 plant and process design. Butterworth-Heinemann.

771 United Nations Environment Programme, 2022. Resolution adopted by the United Nations Envi-
772 ronment Assembly on 2 March 2022, UNEP/EA.5 Res.2, on Sustainable nitrogen management.

773 United States Department of Agriculture, 2009. Agricultural Waste Management Field Handbook.
774 Technical Report. United States Department of Agriculture (USDA).

775 U.S. Census Bureau, 2021. Current versus Constant (or Real) Dollars. [https://www.census.](https://www.census.gov/topics/income-poverty/income/guidance/current-vs-constant-dollars.html)
776 [gov/topics/income-poverty/income/guidance/current-vs-constant-dollars.html](https://www.census.gov/topics/income-poverty/income/guidance/current-vs-constant-dollars.html). [On-
777 line; accessed 23-August-2022].

778 U.S. Department of Agriculture, 2011. Animal Feeding Operations (AFO) and Concentrated
779 Animal Feeding Operations (CAFO). [https://www.nrcs.usda.gov/wps/portal/nrcs/main/](https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/plantsanimals/livestock/afo/)
780 [national/plantsanimals/livestock/afo/](https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/plantsanimals/livestock/afo/). [Online; accessed 10-August-2020].

781 Van Dijk, K.C., Lesschen, J.P., Oenema, O., 2016. Phosphorus flows and balances of the European
782 Union Member States. *Science of the Total Environment* 542, 1078–1093.

783 Van Staden, T., Van Meter, K., Saurette, D., Basu, N., Parsons, C., Van Cappellen, P., 2021.
784 Anthropogenic phosphorus mass balance in ontario counties and watersheds. 10.20383/101.0208.

785 Villalba, G., Liu, Y., Schroder, H., Ayres, R.U., 2008. Global phosphorus flows in the industrial
786 economy from a production perspective. *Journal of Industrial ecology* 12, 557–569.

787 Wang, Y., Zhang, T., Tan, C., Qi, Z., Welacky, T., 2018. Solid cattle manure less prone to
788 phosphorus loss in tile drainage water. *Journal of environmental quality* 47, 318–325.

789 World Integrated Trade Solution, 2022. Data on Export, Import, Tariff. [https://wits.worldbank.](https://wits.worldbank.org/)
790 [org/](https://wits.worldbank.org/). [Online; accessed 13-July-2022].

791 Yang, J., De Jong, R., Drury, C., Huffman, E., Kirkwood, V., Yang, X., 2007. Development of a
792 canadian agricultural nitrogen budget (canb v2. 0) model and the evaluation of various policy
793 scenarios. *Canadian Journal of Soil Science* 87, 153–165.

794 Yokoyama, K., Kubo, H., Mori, K., Okada, H., Takeuchi, S., Nagasaka, T., 2007. Separation and
795 recovery of phosphorus from steelmaking slags with the aid of a strong magnetic field. *ISIJ*
796 *international* 47, 1541–1548.

797 Zagklis, D., Konstantinidou, E., Zafiri, C., Kornaros, M., 2020. Assessing the economic viability of
798 an animal byproduct rendering plant: Case study of a slaughterhouse in greece. *Sustainability*
799 12, 5870.

- 800 Zhang, T., Tan, C., Zheng, Z., Drury, C., 2015. Tile drainage phosphorus loss with long-term
801 consistent cropping systems and fertilization. *Journal of environmental quality* 44, 503–511.
- 802 Zhou, J., Jiao, X., Ma, L., de Vries, W., Zhang, F., Shen, J., 2021. Model-based analysis of
803 phosphorus flows in the food chain at county level in china and options for reducing the losses
804 towards green development. *Environmental Pollution* 288, 117768.