

Phosphorus in Ontario's economic sectors: mapping flows and assessing recovery and recycling potential

Université Laval team^{a,*}, McGill University team^b, University of Waterloo team^c

^a *Université Laval*

^b *McGill University*

^c *University of Waterloo*

Abstract

The dual dimension of the anthropogenic use of phosphorus, i.e., its key role in the food production system and the negative environmental impacts associated with the phosphorus used in intensive agricultural techniques, has been stated by the United Nations Environment Assembly. In addition, phosphorus is a non-renewable material which reserves are concentrated in a few number of regions, making global supply chains vulnerable to regional events and conflicts. As a consequence, the recovery and recycling of phosphorus is not just a desirable but also a necessary approach to assure a sustainable, reliable, and sovereign food production system. In this work we map the phosphorus flows through the economic sectors of the Canadian province of Ontario, and phosphorus recovery and recycling opportunities are identified. These mainly belong to the agricultural sector, including manure (30.5 kt/year) and slaughterhouse waste (3.7 kt/year), although significant amounts of P are also found in food and organic waste, including municipal wastewater (6.4 kt/year). Different scenarios are studied to determine the amount of phosphorus that could be recovered within the province considering according with the technology readiness level of different phosphorus recovery processes, as well as the costs associated with phosphorus recovery [Add some more numbers here](#). Finally, we discuss the implications that would be derived from implementing active phosphorus recovery and recycling approaches regarding phosphorus supply and use in Ontario.

*Corresponding author

Email address: `edgar.martin-hernandez.1@ulaval.ca` (Université Laval team)

8 1. Introduction

9 Phosphorus is an essential for production of food which has been intensively used for crop and
10 livestock production since the development of synthetic fertilizers in the XIX and XX centuries
11 ([Samreen and Kausar, 2019](#)). The combination of synthetic fertilizers with other modern inten-
12 sive agricultural techniques have increased the productivity of agriculture and farming industries
13 ([Pingali, 2012](#)). However, the intensive use of fertilizers in agriculture has resulted in the over-
14 application of phosphorus in many regions worldwide REF, while the run of intensive livestock
15 production facilities, also known as concentrated animal feeding operations (CAFOs) ([U.S. Depart-
16 ment of Agriculture, 2011](#)), result in important difficulties in the management of the large amounts
17 of manure produced, which is often spread in lands in the vicinity of CAFOs, which also leads to
18 the accumulation of phosphorus in soil. Soil acts as a phosphorus reservoir ([Ehlert et al., 2003](#)),
19 building-up a legacy P that can be used for future crops, but also can be transported to waterbodies
20 by erosion and runoff leading to the eutrophication of aquatic ecosystems.

21 The dual dimension of the anthropogenic use of phosphorus, i.e., its key role in the food pro-
22 duction system and the negative environmental impacts associated with the phosphorus used in
23 intensive agricultural techniques, has been stated by the United Nations Environment Assembly in
24 the resolution UNEP/EA.5/Res.2 ([United Nations Environment Programme, 2022](#)). An additional
25 factor to be considered for addressing the phosphorus challenge is the non-renewable nature of
26 phosphorus, since the phosphorus consumed is not replenished by natural means at human time
27 scale, and there is currently no known synthetic substitute for this material ([Cordell et al., 2009](#)).
28 Since the global phosphorus reserves are concentrated in a few number of regions, the supply of
29 phosphorus from a limited number of global supply chains lacks resiliency and it has been proven
30 that it can be globally disrupted by regional events and conflicts, resulting geopolitical tensions
31 ([Food and Agriculture Organization of the United Nations, 2022](#)). As a consequence, the recov-
32 ery and recycling of phosphorus is not just a desirable but also a necessary approach to assure a
33 sustainable, reliable, and sovereign food production system.

34 Although the main uses of phosphorus are in the agri-food sector, phosphorus is also involved in
35 other industrial activities, including steel, chemical, and forestry industries. Henceforth, phosphorus

is a key material for many aspects of human development. As a result, mapping the phosphorus flows involved in human activities to detect opportunities for recovery and recycling is essential for, in a second stage, assess amount of phosphorus that is viable to recover, the economical costs involved, and the enhancement in terms of resiliency of the regional food production system, savings from the reduction of phosphorus imports, and the mitigation of phosphorus pollution on the region implementing strategies for phosphorus recovery and recycling. The quantification of phosphorus flows has been addressed in previous works in the literature for certain sectors such as the agri-food sector (Boh and Clark, 2020; Zhou et al., 2021; Nesme et al., 2018). Additionally, studies on the global phosphorus flows have also been performed (Villalba et al., 2008; Chen and Graedel, 2016), although these studies tend to have a low flow resolution since these are aggregated by major sectors. Additionally, the works quantifying phosphorus often include qualitative recommendations to improve the phosphorus use efficiency and recycling (Van Dijk et al., 2016; Senthilkumar et al., 2012), but often they do not include quantitative assessments on the amount of phosphorus which recovery is feasible along with the costs involved. Conversely, those works focused on estimating the recoverable phosphorus and the associated recovery cost target specific flows, lacking a holistic perspective of the phosphorus flows in the various human activities (Martín-Hernández et al., 2021; Sampat et al., 2018).

In this work, we intend to perform a holistic approach to the opportunities for phosphorus recovery and recycling in the Canadian province of Ontario. In a first stage, we proceed to map the phosphorus flows involved in the economical sectors of Ontario, i.e., the agri-food, industrial, and urban sectors. This data is used in a second stage to identify the flows in which phosphorus recovery is feasible, determining the amount of phosphorus that could be recovered within the province considering different scenarios regarding the technology readiness level of different phosphorus recovery processes, as well as the costs associated with phosphorus recovery. Finally, we discuss the implications that would be derived from implementing active phosphorus recovery and recycling approaches regarding phosphorus supply and use in Ontario.

2. Methods

2.1. Spatial resolution

Phosphorus flows have been mapped within the Canadian province of Ontario, and thus the political borders of Ontario has been considered as the boundaries for the substance flow analysis performed. In those cases where the data was available, the distribution of phosphorus flows within Ontario has also been studied at Census Division level ([Statistics Canada – Statistique Canada, 2017](#)). The database collecting the IDs of Ontario Census Divisions, their names, and geospatial information is taken from [Opendatasoft \(2019\)](#).

ADD MAP WITH CENSUS DIVISIONS???

2.2. Temporal resolution

The study has being performed for year 2019 since the most of data required is available for this year. In addition, the temporal evolution of the largest phosphorus flows, i.e., agricultural and wastewater phosphorus flows, has been studied for a period of 13 years from 2007 to 2019.

2.3. Estimation of phosphorus flows

The estimation of phosphorus flows within the Ontario's agricultural sectors is based on the methodology used in [Pollution Probe \(2022\)](#). It is based on the use open data sources, often from governmental institutions, complemented with information from scientific articles when needed. The particular procedure followed for each flow depends on the information publicly available. In the next sections we depict the main lines of the estimating procedure for each sector, while we refer the reader to [Pollution Probe \(2022\)](#) for a more comprehensive description of the procedure followed for estimating each phosphorus flow.

2.3.1. Agricultural sector

Phosphorus flows in the agricultural sector are estimated based on production data of livestock and crop products, as well as data on fertilizer application.

For those production data were not available, a number of different methods were used to estimate the P flow based on approaches established in the literature. For example, P inflows associated

with synthetic fertilizers could be directly estimated based on application data reported in the Fertilizer Shipments Survey (FSS).³⁷ Conversely, P flows associated with manure were determined indirectly by accounting for the magnitude from which the flow of P could be derived. In this case,

Phosphorus in livestock imports and exports is estimated from livestock trading data REF, multiplying the number of animals by the concentration of phosphorus in the different types of livestock REF,

Phosphorus in livestock feeding and manure is estimated based on the number and type of animals reported for Ontario at Census Division level in the Census of Agriculture REF!, multiplied by the phosphorus feeding requirements REF, and concentration of phosphorus in manure REF. The Census of Agriculture is published by Statistics Canada every five years (i.e., 2001, 2006, 2011, and 2016) for cattle⁵² REF, sheep⁵³ REF, swine⁵⁴ REF, poultry⁵⁵ REF, and other livestock⁵⁶ REF, with the exception of rabbits, where data is not available prior to 2009. The number of animals for the years in between census reporting have been estimated using a linear interpolation. We assumed that the number of animals reported is throughout the year (i.e., the animals culled are replaced by new ones). However, in the case of broilers and turkeys, the number of animals reported by the livestock census have been reduced by a factor of 0.68 (broilers) and 0.80 (turkeys), since these animals have life cycles of 43 and 80 days respectively, meaning barns are empty for 20 days between cycles. ³⁰⁵ REF

Phosphorus contained in meat and slaughterhouse waste is based on the number of animals slaughtered reported by both federally and provincially licensed meat plants.^{59, 60} REF multiplied by the concentration of phosphorus in carcasses REF.

Phosphorus flows associated with the production of milk and eggs is based on provincial production data, multiplying these products by their average phosphorus concentration ^{57, 58} REF.

Phosphorus in fertilizer applied to open fields in Ontario is estimated based on the amount of fertilizer products traded to Ontario's agricultural markets containing P ¹⁰⁰ REF. The distribution of phosphorus fertilizers among the Census Division of the province is based on the fraction of fertilized area of each census division, i.e., dividing the reported area of land fertilized for each census division by the total fertilized area of land in Ontario, removing the areas that correspond

116 with greenhouse crops^{101, 102 103} REF. Regarding manure, we assume that all of the manure
117 generated by livestock is applied in crop fields ⁵⁰ REF.

118 The uptake of phosphorus by crops is determined based on the area used in each Census Division
119 to grow each type of crops by census division^{104, 105, 106} and its yield^{107, 108} multiplied by the
120 specific P content for each crop type.^{109, 110} The phosphorus uptake by crops is divided according
121 to whether it uptake in the grain, fruit or vegetable, or straw and stover components of each type
122 of crop. This is necessary to determine the amount of phosphorus that flows within food or feed
123 (i.e., grains, fruits and vegetables) while straw and stover remain in the field after harvesting as crop
124 residues.

125 A fraction of the phosphorus applied to crop fields as manure of synthetic fertilizer is lost
126 through erosion, runoff, and drainage. This transportation of phosphorus depends on a range
127 of factor, including the amount of phosphorus applied; soil composition, texture, and slope; and
128 precipitation, resulting in a complex and data-intensive process for estimating the phosphorus
129 transported out of the crop fields. As an approximation, we have estimated the phosphorus losses
130 by using export coefficients determined for crop fields in Ontario ¹¹² REF ¹¹³ REF corrected to
131 account for both surface and subsurface runoffs for synthetic fertilizers (1.267 kg/ha/year), and
132 liquid and solid manure (2.548 kg/ha/year and 1.717 kg/ha/year respectively) ¹¹³ REF ([Pollution
133 Probe, 2022](#)).

134 A fraction of the P supplied to crop fields is not taken up by the plants and remains in soil,
135 resulting in the accumulation of P over time as a result of synthetic fertilizer and manure over
136 over sustained periods of time, often applying phosphorus in greater quantities than crops require
137 to ensure satisfactory yields ¹³² REF. This buildup is often referred to as “legacy P”, and it is
138 estimated as the balance between phosphorus inflows to crop fields (application of manure and
139 synthetic fertilizers) and outflows (crop food and feed products, crop residues, and phosphorus
140 losses).

141 Regarding greenhouse crops, the data available was limited, resulting in an estimation of phos-
142 phorus applied as synthetic fertilizers based on the sum of phosphorus uptake by greenhouse crops
143 (i.e., tomatoes, peppers, and cucumbers)¹¹⁹ and the phosphorus releases from greenhouse irrigation

systems (greenhouse nutrient feedwater systems (GNF) REF ONTARIO) systems. The phosphorus uptake by greenhouse crops is determined by multiplying the production of greenhouse crops REF by the phosphorus content of each vegetable type 121, 122 REF. The phosphorus releases from the GNF systems was estimated based on the average concentration of phosphorus in GNF outlet streams of Ontario (33.6 mg/L) 123 REF and the total water discharges from GNF systems 124 REF, assuming that water discharges from GNF systems is equivalent to 25% of the total water applied in greenhouses, which corresponding with the worst-case scenario of no water recirculation in the GNF system. The average water consumption in greenhouses in Ontario was assumed to be 1,000 L/m²/year 125 REF. We have also estimated the phosphorus releases from the seasonal workers live in households in the vicinity of the greenhouses that may use septic systems, considering that the seasonal labour force in Ontario greenhouses is estimated to be 6,699 workers 126 REF, and an average phosphorus load rate of 0.0156 kg P/person/week from septic systems 128 REF.

REVISAR POR SIDNEY Food imports and exports (other than livestock) are estimated scaling each type of food traded in Canada (Statistics Canada - Statistique Canada, 2022c) with the population of Ontario (Statistics Canada - Statistique Canada, 2022b). The phosphorus contained in each type of imported and exported food is estimated multiplying the amount of each type of traded food by its phosphorus content (Health Canada, 202).

2.3.2. Industrial sector

Phosphorus flows through imports, production, exports and waste for the steel, forestry, and food and beverage, industries of Ontario were mapped. The steel industry is the first non-food sector in terms of phosphorus use. The main phosphorus inflows of steel manufacturing are associated with the use of iron ore, coal, and coke, while the main outflow of phosphorus is within slag, which remove most of the impurities from steel, including phosphorus. It must be noted that, although some minor amounts of phosphorus can be desired in steel for making anti-corrosion surface coatings, it is largely considered an impurity in the steel manufacturing process. Phosphorus in these flows is estimated multiplying their average phosphorus content (0.06% P in iron ore, 0.05%

171 P for coal, 0.4% P in slag, and 0.01% in steel) 176 REF by the steel production capacity of the
172 facilities located in Ontario ([Cheminfo Services Inc., 2019](#); [Algoma Steel Inc., 2022](#); [Stelco Inc.,](#)
173 [2022](#); [Pollution Probe, 2022](#)) and the imports and exports of these materials ([World Integrated](#)
174 [Trade Solution, 2022](#); [Statistics Canada - Statistique Canada, 2022a](#)).

175 Phosphorus flows in Ontario's forestry industry includes wood harvesting, wood products man-
176 ufacturing, as well as the production of pulp and paper. The estimation of these phosphorus flows
177 are the result of multiplying the production data of wood, wood products, pulp and paper, and
178 their retrospectives imports, exports, and waste streams ([Canadian Forest Service, 2020](#); [Statistics](#)
179 [Canada - Statistique Canada, 2022a](#)), by the average phosphorus content, which is assumed to be
180 0.01% for wood 181 REF and 0.005% for pulp and paper products REF.

181 Phosphorus in aquaculture are mainly due to supply of feed as part of fish feed the grow of
182 trouts, part of which is uptake by fishes, while the rest of phosphorus is released into aquatic
183 ecosystems since aquaculture effluents are directly discharged to the environment ([Ontario Ministry](#)
184 [of the Environment, Conservation and Parks, 2019](#)). The amount phosphorus uptakes by fishes is
185 calculated multiplying the fish production ([Statistics Canada - Statistique Canada, 2021](#)), by their
186 phosphorus content ([Health Canada, 202](#)), while the phosphorus content in the aquaculture waste
187 effluents of Ontario is estimated to be 10 kg of phosphorus per ton of fish produced ([Bureau et al.,](#)
188 [2003](#)). The sum of phosphorus uptakes by fishes and phosphorus in aquaculture waste effluents
189 result in the phosphorus supplied to aquaculture as fish feed.

190 Regarding other industrial activities which could involve the use of phosphorus, the local pro-
191 duction of phosphorus is assumed to be negligible since phosphorus is not mined or refined in
192 Ontario, and the synthetic phosphorus fertilizer imports are accounted in the agricultural section.
193 The general chemical facilities located in Ontario report 350 t/year of phosphorus as waste REF,
194 in addition of imports and exports of chemical products REF. However, there exist a significant
195 fraction of phosphorus used in the industrial sector that cannot be tracked due to the lack of data.

196 Ask sidney what to do with food industry, and pet feed. My approach is to merge all of them
197 as it is currently in the figure, but confirm with her


198 2.3.3. *Urban sector*

199 In this section we include the phosphorus inflows and outflows through wastewater treatment
200 plants (WWTPs), septic systems, and food and organic waste management facilities (landfills,
201 composting sites, and anaerobic digestion facilities).

202 Jorge do you mind if the purposes of the papaer we stick with just one method? I think it is
203 better, otherwise it becomes lengthy and confusing

204 Phosphorus flows through WWTPs is estimated combining data from the National Pollutant
205 Release Inventory (NPRI) REF, a public database of releases, disposals and transfers of pollutants,
206 including industrial facilities, and data from the Wastewater Systems Effluent Regulations (WSER)
207 database REF. Since the NPRI only contains data of those facilities that meet certain regulatory
208 requirements, the information of this database must be complemented with the data from the
209 WSER database, which includes information of Canadian WWTPs at the federal, provincial, and
210 municipal level. The estimations on phosphorus flows through WWTPs are valitaed using the
211 Municipal Treated Wastewater Effluent (MTWE) database REF, which collects annual data on
212 water quality data and effluent levels for WWTPs in Ontario. We note that this data set only
213 provides information about phosphorus releases from municipal WWTPs, but it does not collect
214 phosphorus disposals and transfers. REVISAR POR JORGE. This methodology is shown in Figure
215 1

216 Ask Jorge if I can make a new Figure



*Note: quantity refers to disposals, releases, and transfers

Figure 1: Methodology used for estimating phosphorus flows through wastewater treatment plants.

However, there exist households that are not connected to any sewer systems. These households are equipped with septic systems to perform a rough treatment of the wastewater produced prior to its release into the environment, which typically consist into a septic tank that separates solid matter from the wastewater, and a drainfield where the effluent is discharged. The estimation of phosphorus releases from septic systems is based on the fraction of households equipped with these systems, estimated on 13% (Statistics Canada - Statistique Canada, 2015), which are inhabited by an average of 2.58 individuals (Statistics Canada - Statistique Canada, 2017), and the average phosphorus load rate from septic systems, which is estimated on 0.81 kg of phosphorus per person per year for the Lake Erie Basin in Ontario by Oldfield et al. (2020).

Phosphorus flows in the form of food and organic waste are based on applying food loss factors for the steps associated with food processing (FAO, 2011), considering the food production and import values estimated in Section 2.3.1.

2.4. Phosphorus recovery techniques (*This section could be Supplementary Material*)

Brief overview of potential P recovery techniques for each sector

231 **3. Results and discussion**

232 *3.1. Phosphorus flows in Ontario*

233 *Showing an overview of the P flows in the province. The use of figures summarizing all the flows*
234 *of the province in the shape of Sankey or network flow figures could be so great*




Figure 2: Phosphorus flows in the province of Ontario (kt/year). The streams within red rectangles denote phosphorus inflows into the province, while those streams within blue rectangles denote phosphorus outflows out of the province.

235 Figure 2 summarizes the phosphorus flows in the province of Ontario. It can be observed that
236 the flow of phosphorus through the anthropogenic activities are divided into 3 independent
237 networks, i.e., the flow of phosphorus involved the production and processing of food (including the
238 treatment of wastewater), the flow of phosphorus used in the steel and chemical industries, and the
239 phosphorus involved in the forestry industry.

240 The production of animal food products exhibits a lower phosphorus use efficiency than the
241 production of plant base products, similarly to the use efficiency of other resources such as water
242 CITE HERE, CALCULAR ENTRA VS SALE!

243 3.1.1. *Agricultural sector*

244 3.1.2. *Industrial sector*

245 3.1.3. *Urban sector*

246 3.2. *Potential of phosphorus recovery in Ontario*

247 Assessment of different scenarios of P recovery in Ontario, P imports that would be saved,
248 reduction of P dependency of the province, etc (all implications related with mass-balances)

249 3.3. *Economic implications of phosphorus recovery in Ontario*

250 Economic costs or saving derived from the recovery of P in the province and all implications
251 related with economy

252 3.4. *Implications on food sovereignty of phosphorus recovery in Ontario*

253 Implications on food production self-sufficiency derived from the (partial) recycling of P. Discus-
254 sion on the improvement of the food production system resiliency against disruptions of the global
255 supply supply chains (e.g., current context derived from the COVID-19 pandemia and the war in
256 Ukraine)

257 3.5. *Gaps of knowledge*

258 4. Conclusions

259 5. Acknowledgments

260 Pollution Probe

261 ECCC

262 References

263 Algoma Steel Inc., 2022. Corporate Profile. <https://algoma.com/about-algoma/corporate-profile/>. [Online; accessed 13-July-2022].

265 Boh, M.Y., Clark, O.G., 2020. Nitrogen and phosphorus flows in Ontario's food systems. Resources, Conservation and Recycling 154, 104639.

267 Bureau, D.P., Gunther, S.J., Cho, C.Y., 2003. Chemical composition and preliminary theoretical estimates of waste outputs of rainbow trout reared in commercial cage culture operations in Ontario. North American Journal of Aquaculture 65, 33–38.

270 Canadian Forest Service, 2020. Statistical data. <https://cfs.nrcan.gc.ca/statsprofile/>. [Online; accessed 13-July-2022].

272 Cheminfo Services Inc., 2019. Economic Assessment of the Integrated Steel Industry. <https://www.canadiansteel.ca/files/resources/Final-Report-Economic-Assessment-of-the-Integrated-Steel-Industry.pdf>. [Online; accessed 13-July-2022].

276 Chen, M., Graedel, T., 2016. A half-century of global phosphorus flows, stocks, production, consumption, recycling, and environmental impacts. Global Environmental Change 36, 139–152.

278 Cordell, D., Drangert, J.O., White, S., 2009. The story of phosphorus: global food security and food for thought. Global environmental change 19, 292–305.

280 Ehlert, P., Morel, C., Fotyma, M., Destain, J.P., 2003. Potential role of phosphate buffering capacity of soils in fertilizer management strategies fitted to environmental goals. Journal of plant nutrition and soil science 166, 409–415.

283 FAO, 2011. Global food losses and food waste - Extent, causes and prevention. [https://www.fao.](https://www.fao.org/3/i2697e/i2697e.pdf)
284 [org/3/i2697e/i2697e.pdf](https://www.fao.org/3/i2697e/i2697e.pdf). [Online; accessed 15-July-2022].

285 Food and Agriculture Organization of the United Nations, 2022. The importance of Ukraine and
286 the Russian Federation for global agricultural markets and the risks associated with the war in
287 Ukraine.

288 Health Canada, 202. Canadian Nutrient File. [https://food-nutrition.canada.ca/cnf-fce/](https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp)
289 [index-eng.jsp](https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp). [Online; accessed 15-July-2022].

290 Martín-Hernández, E., Martín, M., Ruiz-Mercado, G.J., 2021. A geospatial environmental and
291 techno-economic framework for sustainable phosphorus management at livestock facilities. Re-
292 sources, Conservation and Recycling 175, 105843.

293 Nesme, T., Metson, G.S., Bennett, E.M., 2018. Global phosphorus flows through agricultural trade.
294 Global Environmental Change 50, 133–141.

295 Oldfield, L., Rakhimbekova, S., Roy, J.W., Robinson, C.E., 2020. Estimation of phosphorus loads
296 from septic systems to tributaries in the canadian lake erie basin. Journal of Great Lakes Research
297 46, 1559–1569.

298 Ontario Ministry of the Environment, Conservation and Parks, 2019. Provincial policy objectives for
299 managing effects of cage aquaculture operations on the quality of water and sediment in Ontario’s
300 waters. <https://ero.ontario.ca/notice/012-7186>. [Online; accessed 15-July-2022].

301 Opendatasoft, 2019. Census divisions - Canada. [https://public.opendatasoft.com/](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario)
302 [explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario)
303 [en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario](https://public.opendatasoft.com/explore/dataset/georef-canada-census-division/table/?disjunctive.prov_name_en&disjunctive.cd_name_en&sort=year&refine.prov_name_en=Ontario). [Online; accessed
304 29-November-2021].

305 Pingali, P.L., 2012. Green revolution: impacts, limits, and the path ahead. Proceedings of the
306 National Academy of Sciences 109, 12302–12308.

307 Pollution Probe, 2022. Mapping Phosphorus Flows in the Ontario Economy. Exploring Nutrient
 308 Recovery and Reuse Opportunities in a Provincial Context. [https://www.pollutionprobe.org/
 309 mapping-phosphorus-flows-in-the-ontario-economy/](https://www.pollutionprobe.org/mapping-phosphorus-flows-in-the-ontario-economy/). [Online; accessed 13-July-2022].

310 Sampat, A.M., Martin-Hernandez, E., Martín, M., Zavala, V.M., 2018. Technologies and logistics
 311 for phosphorus recovery from livestock waste. *Clean Technologies and Environmental Policy* 20,
 312 1563–1579.

313 Samreen, S., Kausar, S., 2019. Phosphorus fertilizer: The original and commercial sources, in:
 314 Zhang, T. (Ed.), *Phosphorus*. IntechOpen. chapter 6.

315 Senthilkumar, K., Nesme, T., Mollier, A., Pellerin, S., 2012. Conceptual design and quantification
 316 of phosphorus flows and balances at the country scale: The case of France. *Global Biogeochemical
 317 Cycles* 26.

318 Statistics Canada – Statistique Canada, 2017. Census division (CD). [https://www150.
 319 statcan.gc.ca/n1/pub/92-195-x/2016001/geo/cd-dr/cd-dr-eng.htm](https://www150.statcan.gc.ca/n1/pub/92-195-x/2016001/geo/cd-dr/cd-dr-eng.htm). [Online; accessed 23-
 320 November-2021].

321 Statistics Canada - Statistique Canada, 2015. Sewer and septic system connections. House-
 322 holds and the Environment. [https://www150.statcan.gc.ca/n1/pub/11-526-x/2013001/
 323 t059-eng.htm](https://www150.statcan.gc.ca/n1/pub/11-526-x/2013001/t059-eng.htm). [Online; accessed 15-July-2022].

324 Statistics Canada - Statistique Canada, 2017. Estimated number of households and average house-
 325 hold size by domain, Canada. User guide for the survey of household spending. [https:
 326 //www150.statcan.gc.ca/n1/pub/62f0026m/2017002/app-ann-g-eng.htm](https://www150.statcan.gc.ca/n1/pub/62f0026m/2017002/app-ann-g-eng.htm). [Online; accessed
 327 15-July-2022].

328 Statistics Canada - Statistique Canada, 2021. Aquaculture, production and value. [https://www150.
 329 statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210010701](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=3210010701). [Online; accessed 15-July-2022].

330 Statistics Canada - Statistique Canada, 2022a. Interprovincial and international trade flows,

331 basic prices, summary level. [https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1210008801)
332 [1210008801](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1210008801). [Online; accessed 13-July-2022].

333 Statistics Canada - Statistique Canada, 2022b. Population estimates, quarterly. [https://www150.](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901)
334 [statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901](https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1710000901). [Online; accessed 15-July-2022].

335 Statistics Canada - Statistique Canada, 2022c. Trade Data Online. [https://www.ic.gc.ca//app/](https://www.ic.gc.ca//app/scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng)
336 [scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng](https://www.ic.gc.ca//app/scr/tdst/tdo/crtr.html?&productType=HS6&lang=eng). [Online; accessed 15-July-2022].

337 Stelco Inc., 2022. Our Facilities. <https://www.stelco.com/about-us/our-facilities>. [Online;
338 accessed 13-July-2022].

339 United Nations Environment Programme, 2022. Resolution adopted by the United Nations Envi-
340 ronment Assembly on 2 March 2022, UNEP/EA.5 Res.2, on Sustainable nitrogen management.

341 U.S. Department of Agriculture, 2011. Animal Feeding Operations (AFO) and Concentrated
342 Animal Feeding Operations (CAFO). [https://www.nrcs.usda.gov/wps/portal/nrcs/main/](https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/plantsanimals/livestock/afo/)
343 [national/plantsanimals/livestock/afo/](https://www.nrcs.usda.gov/wps/portal/nrcs/main/national/plantsanimals/livestock/afo/). [Online; accessed 10-August-2020].

344 Van Dijk, K.C., Lesschen, J.P., Oenema, O., 2016. Phosphorus flows and balances of the European
345 Union Member States. *Science of the Total Environment* 542, 1078–1093.

346 Villalba, G., Liu, Y., Schroder, H., Ayres, R.U., 2008. Global phosphorus flows in the industrial
347 economy from a production perspective. *Journal of Industrial ecology* 12, 557–569.

348 World Integrated Trade Solution, 2022. Data on Export, Import, Tariff. [https://wits.worldbank.](https://wits.worldbank.org/)
349 [org/](https://wits.worldbank.org/). [Online; accessed 13-July-2022].

350 Zhou, J., Jiao, X., Ma, L., de Vries, W., Zhang, F., Shen, J., 2021. Model-based analysis of
351 phosphorus flows in the food chain at county level in china and options for reducing the losses
352 towards green development. *Environmental Pollution* 288, 117768.