Class 18

Emma Bell (A19247017)

Table of contents

Background 1
CMI-PB project 8
Focus on PT (Pertussis Toxin) Ag 18

Background

Pertussis (aka whooping cough) is a common lung infection caused by the bacteria B. pertussis

This can infect all ages but is most severe for those under 1 years of age

The CDC track the number of reported cases in the US. We can "scrape" this with **datapasta** package.

```
cdc <- data.frame(
  year = c(
    1922L,
    1923L,
    1924L,
    1925L,
    1926L,
    1927L,
    1928L,
    1929L,
    1930L,
    1931L,
    1932L,
    1933L,
    1934L,
    1935L,
```

```
    1936L,
    1937L,
    1938L,
    1939L,
    1940L,
    1941L,
    1942L,
    1943L,
    1944L,
    1945L,
    1946L,
    1947L,
    1948L,
    1949L,
    1950L,
    1951L,
    1952L,
    1953L,
    1954L,
    1955L,
    1956L,
    1957L,
    1958L,
    1959L,
    1960L,
    1961L,
    1962L,
    1963L,
    1964L,
    1965L,
    1966L,
    1967L,
    1968L,
    1969L,
    1970L,
    1971L,
    1972L,
    1973L,
    1974L,
    1975L,
    1976L,
    1977L,
```

```
    1978L,
    1979L,
    1980L,
    1981L,
    1982L,
    1983L,
    1984L,
    1985L,
    1986L,
    1987L,
    1988L,
    1989L,
    1990L,
    1991L,
    1992L,
    1993L,
    1994L,
    1995L,
    1996L,
    1997L,
    1998L,
    1999L,
    2000L,
    2001L,
    2002L,
    2003L,
    2004L,
    2005L,
    2006L,
    2007L,
    2008L,
    2009L,
    2010L,
    2011L,
    2012L,
    2013L,
    2014L,
    2015L,
    2016L,
    2017L,
    2018L,
    2019L,
```

```r
    2020L,
    2021L,
    2022L,
    2023L,
    2024L,
    2025L
  ),
  cases = c(
    107473,
    164191,
    165418,
    152003,
    202210,
    181411,
    161799,
    197371,
    166914,
    172559,
    215343,
    179135,
    265269,
    180518,
    147237,
    214652,
    227319,
    103188,
    183866,
    222202,
    191383,
    191890,
    109873,
    133792,
    109860,
    156517,
    74715,
    69479,
    120718,
    68687,
    45030,
    37129,
    60886,
    62786,
```

```
    31732,
    28295,
    32148,
    40005,
    14809,
    11468,
    17749,
    17135,
    13005,
    6799,
    7717,
    9718,
    4810,
    3285,
    4249,
    3036,
    3287,
    1759,
    2402,
    1738,
    1010,
    2177,
    2063,
    1623,
    1730,
    1248,
    1895,
    2463,
    2276,
    3589,
    4195,
    2823,
    3450,
    4157,
    4570,
    2719,
    4083,
    6586,
    4617,
    5137,
    7796,
    6564,
```

```
    7405,
    7298,
    7867,
    7580,
    9771,
    11647,
    25827,
    25616,
    15632,
    10454,
    13278,
    16858,
    27550,
    18719,
    48277,
    28639,
    32971,
    20762,
    17972,
    18975,
    15609,
    18617,
    6124,
    2116,
    3044,
    7063,
    22538,
    21996
  )
)
```

Q. Make a plot of `year` vs `cases`

```
library(ggplot2)
ggplot(cdc)+
  aes(year, cases) +
  geom_line()
```



Q Add some major milestones including the first wP vaccine roll-out (1946), the switch to the newer aP vaccine (1996), the COVID years (2020).

```
library(ggplot2)
ggplot(cdc)+
  aes(year, cases) +
  geom_line() +
  geom_vline(xintercept=1946, col="maroon", lty=2) +
  geom_vline(xintercept=1996, col="lightpink", lty=2) +
  geom_vline(xintercept=2020, col="magenta", lty=2)
```



> Q. What do you notice after 1946?

There were high case numbers in the pre-1940s. After the roll out of this vaccine, the case numbers decrease dramatically. In the 1970s-90s, cases are very low.

> Q. Describe what happened after the introduction of the aP vaccine? Do you have a possible explanation for the observed trend?

After the introduction of this new vaccine, which is considered safer and with less side effects, cases starts climbing again. The problem could be that no one was getting their boosters. the lag is very consistant that suggests there is a problem with efficiency of this vaccine. The aP protection wanes faster that wP- we do not know why (hence the booster vaccines)

Interestingly, during the COVID years, cases decrease because social distancing measures and lockdown stopped people from catching it.

Why is this vaccine preventable disease on the upswing? To answer this question we need to investigate the mechanisms underlying waning protection against pertussis. This requires evaluation of pertussis-specific immune responses over time in wP and aP vaccinated individuals.

CMI-PB project

Computational Models of Immunity- Pertussis Boost project aimst o provide the scientific community with this very information.

They make their data available via JSON format APIs. We can read this in R with the `read_json()` function from the **jsonlite** package:

```
library(jsonlite)

subject <- read_json("https://www.cmi-pb.org/api/v5_1/subject"
                     , simplifyVector=TRUE)

head(subject)
```

```
  subject_id infancy_vac biological_sex              ethnicity  race
1          1          wP         Female Not Hispanic or Latino White
2          2          wP         Female Not Hispanic or Latino White
3          3          wP         Female                Unknown White
4          4          wP           Male Not Hispanic or Latino Asian
5          5          wP           Male Not Hispanic or Latino Asian
6          6          wP         Female Not Hispanic or Latino White
  year_of_birth date_of_boost      dataset
1    1986-01-01    2016-09-12 2020_dataset
2    1968-01-01    2019-01-28 2020_dataset
3    1983-01-01    2016-10-10 2020_dataset
4    1988-01-01    2016-08-29 2020_dataset
5    1991-01-01    2016-08-29 2020_dataset
6    1988-01-01    2016-10-10 2020_dataset
```

Q4. How many wP and aP individuals are in this `subject` table?

```
table(subject$infancy_vac)
```

```
aP wP
87 85
```

Q5. How many Male and Female subjects/patients are in the dataset?

```
table(subject$biological_sex)
```

```
Female   Male
   112     60
```

Q6. What is the breakdown of race and biological sex (e.g. number of Asian females, White males etc…)? Is this dataset representative of the US population?

```
table(subject$race, subject$biological_sex)
```

```
                                          Female Male
  American Indian/Alaska Native                0    1
  Asian                                       32   12
  Black or African American                    2    3
  More Than One Race                          15    4
  Native Hawaiian or Other Pacific Islander    1    1
  Unknown or Not Reported                     14    7
  White                                       48   32
```

No it is not representative, as there are more women than mena and also, there is not enough representation of some races.

Let's read more databases

```
specimen <- read_json("https://www.cmi-pb.org/api/v5_1/specimen",
                      simplifyVector = TRUE )
ab_titer <- read_json("https://www.cmi-pb.org/api/v5_1/plasma_ab_titer",
                       simplifyVector = TRUE)
```

```
head(specimen)
```

```
  specimen_id subject_id actual_day_relative_to_boost
1           1          1                           -3
2           2          1                            1
3           3          1                            3
4           4          1                            7
5           5          1                           11
6           6          1                           32
  planned_day_relative_to_boost specimen_type visit
1                             0         Blood     1
2                             1         Blood     2
3                             3         Blood     3
4                             7         Blood     4
5                            14         Blood     5
6                            30         Blood     6
```

```r
head(ab_titer)
```

```
  specimen_id isotype is_antigen_specific antigen        MFI MFI_normalised
1           1     IgE               FALSE   Total 1110.21154       2.493425
2           1     IgE               FALSE   Total 2708.91616       2.493425
3           1     IgG                TRUE      PT   68.56614       3.736992
4           1     IgG                TRUE     PRN  332.12718       2.602350
5           1     IgG                TRUE     FHA 1887.12263      34.050956
6           1     IgE                TRUE     ACT    0.10000       1.000000
   unit lower_limit_of_detection
1 UG/ML                 2.096133
2 IU/ML                29.170000
3 IU/ML                 0.530000
4 IU/ML                 6.205949
5 IU/ML                 4.679535
6 IU/ML                 2.816431
```

To analyse this data we need to first "join" (merge/link) the different tables so we have all the data in one place not spread across different tables.

We can use the `*_join()` family of functions from **dplyr** to do this.

```r
library(dplyr)
```

```
Attaching package: 'dplyr'

The following objects are masked from 'package:stats':

    filter, lag

The following objects are masked from 'package:base':

    intersect, setdiff, setequal, union
```

```r
meta <- inner_join(subject, specimen)
```

```
Joining with `by = join_by(subject_id)`
```

```r
head(meta)
```

```
  subject_id infancy_vac biological_sex              ethnicity  race
1          1          wP         Female Not Hispanic or Latino White
2          1          wP         Female Not Hispanic or Latino White
3          1          wP         Female Not Hispanic or Latino White
4          1          wP         Female Not Hispanic or Latino White
5          1          wP         Female Not Hispanic or Latino White
6          1          wP         Female Not Hispanic or Latino White
  year_of_birth date_of_boost      dataset specimen_id
1    1986-01-01    2016-09-12 2020_dataset           1
2    1986-01-01    2016-09-12 2020_dataset           2
3    1986-01-01    2016-09-12 2020_dataset           3
4    1986-01-01    2016-09-12 2020_dataset           4
5    1986-01-01    2016-09-12 2020_dataset           5
6    1986-01-01    2016-09-12 2020_dataset           6
  actual_day_relative_to_boost planned_day_relative_to_boost specimen_type
1                           -3                             0         Blood
2                            1                             1         Blood
3                            3                             3         Blood
4                            7                             7         Blood
5                           11                            14         Blood
6                           32                            30         Blood
  visit
1     1
2     2
3     3
4     4
5     5
6     6
```

```r
abdata <- inner_join(ab_titer, meta)
```

```
Joining with `by = join_by(specimen_id)`
```

```r
head(abdata)
```

```
  specimen_id isotype is_antigen_specific antigen        MFI MFI_normalised
1           1     IgE               FALSE   Total 1110.21154       2.493425
2           1     IgE               FALSE   Total 2708.91616       2.493425
```

```
3           1     IgG                TRUE      PT   68.56614       3.736992
4           1     IgG                TRUE     PRN  332.12718       2.602350
5           1     IgG                TRUE     FHA 1887.12263      34.050956
6           1     IgE                TRUE     ACT    0.10000       1.000000
   unit lower_limit_of_detection subject_id infancy_vac biological_sex
1 UG/ML                 2.096133          1          wP         Female
2 IU/ML                29.170000          1          wP         Female
3 IU/ML                 0.530000          1          wP         Female
4 IU/ML                 6.205949          1          wP         Female
5 IU/ML                 4.679535          1          wP         Female
6 IU/ML                 2.816431          1          wP         Female
               ethnicity  race year_of_birth date_of_boost      dataset
1 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
2 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
3 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
4 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
5 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
6 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
  actual_day_relative_to_boost planned_day_relative_to_boost specimen_type
1                           -3                             0         Blood
2                           -3                             0         Blood
3                           -3                             0         Blood
4                           -3                             0         Blood
5                           -3                             0         Blood
6                           -3                             0         Blood
  visit
1     1
2     1
3     1
4     1
5     1
6     1
```

Q. What Ab isotypes are measured?

```r
table(abdata$isotype)
```

```
  IgE   IgG  IgG1  IgG2  IgG3  IgG4 
 6698  7265 11993 12000 12000 12000 
```

Q. What Ags are reported

```r
table(abdata$antigen)
```

```
    ACT   BETV1      DT   FELD1     FHA  FIM2/3   LOLP1     LOS Measles     OVA 
   1970    1970    6318    1970    6712    6318    1970    1970    1970    6318 
    PD1     PRN      PT     PTM   Total      TT 
   1970    6712    6712    1970     788    6318 
```

Let's focus on the Ags and make a plot of MFI_normalised for all Ag.

```r
igg <- abdata |>
  filter(isotype == "IgG")
head(igg)
```

```
  specimen_id isotype is_antigen_specific antigen        MFI MFI_normalised
1           1     IgG                TRUE      PT   68.56614       3.736992
2           1     IgG                TRUE     PRN  332.12718       2.602350
3           1     IgG                TRUE     FHA 1887.12263      34.050956
4          19     IgG                TRUE      PT   20.11607       1.096366
5          19     IgG                TRUE     PRN  976.67419       7.652635
6          19     IgG                TRUE     FHA   60.76626       1.096457
   unit lower_limit_of_detection subject_id infancy_vac biological_sex
1 IU/ML                 0.530000          1          wP         Female
2 IU/ML                 6.205949          1          wP         Female
3 IU/ML                 4.679535          1          wP         Female
4 IU/ML                 0.530000          3          wP         Female
5 IU/ML                 6.205949          3          wP         Female
6 IU/ML                 4.679535          3          wP         Female
               ethnicity  race year_of_birth date_of_boost      dataset
1 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
2 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
3 Not Hispanic or Latino White    1986-01-01    2016-09-12 2020_dataset
4                Unknown White    1983-01-01    2016-10-10 2020_dataset
5                Unknown White    1983-01-01    2016-10-10 2020_dataset
6                Unknown White    1983-01-01    2016-10-10 2020_dataset
  actual_day_relative_to_boost planned_day_relative_to_boost specimen_type
1                           -3                             0         Blood
2                           -3                             0         Blood
3                           -3                             0         Blood
4                           -3                             0         Blood
5                           -3                             0         Blood
```

```
6                           -3                              0          Blood
  visit
1     1
2     1
3     1
4     1
5     1
6     1
```

```
ggplot(igg) +
  aes(MFI_normalised, antigen) +
  geom_boxplot()
```



Q. Is there a difference for aP vs wP individuals with these values?

```
ggplot(igg) +
  aes(MFI_normalised, antigen) +
  geom_boxplot() +
  facet_wrap(~infancy_vac)
```



OR

```
ggplot(igg) +
  aes(MFI_normalised, antigen, col=infancy_vac) +
  geom_boxplot()
```



Q. Is there a temporal response - ie do values increase or decrease over time

```
ggplot(igg) +
  aes(MFI_normalised, antigen, col= infancy_vac) +
  geom_boxplot() +
  facet_wrap(~visit)
```



Focus on PT (Pertussis Toxin) Ag

```
pt.igg.21 <- igg |> filter(antigen == "PT",
              dataset == "2021_dataset")
```

```
ggplot(pt.igg.21) +
  aes(planned_day_relative_to_boost, MFI_normalised,
      col=infancy_vac,
      group = subject_id) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept= 14, lty=2)
```




```
library(dplyr)

avg_df <- pt.igg.21 %>%
  group_by(infancy_vac, planned_day_relative_to_boost) %>%
  summarise(mean_MFI = mean(MFI_normalised), .groups = "drop")
```

```
ggplot(pt.igg.21) +
  aes(planned_day_relative_to_boost, MFI_normalised,
      color = infancy_vac,
      group = subject_id) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept = 14, lty = 2) +
 stat_summary(aes(group = infancy_vac),
               fun = mean,
               geom = "line",
               linewidth = 1.4)+
  labs(
    title = "CMI-PB 2021 dataset IgG PT",
    subtitle = "Dashed lines at day 0 (pre boost) and day 14 (post boost)",
    x = "Day relative to boost",
    y = "Normalised MFI",
    colour = "infancy_vac"
```

```
)
```


CMI−PB 2021 dataset IgG PT
Dashed lines at day 0 (pre boost) and day 14 (post boost)
Normalised MFI
15
10
5
0
0
25
50
75
100
125
Day relative to boost
infancy_vac
aP
wP