Deriving the Integration Threshold

Information-Theoretic Constraints on $I_{\min}$

Aaron Ben-Shalom Claude

Working Notes — January 2026

Abstract

We explore several approaches to deriving the minimum integration $I_{\min}$ required for self-reference from first principles. If $I_{\min}$ can be determined independently, and if $I_{\min} \sim L_H^2$ (Hubble scale squared), then combined with $\Lambda \leq 1/I_{\min}$, this would predict the cosmological constant. We examine information-theoretic, holographic, computational, and channel-capacity arguments.

Contents

1 The Goal **3**

2 Approach 1: Bekenstein Bound **3**
2.1 The Bound . . . 3
2.2 Application to Self-Reference . . . 3
2.3 Connecting to Integration . . . 3
2.4 Cosmological Application . . . 4
2.5 Revision: Minimal Self-Model . . . 4

3 Approach 2: Holographic Principle **4**
3.1 The Principle . . . 4
3.2 Self-Reference as Boundary-Bulk Correspondence . . . 5
3.3 Integration from Holography . . . 5

4 Approach 3: Channel Capacity **5**
4.1 Information Propagation . . . 5
4.2 Self-Reference Requires Causal Contact . . . 5
4.3 Cosmological Application . . . 6

5 Approach 4: Fixed-Point Stability **6**
5.1 From the Self-Aligning Topologies Paper . . . 6
5.2 Perturbation Analysis . . . 6
5.3 Cosmological Perturbations . . . 6

6 Approach 5: Computational Complexity **7**
6.1 Self-Modeling as Computation . . . 7
6.2 Margolus-Levitin Bound . . . 7
6.3 Self-Modeling Operations . . . 7
6.4 Cosmological Application . . . 7

7 Synthesis: Two Independent Derivations **7**
7.1 Argument 1: Diffusion/Channel Capacity 8
7.2 Argument 2: Computational Capacity 8
7.3 Revised Argument 2 8
7.4 Resolution: Coarse-Grained Self-Models 9

8 Most Promising Direction **9**

9 Open Questions **9**

10 Next Steps **10**

1 The Goal

From the cosmological integration bound, we have:

$$\Lambda \leq \frac{1}{I_{\min}}$$

The observed cosmological constant $\Lambda \approx 10^{-52}$ m^{-2} implies:

$$I_{\min} \lesssim 10^{52} \text{ m}^2 \approx L_H^2$$

where $L_H \approx 10^{26}$ m is the Hubble radius.
Question: Can we derive $I_{\min} \sim L_H^2$ from independent principles?
If yes, this would:

1. Predict Λ rather than just bound it
2. Explain why $\Lambda \sim H^2$ (the coincidence problem)
3. Ground the framework in fundamental physics

2 Approach 1: Bekenstein Bound

2.1 The Bound

Theorem 2.1 (Bekenstein, 1981). *[ESTABLISHED]The maximum entropy (information) that can be contained in a region of space with energy E and radius R is:*

$$S \leq S_B := \frac{2\pi RE}{\hbar c}$$

In Planck units ($\hbar = c = G = 1$):

$$S \leq 2\pi RE$$

2.2 Application to Self-Reference

Conjecture 2.2 (Self-Reference Information Requirement). *[CONJECTURED]A self-referential system must encode at least N bits of information about itself, where N is the minimum complexity for a self-model.*

For a system of size L with energy density ρ:

$$E \sim \rho L^3, \quad S_B \sim \rho L^4$$

If self-reference requires $S \geq N$:

$$\rho L^4 \gtrsim N \implies L \gtrsim \left(\frac{N}{\rho}\right)^{1/4}$$

2.3 Connecting to Integration

The integration measure is $I = 1/\lambda$ where λ is the eigenvalue. For a region of size L:

$$\lambda \sim \frac{1}{L^2} \implies I \sim L^2$$

So:

$$I_{\min} \sim L_{\min}^2 \sim \left(\frac{N}{\rho}\right)^{1/2}$$

2.4 Cosmological Application

For the universe as a whole:

- $\rho \sim \Lambda$ (dominated by dark energy)
- $N \sim ?$ (information content of cosmic self-model)

If we take $N \sim S_{\text{horizon}} \sim L_H^2/\ell_P^2$ (holographic entropy of the cosmic horizon):

$$I_{\min} \sim \left(\frac{L_H^2/\ell_P^2}{\Lambda}\right)^{1/2} \sim \left(\frac{L_H^2}{\ell_P^2 \Lambda}\right)^{1/2}$$

Using $\Lambda \sim 1/L_H^2$:

$$I_{\min} \sim \left(\frac{L_H^4}{\ell_P^2}\right)^{1/2} = \frac{L_H^2}{\ell_P}$$

This is *larger* than L_H^2 by a factor of $1/\ell_P \sim 10^{35}$.
Problem: This gives $I_{\min} \sim 10^{87}$ m^2, not 10^{52} m^2.

2.5 Revision: Minimal Self-Model

Perhaps N should be much smaller—the minimum bits for a self-model, not the full holographic content.

Conjecture 2.3 (Minimal Self-Model Complexity). *[SPECULATIVE]The minimum information for self-reference is $N_{\min} \sim \log_2(\text{states})$ for the simplest self-referential computation, which may be $O(1)$ or $O(\log L_H/\ell_P)$.*

If $N \sim \log(L_H/\ell_P) \sim 100$:

$$I_{\min} \sim \left(\frac{100}{\Lambda}\right)^{1/2} \sim \left(100 \cdot L_H^2\right)^{1/2} \sim 10 L_H$$

This gives $I_{\min} \sim L_H$, still not L_H^2.
Conclusion: The Bekenstein bound approach doesn't naturally give $I_{\min} \sim L_H^2$.

3 Approach 2: Holographic Principle

3.1 The Principle

Theorem 3.1 ('t Hooft, Susskind). *[ESTABLISHED]The maximum entropy in a region is proportional to its boundary area, not volume:*

$$S_{\max} = \frac{A}{4\ell_P^2}$$

For a sphere of radius L:

$$S_{\max} = \frac{4\pi L^2}{4\ell_P^2} = \frac{\pi L^2}{\ell_P^2}$$

3.2 Self-Reference as Boundary-Bulk Correspondence

Conjecture 3.2 (Holographic Self-Reference). *[SPECULATIVE]Self-reference requires the bulk to encode information about itself on the boundary. This is possible only if the bulk dynamics are "simple enough" to be reconstructed from boundary data.*

In AdS/CFT, the bulk can be reconstructed from the boundary. The reconstruction requires:

- Boundary theory has sufficient degrees of freedom
- Bulk fields satisfy appropriate equations (Einstein + matter)
- Entanglement structure matches (Ryu-Takayanagi)

For de Sitter (positive Λ), holography is less understood, but there are proposals:

- dS/CFT correspondence (Strominger)
- Static patch holography
- Stretched horizon approaches

3.3 Integration from Holography

If the integration I measures "how much bulk information is encoded coherently":

Conjecture 3.3 (Holographic Integration). *[SPECULATIVE]*

$$I \sim \frac{(\textit{bulk volume})}{(\textit{boundary area})} \sim \frac{L^3}{L^2} = L$$

for a region of size L.

This would give $I_{\min} \sim L_H$, not L_H^2.

Alternative: If integration measures the square of the coherence length:

$$I \sim L^2$$

then $I_{\min} \sim L_H^2$ directly, but this needs justification.

4 Approach 3: Channel Capacity

4.1 Information Propagation

Self-reference requires information about distant parts of the system to be available locally. This requires communication across the system.

Definition 4.1 (Diffusion Time). On a manifold with Laplacian Δ, information diffuses according to the heat equation. The time to diffuse across a region of size L is:

$$t_{\text{diff}} \sim \frac{L^2}{D}$$

where D is the diffusion coefficient. For the normalized Laplacian, $D \sim 1$, so $t_{\text{diff}} \sim L^2$.

4.2 Self-Reference Requires Causal Contact

Conjecture 4.2 (Causal Self-Reference). *[CONJECTURED]A self-referential system at scale L must be able to "communicate with itself" within its characteristic timescale. This requires:*

$$t_{\textit{self-reference}} \geq t_{\textit{diff}} \sim L^2$$

The integration $I = 1/\lambda$ has units of $(\text{length})^2$, which matches the diffusion time (in units where $D = 1$).

Interpretation: I is the timescale for coherent self-reference. Large I means the system has time to "know itself" before modes decay.

4.3 Cosmological Application

For the universe to be self-referential at the Hubble scale:

$$I_{\min} \sim L_H^2$$

This gives us $I_{\min} \sim L_H^2$ directly!

Proposition 4.3 (Channel Capacity Derivation). *[DERIVED]If self-reference requires information to diffuse across the system, and diffusion time scales as L^2, then:*

$$I_{\min} \sim L_{system}^2$$

For cosmic-scale self-reference:

$$I_{\min} \sim L_H^2$$

This is promising. Let's develop it further.

5 Approach 4: Fixed-Point Stability

5.1 From the Self-Aligning Topologies Paper

The self-aligning topologies work showed that stable eigenvector fixed points require:

1. Non-degenerate spectrum (eigenvalues well-separated)
2. Sufficient spectral gap (first eigenvalue not too small)
3. Resistance to local perturbation

5.2 Perturbation Analysis

Theorem 5.1 (Davis-Kahan). *[ESTABLISHED]If A and $\tilde{A} = A + E$ are self-adjoint with $\|E\| \leq \epsilon$, and λ is a simple eigenvalue of A with eigenvector v, then there exists an eigenvector $\tilde{v}$ of $\tilde{A}$ with:*

$$\sin \angle(v, \tilde{v}) \leq \frac{\epsilon}{gap}$$

where gap is the distance from λ to the rest of the spectrum.

For stability, we need:

$$\text{gap} \gg \epsilon$$

5.3 Cosmological Perturbations

In the real universe, the spatial metric has perturbations $\delta g/g \sim 10^{-5}$ (from CMB observations).

The perturbation to the Laplacian is $\epsilon \sim \delta g/g \cdot \lambda \sim 10^{-5}\lambda$.

For stability:

$$\lambda_2 - \lambda_1 \gg 10^{-5}\lambda_1$$

From the sphere spectrum, $\lambda_2/\lambda_1 = 8/3$, so:

$$\lambda_2 - \lambda_1 = \frac{5}{3}\lambda_1$$

This is $\gg 10^{-5}\lambda_1$, so stability is satisfied.

Conclusion: Perturbation stability doesn't strongly constrain $I_{\min}$; the de Sitter spectrum is robust.

6 Approach 5: Computational Complexity

6.1 Self-Modeling as Computation

Definition 6.1 (Self-Model Computation)**.** A self-referential system computes a function $M : \mathcal{H} \to \mathcal{H}$ where $M(x)$ is the system's model of its own state x. A fixed point $x^* = M(x^*)$ is a consistent self-model.

Conjecture 6.2 (Computational Resources)**.** *[SPECULATIVE]Computing M requires resources (time, space, energy) proportional to the complexity of the system being modeled.*

6.2 Margolus-Levitin Bound

Theorem 6.3 (Margolus-Levitin, 1998)**.** *[ESTABLISHED]The maximum rate of computation is bounded by energy:*

$$\textit{operations per second} \leq \frac{2E}{\pi\hbar}$$

For a system of energy E and lifetime T, the maximum operations is:

$$N_{\text{ops}} \leq \frac{2ET}{\pi\hbar}$$

6.3 Self-Modeling Operations

If self-modeling requires N operations, where N scales with system size:

$$N \sim L^d \quad \text{(for } d\text{-dimensional system)}$$

And the system has energy $E \sim \rho L^3$ and lifetime $T \sim L$ (light-crossing time):

$$N_{\text{ops}} \sim \rho L^4$$

For self-reference to be possible:

$$\rho L^4 \gtrsim L^d$$

In $d = 3$:

$$\rho L^4 \gtrsim L^3 \implies L \gtrsim \rho^{-1}$$

This says the system must be large enough that it has enough computational capacity to model itself.

6.4 Cosmological Application

With $\rho \sim \Lambda$:

$$L_{\min} \sim \Lambda^{-1} \sim L_H^2/L_H = L_H$$

And:

$$I_{\min} \sim L_{\min}^2 \sim L_H^2$$

This also gives $I_{\min} \sim L_H^2$!

7 Synthesis: Two Independent Derivations

We have found two arguments that give $I_{\min} \sim L_H^2$:

7.1 Argument 1: Diffusion/Channel Capacity

Self-reference requires information to propagate across the system. Propagation time scales as L^2 (diffusion). Integration I has units of L^2 and represents the coherence timescale. Therefore:

$$I_{\text{min}} \sim L^2_{\text{system}} \sim L_H^2$$

7.2 Argument 2: Computational Capacity

Self-modeling requires $\sim L^3$ operations. The Margolus-Levitin bound gives $\sim \rho L^4$ available operations. Requiring these to match:

$$\rho L^4 \sim L^3 \implies L \sim \rho^{-1} \sim \Lambda^{-1} \sim L_H^2/L_H$$

Wait, let me redo this. We have $\rho \sim \Lambda \sim L_H^{-2}$:

$$\rho^{-1} \sim L_H^2$$

So $L_{\text{min}} \sim L_H^2$? That doesn't make sense dimensionally. Let me reconsider.

Actually, ρ has units of (mass)/(length)^3 or (length)$^{-2}$ in natural units. So ρ^{-1} has units of (length)^2.

Hmm, the Margolus-Levitin analysis needs more care with units.

7.3 Revised Argument 2

In natural units where $c = \hbar = 1$:

- Energy density: $[\rho] = L^{-4}$
- Cosmological constant: $[\Lambda] = L^{-2}$
- For dark energy dominated: $\rho \sim \Lambda/(8\pi G) \sim \Lambda \cdot M_P^2 \sim \Lambda/\ell_P^2$

The computational capacity in volume L^3 over time T:

$$N_{\text{ops}} \sim \rho L^3 \cdot T \sim \frac{\Lambda}{\ell_P^2} L^3 T$$

For self-modeling requiring $N \sim L^3/\ell_P^3$ operations (one per Planck volume):

$$\frac{\Lambda}{\ell_P^2} L^3 T \gtrsim \frac{L^3}{\ell_P^3}$$

$$\Lambda T \gtrsim \frac{1}{\ell_P}$$

With $\Lambda \sim L_H^{-2}$:

$$T \gtrsim \frac{L_H^2}{\ell_P} \sim 10^{61} \text{ s}$$

This is much longer than the age of the universe ($\sim 10^{17}$ s).

Problem: This suggests self-modeling at the Planck scale is impossible within cosmic time.

7.4 Resolution: Coarse-Grained Self-Models

Perhaps self-reference doesn't require Planck-scale modeling. A coarse-grained self-model with $N \sim (L/L_*)^3$ degrees of freedom, where L_* is some coarse-graining scale:

$$\frac{\Lambda}{\ell_P^2} L^3 T \gtrsim \frac{L^3}{L_*^3}$$

$$T \gtrsim \frac{\ell_P^2}{\Lambda L_*^3}$$

For $T \sim L_H$ (Hubble time) and $\Lambda \sim L_H^{-2}$:

$$L_H \gtrsim \frac{\ell_P^2 L_H^2}{L_*^3} \implies L_*^3 \gtrsim \ell_P^2 L_H$$

$$L_* \gtrsim (\ell_P^2 L_H)^{1/3} \sim 10^{-15} \text{ m}$$

This is around the nuclear scale—perhaps the scale of atoms or nuclei sets the coarse-graining for cosmic self-reference?

8 Most Promising Direction

The **diffusion/channel capacity argument** is cleanest:

Core Argument
Premise 1: Self-reference requires information about the whole system to be available to any part.
Premise 2: Information propagation on a manifold with Laplacian Δ has characteristic timescale $\tau \sim L^2/D$ for distance L and diffusion constant D.
Premise 3: Integration $I = 1/\lambda$ represents the timescale for mode coherence; modes with $I < \tau$ decay before information can propagate.
Conclusion: Self-reference at scale L requires $I \geq L^2$, i.e., $I_{\min} \sim L^2$.
Application: For cosmic self-reference, $L \sim L_H$, so $I_{\min} \sim L_H^2$.

Combined with $\Lambda \leq 1/I_{\min}$:

$$\boxed{\Lambda \lesssim \frac{1}{L_H^2} \sim H^2}$$

This "predicts" $\Lambda \sim H^2$—which is exactly the observed coincidence!

9 Open Questions

1. **Is diffusion the right model?** Information propagation in quantum systems may be faster (Lieb-Robinson bounds). Does this change the analysis?
2. **Why the Hubble scale?** The argument says $I_{\min} \sim L_{\text{system}}^2$. Why is the relevant "system" the Hubble volume rather than, say, a galaxy or a brain?
3. **Tightening the bound:** Can we get $\Lambda = 1/I_{\min}$ (equality) rather than $\Lambda \leq 1/I_{\min}$ (inequality)?
4. **Time evolution:** As the universe expands, L_H grows. Does Λ track this, or is Λ fixed while $I_{\min}$ changes?
5. **Multiple observers:** If there are many self-referential systems at different scales, which one sets $I_{\min}$?

10 Next Steps

1. **Formalize the diffusion argument:** State precisely what "information propagation" means in terms of the Laplacian spectrum. Connect to spectral graph theory results on mixing times.

2. **Justify "cosmic self-reference":** What does it mean for the universe to be self-referential? Is this the same as containing observers, or something stronger?

3. **Compare to other Λ predictions:** How does this compare to other attempts to predict Λ (e.g., Weinberg's anthropic bound, causal set predictions)?

4. **Check consistency with inflation:** During inflation, Λ_{eff} was much larger. Does the framework allow this? (Perhaps observers require post-inflationary conditions.)