

Critical Points

October 2, 2011

We start with the nondimensionalized effective potential,

$$V_{eff} = \frac{1}{\rho^2} \left(n^2 - 2\rho e^{-\rho/\lambda} \right) \quad (1)$$

and attempt to find the points at which the minimum of the effective potential is 0 and where the minimum of the effective potential no longer exists.

For the first, we start by taking the derivative of (1) and setting it to 0, obtaining

$$\frac{dV}{d\rho_0} = -\frac{2n^2}{\rho_0^3} + \frac{2}{\rho_0^2} e^{-\rho_0/\lambda} + \frac{2}{\rho_0 \lambda} e^{-\rho_0/\lambda} \quad (2)$$

$$0 = -2n^2 + 2\rho_0 e^{-\rho_0/\lambda} + 2\frac{\rho_0^2}{\lambda} e^{-\rho_0/\lambda} \quad (3)$$

where ρ_0 is the value of ρ when V_{eff} is minimized. This equation has no general solution, but we can find one for the specific case where $V_{eff}(\rho_0) = 0$ by solving the equation

$$V_{eff}(\rho_0) = 0 = n^2 - 2\rho_0 e^{-\rho_0/\lambda} \quad (4)$$

$$n^2 = 2\rho_0 e^{-\rho_0/\lambda} \quad (5)$$

and plugging it in to the previous, obtaining

$$0 = -2n^2 + n^2 + n^2 \frac{\rho_0}{\lambda} \quad (6)$$

$$= -1 + \frac{\rho_0}{\lambda} \quad (7)$$

$$\rho_0 = \lambda. \quad (8)$$

Plugging this back into the equation for $V_{eff}(\rho_0)$ gives us

$$n = \sqrt{\frac{2}{e}} \sqrt{\lambda} \quad (9)$$

which is pretty close to the naive $n = \sqrt{\lambda}$.

For the second critical point, the one where V_{eff} has no minimum, we rewrite (3) to be

$$n^2 = \left(\rho + \frac{\rho^2}{\lambda} \right) e^{-\rho/\lambda} \quad (10)$$

The right side of this equation will have some maximum, which will be the last point for which the equation can be solved – after that, the value of n^2 will just plain be bigger than the right side. To find this, we differentiate with respect to ρ and set to 0, obtaining

$$0 = (1 + 2\frac{\rho}{\lambda})e^{-\rho/\lambda} - \frac{1}{\lambda}(\rho + \frac{\rho^2}{\lambda})e^{-\rho/\lambda} \quad (11)$$

$$= 1 + \frac{\rho}{\lambda} + \frac{\rho^2}{\lambda^2}. \quad (12)$$

This is a quadratic in ρ and can be solved using the quadratic formula:

$$\rho = \frac{-\frac{1}{\lambda} \pm \sqrt{\left(\frac{1}{\lambda}\right)^2 + 4\frac{1}{\lambda^2}}}{-\frac{2}{\lambda^2}} \quad (13)$$

$$= \frac{\lambda}{2} \mp \frac{\lambda\sqrt{5}}{2} \quad (14)$$

As ρ is a real physical quantity that can't be negative, we can discard one solution, obtaining

$$\rho = \frac{\lambda}{2}(1 + \sqrt{5}) \quad (15)$$

We can plug this in to (3) to find the relationship between λ and n :

$$n^2 = \lambda(1 + \sqrt{5}) \left(\frac{3 + \sqrt{5}}{4} \right) e^{-(1+\sqrt{5})/2} \quad (16)$$