

1. Find the eigenvalues and eigenvectors of the matrices

$$\mathbf{A} = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} -3 & 1 & -3 \\ 20 & 3 & 10 \\ 2 & -2 & 4 \end{bmatrix}$$

2. Determine if either of the following matrices is diagonalizable

$$\mathbf{A} = \begin{bmatrix} -1 & -1 & -2 \\ 8 & -11 & -8 \\ -10 & 11 & 7 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} 1 & -4 & -4 \\ 8 & -11 & -8 \\ -8 & 8 & 5 \end{bmatrix}$$

3. Estimate the eigenvalues of $\mathbf{A} = \begin{bmatrix} 5 & 1 & 1 \\ 0 & 6 & 1 \\ 1 & 0 & -5 \end{bmatrix}$

4. Determine the induced norm $\|\mathbf{A}\|_2$ as well as $\|\mathbf{A}^{-1}\|_2$ for the nonsingular matrix

$$\mathbf{A} = \frac{1}{\sqrt{3}} \begin{bmatrix} 3 & -1 \\ 0 & \sqrt{8} \end{bmatrix}$$

5. Let $\mathbf{A} \in \mathbb{C}^{m \times n}$. Please prove the following statements. (a) $\|\mathbf{A}\|_p = \max_{\|x\|_p=1} \|\mathbf{A}x\|_p$ is a norm for any $p \geq 1$ (b) $\|\mathbf{A}x\|_p \leq \|\mathbf{A}\|_p \|x\|_p$ holds for any $p \geq 1$ (c) $\|\mathbf{A}\mathbf{B}\|_p \leq \|\mathbf{A}\|_p \|\mathbf{B}\|_p$ holds for any $p \geq 1$ (d) $\|\mathbf{Q}\mathbf{A}\mathbf{U}\|_F = \|\mathbf{A}\|_F$ holds for any unitary matrices $\mathbf{Q} \in \mathbb{C}^{m \times m}$ and $\mathbf{U} \in \mathbb{C}^{n \times n}$ (e) $\|\mathbf{Q}\mathbf{A}\mathbf{U}\|_2 = \|\mathbf{A}\|_2$ holds for any unitary matrices $\mathbf{Q} \in \mathbb{C}^{m \times m}$ and $\mathbf{U} \in \mathbb{C}^{n \times n}$

6. Using the induced matrix norm, prove that if \mathbf{A} is nonsingular, then

$$\min_{\|x\|=1} \|\mathbf{A}x\| = \frac{1}{\|\mathbf{A}^{-1}\|}$$

7. Let $\mathbf{A} \in \mathbb{C}^{m \times n}$ is a Vandemonde matrix with distinct roots. Please verify that any collection of r columns of \mathbf{A} , with $r \leq m$, is linearly independent.