

## Note

### On Vector Partition Functions

BERND STURMFELS\*

*Department of Mathematics, University of California, Berkeley, California 94720*

*Communicated by Victor Klee*

Received May 10, 1994

We present a structure theorem for vector partition functions. The proof rests on a formula due to Peter McMullen for counting lattice points in rational convex polytopes. © 1995 Academic Press, Inc.

#### INTRODUCTION

Let  $A = (a_1, \dots, a_n)$  be a  $d \times n$ -matrix of rank  $d$  with entries in  $\mathbb{N}$ , the set of non-negative integers. The corresponding *vector partition function*  $\phi_A: \mathbb{N}^d \rightarrow \mathbb{N}$  is defined as follows:  $\phi_A(u)$  is the number of non-negative integer vectors  $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{N}^n$  such that  $A \cdot \lambda = \lambda_1 a_1 + \dots + \lambda_n a_n = u$ . Equivalently, the function  $\phi_A$  is defined by the formal power series:

$$\prod_{i=1}^n \frac{1}{(1 - t_1^{a_{1i}} t_2^{a_{2i}} \dots t_d^{a_{di}})} = \sum_{u \in \mathbb{N}^d} \phi_A(u_1, \dots, u_d) \cdot t_1^{u_1} t_2^{u_2} \dots t_d^{u_d}. \quad (1)$$

Vector partition functions appear in many areas of mathematics and its applications, including representation theory [9], commutative algebra [14], approximation theory [4] and statistics [5].

It was shown by Blakley [2] that there exists a finite decomposition of  $\mathbb{N}^d$  such that  $\phi_A$  is a polynomial of degree  $n - d$  on each piece. Here we describe such a decomposition explicitly and we analyze how the polynomials differ from piece to piece. Our construction uses the geometric decomposition into chambers studied by Alekseevskaya, Gelfand and Zelevinsky in [1]. Within each chamber we give a formula which refines

\* E-mail address: bernd@math.berkeley.edu.

the results by Dahmen and Micchelli in [4, §3]. The objective of this note is to provide polyhedral tools for the efficient computation of vector partition functions, with a view towards applications, such as the sampling algorithms in [6].

EXAMPLE ( $n = 6$ ,  $d = 3$ ). Consider the vector partition function

$$\phi_A : \mathbb{N}^3 \rightarrow \mathbb{N}, (u, v, w) \mapsto \# \{ \lambda \in \mathbb{N}^6 : A \cdot \lambda = (u, v, w)^t \}$$

associated with the matrix

$$A = \begin{pmatrix} 2 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 2 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 2 \end{pmatrix}.$$

In this instance the value of  $\phi_A$  does not depend on the permutation of  $(u, v, w)$ , so we may assume that  $u \geq v \geq w$ . Also, if  $u + v + w \equiv 1 \pmod{2}$  then  $\phi_A(u, v, w) = 0$ , so we shall assume that  $u + v + w \equiv 0 \pmod{2}$ . Given these assumptions, we distinguish two cases:

Case 1.  $u \geq v + w$ . Then

$$\phi_A(u, v, w)$$

$$\begin{aligned} &= \frac{vw}{2} + \frac{vw^2}{8} - \frac{w^3}{24} \\ &+ \begin{cases} 1 + v/2 + 2w/3 & \text{if } u \equiv 0 \pmod{2} \text{ and } v \equiv 0 \pmod{2}, \\ 1/2 + v/2 + 5w/12 & \text{if } u \equiv 1 \pmod{2} \text{ and } v \equiv 1 \pmod{2}, \\ 1/2 + 3v/8 + 13w/24 & \text{otherwise.} \end{cases} \end{aligned}$$

Case 2.  $u < v + w$ . We set

$$\begin{aligned} \psi &:= -u^2/8 + uv/4 + uw/4 - v^2/8 + vw/4 - w^2/8 \\ &\quad + u^3/48 - u^2v/16 - u^2w/16 + uv^2/16 + uvw/8 + uw^2/16 \\ &\quad - v^3/48 - v^2w/16 + vw^2/16 - w^3/16. \end{aligned}$$

Then

$$\phi_A(u, v, w)$$

$$= \psi + \begin{cases} 1 + u/6 + v/3 + w/2 & \text{if } u \equiv 0 \pmod{2} \text{ and } v \equiv 0 \pmod{2}, \\ 1/2 + u/6 + v/3 + w/4 & \text{if } u \equiv 1 \pmod{2} \text{ and } v \equiv 1 \pmod{2}, \\ 1/2 + u/6 + 5v/24 + 3w/8 & \text{otherwise.} \end{cases}$$

Already this simple example illustrates the main feature of vector partition functions, which is the interplay of a structure of convex polyhedra (as seen in the distinction of cases 1 and 2) with a structure of finite abelian groups (as seen in the “mod”-subcases).

In order to deal with the general case, we introduce some notation. Let  $\text{pos}(A) = \{\sum_{i=1}^n \lambda_i a_i \in \mathbf{R}^n : \lambda_1, \dots, \lambda_n \geq 0\}$ . For  $\sigma \subset [n] := \{1, \dots, n\}$  we consider the submatrix  $A_\sigma := (a_i : i \in \sigma)$ , the polyhedral cone  $\text{pos}(A_\sigma)$ , and the abelian group  $\mathbf{Z}A_\sigma$  spanned by the columns of  $A_\sigma$ . We may assume without loss of generality that  $A$  is surjective over  $\mathbf{Z}$ , that is,  $\mathbf{Z}A = \mathbf{Z}^d$ . This implies that the semigroup  $\mathbf{N}A := \text{pos}(A) \cap \mathbf{Z}A$  is saturated.

The surjectivity assumption does not hold for the  $3 \times 6$ -matrix in our example. In order to apply the results below to such a case, one must choose a rational  $3 \times 3$ -matrix  $B$  which defines an isomorphism from  $\mathbf{Z}A$  onto  $\mathbf{Z}^3$  and then use the formula  $\phi_A(u) = \phi_{BA}(Bu)$ .

A subset  $\sigma$  of  $[n]$  is a *basis* if  $\#(\sigma) = \text{rank}(A_\sigma) = d$ . The *chamber complex* is the polyhedral subdivision of the cone  $\text{pos}(A)$  which is defined as the common refinement of the simplicial cones  $\text{pos}(A_\sigma)$ , where  $\sigma$  runs over all bases. Each *chamber*  $C$  (meaning: maximal cell in the chamber complex) is indexed by the set  $\Delta(C) = \{\sigma \subset [n] : C \subseteq \text{pos}(A_\sigma)\}$ . For each  $\sigma \in \Delta(C)$ , the group  $\mathbf{Z}A_\sigma$  has finite index in  $\mathbf{Z}^d$ ; write  $G_\sigma := \mathbf{Z}^d / \mathbf{Z}A_\sigma$  for the group of residue classes. We say that  $\sigma$  is *non-trivial* if  $G_\sigma \neq \{0\}$ . For  $u \in \text{pos}(A) \cap \mathbf{N}^d$ , let  $[u]_\sigma$  denote the image of  $u$  in  $G_\sigma$ .

In the small example above there are 12 chambers; they are grouped into two equivalence classes with respect to the  $S_3$ -symmetry. The following theorem is our main result.

**THEOREM 1.** *For each chamber  $C$  there exists a polynomial  $P$  of degree  $n - d$  in  $u = (u_1, \dots, u_d)$ , and for each non-trivial  $\sigma \in \Delta(C)$  there exists a polynomial  $Q_\sigma$  of degree  $\#(\sigma) - d$  in  $u$  and a function  $\Omega_\sigma : G_\sigma \setminus \{0\} \rightarrow \mathbf{Q}$  such that, for all  $u \in \mathbf{N}A \cap C$ ,*

$$\phi_A(u) = P(u) + \sum \{ \Omega_\sigma([u]_\sigma) \cdot Q_\sigma(u) : \sigma \in \Delta(C) \text{ and } [u]_\sigma \neq 0 \}.$$

Moreover, the “corrector polynomials”  $Q_\sigma$  satisfy the linear partial differential equations

$$\sum_{i=1}^d a_{ij} \frac{\partial Q_\sigma}{\partial u_i} \equiv 0 \quad \text{for all } j \in \sigma \text{ such that } \sigma \setminus \{j\} \notin \Delta(C).$$

*Remarks.* (1) Thorem 1 provides a generalization of the theory of *denumerants* (the  $d=1$  case) which can be found in Comtet’s book [3, §2.6]. A nice MAPLE package for computing denumerants has been implemented by P. Lisoněk [10].

(2) Another important special case occurs when the matrix  $A$  is *unimodular*, which means that  $G_\sigma = \{0\}$  for each basis  $\sigma$ . In this case  $\phi_A$  is a polynomial function on each chamber [4, Corollary 3.1]. This happens, for instance, in the problem of counting non-negative integer matrices with prescribed row and column sums; see [5] for a general survey and see [13] for the computational state of the art.

(3) The main point of our formula is the “additive decoupling” of the correction term, which generalizes Theorem C in [3, §2.6]. The results of Dahmen and Micchelli in [4, §3] generalize a (somewhat weaker) classical theorem of Bell [3, §2.6, Thm. B]. The computational advantage of the additive decoupling is explained on page 114 in [3].

### THE PROOF

We shall use notation which is standard in the theory of toric varieties; see e.g. [7]. Let  $N$  be a lattice of rank  $m$ ,  $M = \text{Hom}(N, \mathbf{Z})$  its dual lattice, and  $N_{\mathbf{Q}}$  and  $M_{\mathbf{Q}}$  the corresponding rational vector spaces. Suppose we are given a complete simplicial fan  $\Sigma$  in  $N$  having  $n$  rays, and non-zero lattice points  $b_1, \dots, b_n$  on these rays. (The  $b_i$  need not be primitive in  $N$ !) We identify the cones of  $\Sigma$  with subsets of  $\{b_1, \dots, b_n\}$ . For each  $1 \leq l \leq m$  and each cone  $\tau = \{b_{\tau_1}, \dots, b_{\tau_l}\}$ , we let  $H_\tau$  denote the group  $\mathbf{Z}^\tau$  of integer valued functions on  $\tau$  modulo the subgroup of those functions on  $\tau$  which are restrictions from  $M = \text{Hom}(N, \mathbf{Z})$ . We say that  $\tau$  is *non-trivial* if  $H_\tau \neq \{0\}$ .

Consider any convex polytope of the form

$$P_\gamma = \{x \in M_{\mathbf{Q}} : \langle x, b_i \rangle \leq \gamma_i \text{ for } i = 1, \dots, n\}, \quad (2)$$

where  $\gamma = (\gamma_1, \dots, \gamma_n)$  ranges over the set  $C(\Sigma)$  of all vectors in  $\mathbf{Z}^n$  such that the normal fan of  $P_\gamma$  is coarser or equal to  $\Sigma$ . It is well known that there exists a polynomial function  $F = F(\gamma)$  on  $C(\Sigma)$  of degree  $m$  such that  $\#(P_\gamma \cap M) = F(\gamma)$  provided that  $P_\gamma$  is integral, i.e., all vertices of  $P_\gamma$  lie in  $M$ . For a toric proof of this fact see e.g. [8, §5]. In general, however, the polytope  $P_\gamma$  is not integral, since the fan  $\Sigma$  is not assumed to be smooth. The following proposition characterizes the difference between  $\#(P_\gamma \cap M)$  and the polynomial  $F(\gamma)$  in the general case. If  $\gamma \in \mathbf{Z}^n$ , then we write  $[\gamma]_\tau$  for the image of the function  $\tau \rightarrow \mathbf{Z}$ ,  $b_i \mapsto \gamma_i$  in the group  $H_\tau$ .

**PROPOSITION 2.** *For every non-trivial cone  $\tau \in \Sigma$  there exists a polynomial  $R_\tau$  of degree  $m - \#(\tau)$  in the variables  $\gamma = (\gamma_1, \dots, \gamma_n)$  and a function  $\omega_\tau : H_\tau \setminus \{0\} \rightarrow \mathbf{Q}$  such that  $\#(P_\gamma \cap M) - F(\gamma) = \sum \{\omega_\tau([\gamma]_\tau) \cdot R_\tau(\gamma) : \tau \in \Sigma \text{ and } [\gamma]_\tau \neq 0\}$  for all  $\gamma \in C(\Sigma)$ . Moreover, the polynomial  $R_\tau$  depends only on those variables  $\gamma_i$  for which  $\tau \cup \{b_i\} \in \Sigma$ .*

We shall use the following theorem of McMullen. If  $F$  is a face of a polytope  $P \subset M_{\mathbf{Q}}$  then  $v(P, F)$  denotes the cone in  $N$  normal to  $F$  at  $P$ . Let  $\mathcal{L}$  denote the set of all pairs  $(\tau, L)$  where  $\tau$  is a cone in  $N$  and  $L$  is an affine subspace of  $M_{\mathbf{Q}}$  which is a translate of  $\tau^{\perp}$ .

**THEOREM 3** (McMullen [11]). *There exists a function  $\theta: \mathcal{L} \rightarrow \mathbf{Q}$  such that  $\theta(\tau, L) = \theta(\tau, L + m)$  for all  $m \in M$  and*

$$\#(P \cap M) = \sum_{F \text{ face of } P} \theta(v(P, F), \text{aff}(F)) \cdot \text{Vol}(F) \text{ for every polytope } P \text{ in } M_{\mathbf{Q}}.$$

Here “Vol” denotes the standard volume form on the affine span  $\text{aff}(F)$  of the face  $F$ .

*Proof.* This is a special case of Theorem 3 in [11], provided one passes from simple valuations to general valuations using the technique in §3 of [11]. ■

**COROLLARY 4.** *If  $P_{\gamma}$  is an integral polytope then the number of lattice points in  $P_{\gamma}$  equals*

$$F(\gamma) = \sum_{\tau \in \Sigma} \theta(\tau, \tau^{\perp}) \cdot \text{Vol}(P_{\gamma}^{\tau}), \quad (3)$$

where  $P_{\gamma}^{\tau}$  denotes the face of  $P_{\gamma}$  supported by  $\tau$ .

*Proof.* If  $P_{\gamma}$  is integral then  $\text{aff}(P_{\gamma}^{\tau})$  is a lattice translate of the linear subspace  $\tau^{\perp}$ . Therefore  $\theta(\tau, \text{aff}(P_{\gamma}^{\tau})) = \theta(\tau, \tau^{\perp})$ , and the claim follows directly from Theorem 3. ■

We remark that formula (3) is a valid presentation for the polynomial function  $F(\gamma)$  throughout the cone  $C(\Sigma)$ , not just for those special values of  $\gamma$  for which  $P_{\gamma}$  is integral.

*Proof of Proposition 2.* Let  $\tau$  be a cone in  $\Sigma$  and let  $F_{\gamma}$  be the corresponding face of  $P_{\gamma}$ . As  $\gamma$  runs over  $C(\Sigma)$ , the volume of  $F_{\gamma}$  varies as a polynomial in  $\gamma$  of degree  $\dim(F_{\gamma}) = m - \#(\tau)$ . We set  $R_{\tau}(\gamma) := \text{Vol}(F_{\gamma})$ . This function is independent of a support parameter  $\gamma_i$  if the hyperplane  $\langle x, b_i \rangle = \gamma_i$  does not intersect the face  $F_{\gamma}$  for general  $\gamma$ . The latter condition is equivalent to  $\tau \cup \{b_i\}$  not being a cone of  $\Sigma$ . Hence  $R_{\tau}$  has the property asserted in the second part of Proposition 2.

Consider any other vector  $\gamma' \in C(\Sigma)$  and corresponding face  $F_{\gamma'}$  of  $P_{\gamma'}$ . Note that  $\text{aff}(F_{\gamma}) = \{y \in M : \forall b_i \in \tau : \langle y, b_i \rangle = \gamma_i\}$ , and similarly for  $F_{\gamma'}$ . This implies

$$\begin{aligned} [\gamma]_{\tau} = [\gamma']_{\tau} &\Leftrightarrow \exists u \in M : \forall b_i \in \tau : \gamma_i = \gamma'_i + \langle u, b_i \rangle \\ &\Leftrightarrow \exists u \in M : \text{aff}(F_{\gamma}) = \text{aff}(F_{\gamma'}) + u. \end{aligned}$$

We can therefore define a function  $\omega_\tau : H_\tau \setminus \{0\} \rightarrow \mathbf{Q}$  by setting

$$\omega_\tau([\gamma]_\tau) := \theta(\tau, \text{aff}(F_\gamma)) - \theta(\tau, \tau^\perp).$$

Proposition 2 now follows immediately from Theorem 3 and Corollary 4.  $\blacksquare$

*Proof of Theorem 1.* We shall use representation techniques as in [12, §5]. Let  $B = (b_1, \dots, b_n)$  be an integer  $(n-d) \times n$ -matrix whose row space (over  $\mathbf{Z}$ ) equals the kernel of  $A$ . In other words, we construct a short exact sequence of abelian groups

$$0 \longrightarrow \mathbf{Z}^{n-d} \xrightarrow{B^t} \mathbf{Z}^n \xrightarrow{A} \mathbf{Z}^d \longrightarrow 0.$$

We set  $m = n - d$  and  $M = \mathbf{Z}^{n-d}$ , and we consider the polytope  $P_\gamma$  in (2), for an arbitrary  $\gamma \in \mathbf{Z}^n$ . The map  $x \mapsto \gamma - B^t \cdot x$  defines a bijection between the lattice points in  $P_\gamma$  and the set of elements  $\lambda \in \mathbf{N}^n$  such that  $A \cdot \lambda = A \cdot \gamma$ . Therefore we have

$$\phi_A(A \cdot \gamma) = \#(P_\gamma \cap M). \quad (4)$$

We now fix a chamber  $C$  and we consider those vectors  $\gamma \in \mathbf{N}^n$  such that  $A \cdot \gamma$  lies in the interior of  $C$ . This determines the normal fan  $\Sigma$  of  $P_\gamma$  as follows:

$$\Sigma = \{ \{b_{\tau_1}, \dots, b_{\tau_k}\} : [n] \setminus \{\tau_1, \dots, \tau_k\} \in \mathcal{A}(C) \}.$$

Let us now fix  $\sigma \in \mathcal{A}(C)$  and set  $\tau := [n] \setminus \sigma$ . In order to derive the first part of Theorem 1 from Proposition 2, it suffices to show that there exists a group isomorphism  $\delta$  between  $H_\tau$  and  $G_\sigma$ , which takes a class  $[\gamma]_\tau$  in  $H_\tau$  to the class of  $[u]_\sigma$  in  $G_\sigma$ , where  $u := A \cdot \gamma$ . Indeed, in view of (4), we can then simply define  $P(u) := F(\gamma)$ ,  $\Omega_\sigma([u]_\sigma) := \omega_\tau([\gamma]_\tau)$ , and  $Q_\sigma(u) := R_\tau(\gamma)$  to get the desired formula for  $\phi_A(u)$ . (Note that  $\#(\sigma) - d = m - \#(\tau)$ .)

To define the group isomorphism  $\delta$ , we consider the short exact sequence

$$0 \longrightarrow \mathbf{Z}^\sigma \xrightarrow{i} \mathbf{Z}^n \xrightarrow{\pi} \mathbf{Z}^\tau \longrightarrow 0,$$

where  $i$  and  $\pi$  are the obvious coordinate inclusion and projection respectively. We have

$$H_\tau = \text{coker}(\pi \circ B^t) \quad \text{and} \quad G_\sigma = \text{coker}(A \circ i).$$

Consider any element of  $G_\sigma$ , given by a representative  $u \in \mathbf{Z}^d$ . We define  $\delta(u)$  to be  $\pi(\gamma)$ , where  $\gamma$  is any preimage of  $u$  under  $A$ . This defines a unique

element of  $H_\tau$  because  $\gamma$  is well-defined up to  $\text{im}(B') = \ker(A)$ . We have the equivalences

$$\begin{aligned} u \in \mathbf{Z}A_\sigma &\Leftrightarrow u \text{ has a preimage } \tilde{\gamma} \text{ under } A \text{ such that } \pi(\tilde{\gamma}) = 0 \\ &\Leftrightarrow \gamma - \tilde{\gamma} \in \ker(A) = \text{im}(B') \\ &\quad \text{for some } \tilde{\gamma} \in \mathbf{Z}^n \text{ such that } \pi(\tilde{\gamma}) = 0 \\ &\Leftrightarrow \pi(\gamma) = \pi(B' \cdot \lambda) \text{ for some } \lambda \in \mathbf{Z}^{n-d}. \end{aligned}$$

This shows that  $u$  is zero in  $G_\sigma$  if and only if  $\pi(\gamma) = \delta(u)$  is zero in  $H_\tau$ . Therefore the group homomorphism  $\delta$  is injective. But it is also surjective: if  $v \in \mathbf{Z}^\tau$ , then choose any  $w \in \mathbf{Z}$ , consider  $v + w \in \mathbf{Z}^n$  and define  $u = A(v + w)$ . Then  $\delta(u)$  and  $v$  represent the same element of  $H_\tau$ . This completes the proof of the first part of Theorem 1.

To prove the second part we note that an element  $j \in \sigma$  satisfies  $\sigma \setminus \{j\} \notin A(C)$  if and only if  $\tau \cup \{b_j\} \notin \Sigma$ . For such an index  $j$ , we apply the operator  $\partial/\partial\gamma_j$  to the polynomial

$$R_\tau(\gamma) = Q_\sigma(A \cdot \gamma).$$

The result is zero, by Proposition 2, and consequently  $\sum_{i=1}^d a_{ij} (\partial Q_\sigma / \partial u_i) \equiv 0$ , as required. ■

#### ACKNOWLEDGMENTS

I am grateful to Alexander Barvinok for helpful discussions. This project was supported in part by the National Science Foundation and the David and Lucile Packard Foundation.

#### REFERENCES

1. T. V. ALEKSEEVSKAYA, I. M. GEL'FAND, AND A. V. ZELEVINSKY, Arrangements of real hyperplanes and related partition functions, *Doklady Akad. Nauk. SSSR* **297** (1987), 1289–1293.
2. S. BLAKLEY, Combinatorial remarks on partitions of a multipartite number, *Duke Math. J.* **31** (1964), 335–340.
3. L. COMTET, "Advanced Combinatorics," Reidel, Dordrecht/Boston, 1974.
4. W. DAHMEN AND C. A. MICCHELLI, The number of solutions to linear diophantine equations and multivariate splines, *Trans. Amer. Math. Soc.* **308** (1988), 509–532.
5. P. DIACONIS AND A. GANGOLLI, Rectangular arrays with fixed margins, unpublished manuscript, 1994.
6. P. DIACONIS AND B. STURMFELS, Algebraic algorithms for sampling from conditional distributions, *Ann. Statist.*, to appear.
7. W. FULTON, "Introduction to Toric Varieties," Princeton Univ. Press, Princeton, NJ, 1993.

8. W. FULTON AND B. STURMFELS, Intersection theory on toric varieties, *Topology*, to appear.
9. G. J. HECKMAN, Projections of orbits and asymptotic behavior of multiplicities for compact connected Lie groups, *Invent. Math.* **67** (1982), 333–356.
10. P. LISONÉK, “Quasi-Polynomials: A Case Study in Experimental Combinatorics,” Technical report 93-18, RISC-Linz, Austria, 1993.
11. P. McMULLEN, Lattice invariant valuations on rational polytopes, *Arch. Math. (Basel)* **31** (1978), 509–516.
12. P. McMULLEN, Transforms, diagrams and representations, in “Contributions to Geometry” (J. Tölke and J. M. Wills, Eds.), Proceedings of the Geometry Symposium Siegen, Birkhäuser, Basel, 1979.
13. J. MOUNT, “Applications of Convex Sampling to Optimization and Contingency Table Counting,” Ph.D. thesis, Department of Computer Science, Carnegie Mellon University, 1995.
14. R. STANLEY, “Combinatorics and Commutative Algebra,” Birkhäuser, Boston, 1983.