

# CS184 Assignment 2

Gabe Fierro – 20964854 – Section: 103

1. There exists a  $k$  such that  $k \cdot k = 1$  and  $k \cdot j = 0$ .  $k$  can be  $-i$ , and there's only one vector like that in the 2D plane.
2. No, because we're only in 2D space.  $k$  cannot be a new vector orthogonal to both  $j$  and  $i$  if  $j$  and  $i$  are orthogonal to each other.
3. In 3D space there would be more possibilities for  $k$  in part **1**, and the answer to part **2** is yes.

2. For the matrix  $\begin{bmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{bmatrix}$ , the determinant is  $x_1y_2 + y_1x_3 + x_2y_3 - x_3y_2 - x_1y_3 - y_1x_2$ . The area of the triangle given by the three points can be modeled by two vectors  $\begin{bmatrix} x_2 - x_1 \\ y_2 - y_1 \end{bmatrix}$  and  $\begin{bmatrix} x_3 - x_1 \\ y_3 - y_1 \end{bmatrix}$ . The area of the triangle is given as half the cross product, which is  $\frac{1}{2}(x_1y_2 + y_1x_3 + x_2y_3 - x_3y_2 - x_1y_3 - y_1x_2)$ , which is proportional the determinant of the  $3 \times 3$  matrix. If the area of the triangle is 0, then the points are in a line.

3.  $f(x, y) = ax + by + c = 0$ , so we can uniquely determine the line by using the gradient  $\nabla f(x, y) = (a, b)$ . Because  $\begin{bmatrix} a \\ b \end{bmatrix} \cdot \begin{bmatrix} x_1 - x_0 \\ y_1 - y_0 \end{bmatrix} = 0$ , then we can say  $a = y_0 - y_1$  and  $b = x_1 - x_0$ , which gives us

$$f(x, y) = (y_0 - y_1)x + (x_1 - x_0)y + C = 0$$

at which point we can plug in  $(x_0, y_0)$  and solve for  $C$ .

4. For  $\{i, j, k\} = \{1, 2, 3\}, \{2, 3, 1\}, \{3, 1, 2\} : e_i \times e_j = e_k$  so

$$e_1 \times e_2 = \begin{bmatrix} 0 - 0 \\ 0 - 0 \\ 1 - 0 \end{bmatrix} = e_3 \quad e_2 \times e_3 = \begin{bmatrix} 1 - 0 \\ 0 - 0 \\ 0 - 0 \end{bmatrix} = e_1 \quad e_3 \times e_1 = \begin{bmatrix} 0 - 0 \\ 1 - 0 \\ 0 - 0 \end{bmatrix} = e_2$$

For  $\{i, j, k\} = \{1, 3, 2\}, \{3, 2, 1\}, \{2, 1, 3\} : e_i \times e_j = -e_k$  so

$$e_1 \times e_3 = \begin{bmatrix} 0 - 0 \\ 0 - 1 \\ 0 - 0 \end{bmatrix} = -e_2 \quad e_3 \times e_2 = \begin{bmatrix} 0 - 1 \\ 0 - 0 \\ 0 - 0 \end{bmatrix} = -e_1 \quad e_2 \times e_1 = \begin{bmatrix} 0 - 0 \\ 0 - 0 \\ 0 - 1 \end{bmatrix} = -e_3$$

5.  $\begin{bmatrix} x_1 & x_2 & x_3 & x_4 \\ y_1 & y_2 & y_3 & y_4 \\ z_1 & z_2 & z_3 & z_4 \\ 1 & 1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} x_1 & x_2 - x_1 & x_3 - x_1 & x_4 - x_1 \\ y_1 & y_2 - y_1 & y_3 - y_1 & y_4 - y_1 \\ z_1 & z_2 - z_1 & z_3 - z_1 & z_4 - z_1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$ , and the determinant of this matrix is

$\det \begin{pmatrix} x_2 - x_1 & x_3 - x_1 & x_4 - x_1 \\ y_2 - y_1 & y_3 - y_1 & y_4 - y_1 \\ z_2 - z_1 & z_3 - z_1 & z_4 - z_1 \end{pmatrix}$ . If we have the 4 points of the matrix  $p_1, p_2, p_3, p_4$ , then we can define vectors  $v_1 = p_2 - p_1$   $v_2 = p_3 - p_1$   $v_3 = p_4 - p_1$ , and because  $\det(v_1 \ v_2 \ v_3) = \text{vol-}$

ume of the parallelepiped shaped by the vectors, then  $\frac{1}{6}$  of that is the volume of the tetrahedron:

$$\frac{1}{6} \det \begin{pmatrix} x_2 - x_1 & x_3 - x_1 & x_4 - x_1 \\ y_2 - y_1 & y_3 - y_1 & y_4 - y_1 \\ z_2 - z_1 & z_3 - z_1 & z_4 - z_1 \end{pmatrix} = \frac{1}{6} \det \begin{pmatrix} v_1 & v_2 & v_3 \end{pmatrix}$$

6. We can have vectors  $\bar{AB} = \begin{bmatrix} x_2 - x_1 \\ y_2 - y_1 \\ z_2 - z_1 \end{bmatrix}$  and  $\bar{AC} = \begin{bmatrix} x_2 - x_0 \\ y_2 - y_0 \\ z_2 - z_0 \end{bmatrix}$ , and take the cross product to find the

normal vector that helps define the plane:  $\begin{bmatrix} (y_2 - y_0)(z_1 - z_0) - (z_2 - z_0)(y_1 - y_0) \\ (z_2 - z_0)(x_1 - x_0) - (x_2 - x_0)(z_1 - z_0) \\ (x_2 - x_0)(y_1 - y_0) - (y_2 - y_0)(x_1 - x_0) \end{bmatrix}$ .

We can then take this  $(a, b, c)$  and plug into  $f(x, y, z) = ax + by + cz = d$ , then plug in a point then solve for  $D$  to find the full equation of the plane.

7. If  $p_0, p_1, p_2$  are three distinct points in space then the cross product  $n = (p_0 - p_1) \times (p_0 - p_2)$ , then the resulting vector is orthogonal to the vectors  $(p_0 - p_1)$  and  $(p_0 - p_2)$ . If  $p$  is any point on the plane formed by those three points, then  $n$  and  $p - p_0$  are also orthogonal (dot product is 0). So because all points  $p$  in relation to  $p_0$  are orthogonal to  $n$ , then the equation of the plane must be of the form  $n \cdot (p - p_0) = 0$

8.  $M(\theta)^{-1} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^{-1} = \frac{1}{\cos^2 \theta + \sin^2 \theta} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = M(\theta)^T = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$   
so  $M(\theta)$  is orthonormal.

To show that  $M(\theta_1 + \theta_2) = M(\theta_1)M(\theta_2)$ , we consider that  $\sin(\theta_1 + \theta_2) = \sin(\theta_1)\cos(\theta_2) + \cos(\theta_1)\sin(\theta_2)$  and  $\cos(\theta_1 + \theta_2) = \cos(\theta_1)\cos(\theta_2) - \sin(\theta_1)\sin(\theta_2)$  and we can do the matrix multiplication to prove the identity.

If  $\theta_1 = \theta$  and  $\theta_2 = -\theta$ , so  $M(\theta_1 + \theta_2) = M(0) = I$ .  $M(\theta)M(-\theta) = I$ , so  $M(\theta)^{-1} = M(-\theta)$

For  $p = (0, 1)$ , considering  $M(\theta)p$  for every possible  $\theta$  gives you a circle ( $M(\theta)p = \sin \theta + \cos \theta$ , which is the equation for a circle).

9. We have 4 vectors in a plane:  $x_1, x_2, b_1, b_2$  and  $Mx_1 = b_1$  and  $Mx_2 = b_2$ . We need an orthogonal basis, so we define  $y_1 = x_1$  and  $y_2 = x_2 - \frac{x_2 \cdot x_1}{|x_1|} \cdot \frac{x_1}{|x_1|}$ . Then we can write

$$x_3 = ay_1 + by_2 \rightarrow Mx_3 = aMy_1 + bMy_2 = ab_1 + b(b_2 - \frac{x_2 \cdot x_1}{|x_1|} \cdot \frac{b_1}{|x_1|})$$

which gives us  $a = x_3 \cdot \hat{y}_1$  and  $b = x_3 \cdot \hat{y}_2$ , and we can plug these values in to get  $Mx_3 = (x_3 \cdot \hat{y}_1)Mx_1 + (x_3 \cdot \hat{y}_2)M(x_2 - \frac{x_2 \cdot x_1}{|x_1|} \cdot \frac{x_1}{|x_1|})$

10. Parametric eqn for line through  $p_1$  and  $p_2$ :

$$\begin{cases} x(t) = (1-t)x_1 + tx_2 \\ y(t) = (1-t)y_1 + ty_2 \end{cases}$$

For a plane with  $p_1, p_2, p_3$ :

$$\begin{cases} x(t_1, t_2) = (1-t_1-t_2)x_1 + t_1(1-t_2)x_2 + (1-t_1)t_2x_3 \\ y(t_1, t_2) = (1-t_1-t_2)y_1 + t_1(1-t_2)y_2 + (1-t_1)t_2y_3 \\ z(t_1, t_2) = (1-t_1-t_2)z_1 + t_1(1-t_2)z_2 + (1-t_1)t_2z_3 \end{cases}$$

The equalities for

- points on line between  $p_1, p_2$ :  $0 \leq t_1 \leq 1, \quad t_2 = 0$

- triangle between  $p_1, p_2, p_3$ :  $0 \leq t_1 + t_2 \leq 1$ ,  $0 \leq t_1 \leq 1$ ,  $0 \leq t_2 \leq 1$

11.  $\mathbf{M}$  is symmetric  $2 \times 2$  with eigenvalues  $\lambda_0$  and  $\lambda_1$ .

- Show  $0 \leq x^T Mx \leq (\max(\lambda_0, \lambda_1))x^T x$ :

$x = av_1 + bv_2$ , where  $v_1, v_2$  are eigenvalues corresponding to  $\lambda_{0,1}$ . We can then say

$$Mx = Mav_1 + Mbv_2 = \lambda_1 av_1 + \lambda_2 bv_2$$

$$x^T Mx = (av_1 + bv_2)(\lambda_1 av_1 + \lambda_2 bv_2) = \lambda_1 a^2 |v_1|^2 + \lambda_2 b^2 |v_2|^2$$

So, because  $x^T x = a^2 |v_1|^2 + b^2 |v_2|^2$ , then scaling both these terms by  $\max(\lambda_1, \lambda_2)$  will mean that the result will be **at least as big** as  $x^T Mx$ , in which one of the terms will be multiplied by the larger eigenvalue and one by the smaller (or equal).

- If one eigenvalue is 0, then the corresponding eigenvector may not be 0 by definition of an eigenvector. Thus, if  $\lambda_1 = 0$ , then  $Mx = \lambda_1 x = 0$  with a non-zero  $x$ .
- if  $\lambda_0 < 0$  and  $\lambda_1 > 0$  with corresponding eigenvalues  $v_0, v_1$ , then we can say:

$$Mx = Mav_0 + Mbv_1 = \lambda_0 av_0 + \lambda_1 bv_1$$

$$x^T Mx = \lambda_0 a^2 |v_0|^2 + \lambda_1 b^2 |v_1|^2 = 0, \text{ so}$$

$$-\lambda_0 a^2 |v_0|^2 = \lambda_1 b^2 |v_1|^2, \text{ so if } a = 1, \text{ then } b = \sqrt{\frac{-\lambda_0 |v_0|^2}{\lambda_1 |v_1|^2}}, \text{ which we can plug into}$$

$$x = av_0 + bv_1 \neq 0, \text{ but still satisfies } x^T Mx = 0.$$

- The eigenvalues are positive.

12.  $\mathbf{M}$  is a  $2 \times 2$  matrix

$$Mx = \lambda Ix \text{ (where } I \text{ is an identity matrix)}$$

$$Mx - \lambda Ix = 0 = (M - \lambda I)x, \text{ so } \det(M - \lambda I) = 0$$

$$\begin{bmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{bmatrix} = \lambda^2 - (a_{11} + a_{22})\lambda + (a_{11}a_{22} - a_{12}a_{21}) = \lambda^2 - \text{trace}(M)\lambda + \det(M)$$