




Hypothesis tests  
for a single  
proportion

# Overview


Review and continuation of hypothesis tests for a single proportion

# Basic logic of hypothesis tests

We start with a claim about a population parameter

- E.g.,  $\mu = 4$  

This claim implies we should get a certain distribution of statistics





If our observed statistic is highly unlikely, we reject the claim

# Five steps of hypothesis testing

## 1. State $H_0$ and $H_A$

- Assume Gorgias ( $H_0$ ) was right


$$= \sqrt{10.82}$$
$$s_d = 3.29$$




## 2. Calculate the actual observed statistic

## 3. Create a distribution of what statistics would look like if Gorgias is right

- Create the **null distribution** (that is consistent with  $H_0$ )

## 4. Get the probability we would get a statistic more than the observed statistic from the null distribution

- p-value



## 5. Make a judgement

- Assess whether the results are statistically significant




# Are dolphins capable of abstract communication?

Dr. Jarvis Bastian in the 1960's wanted to know whether dolphins are capable of abstract communication

He used an old headlight to communicate with two dolphins (Doris and Buzz)

- Steady light = push button on right to get food
- Flashing = push button on the left to get food

The two dolphins were then separated by a barrier



Buzz got 15 out of 16 trials correct

# Hypothesis testing in 5 easy steps!

## 1. State the null hypothesis... and the alternative hypothesis

- Buzz is just guessing so the results are due to chance:  $H_0: \pi = 0.5$
- Buzz is getting more correct results than expected by chance:  $H_A: \pi > 0.5$

## 2. Calculate the observed statistic

- Buzz got 15 out of 16 guesses correct, or  $\hat{p} = .973$

## 3. Create a null distribution that is consistent with the null hypothesis


- i.e., what statistics would we expect if Buzz was just guessing

## 4. Examine how likely the observed statistic is to come from the null distribution

- What is the probability that the dolphins would guess 15 or more correct?
- The p-value = 0.003

## 5. Make a judgement

- If we have a small p-value, this means that  $\pi = .5$  is unlikely and so  $\pi > .5$
- i.e., we say our results are 'statistically significant'



"Matthews ... we're getting another one of those strange 'aw blah es span yol' sounds."

# Do more than 25% of US adults believe in ghosts?

A telephone survey of 1000 randomly selected US adults found that 31% of them say they believe in ghosts. Does this provide evidence that more than 1 in 4 US adults believe in ghosts?



# Write down/discuss answers to the following questions

1. What are the cases here?
2. What is the variable of interest and is it categorical or quantitative?
3. What is the observed statistic - and what symbols should we use to denote it?
4. What is the population parameter we are trying to estimate - and what symbol should we use to denote it?
5. Do you think that more than 25% of US citizens believe in ghosts?

# 5 steps to null-hypothesis significance testing (NHST)

**Let's go through the 5 steps!**

1. State null and alternative hypotheses
2. Calculate statistic of interest
3. Create a null distribution
4. Calculate a p-value
5. Assess if there is convincing evidence to reject the null hypothesis

# Step 1: State the null and alternative hypotheses

**Null Hypothesis ( $H_0$ ):** Claim that there is no effect or no difference

**Alternative Hypothesis ( $H_a$ ):** Claim for which we seek significant evidence.

# Believing in ghosts study

1. What is the null hypothesis?
2. How would you write it in terms of the population parameter?

$$H_0: \pi = 0.25$$

3. What is the alternative hypothesis?

$$H_A: \pi > 0.25$$

## Step 2: Calculate statistic of interest

For the ghost study, what was the observed statistic?

31% percent of Americans believe in ghosts ( $\hat{p} = .31$ )



## Step 3: Create a null distribution

Q: What is the null distribution?

Answer: A distribution of statistics ( $\hat{p}$ 's) consistent with the null hypothesis ( $H_0: \pi = 0.25$ )

Q: How can we create a null distribution?

Answer: when making inferences on *proportions* we can simulate flipping coins

# Step 3: Create a null distribution

Please answer the following questions for the ghost study

1. How many coins should we flip?

- A: 1,000 people were surveyed, so we want to flip 1,000 coins to simulate each person's answer


2. What should the probability of heads be on each flip?

- A: Our null parameter is  $\pi = 0.25$  so to be consistent with the null hypothesis the probability of heads on each flip should be 0.25

3. How many simulations should we run?

- A: 10,000 simulations should be enough to give us a good sense of the statistics we would get if the null hypothesis was true

# Step 3: Create a null distribution



A null distribution ( $\hat{p}$ 's) based on:

- ***10,000 simulations***
- Each simulation consists of flipping 1,000 coins
- With the probability of getting a head on each flip of 0.25



## Step 4: Calculate a p-value


The **p-value** is the probability, when the null hypothesis is true, of obtaining a statistic as extreme or more extreme than the observed statistic

$$\Pr(\text{Stat} > \text{observed statistic} \mid H_0 = \text{True})$$

The smaller the p-value, the stronger the statistic evidence is against the null hypothesis and in favor of the alternative

## Step 4: Calculate a p-value

$$\hat{p} = 0.31$$



What is the p-value here?

- A: The p-value is close to 0

## Step 5a: Assess if results are statistically significant

When our observed sample statistic is unlikely to come from the null distribution, we say the sample results are **statistically significant**


- i.e., we have a small p-value

‘Statistically significant’ results mean we have convincing evidence against  $H_0$  in favor of  $H_a$

## Step 5b: Make a decision

Are the results seem statistically significant?





NORMAL DISTRIBUTION



PARANORMAL DISTRIBUTION

Let's try the whole ghost example in R...

```
obs_stat <- .31      # step 2: calculate the observed statistic
```

```
# step 3: generate the null distribution
```

```
null_distribution <- do_it(10000) * {  
    rflip_count(1000, prob = .25)/1000  
}
```

```
# step 4: calculate the p-value
```

```
p_value <- pnull(obs_stat, null_distribution, lower.tail = FALSE)
```

# The amazing woman who can smell Parkinson's disease



Joy Milne claimed to have the ability to smell whether someone had Parkinson's disease

To test this claim researchers gave Joy 6 shirts that had been worn by people who had Parkinson's disease and 6 people who did not

Joy identified 11 out of the 12 shirts correctly

# The amazing woman who can smell Parkinson's disease

**Work in pairs to complete the following steps to analyze the data**

1. State Null and Alternative in symbols and words
2. Calculate the observed statistic of interest (`obs_stat`)
3. Create a null distribution using:  

```
null_dist <- do_it(10000) * { rflip_count(num_flips = ... prob = ... ) }
```
4. Calculate a p-value by assessing the probability of getting a statistic as or more extreme than the observed statistic from the null distribution
  - `pnull(obs_stat, null_dist, lower.tail = ...)`
5. Make a decision about whether the results are statistically significant



# Paul the Octopus

In the 2010 World Cup, Paul the Octopus (in a German aquarium) became famous for correctly predicting 11 out of 13 soccer games.



Question: is Paul psychic?

Homework 5!