

The medfate reference book

true

2026-10-06

Contents

Preface	7
How to use this book	7
An collaborative project	7
Funding	9
History and acknowledgements	9
 I Preliminaries	 11
1 Introduction	13
1.1 Purpose	13
1.2 Installation	14
1.3 Main functions in medfate	15
1.4 Main functions in medfateland	17
 2 Model inputs	 21
2.1 Input structures	21
2.2 Latitude and topography	23
2.3 Soil description	24
2.4 Vegetation description	29
2.5 Metereological input	38
2.6 Simulation control	39
 II Basic water balance modelling	 43
3 Basic water balance model	45
3.1 Design principles	45
3.2 State variables	46
3.3 Water balance	46
3.4 Process scheduling	48
3.5 Inputs and outputs	49
3.6 Applications	54
 4 Leaf phenology and radiation extinction	 55
4.1 Leaf phenology	55
4.2 Radiation extinction	58
 5 Forest hydrology	 61
5.1 Snow and rainfall	61
5.2 Snow pack dynamics	61
5.3 Rainfall interception loss	62

5.4	Bare soil evaporation	65
5.5	Transpiration of the herbaceous layer	65
5.6	Infiltration	66
5.7	Soil water flows	68
6	Plant transpiration, photosynthesis and drought stress	77
6.1	Plant transpiration and photosynthesis	77
6.2	Plant drought stress and cavitation	82
III	Advanced water balance modelling	85
7	Advanced water balance model	87
7.1	About Sureau	87
7.2	Design principles	88
7.3	State variables	89
7.4	Water and energy balances	90
7.5	Process scheduling	91
7.6	Inputs and outputs	93
7.7	Applications	100
8	Sub-daily atmospheric temperature and radiation variation	101
8.1	Above-canopy air temperature	101
8.2	Incoming diffuse and direct short-wave radiation	101
8.3	Incoming long-wave radiation	101
9	Radiation transfer	103
9.1	Short-wave radiation	103
9.2	Long-wave radiation	108
10	Plant hydraulics	111
10.1	Hydraulic architecture	111
10.2	Vulnerability curves	112
10.3	Water content of plant tissues	118
10.4	Sperry's sub-model of plant hydraulics	120
10.5	Sureau's sub-model of plant hydraulics	131
11	Leaf/canopy photosynthesis	137
11.1	Leaf energy balance, gas exchange and photosynthesis	137
11.2	Crown/canopy photosynthesis	140
12	Stomatal regulation, transpiration and photosynthesis	143
12.1	Stomatal regulation, transpiration and photosynthesis under Sperry's sub-model	143
12.2	Stomatal regulation, transpiration and photosynthesis under Sureau's sub-model	153
12.3	Scaling of transpiration, soil water uptake and photosynthesis rates	156
12.4	Stomatal regulation and transpiration of hemi-parasitic plants	156
13	Closing energy balances	159
13.1	Single-layer canopy	159
13.2	Multi-layer canopy	161
14	Day-level results	165
14.1	Photosynthesis and transpiration	165
14.2	Plant water potentials and relative water contents	165
14.3	Plant drought stress	166

14.4 Cavitation and conduit refilling	166
14.5 Cavitation-induced defoliation and recovery	167

IV Forest growth modelling 169

15 Forest growth model 171

15.1 Design principles	171
15.2 State variables	183
15.3 Mass and energy balances	186
15.4 Process scheduling	190
15.5 Inputs and outputs	191

16 Plant carbon pools and components of the labile carbon balance 199

16.1 Size of plant carbon pools	199
16.2 Components of labile carbon balance	202

17 Growth, senescence and mortality 205

17.1 Growth	205
17.2 Senescence	209
17.3 Update of plant traits, structural variables and allocation targets	210
17.4 Plant mortality	214
17.5 Fire severity	220

18 Decomposition and soil biogeochemistry 223

18.1 Carbon pools, losses and fluxes	223
18.2 Decomposition rates	224
18.3 Decomposition process of leaf and fine root litter, coarse woody debris and standing dead wood	226
18.4 Decomposition in soil organic carbon pools	229

V Forest dynamics modelling 233

19 Forest dynamics model 235

19.1 Design principles	235
19.2 State variables	237
19.3 Process scheduling	237
19.4 Inputs and outputs	237
19.5 Applications	240

20 Regeneration 241

20.1 Seed production and seed bank dynamics	241
20.2 Recruitment from seeds	241
20.3 Resprouting	242

VI Landscape/regional modelling 243

21 Watershed hydrology 245

21.1 Design principles	245
21.2 State variables	246
21.3 Water balance	246
21.4 Process scheduling	247
21.5 Inputs and outputs	248
21.6 Process details	250

22 Seed dispersal	253
22.1 Design principles	253
22.2 Process scheduling	253
22.3 Process details	255
 VII Static modules	 257
23 Allometric models	259
23.1 Input data	259
23.2 Allometric relationships	260
24 Post-simulation procedures	263
24.1 Water use efficiency	263
25 Wind extinction	265
25.1 Wind speed at the top of the canopy	265
25.2 Wind extinction profile	265
26 Fuel characteristics and fire behaviour	267
26.1 Overview	267
26.2 Input data	267
26.3 Fuel characteristics	269
26.4 Surface fire behavior	276
26.5 Crown fire behavior	282
26.6 Fire potentials	283
26.7 Unit conversion of outputs	284
26.8 Applications	285
 A Inbuilt parameter estimation	 287
A.1 Introduction	287
A.2 Strict, scaled and imputable parameters	287
A.3 Imputation of missing values	292
A.4 Scaling size-related parameters	303
 B Symbols	 313
B.1 Dimensions	313
B.2 Soils	313
B.3 Plant/leaf classification	314
B.4 Vegetation	314
B.5 Meteorology	315
B.6 Anatomy	316
B.7 Radiation	316
B.8 Water balance	317
B.9 Energy balance	318
B.10 Forest hydrology	318
B.11 Plant hydraulics	318
B.12 Photosynthesis	319
B.13 Plant water content	320
B.14 Stomatal regulation	321
B.15 Carbon pools	321
B.16 Carbon balance	321
B.17 Growth, senescence and mortality	322

Preface

This is a reference book for the models implemented in **medfate** and **medfateland**, two R packages that provide functions for the simulation of functioning and dynamics of Mediterranean forests and landscapes.

How to use this book

This *reference book* is meant to help you understand the static and dynamic models included in packages **medfate** (ver. **5.2.0**) and **medfateland** (ver. **3.0.1**). Hands-on user guides to run simulation functions and static functions can be found in *articles* of the package website and training course materials can be found here. As any reference book, you are not expected to read the book linearly, but to jump to specific book sections whenever you have doubts about the design or implementation of certain processes or calculations.

The first part of the book presents the main functions of the package and describes the soil, vegetation and weather inputs that apply to all simulation models. After that, the book presents in detail each simulation model using a set of chapters. The first of those provides an overview of the model design and the remaining ones describe the details of formulations by groups of related processes. Process formulations that are common to more than one model are described only once, to avoid repeating material already presented, but links are provided whenever needed. Static models and their corresponding R functions are presented grouped at the end of the book, before the appendices, which include parameter estimation information and a list of all symbols for reference.

In this book we use **objectname** or **variablename** to refer to an R objects or a variable within data frames, and **functionname()** to refer to a package function. Whenever relevant, we indicate the correspondence between mathematical symbols, their units and the names used within the R package.

Our aim is to continuously update this reference book along with package developments, so that users have detailed and up-to-date information about the models at the time functions are run. Since the book evolves with the package, after any model application we recommend users to store a **PDF version** of the reference book to be sure it matches the version the package reported in their application report or article.

An collaborative project

You should look at **medfate**, **medfateland** and associated R packages as an ongoing research project for the development of forest ecosystem modelling tools. The models included in the package may contain bugs, and provide wrong predictions (all models are *wrong*, remember?). Package functions may evolve in the future, so that you may have to modify your scripts after package updates (hopefully not too often). Nevertheless, we feel the tool is mature enough to become useful for applications and that is why we made an effort to document it thoroughly.

The set of R packages are developed and maintained by the *Ecosystem Modelling Facility* unit at *CREAF* (in Spain), in close collaboration with researchers from *URFM-INRAE* (in France) and *CTFC* (in Spain):



At the individual level, the R packages are the result of collaborative work between modelers and experts in different disciplines. The following is a list of researchers that have contributed to the project with ideas, data or code:

- Jordi Martínez-Vilalta (CREAF-UAB, Spain)
- Maurizio Mencuccini (CREAF-ICREA, Spain)
- Juli G. Pausas (CIDE-CSIC, Spain)
- Pilar Llorens (CSIC, Spain)
- Rafa Poyatos (CREAF, Spain)
- Lluís Brotons (CREAF-CSIC, Spain)
- Antoine Cabon (WSL, Switzerland)
- Roberto Molowny (EMF-CREAF, Spain)
- Victor Granda (EMF-CREAF, Spain)
- Adriana Tovar (EMF-CREAF, Spain)
- Alicia Forner (MNCN-CSIC, Spain)
- Lluís Coll (UdL, Spain)
- Pere Casals (CTFC, Spain)
- Mario Beltrán (CTFC, Spain)
- Jean-Marc Limousin (CEFE, France)
- Aitor Améztegui (UdL, Spain)
- Nicolas Martin-StPaul (INRAE, France)
- Hervé Cochard (INRAE, France)
- Julien Ruffault (INRAE, France)
- François Pimont (INRAE, France)
- Shengli Huang (USDA, USA)
- Enric Batllori (UB-CREAF, Spain)
- Santi Sabaté (UB-CREAF, Spain)
- Daniel Nadal-Sala (UB, Spain)
- Maria González Sanchis (UVA, Spain)
- Daniel Caviedes-Voullième (Forschungszentrum Jülich, Germany)
- Mario Morales-Hernández (U Zaragoza, Spain)
- Mario Morales (UdZ, Spain)
- Raúl García Valdés (URJC, Spain)
- Josep M^a Espelta (CREAF, Spain)
- Francisco Lloret (CREAF, Spain)
- Teresa Valor (CTFC, Spain)

If your own expertise includes any of the processes that are modeled in the package and you feel that your expertise could be helpful to improve the package, you are more than welcome to contact us!

Funding

I have been intensively working on **medfate** since 2013, when I obtained a *Ramon y Cajal* research fellowship from the Spanish government (Reference: RYC-2012-11109; Amount: 308.600 €). An investment on a long-term modelling project could not have started with this initial financial help and that of the following competitive research projects:

- **Title:** Forecasting of ecosystem services on forests: impacts and adaptation to extreme climate events (FORESTCAST). **Financial Entity:** Ministerio de Economía y Competitividad (CGL2014-59742-C2-2-R). **Amount:** 152.460 €. **Duration from:** 01/01/2015 **to:** 31/12/2017. **PI:** Lluís Brotons/Miquel De Cáceres.
- **Title:** Modeling plant water use and drought resistance strategies across scales (DRESS). **Financial Entity:** Ministerio de Economía y Competitividad (CGL2017-89149-C2-2-R). **Amount:** 93.170 €. **Duration from:** 01/01/2018 **to:** 31/12/2021. **PI:** Miquel De Cáceres.
- **Title:** Boosting process-based models to project forest dynamics and associated ecosystem services at stand-to-regional scales (BOMFORES). **Financial Entity:** Ministerio de Ciencia e Innovación (PID2021-126679OB-I00). **Amount:** 75.020 €. **Duration from:** 01/09/2022 **to:** 31/08/2024. **PI:** Miquel De Cáceres.
- **Title:** Improving the modelling of key forest dynamic processes to forecast long-term changes in Mediterranean forests under climate change (IMPROMED). **Financial Entity:** Ministerio de Ciencia e Innovación (PID2023-152644NB-I00). **Duration from:** 01/09/2024 **to:** 31/08/2025. **PI:** Miquel De Cáceres/Josep M^a Espelta.

Besides this specific funding, this model developments could not have occurred without the institutional support of the Forest Science and Technology Center of Catalonia (CTFC), where I worked until March 2021, and Center for Ecological Research and Forestry Applications (CREAF), where I currently coordinate its Ecosystem Modelling Facility (EMF).

History and acknowledgements


The idea of an R package devoted to Mediterranean vegetation evolved after a previous attempt at modelling the dynamics of plant cohorts within a landscape model that tried to extend the MEDFIRE landscape model (Brotons et al., 2013) by incorporating drought effects and a simple representation of vegetation structure and composition. The word ‘MEDFATE’ originated from MEDiterranean vegetation and the ‘FATE’ model (Moore and Noble, 1990), which inspired me in its approach to model vegetation based on vital attributes. I made these initial efforts when I was still a post-doc at CTFC under the supervision of Lluís Brotons, to whom I am grateful for having greatly expanded my view of ecology and its applications. Undertaking this initial endeavor made us contact two excellent researchers in fire and drought ecology respectively, Juli G. Pausas (CIDE-CSIC, Spain) and Jordi Martínez-Vilalta (CREAF-UAB, Spain). While interacting with Juli was rewarding and fostered my interest in wildfires and fire ecology, Jordi had a profound influence on the project. Collaborating with him caused a slow but steady drift of my research interests towards mechanisms around soil water balance, plant hydraulics and physiological responses to drought. His influence on my work was not so apparent when we published water balance model (De Cáceres et al., 2015) and launched the R package, with the help of the forest hydrologist Pilar Llorens (CSIC, Spain) and other researchers (Rafa Poyatos, Lluís Coll and Pere Casals). However, it became undeniable when I started studying plant hydraulics, trying to understand and implement Sperry’s model in the package. Jordi made me interact with people in his research group and introduced me to Maurizio Mencuccini, another great ecophysiologicalist with whom he had been working for a long time. I have to thank Maurizio and Jordi for encouraging me to continue working on this development project and overcome the difficulty that entails learning scientific disciplines quite far from my previous experience. All this efforts crystallized in the advanced version of the model, which we presented in De Cáceres et al. (2021) as a tool to study species interactions.

The development of the package has benefited from many colleagues who have provided me with ideas or have struggled to use the models within it. Asaf Karavani was among the first package users besides me. Together, we used the package to predict soil moisture dynamics in forest plots where mushroom

production was being monitored. Antoine Cabon (CTFC, Spain) was also an early user of the package. I have enjoyed endless discussions with him about several processes and how to implement them. While he was the Ph.D. student and I one of his advisors, I felt we both learned together. Moreover, Antoine has contributed with some functions to the package (and I hope he will contribute more!). Another key person for the development of **medfate** was working with Victor Granda (Ecosystem Modelling Facility at CREAM, Spain), whose great programming skills and profound knowledge about R technologies has encouraged me to embrace new programming habits and skills. I owe the idea of separating packages **meteoland** and **medfate** to Roberto Molowny (Ecosystem Modelling Facility at CREAM, Spain) an excellent modeller who has also contributed to the design of dispersal in **medfateland**. More recently, collaboration with another ecophysiologicalist, Nicolas Martin-StPaul (INRA, France), has stimulated improvements both in **meteoland** and **medfate**. In fact, Nicolas has become another key person in my research, with whom I share the interest for one of the interfaces between fire and drought, namely fuel moisture dynamics. Thanks to him and other colleagues (Julien Ruffault and François Pimont), since ver. 3.0 **medfate** includes an additional sub-model of plant hydraulics, based on SurEau-ECOS. Collaboration with Shengli Huang (Forest Service, US) on adapting the package for the US Forest Inventory and Analysis (FIA) program has expanded the horizons of its application. Collaboration with Josep Maria Espelta (CREAF) and Francisco Lloret (UAB-CREAM) has been key to the design of regeneration processes. The soil hydrology processes in the package, as well as recent coupling developments would be not possible without María Gonzalez Sanchís (UPV, Spain), Daniel Caviedes-Voullième (Forschungszentrum Jülich, Germany) and Mario Morales-Hernández (U. Zaragoza, Spain). The package have also benefited from the feed back of other users, such as Francesco d'Adamo (CREAF, Spain), Vincenzo Saponaro (Free University of Bozen-Bolzano, Italy) and Paula Martín (CTFC, Spain).

I would not like to finish the acknowledgements without thanking my former colleagues at CTFC (located in the small town of Solsona, Catalonia, Spain), for their collaboration and support. Particularly, Lluís Coll, Aitor Ameztegui and Pere Casals have been continuously supporting my efforts and finding new model applications. Moreover, they have contributed to the knowledge within the package with their expertise in forest dynamics, forest management, soils and fire issues. I also thank Mario Beltrán and Assu Gil for teaching me the basics of forest management.

Finally, I would like to thank José Manuel, who takes care of me and has to endure my endless talking about the package and its models, to my family for being there at difficult times, including my three nephews (Aran, Jan and Nil) that constantly remind me the joy of life.



Part I

Preliminaries

Chapter 1

Introduction

This chapter provides an overview of the purpose of developing the **medfate** and **medfateland** R packages, their main simulation functions and expected applications.

1.1 Purpose

Being able to anticipate the impact of global change on forest function and dynamics is one of the major environmental challenges in contemporary societies. However, uncertainties in the mechanisms underpinning forest function and practical constraints in how to integrate available information still hinder the availability of robust and reliable predictive models. Despite the amount of knowledge accumulated about the function and dynamics of forest ecosystems and the plethora of observations and modelling tools already available, further efforts are necessary to achieve the challenge of integrating different global change drivers into simulation frameworks useful for research and applications.

1.1.1 Package **medfate**

The R package **medfate** has been designed as a platform to simulate the function and dynamics of individual forest stands at temporal scales from days to years. Climate is the main environmental driver forcing model predictions, and the ecological, biophysical, hydrological and physiological processes surrounding these are fundamental for the simulation models included in the package. In particular, the package allows the simulation of daily energy, water and carbon balances within forest stands. It also allows simulating the growth, mortality and regeneration processes of a set of plant cohorts competing for light and water within a forest stand. Finally, additional package functions allow relating the amount of plant biomass and the water status of plant live and dead tissues to fuel characteristics and, hence, fire hazard.

The version of the reference manual that you are reading is intended to reflect **version 5.2.0** of the package.

1.1.2 Package **medfateland**

The R package **medfateland** has been designed to extend the capabilities of **medfate** to a spatial context, most usually at the landscape or regional level. It allows running the stand-level models available in **medfate** on a set of spatial units (normally points or cells) within a target area. The package allows coordinating the dynamics of multiple stands via the evaluation of demand-based management scenarios. Additionally, the package allows considering lateral water transfer processes of forested watersheds. Hence, **medfateland** can also be used as a tool for eco-hydrological applications.

The version of the reference manual that you are reading is intended to reflect **version 3.0.1** of the package.

1.1.3 Companion packages

Packages **medfate** and **medfateland** are more easily used in conjunction with three other packages, called **meteoland**, **traits4models** and **forestables**. Package **meteoland** assists the estimation of weather variables over landscapes, whereas **traits4models** provides functions to assist the harmonization of plant trait data bases with the aim to create species parameter tables for simulation models, and **forestables** allows reading and harmonizing forest inventory data. Together the five R packages conform a modelling framework designed to help simulating the function and dynamics of forest ecosystems, with a particular focus on drought impacts under Mediterranean conditions (Fig. 1.1).




Figure 1.1: Set of packages conforming the medfate modelling framework. Black arrows indicate library requirements, whereas gray arrows indicate model parameter or data provision.

Although **meteoland** and **forestables** are part of the forest modelling framework, they can be used independently for many other purposes. Similarly, package **traits4models** can be used to generate parameter inputs for other models or frameworks.

1.2 Installation

1.2.1 Requirements

Both **medfate** and **medfateland** have a user interface in R, but models have been coded in **C++ language**, and the resulting simulation functions are linked to R via the **Rcpp** package. This means that whenever installation is done from sources you will need a C++ compiler (i.e. Rtools in Windows). Note that compilers will always be a requirement when installing packages from GitHub.

1.2.2 Installing medfate

Package **medfate** can be found at CRAN, where it is updated every few months. Hence, it can be installed as any other R package using:

```
install.packages("medfate")
```

Users can also download and install the latest stable versions GitHub as follows (required package **remotes** should be installed/updated first):

```
remotes::install_github("emf-creaf/medfate")
```

Among other dependencies, installing **medfate** requires package **meteoland**, as it links to some of its C++ functions. Note that some package dependencies are only *suggested*, so that they are not automatically installed when installing **medfate**.

1.2.3 Installing medfateland

Since both packages evolve together, installing **medfateland** normally requires an up-to-date version of **medfate**. Package **medfateland** can be found at CRAN, where it is updated every few months. Hence, it can be installed as any other R package using:

```
install.packages("medfateland")
```

Users can also download and install the latest stable versions GitHub as follows:

```
remotes::install_github("emf-creaf/medfateland")
```

Among other dependencies, installing **medfateland** requires packages **medfate** and **meteoland**, as it links to some of their C++ functions.

1.3 Main functions in medfate

1.3.1 Dynamic simulation functions

Dynamic simulation functions in **medfate** include three nested simulation levels (Fig. 1.2):

1. Forest water and energy balance can be studied for a given forest stand using function **spwb()** (soil-plant-water-balance). The same function is used to run three different models (**Granier**, **Sperry** or **Sureau**), that correspond to two levels of complexity (*basic* or *advanced*; see Fig. 1.2). The user should choose one model or another depending on the intended application and data availability. Water balance simulations include hydrological processes (rainfall interception, soil infiltration, percolation and evapotranspiration) and plant physiological processes related to transpiration (hydraulics, photosynthesis and stomatal regulation). Most processes are implemented at the daily scale, although some operate at subdaily time steps for *advanced* models. Function **spwb()** has a closely related function **spwb_day()** that allows single-day simulation.
2. Changes in primary (leaf area) and secondary (sapwood) growth are key to evaluate the influence of climatic conditions on plant and forest structure and function. Function **growth()** extends the previous models because it allows simulating carbon balance, tissue growth and mortality of a set of plant cohorts competing for light and water in a given forest stand. Carbon balance is performed at the plant level, at the cohort level (once mortality is included), and at the stand level, after accounting for heterotrophic respiration due to decomposition processes. Analogously to the simulation of water balance, **growth()** has a closely related function **growth_day()** that allows focusing on sub-daily processes (i.e. carbon balance).
3. Finally, function **fordyn()** complements growth and mortality processes with recruitment (from seeds) and resprouting, thus completing the minimum set of demographic processes needed to simulate Mediterranean forest dynamics. The function splits the period to be simulated by years and makes internal calls to **growth()** for the simulation of growth and mortality. **fordyn()** is suited to simulate the inter-annual variation in forest structure and composition, while accounting for the biophysical and physiological processes provided by the former models. In addition, it allows simulating human interventions (i.e. tree or shrub cutting or planting) on forest stands.



1.3.2 Sub-model functions

Many of the sub-models included in **medfate** are implemented as C++ functions and internally called by the simulation functions implementing top-level models. Most sub-models have, however, their corresponding R


Processes

User-level functions

Water balance (daily)



Carbon balance, growth/senescence and mortality (daily)



Regeneration and management (yearly)




Figure 1.2: Nested relationships between medfate simulation functions and processes included in each model

function to made them directly available to the user. This facilitates understanding the different sub-models and a more creative use of the package. However, most users will not normally use them in their simulation workflows. Sub-model functions are grouped by *subject*, which is included in the name of the function. The different sub-model functions are (by subject):

Group	Description
<code>biophysics_*</code>	Physics and biophysics
<code>carbon_*</code>	Biomass estimation and carbon balance
<code>decomposition_*</code>	Dynamics of litter and soil organic carbon pools
<code>fuel_*</code>	Fuel properties
<code>fire_*</code>	Fire behaviour and severity
<code>hydraulics_*</code>	Plant hydraulics
<code>hydrology_*</code>	Canopy and soil hydrology
<code>light_*</code>	Light extinction and absorption
<code>moisture_*</code>	Live tissue moisture
<code>mortality_*</code>	Plant mortality
<code>pheno_*</code>	Leaf phenology
<code>photo_*</code>	Leaf photosynthesis
<code>regeneration_*</code>	Regeneration from seed germination or resprouting
<code>root_*</code>	Root distribution and conductance calculations
<code>soil_*</code>	Soil hydraulics and thermodynamics
<code>transp_*</code>	Stomatal regulation, transpiration and photosynthesis
<code>wind_*</code>	Canopy turbulence

1.3.3 Static functions

Package **medfate** includes a number of functions to examine static properties of plants conforming forest stands, summary functions at the species- and stand- levels:

- `plant_*`: Cohort-level information (species name, id, leaf area, height, etc.).
- `species_*`: Cohort-level attributes aggregated by species (e.g. basal area, cover, density, etc.).
- `herb_*`: Attributes of the (aggregated) herbaceous layer (e.g. fuel density or leaf area index).
- `stand_*`: Stand-level attributes (e.g. stand basal area, fuel loading, etc.).

Vegetation functioning and dynamics have strong, but complex, effects on fire hazard. On one hand, growth and death of organs and individuals changes the amount of standing live and dead fuels, as well as downed dead fuels. On the other, day-to-day changes in soil and plant water content changes the physical properties of fuel, notably moisture content of live and dead fuels. Package **medfate** provides functions to estimate fuel properties and potential fire behaviour in forest stands. Specifically, `fuel_FCCS()` calculates fuel characteristics for a set of fuel strata; and a fire behaviour model is implemented in function `fire_FCCS()` to calculate the intensity of surface fire reaction and the rate of fire spread of surface fires assuming a steady-state fire. Fuel and fire behavior functions allow obtaining the following:

1. Fuel characteristics by stratum.
2. Surface fire behavior (i.e. reaction intensity, rate of spread, fireline intensity and flame length).
3. Crown fire behavior.
4. Fire potential ratings of surface fire behavior and crown fire behavior.

1.4 Main functions in medfateland

Package **medfateland** allows simulating forest functioning and dynamics on sets forests stands distributed across space, with or without spatial processes (Fig. 1.3).

When spatial processes are omitted, the package offers a set of simulation functions that are analogous to those of package **medfate** but allow processing multiple forest stands sequentially or in parallel:

- Forest water and energy balance can be studied for a set of forest stands using functions `spwb_spatial()` or `spwb_spatial_day()`.
- Growth and mortality of a set of plant cohorts competing for light and water can be simulated for a set of forest stands using functions `growth_spatial()` or `growth_spatial_day()`.
- Forest dynamics arising from competition for light and water can be simulated for a set of forest stands using function `fordyn_spatial()`.

The package offers a simulation function where the dynamics of forest stands are related by management decisions:

- Function `fordyn_scenario()` allows simulating `fordyn()` on forest stands, while applying a demand-based forest management scenario. Simulations with this function also include seed dispersal among forest stands.

Finally, there are three functions that are meant to simulate watersheds, including spatial processes:

- Functions `spwb_land()` and `growth_land()` include distributed hydrological models that allow simulating daily water balance or growth processes on the cells of a watershed while accounting for overland runoff, subsurface flow and groundwater flow between cells.
- Forest dynamics arising from competition for light and water can be simulated in a watershed using function `fordyn_land()`. Simulations with this function also include seed dispersal among grid cells.




Figure 1.3: Overview of simulation functions in medfateland and the processes available for each of them. Colored squares below each simulation function indicate the processes included

Chapter 2

Model inputs

This chapter describes the inputs required by simulation models of **medfate** and **medfateland** packages. First, we describe the R structures required for the two packages. Then, we provide the details for four different kinds of inputs:

- (1) *Vegetation existing in the forest stand.* Information of the living plants includes not only species composition and/or structure, but also functional parameters of a set of plant cohorts. Vegetation can include information about dead plant elements (e.g. snags, leaf litter, large wood) or soil carbon pools. Vegetation is considered static in **spwb()** calls, except for leaf area variation, but becomes dynamic in calls to **growth()** or **fordyn()**. Hence, in these functions the input vegetation describes the starting point.
- (2) *Information on the physical structure of soil.* Soil information is also required but, unlike vegetation, the soil physical structure and its hydraulic properties are considered static in the current version.
- (3) *Climatic forcing.* Daily weather series are always required, defining the temporal extent of the simulation and the temporal variation of environmental conditions in which physical and biological processes occur.
- (4) *Topographic information and geographic location.* Topography and latitude are also required, because internal calculations of some physical variables require them.

The last section (2.6) describes additional input parameters that are used to specify global options of simulation functions.

Since preparing inputs for simulations can be daunting for most users, package **medfateland** provides a set of utility functions aimed at facilitating the preparation of vegetation and soil inputs, whereas package **meteoland** provides functions to assess daily weather over landscapes.

2.1 Input structures

2.1.1 Input structures in medfate

Package **medfate** deals with simulations at stand level and, hence, the data structures defined in the package are meant to encapsulate information about a target forest stands and the properties of the soil where it grows:

- S3 class **soil** is a *data frame* that is initialized using a data frame of soil physical properties, but the table includes also soil hydraulic parameters and state variables. Soil objects are described in subsection 2.3.3.
- S3 class **forest** is a *list* of forest inventory data corresponding to a single forest stand. It mainly refers to tree and shrub measurements, but can include many other information such as herbaceous

vegetation, seedling bank, seed bank, tree snags or litter. Forest plot objects are described in subsection 2.4.2.

- S3 classes `spwbInput` and `growthInput` are lists containing all the vegetation parameters and state variables necessary to run water-balance simulations (function `spwb()` and alike) and forest growth simulations (function `growth()` and alike). Objects of classes `spwbInput` and `growthInput` are created using functions whose input includes a `forest` object and a `soil` object.

The figure below (Fig. 2.1) can be used to illustrate the workflow of soil/forest input creation for the main simulation functions in **medfate**. Note the different inputs in the case of `fordyn()`.




Figure 2.1: Schematic workflow of input creation for simulation functions in medfate

2.1.2 Input structures in medfateland

Spatial information for **medfateland** needs to be provided in `sf` objects (i.e. data frames with geometry, see package `sf`). For each forest stand to be simulated, these object must include a geometry, topographic features as well as a `forest` object and `soil` object, as illustrated in the figure below:




Figure 2.2: Relationship between data structures in medfate and medfateland

Non-forest (i.e. croplands or urban areas) patches may be also described, and then `forest` or `soil` objects may not be required. The actual columns required in a `sf` depend on the simulation function to be used, as described in the following table:

Column	Description	Use
<code>geom</code>	Spatial geometries (points, polygons, ...)	Always required
<code>id</code>	Stand identifiers	Required for <code>*_spatial()</code> , <code>*_spatial_day()</code> and <code>fordyn_scenario()</code>
<code>elevation</code>	Elevation values above sea level (in m)	Always required
<code>slope</code>	Slope values (in degrees)	Always required
<code>aspect</code>	Aspect values (in degrees)	Always required
<code>land_cover_type</code>	Land cover type: <i>wildland</i> , <i>agriculture</i> , <i>rock</i> , <i>artificial</i> or <i>water</i>	Required for <code>*_land()</code> , optional for <code>*_spatial()</code> and <code>*_spatial_day()</code>
<code>forest</code>	Objects of class <code>forest</code>	Always required
<code>soil</code>	Objects of class <code>soil</code>	Always required
<code>state</code>	Objects of class <code>spwbInput</code> or <code>growthInput</code>	Optional to recover the final state of a previous simulation
<code>meteo</code>	Weather data frames (see 2.5)	Optional
<code>crop_factor</code>	Crop evapo-transpiration factor	Only required for <i>agriculture</i> land cover type
<code>management_arguments</code>	Management arguments for each stand	Optional for <code>fordyn_spatial()</code>
<code>management_unit</code>	Management unit corresponding to each stand	Required for <code>fordyn_scenario()</code>
<code>represented_area</code>	Area represented by each stand, in hectares	Required for <code>fordyn_scenario()</code>
<code>depth_to_bedrock</code>	Depth to bedrock (mm).	Required for <code>*_land()</code>
<code>bedrock_conductivity</code>	Bedrock (saturated) conductivity (in $\text{m} \cdot \text{day}^{-1}$).	Required for <code>*_land()</code>
<code>bedrock_porosity</code>	Bedrock porosity ([0-1]).	Required for <code>*_land()</code>
<code>snowpack</code>	Snow water equivalent content of the snowpack (mm).	Required for <code>*_land()</code>
<code>aquifer</code>	Water content of the aquifer in each cell.	Required for <code>*_land()</code>
<code>deep_aquifer_loss</code>	A numeric vector with the maximum daily loss to a deeper aquifer (in $\text{mm} \cdot \text{day}^{-1}$). If missing all cells take their value from watershed control.	Optional for <code>*_land()</code>
<code>channel</code>	A logical (or binary) vector indicating overland channel routing.	Optional for <code>*_land()</code>
<code>outlet_backlog</code>	A list vector indicating channel backlog of outlet cells from a previous simulation.	Optional for <code>*_land()</code>

2.2 Latitude and topography

Topography has a relatively small direct influence in **medfate** simulations, because most topographic effects are assumed to be taken into account in the estimation of weather inputs. Nonetheless, *elevation* (in meters) is needed to estimate atmospheric pressure and air density (see 2.5), whereas *aspect* (in degrees from North) and *slope* (also in degrees) are needed in addition to *latitude* (also in degrees North) to estimate sunrise/sunset hours and potential radiation. Topographic variables should be readily accessible for users having access to a digital elevation model of the target area.

2.3 Soil description

Soils are described in **medfate** using $S > 1$ discrete soil layers ($s \in \{1, 2, \dots, S\}$). Here the word *soil* refers to the volume comprised between forest floor and the depth that plant rooting systems can reach (i.e., the critical zone), including cracks within the bedrock. The number and size of soil layers may correspond to changes in soil properties along depth, but also can be chosen to reflect differences in plant rooting depths. The physical properties of the soil are needed to estimate soil hydraulic properties and are considered **static soil parameters** in simulations. In contrast, soil moisture content, other variables that depend on moisture (e.g. soil water potential and soil conductivity) and soil temperature, if the soil energy balance is considered, are all **soil state variables**.

2.3.1 Soil physico-chemical properties

Soil physico-chemical properties, such as *texture* (i.e. volume percent of sand, silt and clay), *organic matter content*, *bulk density* or *rock fragment content*, can differ between soil layers, and this has important consequences for soil water retention capacity and soil hydraulics. Specifying layers with an elevated rock fragment content (e.g. layers of weathered rock where soil particles are scarce) may be important in arid or seasonally-dry climates like the Mediterranean, because plants often extend their roots into cracks existing in the parent rock to access water during summer (Ruffault et al., 2013).

For each soil layer s , the following physico-chemical parameters are accepted (some of them are only optional):

Symbol	Units	R	Description	Required
d_s	mm	widths	Width of soil layer s	Yes
$P_{clay,s}$	%	clay	Percent volume of clay (within soil particles fraction) in layer s	Yes
$P_{sand,s}$	%	sand	Percent volume of sand (within soil particles fraction) in layer s	Yes
OM_s	%	om	Percentage of organic matter per dry weight (can be missing) in layer s	No
N_s	g/kg	nitrogen	Sum of total nitrogen (ammonia, organic and reduced nitrogen) in layer s	No
pH_s	[0-14]	ph	pH in water of layer s	No
BD_s	$g \cdot cm^{-3}$	bd	Bulk density in layer s	Yes
$P_{rocks,s}$	%	rfc	Rock fragment content as percent volume (within whole soil) in layer s	Yes

The percent volume of silt is 100% minus the percent volume of clay and sand. Whenever possible, soil physico-chemical properties should be measured in soil profiles conducted at the target forest plot (although soil profiles rarely reach rooting depths of deeply-rooted plants). Soil input data should be arranged in a `data.frame` with soil layers in rows and physico-chemical variables in columns (see function `defaultSoilParams()`). The following is an example with four soil layers:

```
soildf <- defaultSoilParams()
soildf
```

```
##   widths clay sand om nitrogen ph   bd rfc
## 1    300   25   25 NA        NA NA  1.5  25
## 2    700   25   25 NA        NA NA  1.5  45
## 3   1000   25   25 NA        NA NA  1.5  75
## 4   2000   25   25 NA        NA NA  1.5  95
```

Package **medfateland** includes a function `add_soilgrids()` to fetch soil information from SoilGrids.org, a global soil database currently providing soil data at 250m scale. This can be helpful to users lacking local

soil measurements, but the uncertainty of SoilGrids estimates can be high for some areas and soil properties, especially soil depth and rock fragment content.

The depth of a given soil layer s (Z_s ; in mm) is defined as the sum of layer widths from the surface to the target layer:

$$Z_s = \sum_{i=1}^s d_s \quad (2.1)$$

And the overall *soil depth* (Z_{soil} ; in mm) is the sum of widths across all soil layers. As the soil may include deep rocky layers, the value of Z_{soil} may be larger than the soil depth reported in soil profiles.

2.3.2 Water retention curves

The **water retention curve** of a soil is the relationship between *volumetric soil moisture content* (θ , in $m^3 \cdot m^{-3}$ of soil, excluding rock fragments) and the corresponding *soil water potential* (Ψ , in MPa), i.e. the amount of work that must be done per unit quantity of pure water in order to transport reversibly and isothermally an infinitesimal quantity of water from a reference pool. The shape of the water retention curve (also called the *soil moisture characteristic curve*) depends on physical properties (mainly texture and bulk density, but also organic matter content). Since soil layers usually differ in their physical properties they also normally have different water retention curves.

Two water retention curve models are available in **medfate** (fig. 2.3):

1. *Saxton model*: In this model, volumetric soil moisture $\theta(\Psi)$ corresponding to a given water potential Ψ (in MPa) below field capacity (i.e., $\Psi < \Psi_{fc}$) is calculated using:

$$\theta(\Psi) = (\Psi/A)^{(1/B)} \quad (2.2)$$

where A and B depend on the texture and, if available, organic matter in the soil. If organic matter is available, A and B are calculated from P_{clay} , P_{sand} and OM following Saxton and Rawls (2006). Otherwise, they are calculated from P_{clay} and P_{sand} as indicated in Saxton et al. (1986). Volumetric changes between field capacity and saturation (i.e., $\Psi_{fc} \leq \Psi < 0$) are estimated using a linear function.

2. *Van Genuchten model*: The well known van Genuchten (1980) model is:

$$\theta(\Psi) = \theta_{res} + \frac{\theta_{sat} - \theta_{res}}{[1 + (\alpha \cdot \Psi)^n]^{1-1/n}} \quad (2.3)$$

where $\theta(\psi)$ is the water retention, θ_{sat} is the saturated water content, θ_{res} is the residual water content, α is related to the inverse of the air entry pressure, and n is a measure of the pore-size distribution.

Hydraulic conductivity governs the rate of water flow for a unit gradient in potential. It decreases sharply as soil becomes dry, and the relationship also depends on soil texture. The Saxton and Van Genuchten models also describe and parameterize the relationship between conductivity and soil water potential.

2.3.3 Soil initialization

Soil initialization is needed to estimate hydraulic parameters for each soil layer s from its physico-chemical attributes. Soil initialization can be done using function `soil()`, which returns a data frame of a S3 class with the same name (see 2.1.1), but most often is performed implicitly when calling functions `spwbInput()` or `growthInput()` with a data frame of physical properties.

Soil initialization adds the following information to the physico-chemical soil description:

Symbol	Units	R	Description
P_{macro}	%	macro	Percentage of macroporosity corresponding to each soil layers

Symbol	Units	R	Description
K_{sat}	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	Ksat	Saturated soil conductivity for each soil layer
α	MPa^{-1}	VG_alpha	Parameter α of the Van Genuchten (1980) model for each soil layer
n		VG_n	Parameter n of the Van Genuchten (1980) model for each soil layer
θ_{sat}	$m^3 \cdot m^{-3}$	VG_theta_sat	Volumetric soil moisture at saturation for each soil layer
θ_{res}	$m^3 \cdot m^{-3}$	VG_theta_res	Residual volumetric soil moisture for each soil layer

Macroporosity values are calculated using the equations given in Stolf et al. (2011), while γ_{soil} is derived from texture. Parameters of the van Genuchten model for each layer are estimated from the physical description of the layer using one of two pedotransfer functions (see help for `soil()`):

1. Using the USDA texture classification and the average texture class parameters given by Carsel and Parrish (1988).
2. Directly from the soil texture, organic matter and bulk density, using the pedotransfer functions in Tóth et al. (2015).

Parameter α of the Van Genuchten model is given in MPa^{-1} , but it can be transformed to cm^{-1} using the relationship:

$$1 \text{ cm} = 0.00009804139432 \text{ MPa}, \quad (2.4)$$

which can also be applied to water potential units. Soil layer saturated conductivity (K_{sat}) is estimated using Saxton and Rawls (2006) or Saxton et al. (1986), depending on whether an estimate of organic matter is available. K_{sat} is given in physiological units $mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$, but they can be transformed into $cm \cdot day^{-1}$ using:

$$1 \text{ cm} \cdot day^{-1} = 655.2934 \text{ mmol} \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1} \quad (2.5)$$

Saturated conductivity (K_{sat}) is corrected for bulk density (BD) of the soil layer, because compact soils have lower conductivity (Rahimi et al., 2011):

$$K_{sat} = K_{sat,ref} \cdot \left(\frac{\rho_s - BD}{\rho_s - BD_{ref}} \right)^3 \quad (2.6)$$

where $K_{sat,ref}$ is the reference conductivity according to soil texture, $\rho_s = 2.73 \text{ g} \cdot cm^{-3}$ is density of soil particles, $BD_{ref} = 1.2 \text{ g} \cdot cm^{-3}$ is the reference bulk density.

Below is an example of soil initialization:

```
soilinit <- soil(soildf)
soilinit
```

```
## widths sand clay usda om nitrogen ph bd rfc macro Ksat VG_alpha
## 1 300 25 25 Silt loam NA NA NA 1.5 25 0.0485 5401.471 89.16112
## 2 700 25 25 Silt loam NA NA NA 1.5 45 0.0485 5401.471 89.16112
## 3 1000 25 25 Silt loam NA NA NA 1.5 75 0.0485 5401.471 89.16112
## 4 2000 25 25 Silt loam NA NA NA 1.5 95 0.0485 5401.471 89.16112
## VG_n VG_theta_res VG_theta_sat W Temp
## 1 1.303861 0.041 0.423715 1 NA
## 2 1.303861 0.041 0.423715 1 NA
## 3 1.303861 0.041 0.423715 1 NA
## 4 1.303861 0.041 0.423715 1 NA
```




Figure 2.3: Water retention curves under the Saxton and Van Genuchten models, for the same physical attributes (25% sand, 25% clay, 50% silt and bulk density of $1.5g \cdot cm^{-3}$). Parameters of the Van Genuchten curve have been estimated using pedotransfer functions given in Tóth et al. (2015).

Besides defining soil parameters, function `soil()` also initializes *soil state variables* for each soil layer:

Symbol	Units	R	Description	Initial value
$\theta(\Psi_s)/\theta(\Psi_{fc,s})$		W	Soil moisture content with respect to field capacity for layer s	1
$T_{soil,s}$	Celsius	Temp	Soil temperature for layer s	NA

Users can edit the `soil` object manually (it is actually a `list`) before simulations, for example to provide specific parameters of the Van Genuchten retention curve calibrated from soil measurements or to modify the initial values of state variables.

2.3.4 Water content and depth to saturation

The *volume of water* (V_s in $mm = l \cdot m^{-2}$ of ground area) in a soil layer s is calculated from its water potential (Ψ_s) and water retention curve using:

$$V_s(\Psi_s) = d_s \cdot ((100 - P_{rocks,s})/100) \cdot \theta_s(\Psi_s) \quad (2.7)$$

where d_s is the depth of the soil layer (in mm) and $P_{rocks,s}$ is the percentage of rock fragments. The overall volume of water in the soil (accessible by roots) is simply the sum of water content over soil layers:

$$V_{soil} = \sum_{s=1}^S V_s(\Psi_s) \quad (2.8)$$

A number of fixed water volumes (and their corresponding moisture) are important to remember:

- *Water holding capacity* ($V_{fc,s}$, in mm) of the soil layer s is defined as the volume of water at *field capacity*, i.e. at $\Psi_{fc} = -0.033$ MPa, the amount of water held in the soil after gravitational water has drained away:

$$V_{fc,s} = V_s(-0.033) = d_s \cdot ((100 - P_{rocks,s})/100) \cdot \theta_{fc,s} \quad (2.9)$$

where $\theta_{fc,s} = \theta_s(-0.033)$ is the corresponding moisture at field capacity.

- *Water content at saturation* ($V_{sat,s}$, in mm) is calculated analogously, but replacing $\theta_{fc,s}$ by $\theta_{sat,s} = \theta_s(0)$, the moisture at *saturation*, the amount of soil moisture when all easily drained spaces between soil particles (i.e. macropores) are also filled with water.
- *Water content at wilting point* ($V_{wp,s}$, in mm) is calculated replacing $\theta_{fc,s}$ by $\theta_{wp,s} = \theta_s(-1.5)$, the moisture at *wilting point*, i.e. at $\Psi_{wp} = -1.5$ MPa, a conventional amount of soil moisture beyond which plants are assumed have problems extracting water and start to wilt. However, plants can often extract water from much drier soils.
- The *amount of extractable water* (in mm) is the difference between water content at field capacity and the water content at a conventional minimum water potential, which can be set at wilting point or lower values (by default at -5 MPa in **medfate**).

The *saturated water depth* (Z_{sat} , in mm) is not defined (i.e. missing) when all soil layers are below field capacity. When some layers are between field capacity and saturation, depth of the saturated water is calculated as:

$$Z_{sat} = \sum_s d_s \cdot \min [1, (\theta_{sat,s} - \theta(\Psi_s)) / (\theta_{sat,s} - \theta_{fc,s})] \quad (2.10)$$

Function `summary.soil()` can be used to describe the properties and water content of the different soil layers, after initialization:

```
summary(soilinit)
```

```
## Soil depth (mm): 4000
##
## Layer 1 [ 0 to 300 mm ]
##   clay (%): 25 silt (%): 50 sand (%): 25 usda: Silt loam
##   organic matter (%): NA nitrogen (g/kg): NA ph (0-14): NA
##   rock fragment content (%): 25 macroporosity (%): 5
##   Theta WP (%): 14 Theta FC (%): 30 Theta SAT (%): 49 Theta current (%) 30
##   Vol. WP (mm): 32 Vol. FC (mm): 68 Vol. SAT (mm): 111 Vol. current (mm): 68
##   Temperature (Celsius): NA
##
## Layer 2 [ 300 to 1000 mm ]
##   clay (%): 25 silt (%): 50 sand (%): 25 usda: Silt loam
##   organic matter (%): NA nitrogen (g/kg): NA ph (0-14): NA
##   rock fragment content (%): 45 macroporosity (%): 5
##   Theta WP (%): 14 Theta FC (%): 30 Theta SAT (%): 49 Theta current (%) 30
##   Vol. WP (mm): 55 Vol. FC (mm): 117 Vol. SAT (mm): 190 Vol. current (mm): 117
##   Temperature (Celsius): NA
##
## Layer 3 [ 1000 to 2000 mm ]
##   clay (%): 25 silt (%): 50 sand (%): 25 usda: Silt loam
##   organic matter (%): NA nitrogen (g/kg): NA ph (0-14): NA
##   rock fragment content (%): 75 macroporosity (%): 5
##   Theta WP (%): 14 Theta FC (%): 30 Theta SAT (%): 49 Theta current (%) 30
##   Vol. WP (mm): 36 Vol. FC (mm): 76 Vol. SAT (mm): 123 Vol. current (mm): 76
##   Temperature (Celsius): NA
##
## Layer 4 [ 2000 to 4000 mm ]
##   clay (%): 25 silt (%): 50 sand (%): 25 usda: Silt loam
```

```
##      organic matter (%): NA nitrogen (g/kg): NA ph (0-14): NA
##      rock fragment content (%): 95 macroporosity (%): 5
##      Theta WP (%): 14 Theta FC (%): 30 Theta SAT (%): 49 Theta current (%) 30
##      Vol. WP (mm): 14 Vol. FC (mm): 30 Vol. SAT (mm): 49 Vol. current (mm): 30
##      Temperature (Celsius): NA
##
## Total soil saturated capacity (mm): 473
## Total soil water holding capacity (mm): 291
## Total soil extractable water (mm): 183
## Total soil current Volume (mm): 291
## Saturated water depth (mm): NA
```

2.4 Vegetation description

The representation of vegetation in a target forest stand is done primarily by means of **cohorts of living plants** (i.e. trees, shrubs or herbs). This representation was chosen so that package functions can be easily applied to **forest inventory data**, for example from national forest surveys. Note that one limitation of the representation of vegetation in **medfate** is that it is not spatially-explicit (i.e. plants cannot have explicit coordinates within forest stands) and hence spatial (horizontal) interactions between plant cannot be taken into account explicitly (but see 2.4.3.5).

Apart from its living plant cohorts, vegetation description can also include other elements, such as **cohorts of standing dead wood** (i.e. snags), or **leaf/branch/root litter** lying on the ground or found below the surface, or even the **seedling and seed banks**. All vegetation elements are grouped into a single structure, the **forest** object, which is described in section 2.4.2.

2.4.1 Cohorts of living plants

A living plant cohort represents a set of plants that belong to the same species (or functional group) and are more or less of the same *size*. For example, trees in a stand may be grouped by species and diameter class, whereas shrubs or herbs may be grouped by species and height. In practice, however, plant cohorts are entities that are treated separately when modeling the forest stand. The description of plant cohorts includes a measure of *abundance* of the cohort in the stand, which in the case of trees is *density* as stems per hectare and for shrubs or herbs is *percent cover*. The difference in the kind of measurements made for abundance is the main motivation for distinguishing between tree cohorts and shrub/herbs cohorts, rather than their size or growth form. Herbaceous cohorts are different from shrub cohorts for functional reasons. While shrubs are woody (often multi-stemmed) plants with secondary growth, herbs are non-woody, lacking secondary growth and with crowns starting at the ground level. Although herbs can be described as plant cohorts, similar to shrubs, the description of vegetation in the forest stand may include the cover and mean height of a *herbaceous layer* defined collectively, for practical reasons.

2.4.1.1 Structural attributes of living plant cohorts

A plant cohort i is either a *tree cohort*, a *shrub cohort* or a *herb cohort*; and is formally defined using a set of structural attributes shown in the following table, where columns **spwb**, **growth** and **fordyn** indicate whether attributes are required by simulation functions **spwb()**, **growth()** and **fordyn()**, respectively:

Symbol	Units	R	Description	spwb	growth	fordyn
<i>SP</i>		Species	Species identity	Y	Y	Y
<i>H</i>	<i>cm</i>	Height	Average plant height	Y	Y	Y
<i>CR</i>		CR	Crown ratio (i.e. ratio between crown length and plant height)	Y	Y	N
<i>N</i>	<i>ind·ha⁻¹</i>	N	Density of individuals	N	Y	Y
<i>DBH</i>	<i>cm</i>	DBH	Tree diameter at breast height	N	Y	Y

Symbol	Units	R	Description	spwb	growth	fordyn
<i>Cover</i>	%	Cover	Projected percent cover	N	Y	Y
<i>LAI_{live}</i>	$m^2 \cdot m^{-2}$	LAI_{live}	(Maximum) leaf area index	Y	Y	N
<i>LAI_{dead}</i>	$m^2 \cdot m^{-2}$	LAI_{dead}	(Standing) dead leaf area index	Y	Y	N
<i>LAI_φ</i>	$m^2 \cdot m^{-2}$	LAI_{expanded}	expanded leaf area index	Y	Y	N
<i>LAI_{NC}</i>	$m^2 \cdot m^{-2}$	LAI_{noComp}	leaf area index assuming no competition with other plants	N	Y	N
<i>Z₅₀</i>	mm	Z50	Depth above which 50% of the fine root mass is located	Y	Y	Y
<i>Z₉₅</i>	mm	Z95	Depth above which 95% of the fine root mass is located	Y	Y	Y

Height (H) values refer to average height of individuals included in the cohort, and the same for crown ratio (CR) and diameter at breast height (DBH). While plant size (i.e. height or diameter) is relatively easy to tally, other measurements are not usually made in the field. Package **medfate** includes utility functions that provide estimates CR and LAI from forest inventory data (e.g. heights, DBH and density for measured trees), using allometric relationships calibrated for Catalonia, Spain (see chapter 23). As mentioned above, herbaceous cohorts are assumed to have $CR = 1$.

2.4.1.2 Collective herbaceous layer

As stated above, the herbaceous layer of the stand can be described collectively, without distinguishing among plant cohorts. Two attributes, cover and height, are required in this case:

Symbol	Units	R	Description
<i>H_{herb}</i>	cm	herbHeight	Average height of herbaceous vegetation
<i>Cover_{herb}</i>	%	herbCover	Percent cover of herbaceous vegetation
<i>LAI_{herb}</i>	$m^2 \cdot m^{-2}$	LAI_{herb}	Leaf area index of herbaceous vegetation

Leaf area index of herbaceous layer (LAI_{herb}), is normally estimated from H_{herb} and $Cover_{herb}$, but could be directly prescribed by the user. Rooting depth parameters are not needed for herbaceous vegetation, which is assumed to extract water from the topmost soil layer. Note that neither phenology or growth/senescence processes are modelled when herbaceous vegetation described in this way.

2.4.2 Forest objects

Forest data input in the **medfate** package is facilitated through the adoption of a data structure that follows closely forest inventory plot descriptions. Each forest stand is represented in an object of class **forest**, a *list* that contains several elements. Here we display an example of forest stand provided with the package:

```
data(exampleforest)
exampleforest

## $treeData
##      Species  DBH Height    N Z50  Z95
## 1 Pinus halepensis 37.55   800 168 100  300
## 2   Quercus ilex 14.60   660 384 300 1000
##
## $shrubData
##      Species Height Cover Z50  Z95
## 1 Quercus coccifera    80  3.75 200 1000
##
```

```
## attr("class")
## [1] "forest" "list"
```

The most important items of **forest** are the two data frames describing woody plants: **treeData** (for trees) and **shrubData** for shrubs. While not required, an additional **herbData** item can be used to describe the herbs, with the same structure as for shrubs. With the aim to help users in the task of constructing **forest** objects, package **medfate** includes functions to map user data in into tables **treeData** and **shrubData**.

forest objects are a convenient format to start calculations with **medfate**, because there are many static functions that take forest objects as input. For example, a **summary.forest()** function provides the basal area, density, cover and leaf area index of the forest stand, and its different components:

```
summary(exampleforest, SpParamsMED)
```

```
## Tree BA (m2/ha): 25.0333016  adult trees: 25.0333016  saplings: 0
## Density (ind/ha) adult trees: 552  saplings: 0  shrubs (estimated): 749.4923076
## Cover (%) adult trees: 100  saplings: 0  shrubs: 3.75  herbs: 0
## LAI (m2/m2) total: 1.4977163  adult trees: 1.4449762  saplings: 0  shrubs: 0.0527401  herbs: 0  mist:
## Fuel loading (kg/m2) total: 0.5423816  adult trees: 0.5255004  saplings: 0  shrubs: 0.0168812  herbs:
## PAR ground (%): 45.7239003  SWR ground (%): 56.0085428
```

2.4.2.1 Single-cohort forests

Although **medfate** has been designed to perform simulations on multi-cohort forests, it can also handle simulations where vegetation is described using a single cohort. Functions **tree2forest()** and **shrub2forest()** allow defining single-cohort forests from attributes. For example a holm oak (*Quercus ilex*) forest of 4-m height and having a leaf area index of $2\text{ m}^2 \cdot \text{m}^{-2}$ can be defined using:

```
oak_forest <-tree2forest("Quercus ilex", Height= 400, LAI = 2)
```

The function will return a **forest** object where most attributes are empty:

```
oak_forest
```

```
## $treeData
##      Species DBH Height  N Z50 Z95 LAI
## 1 Quercus ilex  NA   400 NA  NA  NA   2
##
## $shrubData
## [1] Species Height  Cover   Z50     Z95
## <0 files> (o «row.names» de longitud 0)
##
## attr("class")
## [1] "forest" "list"
```

Nevertheless, the object is well defined, for example we can summarize it using:

```
summary(oak_forest, SpParamsMED)
```

```
## Tree BA (m2/ha): 0  adult trees: 0  saplings: 0
## Density (ind/ha) adult trees: 0  saplings: 0  shrubs (estimated): 0
## Cover (%) adult trees: 0  saplings: 0  shrubs: 0  herbs: 0
## LAI (m2/m2) total: 2  adult trees: 0  saplings: 0  shrubs: 0  herbs: 0  mistletoe: 0
## Fuel loading (kg/m2) total: 0.6473467  adult trees: 0  saplings: 0  shrubs: 0  herbs: 0
## PAR ground (%): 33.2871084  SWR ground (%): 44.2721305
```

Since density and diameter have not been provided, simulations in this case will be restricted to function **spwb()**. Moreover, note that when defining single-cohort forests all possible interactions with functionally distinct plants are neglected.

2.4.2.2 Optional variables for living plant cohorts in forest objects

The structure presented above for **forest** objects corresponds to the required data elements. A forest object can contain additional information when this is available. In particular, data frames **treeData**, **shrubData** and **herbData** can contain additional columns:

- **LAI**: Leaf area index ($m^2 \cdot m^{-2}$), corresponding to LAI_{live} , avoiding the use of allometric relationships for leaf biomass in water balance simulations.
- **LAIMistletoe**: Leaf area index ($m^2 \cdot m^{-2}$) of hemi-parasitic (mistletoe) plants.
- **FoliarBiomass**: Standing dry biomass of leaves ($kg \cdot m^{-2}$), which together with SLA will be used to estimate LAI_{live} , avoiding the use of allometric relationships for leaf biomass in water balance simulations.
- **FuelLoading**: Fine fuel loading ($kg \cdot m^{-2}$), to avoid its calculation from allometric relationships.
- **CrownRatio**: The ratio between crown length and total height (between 0 and 1), avoiding the use of allometric relationships for crown ratio in water balance simulations.
- **Z100**: Depth (in *mm*) corresponding to 100% of fine roots (to specify a truncated root distribution).
- **Age**: A numeric vector indicating age of cohorts in years. Used to track the age of plant cohorts in simulations of forest dynamics.
- **ObsID**: A string identifying plant cohorts at the stage of forest sampling. Used to track the fate of particular plant cohorts in simulations of forest dynamics.

2.4.2.3 Seedling and seed banks

Forest objects can include the description of the seed bank and the seedling bank in forest stands, which are used to model regeneration from seeds in simulations with function **fordyn()** (see chapter 20).

The optional data frame **seedBank** contains information about the **seed bank**, with the following columns:

R column	Description
Species	String with species (taxon) name
Percent	Amount of seeds in relation to a full seed bank (%)

Similarly, the optional data frame **seedlingBank** contains information about the **seedling bank**, with the following columns:

R column	Description
Species	String with species (taxon) name
Percent	Amount of seedling in relation to a full seedling bank (%)
Age	The age of seedlings/saplings in years
Z50	Depth (in <i>mm</i>) corresponding to 50% of fine roots
Z95	Depth (in <i>mm</i>) corresponding to 95% of fine roots.

2.4.2.4 Dead plant elements

Forest objects can include the description of dead vegetation elements of forest stands. Dead elements are grouped into *standing dead wood* (e.g. tree snags), *downed litter* (e.g. branches, logs or leaf litter) and *belowground litter* (i.e. fine roots or coarse roots). They are used to model organic matter decomposition of standing or downed plant organs (see chapter 18) in simulations with functions **growth()** or **fordyn()**.

The optional data frame **snagData** contains information about the **standing dead wood** of dead plant cohorts, with the following columns:

R column	Description
Species	String with species (taxon) name
DBH	Diameter at breast height (in cm) of the original living cohort, for tree snags only
Height	Height of the original living cohort (tree or shrub) (in cm)
DeathAge	Years passed since plant death (in yrs)
SmallBranches	Necromass of standing small dead branches (diameters between 0.635 cm and 7.5 cm; 10h and 100h fuels; in g C/m ²)
LargeWood	Necromass of standing large dead wood (diameters > 7.5 cm; 1000h and 10000h fuels; in g C/m ²).

Dead plant cohorts are characterized by species, diameter (for trees), height and age since death. Note that the description of dead cohorts quantifies the abundance of branches and large wood (i.e. stems), but not leaves (or twigs), which are assumed to become litter before or soon after plant death.

Elements corresponding to **downed litter** or **belowground litter** are described in an additional data frame called **litterData**, which contains the following columns:

R column	Description
Species	String with species (taxon) name.
Leaves	Necromass of leaf litter (in g C/m ²).
Twigs	Necromass of twig litter (diameters < 0.635; in g C/m ²).
SmallBranches	Necromass of downed small dead branches (diameters between 0.635 cm and 7.5 cm; 10h and 100h fuels; in g C/m ²).
LargeWood	Necromass of (downed) large dead wood (diameters > 7.5 cm; 1000h and 10000h fuels; in g C/m ²).
CoarseRoots	Necromass of dead coarse roots (in g C/m ²).
FineRoots	Necromass of fine root litter (in g C/m ²).

Note that litter elements are characterized by species, so that the model can decompose litter elements at different rates depending on species identity (see chapter 18).

2.4.2.5 Soil organic carbon pools

Forest objects can include the description of soil organic matter pools, in an element called **SOCDData**. Soil organic matter pools will depend on the decomposition model employed, being CENTURY the only available at present. **SOCDData** for CENTURY is a named numerical vector with the following items:

R items	Description
SurfaceMetabolic	Metabolic carbon in the surface (in g C/m ²).
SoilMetabolic	Metabolic carbon in the soil (in g C/m ²).
SurfaceActive	Active decomposition carbon pool in the surface (in g C/m ²).
SoilActive	Active decomposition carbon pool in the soil (in g C/m ²).
SurfaceSlow	Slow decomposition carbon pool in the surface (in g C/m ²).
SoilSlow	Slow decomposition carbon pool in the soil (in g C/m ²).
SoilPassive	Passive decomposition carbon pool in the soil (in g C/m ²).

2.4.3 Leaf and root distribution

2.4.3.1 Leaf area index of a plant cohort

LAI variables refer to *one-sided leaf area* of plants per surface area of the stand. Only living plant cohorts (or the collective herbaceous layer) contribute to the leaf area of the forest stand (i.e. seedlings or dead snags

are assumed to not contribute significantly to leaf area). Leaves standing on branches of living plant cohorts can be alive or dead. We will call *total leaf area index* of a plant cohort i ($LAI_{all,i}$) to the sum of its dead and live (and unfolded) leaf area:

$$LAI_{all,i} = LAI_{\phi,i} + LAI_{dead,i} \quad (2.11)$$

For simulations not involving growth, winter-deciduous phenology is a prognosed variable, and the current level of leaves unfolded may be lower than the maximum (live) amount of leaves. In simulations involving growth, both LAI_{ϕ} and LAI_{live} can change, the former due to leaf phenology (unfolding is simulated explicitly and has carbon costs) and the latter due to leaf allocation limits during plant development.

2.4.3.2 Leaf area index of the forest stand

If there are c living plant cohorts, the leaf area index corresponding to living leaves, unfolded leaves, dead leaves and leaves of hemi-parasitic (mistletoe) plants can be aggregated across cohorts using:

$$LAI_{live} = \sum_{i=1}^c LAI_{live,i} \quad (2.12)$$

$$LAI_{\phi} = \sum_{i=1}^c LAI_{\phi,i} \quad (2.13)$$

$$LAI_{dead} = \sum_{i=1}^c LAI_{dead,i} \quad (2.14)$$

$$LAI_{mistletoe} = \sum_{i=1}^c LAI_{mistletoe,i} \quad (2.15)$$

We can also estimate the overall leaf area index of living plant cohorts (LAI_{all}), including dead leaves:

$$LAI_{all} = \sum_{i=1}^c [LAI_{\phi,i} + LAI_{dead,i}] = LAI_{\phi} + LAI_{dead} \quad (2.16)$$

Finally, we can also estimate the overall leaf area index of the whole stand ($LAI_{all,stand}$), which also takes into account the leaf area of mistletoes ($LAI_{mistletoe}$) and the collective herbaceous layer (LAI_{herb}), if defined:

$$LAI_{all,stand} = LAI_{all} + LAI_{herb} = LAI_{\phi} + LAI_{dead} + LAI_{mistletoe} + LAI_{herb} \quad (2.17)$$

2.4.3.3 Vertical leaf distribution

The vegetation input structures allow the package to determine the vertical distribution of leaves in the stand. The leaf area of any woody plant cohort is assumed to be distributed vertically following a **truncated Gaussian function** whose standardized values -1.5 and 1.5 correspond to crown base height (H_{crown} ; in *cm*) and total plant height (H), respectively. Crown base height is defined as the height corresponding to the first living branch. It is calculated from the crown ratio of the cohort (CR ; a proportion between 0 and 1), which in turn can be estimated as explained in (23.11).

Simulation models in **medfate** divide the vertical dimension into vertical layers (by default are 100 cm width, but see control parameter **verticalLayerSize**). Let us define $LAI_{all,i,j} = LAI_{\phi,i,j} + LAI_{dead,i,j}$ as the leaf area index of cohort i in layer j including both functional leaves and dead leaves standing on branches. The truncated Gaussian distribution defines the $LAI_{\phi,i,j}$ and $LAI_{dead,i,j}$ for all plant cohorts and vertical layers.




Figure 2.4: Leaf area density distribution of woody vegetation in a forest stand. The left panel shows the distribution for each plant cohort separately, whereas the right panel shows the overall density distribution.

Dividing the leaf area of a given layer by its width, one obtains the *leaf area density* (LAD in $m^2 \cdot m^{-3}$). Figure 2.4 illustrates the leaf area density profile (see function `plot.forest()`) corresponding to the forest stand described in the example `forest` object that we showed above.

The leaf area density profile determines the light extinction rates through the forest canopy. The same truncated Gaussian distribution is used to distribute leaf and small branch biomass along the vertical dimension.

2.4.3.4 Vertical root distribution

Roots can be distinguished functionally into two different organs: **coarse roots** are responsible for mechanical support and transport of water and nutrients towards the stem, whereas **fine roots** are responsible for absorbing water and nutrients.

Analogously to the aboveground vegetation parameters, inputs describing depths corresponding to 50% and 95% of fine roots ($Z_{50,i}$ and $Z_{95,i}$) are used to completely specify the distribution of roots across soil layers. The root system of each plant cohort i is described using $FRP_{i,s}$, the proportion of fine roots (with respect to the whole plant) in each soil layer s . $FRP_{i,s}$ values are normally defined using the linear dose response model (Schenk and Jackson, 2002; Collins and Bras, 2007):

$$FRP_{i,s} = Y_i(Z_s) - Y_i(Z_{s-1}) \quad (2.18)$$

$$Y_i(z) = \frac{1}{1 + (z/Z_{50,i})^{c_i}} \quad (2.19)$$

where $Y_i(z)$ is the cumulative fraction of fine root mass located between surface and depth z ; $Z_{50,i}$ is the depth (in mm) above which 50% of the root mass is located; and c_i is a shape parameter related to $Z_{50,i}$ and $Z_{95,i}$ as $c_i = 2.94 / \ln(Z_{50,i}/Z_{95,i})$.

The linear dose response model is quite flexible with respect to fine root distribution (see internal function `root_ldrDistribution()`). The following figure shows the fine root distribution profile of the same cohorts as in fig. 2.4 (see function `plot.forest()`):




Figure 2.5: Examples of root density profile according to the linear dose response model.

The vertical distribution of fine roots can be effectively truncated to a maximum depth by specifying a value for Z_{100} (or by allowing the model to estimate it using `truncateRootDistribution = TRUE` in control parameters).

The actual input to simulation functions `spwb()`, `growth()` and alike, describes fine root distribution using the matrix of $FRP_{i,s}$ values - not $Z_{50,i}$ and $Z_{95,i}$ -, which means that the proportions of fine roots can be modified manually or be defined by another model. In fact, **medfate** also allows calculating $FRP_{i,s}$ values assuming a conic distribution of fine roots (see internal function `root_conicDistribution()`).

2.4.3.5 Water pools and root horizontal distribution

By default, models in **medfate** assume that soil moisture under all plant cohorts is the same (i.e. water sources corresponding to vertical soil layers are shared among cohorts). Therefore, it neglects spatial variation in soil moisture, against moisture variation patterns observed in mixed stands (Schume et al., 2004). However, variations in soil moisture beneath plant cohorts (and, implicitly, horizontal variation of soil moisture) can also be simulated if required by the user (see parameter `rhizosphereOverlap` in 2.6). This involves considering that a given plant cohort will perform water uptake from the **water pool** surrounding its roots, whereas it may not have access to the water beneath other plants. However, there can exist some degree of horizontal overlap between water pools exploited by different plants.

Considering water pools involves partitioning the stand area into fractions corresponding to the abundance of each plant cohort. More specifically, the model defines as many water pools as plant cohorts, with proportions defined by their LAI values:

$$f_{pool,i} = \frac{LAI_{live,i}}{LAI_{live}} \quad (2.20)$$

Models assume that the rhizosphere of each plant cohort occupies its own water pool but may extend into the water pools under other plant cohorts. In other words, the root systems of different cohorts may overlap horizontally. Moreover, the horizontal overlap of root systems will vary across soil layers. A given plant cohort i will have its roots in layer s partitioned among different water pools. We thus need to define $fr_{i,s,j}$, the (horizontal) proportion of fine roots of cohort i in layer s of the water pool j , with the restriction that:

$$\sum_j fr_{i,s,j} = 1 \quad \forall i, s \quad (2.21)$$

A simplifying assumption is to force a complete independence of water pools, which is equal to say that $fr_{i,s,j} = 1$ for $i = j$ and $fr_{i,s,j} = 0$ when $i \neq j$ (this is the simplification adopted when using `rhizosphereOverlap = "none"`). In the general case of partial overlap (`rhizosphereOverlap = "partial"`), it is important to realize that proper estimation of $fr_{i,s,j}$ is challenging when we do not have explicit plant coordinates, root lateral widths, etc. At this point, let us assume we have a reasonable estimate of Vol_i , the **soil volume explored by coarse roots** of an individual of cohort i . Assuming that the proportion of fine roots in each layer is proportional to the proportion of total soil volume explored by coarse roots that corresponds to the same layer we have that $Vol_{i,s}$, the soil volume explored by coarse roots in layer s is:

$$Vol_{i,s} = FRP_{i,s} \cdot Vol_i \quad (2.22)$$

In **medfate**, coarse roots are represented by *axial* and *radial* components. The length of axial component of a given layer ($L_{axial,i,s}$) is simply equivalent to its bottom depth, whereas the radial component ($L_{radial,i,s}$) is unknown. However, we can assume that the soil volume explored by coarse roots corresponds to a cylinder:

$$Vol_{i,s} = \pi \cdot L_{radial,i,s}^2 \cdot L_{axial,i,s} \quad (2.23)$$

and substituting the soil volume $Vol_{i,s}$ in the equation above we can isolate $L_{radial,i,s}$ (see internal function `root_coarseRootLengthsFromVolume()`). Once we have the radial component we can estimate the area (m^2) covered by coarse roots of an individual of the cohort in a given layer s :

$$Area_{i,s} = \pi \cdot L_{radial,i,s}^2 \quad (2.24)$$

We can then compare $Area_{i,s}$ to the area of the water pool for an individual of cohort i , which in m^2 is:

$$AreaPool_i = 10000 \cdot f_{pool,i} / N_i \quad (2.25)$$

If $Area_{i,s} \leq AreaPool_i$, then the roots of the individual do not exceed the area of the pool, in other words $fr_{i,s,i} = 1$ and $fr_{i,s,j} = 0$ for all $j \neq i$. If $Area_{i,s} > AreaPool_i$ then the excess should be counted as overlap with the other pools. Assuming that plants are randomly distributed (i.e. no clumping) the probability of finding the pool of a cohort is equal to the proportion of its water pool (see internal function `root_horizontalProportions()`). All these calculations were done assuming Vol_i were known. Estimation of Vol_i depends on the complexity of the water balance model and is explained in A.4.6.

2.4.4 Functional traits and vegetation initialization

Besides the physical representation of vegetation, forest ecosystem models require information regarding plant functional traits, because this influences the outcome of physical, physiological and hydrological processes. In the following chapters, we will indicate for each simulated process its required plant functional parameters. A description of all parameters is provided in data frame `SpParamsDefinition` included in the package.

Normally, functional traits are described at the species level because infra-specific parameters are hard to get. Even at the species level, many functional (e.g. physiological) traits are hard to obtain, so in **medfate** we provide a default species parameter table for tree/shrub species found in Spain, with many of them occurring elsewhere in the Mediterranean Basin (`SpParamsMED`). User can modify this data frame to account for intra-specific trait variation (see function `modifySpParams()`).

The package provides functions `spwbInput()` and `growthInput()` that prepare the **forest** vegetation input for simulation functions `spwb()` and `growth()`, respectively (see 2.1.1). This initialization mostly consists

in compiling the necessary functional traits from the species parameter table, following the SP_i attribute of plant cohorts. Some new model parameters are estimated from the initial ones, while taking into account the structural attributes of cohorts (an example of these derived quantities is stem conductance, which estimated from stem xylem conductivity, Huber value and plant height). Users can take the output of these initialization functions and replace parameter values for specific plant cohorts (see function `modifyCohortParams()`).

2.4.5 Update of vegetation during simulations

Vegetation characteristics stay constant during simulations using function `spwb()`, although the actual (unfolded) leaf area and dead leaf area may vary depending on leaf phenology.

In contrast, growth simulation requires updating the structure of vegetation, i.e. plant heights, tree diameters, tree density and shrub cover. Therefore, function `growth()` can modify any of the structural attributes of living plant cohorts.

Finally, function `fordyn()` modifies not only structural variables of plants cohorts but also involves many other changes: (a) the removal of dead (or cut) cohorts; (b) the modification of the seed bank and the seedling bank; (c) the recruitment of new cohorts during simulations; (d) the modification of snag data and litter data.

2.5 Meteorological input

Weather input data must include variables calculated at the **daily** scale. Weather data should be arranged in a data frame with days in **rows** and variables in **columns**.

Since it allows producing data frames with appropriate variable units and row/column names, we recommend meteorological input to be generated using package **meteoland** (De Cáceres et al., 2018), but other sources are possible.

2.5.1 Date format

Dates are needed in **medfate** to estimate parameters like solar declination or the day of the year (*DOY*). Dates can be specified in the input data frame either:

- As data frame **row names** (in string format *year-month-day*)
- As **Date** elements in a column called "**dates**".

2.5.2 Required weather variables

The following table indicates the symbols, units, definitions and the variable name in R of input weather variables that are always **required** in the input data frame:

Symbol	Units	R param	Description	Optional	Missing allowed
T_{min}	°C	MinTemperature	Minimum temperature	No	No
T_{max}	°C	MaxTemperature	Maximum temperature	No	No
RH_{min}	%	MinRelativeHumidity	Minimum relative humidity	No	Yes
RH_{max}	%	MaxRelativeHumidity	Maximum relative humidity	No	Yes
P	$L \cdot m^{-2} = mm$	Precipitation	Precipitation (either rainfall or snow).	No	No
Rad	$MJ \cdot m^{-2}$	Radiation	Solar radiation after accounting for clouds	No	Yes

Some weather variables (temperature, precipitation) cannot contain missing values, whereas for others (radiation, relative humidity) the package can impute values when missing, thanks to its dependency with package **meteoland**.

2.5.3 Optional weather variables

The following table lists the same attributes for input weather variables that are **optional**:

Symbol	Units	R param	Description	Optional
u	$m \cdot s^{-1}$	WindSpeed	Wind speed	Yes
R_{int}	$mm \cdot h^{-1}$	RainfallIntensity	Rainfall intensity	Yes
C_{atm}	ppm	CO2	Atmospheric (above-canopy) CO_2 concentration	Yes
P_{atm}	kPa	Patm	Surface atmospheric pressure	Yes
P_{fire}	[0-1]	FireProbability	Probability of wildfire occurrence	Yes

Missing wind speed (u) and CO_2 values will be replaced by defaults defined in control parameters. As an alternative to providing daily atmospheric CO_2 concentration values, simulation functions allow the user to provide CO_2 concentration values in **annual** time steps using a separate vector.

When missing, atmospheric pressure (P_{atm}) is also derived from elevation using an utility function available in **meteoland**.

Missing rainfall intensity (R_{int}) values will be estimated from the current month using function `hydrology_rainfallIntensity()` and the seasonality of rainfall intensity given in control parameters (see `defaultRainfallIntensityPerMonth` in 3.5.3).

2.5.4 Derived weather variables

While the weather variables listed in the tables above are provided as inputs, **medfate** derives other atmospheric variables from them, using also topographic information (see utility functions of the **meteoland** reference manual):

Symbol	Units	R param	Description
ρ_{air}	$kg \cdot m^{-3}$		Air density
e_{atm}	kPa		Atmospheric water vapor pressure
T_{mean}	$^{\circ}C$	MeanTemperature	Mean daily temperature
PET	$L \cdot m^{-2} = mm$	PET	Potential evapotranspiration, calculated using Penman's (1948) equation

2.6 Simulation control

Simulation control parameters are a list of global parameter values, initialized using function `defaultControl()`, that the user can modify to change the general behavior of simulation functions. Here is the set of global control parameters currently accepted in **medfate**:

```
names(defaultControl())
```

```
## [1] "fillMissingRootParams"      "fillMissingSpParams"
## [3] "fillMissingWithGenusParams" "verbose"
## [5] "subdailyResults"           "standResults"
## [7] "soilResults"                "soilPoolResults"
## [9] "snowResults"               "plantResults"
## [11] "labileCarbonBalanceResults" "plantStructureResults"
```

## [13] "growthMortalityResults"	"decompositionPoolResults"
## [15] "leafResults"	"temperatureResults"
## [17] "fireHazardResults"	"fireHazardStandardWind"
## [19] "fireHazardStandardDFMC"	"transpirationMode"
## [21] "soilDomains"	"rhizosphereOverlap"
## [23] "truncateRootDistribution"	"fullRhizosphereOverlapConductivity"
## [25] "soilFunctions"	"VG_PTF"
## [27] "ndailysteps"	"max_nsubsteps_soil"
## [29] "defaultWindSpeed"	"defaultCO2"
## [31] "defaultRainfallIntensityPerMonth"	"leafPhenology"
## [33] "bareSoilEvaporation"	"unlimitedSoilWater"
## [35] "interceptionMode"	"infiltrationMode"
## [37] "infiltrationCorrection"	"verticalLayerSize"
## [39] "windMeasurementHeight"	"stemCavitationRecovery"
## [41] "leafCavitationRecovery"	"lfmcComponent"
## [43] "mistletoeParams"	"defoliationParams"
## [45] "hydraulicRedistributionFraction"	"nsubsteps_canopy"
## [47] "taper"	"multiLayerBalance"
## [49] "leafConductanceEstimation"	"rootConductivityEstimation"
## [51] "sapFluidityVariation"	"TPhase_gmin"
## [53] "Q10_1_gmin"	"Q10_2_gmin"
## [55] "rootRadialConductance"	"averageFracRhizosphereResistance"
## [57] "thermalCapacityLAI"	"boundaryLayerSize"
## [59] "cavitationRecoveryMaximumRate"	"sunlitShade"
## [61] "numericParams"	"leafCavitationEffects"
## [63] "stemCavitationEffects"	"stomatalSubmodel"
## [65] "plantCapacitance"	"cavitationFlux"
## [67] "leafCuticularTranspiration"	"stemCuticularTranspiration"
## [69] "C_SApoInit"	"C_LApoInit"
## [71] "k_SSym"	"fractionLeafSymplasm"
## [73] "gs_NightFrac"	"JarvisPAR"
## [75] "fTRBToLeaf"	"subdailyCarbonBalance"
## [77] "allowDessication"	"allowStarvation"
## [79] "sinkLimitation"	"shrubDynamics"
## [81] "herbDynamics"	"allocationStrategy"
## [83] "phloemConductanceFactor"	"nonSugarConcentration"
## [85] "equilibriumOsmoticConcentration"	"minimumRelativeStarchForGrowth"
## [87] "constructionCosts"	"senescenceRates"
## [89] "maximumRelativeGrowthRates"	"organogenesisEfficiencyModulator"
## [91] "elongationEfficiencyModulator"	"mortalityMode"
## [93] "mortalityBaselineRate"	"mortalityRelativeSugarThreshold"
## [95] "mortalityRWCThreshold"	"recrTreeDBH"
## [97] "recrTreeDensity"	"ingrowthTreeDBH"
## [99] "ingrowthTreeDensity"	"decompositionAnnualBaseRates"
## [101] "decompositionAnnualTurnoverRate"	"allowSeedBankDynamics"
## [103] "allowRecruitment"	"allowResprouting"
## [105] "recruitmentMode"	"removeEmptyCohorts"
## [107] "minimumTreeCohortDensity"	"minimumShrubCohortCover"
## [109] "dynamicallyMergeCohorts"	"keepCohortsWithObsID"
## [111] "seedRain"	"seedProductionTreeHeight"
## [113] "seedProductionShrubHeight"	"probRecr"
## [115] "minTempRecr"	"minMoistureRecr"
## [117] "minFPARRecr"	"recrAge"
## [119] "recrTreeHeight"	"recrShrubCover"


```
## [121] "recrShrubHeight"
```

Control parameters are employed when initializing vegetation inputs and state variables using functions `spwbInput()` and `growthInput()`, since they are needed also to make decisions at the point of initialization. Control parameters are also stored in the result of these functions, so the user does not need to specify again control parameters when calling simulation functions such as `spwb()` or `growth()`. Function `fordyn()` is different than the other simulation functions, because the list control parameters is directly passed to the function (see 2.1.1).

Even if the list of control parameters is long, not all control parameters are relevant to all simulation functions (this will be indicated in the following chapters), and most of them should not be altered from their default values. However, there are a set of parameters that are worth learning:

- **verbose** [=TRUE]: Whether extra console output is desired during simulations.
- **transpirationMode** ["Granier"] is an important control parameter, because it allows the user to switch between the *basic* (i.e. "Granier") and *advanced* (i.e., either "Sperry" or "Sureau") water balance models, corresponding to chapters 3 and 7, respectively.
- **soilDomains** ["buckets"] is an important control parameter, because it specifies how water moves between soil layers. Accepted values are "buckets" (for multi-bucket model), "single" (for single-domain Richards model) and "dual" (for dual-permeability model). See 5.7.
- **rhizosphereOverlap** ["partial"] is also a relevant parameter, since changing its value allows considering that plants extract soil water from the same pools, or partially or totally unrelated water pools, as explained in 2.4.3.5.

Some parameters are only relevant for parameter estimation at the stage of initialization:

- **fillMissingRootParams** [= TRUE]: Boolean flag to indicate that initialization functions should provide estimates for Z50 and Z95 if these are lacking in the forest data. Note that if **fillMissingRootParams** is set to FALSE then simulations may fail if the user does not provide values for Z50 and Z95 in tree or shrub data.
- **fillMissingSpParams** [=TRUE]: Boolean flag to indicate that initialization functions should provide estimates for functional parameters if these are lacking in the species parameter table **SpParams**. Note that if **fillMissingSpParams** is set to FALSE then simulations may fail if the user does not provide values for required parameters.
- **fillMissingWithGenusParams** [=TRUE]: Boolean flag to indicate that initializing functions should provide estimates from genus value, if species-level values are missing in the species parameter table **SpParams** but genus-level ones are not.

Part II

Basic water balance modelling

Chapter 3

Basic water balance model

This chapter provides an overview of the basic soil and plant water balance model included in **medfate**. The model allows simulating vertical water fluxes in a forest stand, and performs soil and plant water balance on a daily step basis for the period corresponding to input weather data.

The model is run using function `spwb()`, for a set of days, or function `spwb_day()`, for a single day. The basic water balance model corresponds to the control variable `transpirationMode = "Granier"`.

The description of the model presented here conforms to a large extent to the description given in De Cáceres et al. (2015), but includes several modifications that have been implemented since the publication of this scientific article. The information provided in this chapter should be enough to understand what the model does and when it can be useful, but the reading the following ones should provide a more detailed understanding on how the different processes are formulated.

3.1 Design principles

The water balance model considers only the vertical spatial dimension of the stand and not the horizontal distribution of plants within it. In other words, the model is not spatially explicit (i.e., plants do not have horizontal coordinates nor interact for space explicitly). Living plants in the stand are represented in groups, here referred to as *plant cohorts*, of different species, height and leaf area index (*LAI*). Height and *LAI* values determine competition for light. *LAI* values also drive competition for soil water, along with fine root distribution. The model can also consider a collective layer herbaceous vegetation (see 2.4.1.2), which also has a height value and *LAI*, but where plant cohorts are not distinguished and with fine roots confined to the topmost layer.

The soil water balance initially followed mostly the design principles of SIERRA (Mouillot et al., 2001; Ruffault et al., 2014, 2013) and BILJOU (Granier et al., 1999; 2007), although nowadays it has evolved substantially. Hydrological processes include rainfall interception loss (Gash et al., 1995; Liu, 2001), plant transpiration, evaporation from bare soil (Ritchie, 1972), snow melt, infiltration into the soil matrix (Boughton, 1989; Green and Ampt, 1911) and vertical water movement within the soil, the latter following either a single-domain model (Bonan, 2019) or a dual-permeability model (Jarvis et al., 1991; Larsbo et al., 2005), where total porosity is partitioned into two separate flow regions (matrix and macropores), each with its own degree of saturation, conductivity and flux rate. Potential evapo-transpiration (*PET*) determines maximum herbaceous and maximum canopy transpiration ($Tr_{\max}$) using an empirical relationship between the *LAI* of the stand and the ratio $Tr_{\max}/PET$ (Granier et al., 1999). Actual plant transpiration is estimated using a function that depends not only on $Tr_{\max}$ but also on current soil moisture levels, cohort-specific drought resistance, fine root distribution and the degree of shading of the target plant cohort. When soil water deficit progresses, transpiration rates are consequently reduced, but minimum transpiration rates of plant cohorts are not zero, because of the role of cuticular transpiration and imperfectly closed

stomata (Duursma et al., 2018). Plant water status (i.e. plant water potential and, hence, plant water content) follows variations of soil moisture in the soil layers where fine roots are located.

By default, the model assumes that soil moisture under all plant cohorts is the same (i.e. water sources corresponding to vertical soil layers are shared among cohorts). However, 3D variations in soil moisture beneath plant cohorts can be simulated in spatially-explicit forest ecosystem models (Manoli et al., 2017; Rötzer et al., 2017). The water balance model does not simulate horizontal spatial processes explicitly, but allows considering more or less independent water pools for plant cohorts. If partial or completely independent water pools are considered, hydrological processes are replicated for the fraction of soil corresponding to each plant cohort. Transpiration of each plant cohort depends on the soil moisture beneath itself and, depending on the degree of rhizosphere overlap, on the soil moisture beneath other plant cohorts.

Optionally, the model can simulate additional transpiration caused by hemi-parasitic plants (mistletoe) attached to specific plant cohorts. This causes water update from the corresponding soil water pools to be higher than cohort transpiration. Mistletoe transpiration flow is not extracted from plant tissues directly, as in Zweifel et al. (2012), but added to the water uptake from the soil layers (or soil water pools) where the plant is connected. Therefore, increased stress effects on the host plant are better represented when soil water pools are considered.

3.2 State variables

The following are state variables in the model under all simulations:

- Cumulative degree days (a) to budburst, (b) to complete unfolding or (c) to senescence ($S_{eco,d}$, $S_{unf,d}$ or $S_{sen,d}$; all in $^{\circ}C$), are tracked by the model to determine leaf phenological status (see 4.1).
- Soil moisture content dynamics on each layer s are tracked daily using $W_s = \theta_s(\Psi)/\theta_{fc,s}$, the *proportion of soil moisture in relation to field capacity*, where moisture at field capacity, $\theta_{fc,s}$, is assumed to correspond to $\Psi_{fc} = -0.033$ MPa. Note that W_s values larger than one are possible if the soil is between field capacity and saturation (which can happen if deep drainage is not allowed).
- Plant water potential ($\Psi_{plant,i}$) of each plant cohort i are also tracked daily and follows the soil water potential of layers (or water pools) where fine roots are located.
- Leaf area index ($LAI_{act,i}$) of each plant cohort i will change during simulations according to phenology if the species is winter-deciduous or evergreen with autumn leaf senescence.

Additional state variables depend on the activation of specific control flags:

- If *snow accumulation/melting is considered*, the model tracks S_{snow} , the snow water equivalent (in mm) of the snow pack storage over the surface.
- If *stem cavitation is not completely reversible*, the model tracks $PLC_{stem,i}$ and $PLC_{leaf,i}$, the proportion of conductivity lost the stem and leaves, respectively, of plants of cohort i .
- If *the degree of rhizosphere overlap between plant cohorts is not total* (i.e. if `rhizosphereOverlap` = "none" or `rhizosphereOverlap` = "partial"), the model also tracks $W_{i,s}$, the proportion of soil moisture in relation to field capacity for layer s within the areal fraction of the stand covered by the plant cohort i .
- If *cavitation-induced defoliation is considered* (i.e. `cavitationInducedDefoliation` = TRUE), then leaf area index ($LAI_{act,i}$) will become a state variable for all plant cohorts (see 6.2.3).

3.3 Water balance

Water balance is performed for both the snow pack and the soil (fig. 3.1). Precipitation (P) is considered be snow precipitation (Ps) or rainfall (Pr), depending on temperature. Thus, we have:

$$P = Pr + Ps \quad (3.1)$$

The water balance of the snow pack (ΔS_{snow} in mm of snow water equivalents) is simply defined as:

$$\Delta S_{snow} = Ps - Sm \quad (3.2)$$

where Ps is precipitation as snowfall and Sm is snow melt. Interception on canopy surfaces is only modelled for liquid precipitation, which leads to a partitioning of Pr into interception loss (In ; i.e. water evaporated after being intercepted by the canopy surfaces) and net rainfall (Pr_{net} ; the sum of throughfall and stem flow):

$$Pr = Pr_{net} + In \quad (3.3)$$

Several water sources can contribute to soil infiltration, namely net rainfall (Pr_{net}), surface runoff from nearby locations (Ro , if provided) and snowmelt (Sm). Given that not all water that reaches the soil surface actually infiltrates, the amount of water entering the first soil layer via infiltration (If) is:

$$If = Pr_{net} + Sm + Ro - Ie \quad (3.4)$$

where Ie is the infiltration excess that may occur. Other water inputs into the soil can come via capillarity rise (Cr) from the water table (if present) or via lateral interflow from nearby locations (Lw ; if provided, assumed positive in the direction of entering the local soil). There are multiple ways water can leave the soil. Regarding purely physical processes we have the evaporation from soil surface (Es), saturation excess (Se) and deep drainage (Dd). Fluxes mediated by plants are the transpiration (Tr) made by herbaceous plants (Tr_{herb}), by plant cohorts ($Tr_{cohorts}$) and (sometimes) by hemi-parasitic (mistletoe) plants ($Tr_{mistletoe}$):

$$Tr = Tr_{herb} + Tr_{cohorts} + Tr_{mistletoe} \quad (3.5)$$




Figure 3.1: Schematic representation of soil and snow pack water balance components. Subsurface lateral source/sinks and surface runoff are omitted for simplicity. Green and blue boxes represent outflows corresponding to green and blue water respectively.

Taking all input and output terms, daily variations in soil water content (ΔV_{soil} in mm) are then summarized as:

$$\Delta V_{soil} = (If + Lw + Cr) - (Dd + Se + Es + Tr) \quad (3.6)$$

The water leaving the soil via surface runoff (Ru) is defined as the sum of infiltration excess (Ie) and saturation excess (Se):

$$Ru = Ie + Se \quad (3.7)$$

If we consider soil and snow pack altogether, the water balance of the whole forest becomes:

$$\Delta V_{soil} + \Delta S_{snow} = (P + Ro + Lw + Cr) - (Dd + Ru + In + Es + Tr) \quad (3.8)$$

Among inputs, precipitation (P) will be the most relevant inflow on most instances. The liquid water fluxes contributing to **blue water** outflow (i.e. entering river networks or leading to ground water) are runoff (Ru) and deep drainage (Dd). Water vapor fluxes (Es , $Tr_{cohorts}$, Tr_{herb} , Tr_{mist} , In) contribute to total evapotranspiration, or **green water** (i.e. water vapor returning to the atmosphere). These are represented in Fig. 3.1 using blue/green boxes.

While the above describes the default water balance equations, variations can occur depending on specific control flags:

- a. *If snow dynamics are considered* [`snowpack`], evaporation from bare soil cannot occur if there is a snow pack over the soil (i.e., if $S_{snow} > 0$ then $Es = 0$).
- b. *If plant water pools are considered* [`rhizosphereOverlap`], the soil water balance equation applies not only to the soil of the overall stand but also to the soil beneath each plant cohort. The fraction of stand area corresponding to the water pool of each cohort is used to keep the water balance at the two scales aligned. Moreover, the water balances of soils beneath the different plant cohort are more or less correlated depending on the degree of rhizosphere overlap.
- c. *If soil flows are simulated using a dual-permeability domain* [`soilDomains`] there are water balance equations for the soil matrix and the macropore domain. Moreover, it affects the infiltration amounts (and hence runoff), in the latter including infiltration via macropores.

3.4 Process scheduling

For every day to be simulated, the model performs the following steps:

1. Update leaf area values according to the phenology of species and recalculate radiation extinction (sections 4.1 and 4.2).
2. Determine rainfall interception loss (In) and net rainfall (Pr_{net}) (section 5.3). The amount of water reaching the soil surface from the rainfall event is the sum of net rainfall (Pr_{net}) and surface runoff (Ro), if given. If snow dynamics are included, increase snow pack from snow precipitation (Ps) and decrease it following snow melt (Sm) (section 5.2).
3. Determine bare soil evaporation (Es), if snow is not present (section 5.4), and herbaceous layer transpiration (Tr_{herb} ; section 5.5) if required.
4. Determine plant cohort transpiration ($Tr_{cohorts}$), including hydraulic redistribution, and photosynthesis (chapter 6). Update plant water potential (Ψ_{plant}) and plant drought stress for each plant cohort. If necessary, determine mistletoe transpiration ($Tr_{mistletoe}$) and the corresponding additional soil water uptake (section 6.1.6).
5. Determine rainfall infiltration (section 5.6) and vertical water movement within the soil (section 5.7), while considering soil layer source/sinks from snow melt, bare soil evaporation and herbaceous layer transpiration and water uptake/redistribution by plant cohorts. This updates soil moisture and returns infiltration (If), runoff (Ru) and deep drainage (Dd).
6. If fire hazard calculations are requested, estimate live and dead fuel moisture and potential fire behaviour (chapter 26).

The set of steps corresponds to simulations if plant water pools *are not* considered, so that only the water balance of the overall soil matters. If plant water pools *are* considered, steps 1, 2 and 6 are the same, because they apply to the whole forest stand. However, step 3 is performed for each water pool separately because it depends on antecedent soil moisture. Step 4 also involves defining plant water uptake from different plant water pools depending on the proportion of fine roots of each plant cohort in each pool (see 6.1.5).

Step 5 is also performed on the soil corresponding to each pool, and the resulting infiltration, runoff and drainage estimates are averaged at the stand level. Finally, the soil moisture of the overall soil is estimated by averaging the soil moisture of water pools.

3.5 Inputs and outputs

3.5.1 Soil, vegetation and meteorology

Soil

Soil input requirements are fully described in section 2.3.

Vegetation

Vegetation input requirements are fully described in section 2.4. Tree, shrub and herb cohorts do not need to be characterized with different structural variables in soil water balance calculations, as long as all plant cohorts have a LAI value. In most cases, users only need to estimate the leaf area index corresponding to live leaves, i.e. LAI_{live} , because normally at the starting point all leaves are expanded (i.e. $LAI_{act} = LAI_{live}$) and one can assume no dead leaves in the canopy (i.e., $LAI_{dead} = 0$). Vegetation characteristics stay constant during simulations using function `spwb()`, although the actual expanded leaf area (LAI_{act}) and dead leaf area (LAI_{dead}) may vary if the species is winter deciduous. For each plant cohort with hemi-parasitic (mistletoe) plants living on crowns, a corresponding value of mistletoe leaf area index ($LAI_{mistletoe}$) can be defined.

Meteorology

The minimum weather variables required to run the model are min/max temperatures (T_{min} and T_{max}), min/max relative humidity (RH_{min} and RH_{max}), precipitation (P) and solar radiation (Rad). Wind speed (u) is also needed, but the user may use missing values if not available (a default value will be used in this case). Wind speed is assumed to have been measured at a specific height above the canopy (by default at 2 m). Atmospheric CO_2 concentration (C_{atm}) may also be specified, but if missing a default constant value is assumed, which is taken from the control parameters. Definitions and units of these variables were given in section 2.5.

3.5.2 Plant functional parameters

The following sets of functional parameters should be supplied for each plant cohort. In practice, they are normally estimated from the species-specific parameter table and imputation of missing values (see A). Functional attributes are filled for each cohort i by function `spwbInput()` from species identity (i.e. SP_i). However, different parameters can be specified for different cohorts of the same species if desired (see section 2.4.4).

A first set of functional parameters refers to leaf phenology (`paramsPhenology`):

Symbol	Units	R	Description
		<code>PhenologyType</code>	Phenology type ("evergreen", "winter-deciduous", or "winter-semideciduous").
		<code>SenescencePeriod</code>	Senescence for evergreen plants ("whole-year", "spring" or "autumn").
		<code>GrowthPeriod</code>	Period of primary growth for evergreen plants ("whole-year" or "spring").
		<code>GrowthDeterminacy</code>	Growth determinacy ("determined", "indetermined" or "intermediate"), only relevant for the growth model (see 15.1.3).
LD	years	<code>LeafDuration</code>	Duration of leaves (in years).
BFD	days	<code>BudFormationDays</code>	Number of days required to complete bud formation (only relevant for the growth model (see 15.1.3)).
$t_{0,eco}$	days	<code>t0gdd</code>	Date to start the accumulation of degree days.

Symbol	Units	R	Description
S_{eco}^*	$^{\circ}\text{C}$	Sgdd	Cumulative degree days required for leaf budburst (see section 4.1.3).
S_{unf}^*	$^{\circ}\text{C}$	Ugdd	Cumulative degree-days required to complete leaf unfolding after budburst has occurred (see section 4.1.3).
T_{eco}	$^{\circ}\text{C}$	Tbgdd	Base temperature for the calculation of degree days to leaf budburst and unfolding (see section 4.1.3).
S_{sen}^*	$^{\circ}\text{C}$	Ssen	Cumulative degree days corresponding to the start of leaf senescence (see section 4.1.4).
F_{sen}^*	$^{\circ}\text{C}$	Fsen	Cumulative degree days corresponding to complete leaf senescence (see section 4.1.4).
Ph_{sen}	hours	Phsen	Photoperiod corresponding to start counting senescence degree-days (see section 4.1.4).
T_{sen}	$^{\circ}\text{C}$	Tbsen	Base temperature for the calculation of degree days to leaf senescence (see section 4.1.4).
x_{sen}	{0,1,2}	xsen	Discrete values, to allow for any absent/proportional/more than proportional effects of temperature on senescence (see section 4.1.4).
y_{sen}	{0,1,2}	ysen	Discrete values, to allow for any absent/proportional/more than proportional effects of photoperiod on senescence (see section 4.1.4).

A second set of parameters relate to light extinction and water interception (**paramsInterception**):

Symbol	Units	R	Description
k_{PAR}		kPAR	PAR extinction coefficient (see section 4.2).
s_{water}	$\text{mm } H_2O \cdot LAI^{-1}$	g	Crown water storage capacity (i.e. depth of water that can be retained by leaves and branches) per LAI unit (see section 5.3).

A third set includes parameters related to plant anatomic and morphological attributes (**paramsAnatomy**):

Symbol	Units	R param	Description
$1/H_v$	$\text{m}^2 \cdot \text{m}^{-2}$	A12As	Ratio of leaf area to sapwood area
RLR	$\text{m}^2 \cdot \text{m}^{-2}$	Ar2A1	Fine root area to leaf area ratio
SLA	$\text{m}^2 \cdot \text{kg}^{-1}$	SLA	Specific leaf area
ρ_{leaf}	$\text{g} \cdot \text{cm}^{-3}$	LeafDensity	Leaf tissue density
ρ_{wood}	$\text{g} \cdot \text{cm}^{-3}$	WoodDensity	Wood tissue density
$\rho_{fineroot}$	$\text{g} \cdot \text{cm}^{-3}$	FineRootDensity	Fine root tissue density
SRL	$\text{cm} \cdot \text{g}^{-1}$	SRL	Specific root length
RLD	$\text{cm} \cdot \text{cm}^{-3}$	RLD	Fine root length density (i.e. density of root length per soil volume)
$r_{6.35}$		r635	Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm

A fourth set of parameters are related to transpiration and photosynthesis (**paramsTranspiration**):

Symbol	Units	R	Description
g_{swmin}	$\text{mol } H_2O \cdot \text{s}^{-1} \cdot \text{m}^{-2}$	Gwmin	Minimum stomatal conductance to water vapour
$T_{max,LAI}$		Tmax_LAI	Empirical coefficient relating LAI with the ratio of maximum transpiration over potential evapotranspiration (see section 6.1.1).

Symbol	Units	R	Description
$T_{max,sqLAI}$		Tmax_LAI	Empirical coefficient relating squared LAI with the ratio of maximum transpiration over potential evapotranspiration (see section 6.1.1).
$\Psi_{extract}$	MPa	Psi_Extract	Root water potential at which plant transpiration is 50% of its maximum (see section 6.1.2).
$c_{extract}$		Exp_Extract	Parameter of the Weibull function regulating transpiration reduction (see section 6.1.2).
c_{leaf}, d_{leaf}	(unitless), MPa	VCleaf_d	Parameters of the vulnerability curve for leaf xylem
c_{stem}, d_{stem}	(unitless), MPa	VCstem_d	Parameters of the vulnerability curve for stem xylem
WUE_{max}	$g \cdot C \cdot mm^{-1}$	WUE	Water use efficiency at VPD = 1kPa and without light or CO2 limitations (see section 6.1.7).
WUE_{PAR}		WUE_par	Coefficient describing the progressive decay of WUE with lower light levels (see section 6.1.7).
WUE_{CO2}		WUE_co2	Coefficient for WUE dependency on atmospheric CO2 concentration 6.1.7).
WUE_{VPD}		WUE_vpd	Coefficient for WUE dependency on vapor pressure deficit 6.1.7).

A fifth (final) set of parameters are related to water storage and water relations in plant tissues (**paramsWaterStorage**):

Symbol	Units	R	Description
$LFMC_{max}$	%	maxFMC	Maximum live fuel moisture content, corresponding to fine fuels (< 6.35 mm twigs and leaves).
ϵ_{leaf}	MPa	LeafEPS	Modulus of elasticity of leaves
ϵ_{stem}	MPa	StemEPS	Modulus of elasticity of symplastic xylem tissue
$\pi_{0,leaf}$	MPa	LeafPIO	Osmotic potential at full turgor of leaves
$\pi_{0,stem}$	MPa	StemPIO	Osmotic potential at full turgor of symplastic xylem tissue
$f_{apo,leaf}$	[0-1]	LeafAF	Apoplastic fraction in leaf tissues
$f_{apo,stem}$	[0-1]	StemAF	Apoplastic fraction in stem tissues
V_{leaf}	$l \cdot m^{-2}$	Vleaf	Leaf water capacity per leaf area unit
$V_{sapwood}$	$l \cdot m^{-2}$	Vsapwood	Sapwood water capacity per leaf area unit

3.5.3 Control parameters

Control parameters modulate the overall behaviour of water balance simulations (see section 2.6). Most importantly, **transpirationMode** defines the transpiration model, which in turn defines the complexity of the water balance model. If **transpirationMode** = "**Granier**" (the default value) then the basic water balance model is used.

The other relevant control parameters for the basic water balance model are:

- **soilFunctions** [= "SX"]: Soil water retention curve and conductivity functions, either 'SX' (for Saxton) or 'VG' (for Van Genuchten).
- **truncateRootDistribution** [= FALSE]: Boolean flag to indicate that fine root distribution is to be truncated (i.e. Z_{100} is estimated from Z_{95} when not provided).
- **ndailysteps** [= 24]: Number of steps into which each day is divided for determination of soil water balance (24 equals 1-hour intervals).
- **max_nsubsteps_soil** [= 300]: Maximum number of substeps for soil water balance solving.
- **defaultWindSpeed** [= 2.5]: Default value for wind speed (in $m \cdot s^{-1}$) when this is missing (only used for leaf fall, see section 4.1).

- **defaultCO2** [=386]: Default atmospheric (abovecanopy) CO_2 concentration (in micromol $CO_2 \cdot mol^{-1}$ = ppm). This value will be used whenever CO_2 concentration is not specified in the weather input.
- **defaultRainfallIntensityPerMonth** [= c(1.5, 1.5, 1.5, 1.5, 1.5, 1.5, 5.6, 5.6, 5.6, 5.6, 5.6, 1.5)]: A vector of twelve values indicating the rainfall intensity (mm/h) per month. By default synoptic storms (1.5 mm/h) are assumed between December and June, and convective storms (5.6 mm/h) are assumed between July and November (Miralles et al., 2010).
- **snowpack** [= TRUE]: Whether dynamics of snow pack are included (see section 5.2).
- **leafPhenology** [= TRUE]: Whether leaf phenology is simulated (see section 4.1). If FALSE then all species are assumed to be evergreen.
- **bareSoilEvaporation** [= TRUE]: Whether evaporation from the soil surface is simulated (see section 5.4).
- **unlimitedSoilWater** [=FALSE]: Boolean flag to indicate the simulation of plant transpiration assuming that soil water is always at field capacity.
- **interceptionMode** [= "Gash1995"]: Infiltration model, either "Gash1995" or "Liu2001"; see section 5.3.
- **infiltrationMode** [= "GreenAmpt1911"]: Infiltration model, either "GreenAmpt1911" or "Boughton1989"; see section 5.6.
- **soilDomains** [= "buckets"]: Either "buckets" (for multi-bucket model), "single" (for single-domain Richards model) and "dual" (for dual-permeability model). See 5.7.
- **max_nsubsteps_soil** [= 300]: Maximum number of substeps for soil water balance solving.
- **fireHazardResults** [= FALSE]: Boolean flag to force calculation of daily fire hazard.
- **rhizosphereOverlap** [= "partial"]: A string indicating the degree of rhizosphere spatial overlap between plant cohorts:
 - "none" - no overlap (independent water pools).
 - "partial" - partial overlap determined by soil moisture.
 - "total" - total overlap (plants extract from common soil pools).
- **verticalLayerSize** [= 100]: Size of vertical layers (in cm) for the calculation of light extinction (see section 2.4.3.3).
- **windMeasurementHeight** [= 200]: Height (in cm) over the canopy corresponding to wind measurements.
- **stemCavitationRecovery/leafCavitationRecovery** [= "rate"]: A string indicating how to model recovery of xylem conductance (see section 14.4). Allowed values are:
 - "none" - no recovery.
 - "annual" - complete recovery every first day of the year.
 - "total" - instantaneous complete recovery.
 - "rate" - following a rate of new leaf or sapwood formation.
- **cavitationRecoveryMaximumRate** [= 0.05]: Maximum rate of conductance recovery (see section 6.2.2).
- **lfmcComponent** [= "fine"]: Plant component used to estimate LFMC, either "leaf" or "fine" (for fine fuel) (see section 10.3.4).
- **mistletoeParams**: A list with the following elements:
 - **kPAR** [= 0.5]: Mistletoe light extinction coefficient.
 - **g** [= 0.8]: Mistletoe water storage capacity per LAI unit.
 - **Tmax_LAI** [= 0.134]: Empirical coefficient relating mistletoe LAI with the ratio of maximum transpiration over potential evapotranspiration.
 - **Tmax_LAI^{sq}** [= -0.006]: Empirical coefficient relating squared mistletoe LAI with the ratio of maximum transpiration over potential evapotranspiration.
 - **Gs_P50_Baldocchi** [= -2.5]: Water potential corresponding to 50% reduction of mistletoe stomatal conductance (i.e. $\Psi_{50_{gs,bald,mistletoe}}$).
 - **Gs_slope_Baldocchi** [= 30]: Rate of decrease in mistletoe stomatal conductance at Gs_P50 (i.e. $slope_{gs,bald,mistletoe}$).
- **defoliationParams**: A list with the following elements:
 - **cavitationInducedDefoliation** [= TRUE]: Whether leaf hydraulic impairment induces defoliation.

- `criticalLeafPLC` [= 0.88]: Level of PLC (a proportion) corresponding to 50 % crown defoliation.
- `cvLeafP50` [= 10.0]: Coefficient of variation (in percent) of leaf P50 within crown leaves.

3.5.4 Model output

Function `spwb()` returns a list of a class with the same name. The first four elements of this list (i.e., `latitude`, `topography`, `weather` and `spwbInput`) are simply copies of model inputs. The next element is `spwbOutput`, which contains the state of the input object at the end of the simulation (this can be used to perform further simulations starting with current values of state variables). The next five elements correspond to *water balance flows*, *soil state* variables, *stand-level* variables, *plant-level* and *fire hazard* results:

Element	Description
WaterBalance	Climatic input and water balance flows of eq.(3.6) (i.e. net precipitation, infiltration, runoff, transpiration...). All of them in $mm = l \cdot m^{-2}$.
Soil	Soil variables for each soil layer: Moisture relative to field capacity (W_s), water potential (Ψ_s) and volumetric water content (V_s). Only returned if <code>soilResults</code> = TRUE.
Stand	Stand-level variables, such as LAI_{act} , LAI_{live} , LAI_{dead} , the water retention capacity of the canopy (S_{canopy}) or the fraction of light reaching the ground (L_{ground}^{PAR} and L_{ground}^{SWR}). Only returned if <code>standResults</code> = TRUE.
Plants	Variables defined at the level of plant cohort, such as expanded leaf area ($LAI_{act,i}$), transpiration, photosynthesis, water potential, etc. Only returned if <code>plantResults</code> = TRUE.
FireHazard	Fire hazard variables (fuel moisture, rate of spread, fire potentials, etc). Only returned if <code>fireHazardResults</code> = TRUE.

Elements `WaterBalance`, `Soil`, `Stand` and `FireHazard` are data frames with dates in rows and variables in columns, whereas `Plants` is itself a list with several data frames, with dates in rows and plant cohorts in columns:

Element	Symbol	Units	Description
LAI	LAI_{act}	$m^2 \cdot m^{-2}$	Actual leaf area index (expanded).
LAIlive	LAI_{live}	$m^2 \cdot m^{-2}$	Leaf area index (live) without accounting for phenology or drought-induced defoliation.
Phenology			Daily phenological status for each plant cohort (bud formation, bud dormancy, leaf elongation, leaf senescence), codified into an integer value.
FPAR		[0-1]	Fraction of of PAR at the canopy level of each plant cohort.
AbsorbedSWRFraction		[0-1]	Fraction of shortwave radiation absorbed.
Transpiration	Tr	mm	Transpiration per unit ground area.
GrossPhotosynthesis		$gC \cdot m^{-2}$	Gross photosynthesis per unit ground area.
PlantPsi	Ψ_{plant}	MPa	Plant water potential.
LeafPLC	PLC_{leaf}	[0-1]	Degree of leaf embolisation.
StemPLC	PLC_{stem}	[0-1]	Degree of stem embolisation.
PlantWaterBalance		mm	Internal daily plant water balance (balance of soil extraction and transpiration).
LeafRWC	RWC_{leaf}	%	Mean leaf relative water content.
StemRWC	RWC_{stem}	%	Mean stem relative water content.
LFMC	$LFMC$	%	Live fuel moisture content (as percent of dry weight), corresponding to fine fuels (< 6.35 mm twigs and leaves).
PlantStress	DDS	[0-1]	Drought stress level suffered by each plant cohort (relative whole-plant transpiration).

The output of simulations can be inspected using `plot`, `shinyplot` and `summary` functions specific to `spwb` objects (examples are given in the corresponding package *vignette*).

As it simulates water balance for only one day, function `spwb_day()` returns a much more reduced output. This function is most useful with advanced water balance modelling (chapter 7).

3.6 Applications

The basic water balance model is easier to parameterize and is faster than the advanced water balance model. These features make it appropriate for applications that do not seek a high predictive capacity of water status at the plant level, but require robust estimates of water fluxes at the stand, landscape or regional levels. In our opinion the basic water balance model may be useful to monitor or forecast temporal variation in hydrological fluxes (i.e. evapotranspiration vs. drainage/runoff) and soil moisture in particular stands or in multiple stands across a landscape or region. Similarly, it can be used to monitor or forecast temporal variation of plant drought stress, fuel moisture and potential fire behavior in particular stands or in multiple stands across a landscape or region.

For example, Karavani et al. (2018) employed the water balance to provide sound estimates of soil moisture dynamics to predict mushroom production in a network forest stands. The model is particularly interesting to test the relationship between forest structure/composition and the water balance at the stand level or the drought stress suffered by plants, as done in Ameztegui et al. (2017) or Rolo and Moreno (2019). Finally, the model has been used to estimate green water and blue water provision at landscape to regional scales (Morán-Ordóñez et al., 2020; Roces-Díaz et al., 2021; Sánchez-Dávila et al., 2024, 2026).

Chapter 4

Leaf phenology and radiation extinction

Leaf phenology determines seasonal variations in the surface area of leaves, hence influencing the energy and water exchange between plants and the atmosphere. Radiation extinction is an important process to consider in a model where plants have different heights.

4.1 Leaf phenology

4.1.1 Leaf phenology and phenological models

The following three phenological types are currently distinguished (see parameter **PhenologyType**):

- *Evergreen*: Evergreen plants involve continuous presence of leaves throughout the year, although the amount of leaves can vary seasonally.
- *Winter-deciduous*: In winter-deciduous plants, all leaves are shed in autumn and buds remain under dormancy until spring, when budburst and elongation occurs. Therefore, the plant has a non-zero leaf area during the period limited by leaf elongation and leaf shedding events.
- *Winter semi-deciduous*: Winter semi-deciduous plants are like deciduous ones, with the difference that they retain dead leaves in the plant until next season elongation period, so that leaf abscission is retarded.

Whereas leaves of deciduous plants always undergo senescence in autumn and new leaves are flushed in spring, different strategies can be distinguished with respect to seasonality in evergreen plants. In particular, we distinguish two possibilities for leaf growth period (parameter **GrowthPeriod**) - i.e. *whole-year growth* vs. *spring growth* - and three possibilities for leaf senescence period (parameter **SenescencePeriod**) - i.e. *whole-year senescence* vs *autumn senescence* or *spring senescence* - .

The most usual combinations of phenology type and growth/senescence periods are shown in Fig. 4.1 (see also primary growth phenology in 15.1):

Evergreen plants with whole-year leaf senescence and growth are typical of tropical regions and they keep the same amount of leaves constant throughout the year (unless there is cavitation-induced defoliation). Evergreen plants with spring senescence and spring growth result also in the same amount of leaves more or less constant throughout the year. In contrast, evergreen plants with spring growth and autumn senescence result in lower amount of leaves during winter, determined from their leaf lifespan.

Legend

- Live leaves present
- Budburst and leaf elongation
- Leaf abscission




Figure 4.1: Schematic representation of processes and timings for different leaf phenology types

4.1.2 Leaf phenological models

In winter-deciduous leaf phenology (or evergreens with autumn leaf senescence and spring growth), leaf-expanded status is updated daily, represented by ϕ_i , the fraction of maximum leaf area for cohort i :

$$LAI_{act,i} = LAI_{live,i} \cdot \phi_i \quad (4.1)$$

whereas for evergreen species $LAI_{act,i} = LAI_{live,i}$ (or equivalently $\phi_i = 1$ at all times).

The general structure of process based phenological models explained in Chuine et al. (2013). These models aim to determine, for a given phenological phase n (endodormancy, ecodormancy, maturation, etc), the day of its finalisation d_n , such that the following equation holds:

$$S_{n,d} = \sum_{d_{n-1}}^{d_n} R_{n,d} = S_n^* \quad (4.2)$$

where $S_{n,d}$ is the state of development on day d in phase n , and d_{n-1} is the end of the previous phase. $R_{n,d}$ is the rate of development during phase n on day d , which depends on environmental variables (temperature, photoperiod,...) and S_n^* is the critical threshold to achieve the change of phase.

In the water balance model, leaf area index (LAI) values of winter (semi-)deciduous plants are adjusted for leaf phenology following a leaf development model in spring and a leaf senescence model in autumn. Similarly, evergreen species whose leaf growth period is in spring but leaf senescence occurs in autumn also have a LAI that varies according to those models. Function `pheno_updateLeaves()` updates the status of expanded leaves and dead leaves in a simulation object. Regardless of leaf phenology type, when control parameter `cavitationInducedDefoliation` = TRUE the function `pheno_updateLeaves()` will take into account the degree of cavitation (i.e. using PLC_{leaf}) to determine the degree of defoliation (see 6.2.3).

4.1.3 Bud burst and leaf unfolding

Transition from ecodormancy to bud burst is estimated using a very simple one-phase ecodormancy model (also called the spring warming model) implemented in function `pheno_leafDevelopmentStatus()`. Given a base temperature (T_{eco} , in °C), the rate of development during the ecodormancy phase (R_{eco} , in °C) is zero for those days where mean temperature T_{mean} is below T_{eco} and $T_{mean} - T_{eco}$ for those days where temperatures become warmer than this threshold:

$$R_{eco,d}(T_d) = \begin{cases} 0 & T_d \leq T_{eco} \\ T_d - T_{eco} & T_d > T_{eco} \end{cases} \quad (4.3)$$

Degree accumulation starts after the year date surpasses parameter $t_{0,eco}$. If $DOY > t_{0,eco}$, daily $R_{eco,d}$ values are added to the cumulative sum S_{eco} and budburst occurs the first day that $S_{eco} \geq S_{eco}^*$ (see Fig. 4.1). Both T_{eco} and S_{eco}^* are plant-specific parameters in `spwbInput`.

After budburst, we model progressive leaf unfolding and elongation in a similar manner, by accumulating degree days using the same definition as for $R_{eco,d}$ and then defining an unfolding development status $S_{unf} = S_{eco} - S_{eco}^*$, which we use to determine ϕ_i , the degree of leaf elongation of cohort i :

$$\phi_i = S_{unf}/S_{unf}^* = (S_{eco} - S_{eco}^*)/S_{unf}^* \quad (4.4)$$

where S_{unf}^* is another plant-specific parameter of `spwbInput`. The leaf elongation process continues until $S_{unf} \geq S_{unf}^*$ (equivalently, when $S_{eco} \geq S_{eco}^* + S_{unf}^*$, see Fig. 4.1). By design, leaf elongation cannot extend beyond $DOY = 212$ (corresponding to 31st July in the northern hemisphere), i.e. the phenology submodel forces $\phi_i = 1$ at this date.

4.1.4 Leaf senescence

Leaf senescence for deciduous or evergreen plants with autumn senescence extends the design of the models developed by Delpierre et al. (2009) for leaf colouring. The daily rate of senescence forcing $R_{sen,d}$ is defined on the basis of daily photoperiod (Ph_d , in hours) and temperature (T_d , in °C):

$$R_{sen,d}(Ph_d, T_d) = \begin{cases} 0 & T_d \geq T_{sen} \text{ or } Ph_d > Ph_{sen} \\ (T_{sen} - T_d)^{x_{sen}} \cdot (Ph_d/Ph_{sen})^{y_{sen}} & T_d < T_{sen} \text{ and } Ph_d < Ph_{sen} \end{cases} \quad (4.5)$$

where Ph_{sen} is the maximum photoperiod to start counting the senescence; T_{sen} is the base temperature for the calculation of degrees of forcing senescence; x_{sen} and y_{sen} are exponents regulating the importance of temperature and photoperiod on leaf senescence. The state of development of senescence S_{sen} is defined as the sum of senescence forcing:

$$S_{sen} = \sum R_{sen,d} \quad (4.6)$$

The ratio S_{sen}/S_{sen}^* can be conceptually understood as the proportion of chlorophyll degradation and nutrient translocation in crown leaves (but at present it is not used to downregulate transpiration nor photosynthesis). When $S_{sen} \geq S_{sen}^*$ then leaves are assumed to have lost their green color (i.e. turned yellow or brown) and be ready for the terminal senescence phase (see Fig. 4.1), which involves cell death and separation in the abscission zone. From this point on, the fraction of crown leaves that are considered ready to abscission and fall increases due to further forcing accumulation, until $S_{sen} \geq S_{sen}^* + F_{sen}^*$ (see Fig. 4.1). The senescence submodel is implemented in function `pheno_leafSenescenceStatus()` and returns $\phi_{i,d} = 1$ for all days until $S_{sen} \geq S_{sen}^*$. After that, the function returns progressively lower values, dictated by $\phi_{i,d} = (S_{sen} - S_{sen}^*)/F_{sen}$. Parameters Ph_{sen} , T_{sen} , x_{sen} , y_{sen} , S_{sen}^* and F_{sen}^* are plant-specific parameters in `spwbInput`.

4.1.5 Leaf abscission and fall

The decrease in ϕ_i causes live expanded leaves to become dead leaves, still attached to branches but ready to be detached by wind action. Dead leaves are kept in the canopy and they are reduced daily using a negative exponential function of wind speed:

$$LAI_{dead,i,d} = LAI_{dead,i,d-1} \cdot e^{-u/10} \quad (4.7)$$

where u is wind speed in ($m \cdot s^{-1}$), $LAI_{dead,i,d-1}$ is the dead leaf area index of the previous day and $LAI_{dead,i,d}$ is the dead leaf area index of the current day.

4.2 Radiation extinction

The proportion of photosynthetically active radiation (PAR) and short-wave radiation (SWR; 400-3000 nm) decreases through the canopy following the Beer-Lambert's light extinction equation. L_{herb}^{PAR} , the proportion of PAR that reaches the herbaceous layer, is calculated as:

$$L_{herb}^{PAR} = \exp \left(- \sum_{i=1}^c k_{PAR,i} \cdot LAI_{all,i} + k_{PAR,mistletoe} \cdot LAI_{mistletoe,i} \right) \quad (4.8)$$

where $k_{PAR,i}$ is the PAR extinction coefficient of plant cohort i and $k_{PAR,mistletoe}$ is the PAR extinction coefficient of mistletoe. If we add the extinction caused by the herbaceous layer, we have that L_{ground}^{PAR} , the proportion of PAR that reaches the ground, is calculated as:

$$L_{ground}^{PAR} = L_{herb}^{PAR} e^{-0.5 \cdot LAI_{herb}} \quad (4.9)$$

Where herbaceous vegetation is assumed to have a fixed extinction coefficient.

We can also define L^{PAR} as the proportion of PAR available for a given plant cohort, which we associate to the PAR available at a height corresponding to half of the cohort's crown. This height will include some self-shading of the target cohort.

The proportion of short-wave radiation (SWR) energy absorbed by each plant cohort needs to be calculated to divide the transpiration of the stand among cohorts (chapter 6), and the radiation absorbed by the soil is needed to calculate soil evaporation (section 5.4). Foliage absorbs a higher proportion of PAR than SWR; thus, the extinction coefficient is higher for PAR than for SWR. However, values for the ratio of extinction coefficients are rather constant. Following Friend et al. (1997) it is assumed that the extinction coefficient for PAR is 1.35 times larger than that for SWR (i.e. $k_{SWR,i} = k_{PAR,i}/1.35$). Figure 4.2 shows the PAR and SWR extinction profiles (see functions `vprofile_PARExtinction()` and `vprofile_SWRExtinction()`) corresponding to the leaf area density distribution of Fig. 2.4.




Figure 4.2: Light extinction in a forest stand. Note the sharper decrease of PAR (left panel) in comparison to SWR (right panel)

To calculate radiation absorption, where the vertical dimension of the plot is divided into layers (as explained in 2.4.3.3), and the SWR absorbed is calculated for each plant cohort in each layer. Let l be the number of vertical layers. The fraction of radiation incident on layer j that is absorbed in the same layer is:

$$f_j = 1 - \exp \left(- \sum_{i=1}^c k_{SWR,i} \cdot LAI_{all,i,j} + k_{SWR,mistletoe} \cdot LAI_{mistletoe,i,j} \right) \quad (4.10)$$

where $LAI_{all,i,j}$ is the leaf area index of cohort i in layer j and $LAI_{mistletoe,i,j}$ is the leaf area index of mistletoe of cohort i in layer j . Hence, the fraction transmitted is $(1 - f_j)$. The fraction of radiation incident on layer j that is absorbed by expanded leaves of plant cohort i in that layer (f_{ij}) is calculated from the relative contribution of these leaves to the total absorption in the layer:

$$f_{ij} = f_j \cdot \frac{k_{SWR,i} \cdot LAI_{act,i,j}}{\sum_{h=1}^c k_{SWR,h} \cdot LAI_{all,h,j} + k_{SWR,mistletoe} \cdot LAI_{mistletoe,h,j}} \quad (4.11)$$

The fraction of canopy radiation absorbed by a plant cohort i across all layers is found by adding the fraction absorbed in each layer:

$$f_i = \sum_{j=1}^l f_{ij} \cdot \prod_{h>j}^l (1 - f_h) \quad (4.12)$$

where for each layer the fraction of the radiation incident in the canopy that reaches the layer is found by multiplying the transmitted fractions across the layers above it. For example, the fraction of SWR absorbed by each of the three cohorts of the example in Fig. 2.4 would be (see function `light_cohortAbsorbedSWRFraction()`):

```
##      T1_158      T2_179      S1_176
## 0.24545010 0.18229985 0.01216462
```

Chapter 5

Forest hydrology

Forest hydrological processes are key for soil and plant water balances. This chapter details the design and implementation of most of the vertical hydrological processes included in package **medfate**. These processes determine the value of the water flows mentioned in eq. (3.6). The only process that is not described here is plant cohort transpiration, as it will be covered in detail in chapter 6.

Although each process can be simulated in specific sub-model functions, function `hydrology_verticalInputs()` allows simulating soil water input processes altogether, including canopy water interception, snow accumulation/melt, soil infiltration and percolation.

5.1 Snow and rainfall

Precipitation (P) is considered be snow precipitation (Ps) when $T_{mean} < 0$, and is considered rainfall (Pr) otherwise. Thus, we have:

$$P = Pr + Ps \quad (5.1)$$

Interception of snow by the canopy is neglected, and all snow is assumed to accumulate in a single storage compartment S_{snow} over the soil (i.e. canopy snow storage capacity is neglected). Rainfall interception is described in section 5.3.

5.2 Snow pack dynamics

A very simple snow submodel is used for snow pack dynamics (accumulation and melt), taken from Kergoat (1998). When mean air temperature is above 0 Celsius ($T_{mean} > 0$), a simple energy budget relates snow melt, Sm (mm), to air temperature and soil-level radiation (see function `hydrology_snowMelt()`):

$$Sm = \frac{Rad \cdot L_{ground}^{SWR} \cdot (1 - \alpha_{ice}) + \tau_{day} \cdot T_{mean} \cdot \rho_{air} \cdot C_p / r_s}{\lambda_{ice}} \quad (5.2)$$

where Rad is solar radiation ($MJ \cdot m^{-2}$), L_{ground}^{SWR} is the fraction of (short-wave) radiation reaching the ground, $\alpha_{ice} = 0.9$ is the albedo of snow, $\tau_{day} = 86400$ is the day duration in seconds, ρ_{air} is the air density ($kg \cdot m^{-3}$), depending on temperature and elevation (see utility functions of the **meteoland** reference manual), $C_p = 1013.86 \cdot 10^{-6} MJ \cdot kg^{-1} \cdot C^{-1}$ is the specific heat capacity of the air and $r_s = 100 s \cdot m^{-1}$ is the snow aerodynamic resistance and $\lambda_{ice} = 0.33355 MJ \cdot kg^{-1}$ is the (latent) heat of fusion of snow.

5.3 Rainfall interception loss

As mentioned above, interception loss is only modelled for liquid precipitation (i.e. snow interception is not modelled). Rainfall interception loss, In , is estimated using either the Gash et al. (1995) or Liu (2001) models.

5.3.1 Gash (1995) model

In the Gash et al. (1995) analytical interception model for sparse canopies, rain is assumed to fall in a single event during the day. First, the amount of rainfall needed to saturate the canopy is calculated:

$$P_G = -\frac{S_{canopy}/C_{canopy}}{ER_{ratio}} \cdot \ln(1 - ER_{ratio}) \quad (5.3)$$

where S_{canopy} is the canopy water storage capacity (in mm) – i.e. the minimum amount of water needed to saturate the canopy –, C_{canopy} is the canopy cover and ER_{ratio} is the ratio of evaporation rate to rainfall intensity during the rainfall event (R_{int} ; in $mm \cdot h^{-1}$). Although interception models are normally applied to single-canopy stands, we apply the sparse Gash model to the whole stand (including shrubs and the herbaceous layer). Moreover, in our implementation stem interception is lumped with crown interception, so that S_{canopy} represents both.

Following Watanabe and Mizutani (1996) we estimate S_{canopy} , the canopy water storage capacity, from adjusted LAI values:

$$S_{canopy} = LAI_{herb} + \sum_i s_{water,i} \cdot LAI_{act,i} \quad (5.4)$$

where $s_{water,i}$ is the depth of water that can be retained by leaves, branches and stems of cohort i per unit of leaf area index ($mm \cdot LAI^{-1}$). To estimate the stand cover, C_{canopy} , we use the complement of the percentage of PAR that reaches the ground, i.e. $C_{canopy} = 1 - L_{ground}^{PAR}$ (Deguchi et al., 2006).

The amount of water evaporated from interception (In , in mm), is calculated as:

$$In = C_{canopy} \cdot P_G + C_{canopy} \cdot ER_{ratio} \cdot (Pr - P_G) \text{ if } Pr > P_G \quad (5.5)$$

$$In = C_{canopy} \cdot Pr \text{ if } Pr \leq P_G \quad (5.6)$$

where Pr is the daily gross rainfall (in mm). Net rainfall (Pr_{net} , also in mm) includes throughfall and stemflow, and is calculated as the difference between gross rainfall and interception loss. Fig. 5.1 below shows examples of relative throughfall (including stemflow), calculated according to the Gash et al (1995) interception model, under different situations (see function `hydrology_rainInterception()`).

5.3.2 Liu (2001) model

In the single-storm form of the Liu (2001) model is similar to the Gash et al. (1995) model in terms of its inputs. Rainfall interception during a single storm can be predicted by gross rainfall amount P_G , the ratio of evaporation rate to rainfall rate during the rainfall event, ER_{ratio} , canopy gap fraction (C_{canopy}) and storage capacity (S_{canopy}):

$$In = S_{canopy} \cdot \left[1.0 - \exp \left(-P_G \cdot \left(\frac{C_{canopy}}{S_{canopy}} \right) \right) \right] \cdot (1.0 - (ER/C_{canopy})) + (ER \cdot P_G) \quad (5.7)$$

Fig. 5.2 below shows examples of relative throughfall (including stemflow), calculated according to the Liu et al (2001) interception model, under different situations (see function `hydrology_rainInterception()`).




Figure 5.1: Examples of canopy interception with different S_{canopy} (canopy water storage capacity), ER_{ratio} (ratio between evaporation and rainfall rates) and p (throughfall coefficient; $p = 1 - C_{canopy}$), using the Gash (1995) model.




Figure 5.2: Examples of canopy interception with different S_{canopy} (canopy water storage capacity), ER_{ratio} (ratio between evaporation and rainfall rates) and p (throughfall coefficient; $p = 1 - C_{canopy}$), using the Liu (2001) model.

5.4 Bare soil evaporation

Evaporation from the soil surface is the last component of the soil water balance to be calculated before calculating plant transpiration. Bare soil evaporation cannot happen if there is snow over the soil surface (i.e., if $S_{snow} > 0$).

Potential evaporation from the soil (PE_{soil} ; in $mm \cdot day^{-1}$) is defined as the product between PET and L_{ground}^{SWR} , the proportion of SWR absorbed by the ground:

$$PE_{soil} = PET \cdot L_{ground}^{SWR} \quad (5.8)$$

Actual evaporation from the soil surface is modeled as in Mouillot et al. (2001), who in turn followed Ritchie (1972). First, the model determines t_{soil} , the time needed to evaporate the current water deficit (difference between field capacity and current moisture) in the surface soil layer:

$$t_{soil} = \left\{ \frac{V_{fc,1} \cdot (1 - W_1)}{\gamma_{soil}} \right\}^2 \quad (5.9)$$

where $V_{fc,1}$ is the water retention capacity of layer 1, W_1 is the proportion of moisture in relation to field capacity of layer 1 and γ_{soil} is the maximum daily evaporation ($mm \cdot day^{-1}$). The calculated time is used to determine the ‘supplied’ evaporation, SE_{soil} :

$$SE_{soil} = \gamma_{soil} \cdot (\sqrt{t_{soil} + 1} - \sqrt{t_{soil}}) \quad (5.10)$$

The amount of water actually evaporated from the soil, Es , is then calculated as the minimum between supply, SE_{soil} , and demand (Federer, 1982), i.e. PE_{soil} (see function `hydrology_soilEvaporationAmount`):

$$Es = \min(PE_{soil}, SE_{soil}) \quad (5.11)$$

Figure 5.3 shows the cumulative evaporation from soils for different values of maximum evaporation rate.

5.5 Transpiration of the herbaceous layer

Transpiration of herbaceous layer is modelled analogously to the transpiration of plant cohorts. First, potential evapotranspiration from the herbaceous layer (PET_{herb} ; in $mm \cdot day^{-1}$) is defined as the product between PET and L_{herb}^{SWR} , the proportion of SWR reaching the herbaceous layer:

$$PET_{herb} = PET \cdot L_{herb}^{SWR} \quad (5.12)$$

Thus, the more dense the canopy of plant cohorts, the lower herbaceous layer transpiration will be. Maximum herbaceous transpiration $Tr_{herb,max}$ depends on both PET_{herb} and the amount of transpiring surface, i.e. LAI_{herb} . To estimate Tr_{max} the model uses the empirical equation of Granier et al. (1999), as in section 6.1.1:

$$\frac{Tr_{herb,max}}{PET} = -0.006 \cdot (LAI_{herb})^2 + 0.134 \cdot LAI_{herb} \quad (5.13)$$

Finally, actual herbaceous transpiration for a given soil layer s ($Tr_{herb,s}$) is reduced according to the soil water potential (Ψ_s), assuming that transpiration reduction follows Weibull function with fixed coefficients:

$$Tr_{herb,s} = FRP_{herb,s} \cdot Tr_{herb,max} \cdot \exp \left\{ \ln(0.5) \cdot \left[\frac{\Psi_i}{-1.5} \right]^2 \right\} \quad (5.14)$$

where $FRP_{herb,s}$ is the proportion of fine roots in layer s , defined using $Z_{50} = 50mm$ and $Z_{95} = 500mm$ (see 2.4.3.4).




Figure 5.3: Cumulative bare soil evaporation for different values of maximum evaporation rate (γ_{soil}). Topsoil layer (0 – 30 cm) is initialized at field capacity ($V_1 = 50mm$). PE_{soil} was assumed not to be limiting.

5.6 Infiltration

The amount of rainfall event water that reaches the soil is the sum of net rainfall (Pr_{net}) and runoff (Ro , in mm), if any. The amount of rainfall event water infiltrating into the soil matrix (If) is estimated using either the Boughton (1989) or the Green and Ampt (1911) models.

5.6.1 Boughton (1989) model

Surface runoff Ru can be estimated using the USDA SCS curve number method, following Boughton (1989):

$$Ru = \frac{(Pr_{net} + Ro - 0.2 \cdot V_{fc,soil})^2}{(Pr_{net} + Ro - 0.8 \cdot V_{fc,soil})} \quad (5.15)$$

where $V_{fc,soil}$ (in mm) is the overall soil water retention capacity. The Boughton (1989) model is available in function `hydrology_infiltrationBoughton()` and examples of its behavior are given in the figure below.

5.6.2 Green-Ampt (1911) model

The description of the Green and Ampt (1911) model has been adapted from chapter 9 of Bonan (2019). The Green-Ampt equation uses the concept of a horizontal wetting front at some distance L_w into the soil. The soil above L_w is saturated (i.e. θ_{sat}), whereas soil below the wetting front has a uniform content θ_{dry} corresponding to conditions prior to the rainfall event. Darcy's law provides the flux of water i_c infiltrating into the soil from the surface to the wetting front at depth $z = -L_w$:

$$i_c = K_{sat} \left[\frac{|\Psi_w|}{L_w} + 1 \right] \quad (5.16)$$




Figure 5.4: Examples of infiltration/runoff calculation using the Boughton (1989) model for different values of net rainfall and overall retention capacity, V_{soil} , calculated from different soil depths (topsoil+subsoil) and assuming that soil texture is 15% clay and 25% sand. Rock fragment content was 25% and 40% for the topsoil and subsoil, respectively.

where Ψ_w is the matric potential at the wetting front. The total cumulative infiltrated water I_c at time t is the time integral of i_c , and is also equal to the L_w multiplied by the change in water, i.e. $\Delta\theta = \theta_{sat} - \theta_{dry}$:

$$I_c = \Delta\theta L_w \quad (5.17)$$

The infiltration rate can also be expressed as the time derivative of I_c , i.e.:

$$i_c = \frac{dI_c}{dt} = \Delta\theta \frac{dL_w}{dt} \quad (5.18)$$

Combining the two equations for i_c we obtain:

$$\Delta\theta \frac{dL_w}{dt} = K_{sat} \left[\frac{|\Psi_w|}{L_w} + 1 \right] \quad (5.19)$$

and substituting $I_c/\Delta\theta$ for L_w gives:

$$\frac{dI_c}{dt} = K_{sat} \left[\frac{|\Psi_w| \Delta\theta}{I_c} + 1 \right] \quad (5.20)$$

By rearranging terms and integrating we obtain an expression for $I_c(t)$:

$$I_c(t) = K_{sat} t + |\Psi_w| \Delta\theta \ln \left[1 + \frac{I_c(t)}{|\Psi_w| \Delta\theta} \right] \quad (5.21)$$

This equation defines the cumulative infiltration to time t as a function of K_{sat} , Ψ_w , θ_{sat} and θ_{dry} . Because I_c appears on both sides of the equation, it must be found using numerical root-finding (the Newton-Raphson method in our case). To estimate Ψ_w the following equation is used:

$$\Psi_w = \left(\frac{2b+3}{2b+6} \right) \Psi_{sat} \quad (5.22)$$

Parameter θ_{dry} is the volumetric content of the topmost soil layer (antecedent soil conditions), and parameters θ_{sat} , Ψ_{sat} , K_{sat} and b are estimated using Campbell's water retention curves from the texture of the topmost layer. The infiltration time t (in h) is estimated from the sum of net rainfall (Pr_{net}) and runoff (Ro) (both in mm) using rainfall intensity of the rainfall event (R_{int} ; in mm/h). The Green and Ampt (1911) model is available in function `hydrology_infiltrationGreenAmpt()` and examples of its behavior are given in the figure below.




Figure 5.5: Examples of infiltration using the Green-Ampt (1911) model for different values of net rainfall. The left panel shows the infiltration for different soil textural types, assuming a $\theta_{dry} = 0.3$. The right panel shows the infiltration for different values of θ_{dry} , assuming a silt loam soil. In both cases rainfall intensity is assumed $R_{int} = 6 mm/h$.

5.7 Soil water flows

5.7.1 Design of soil water flows

In this chapter we describe water movement into, out of, and within the soil. The water input to the soil is the sum of net rainfall (Pr_{net} , i.e. rainfall minus interception), snow melt (Sm) and (if simulations are done in a spatially-explicit context) runoff from an adjacent location (Ro). Water movement can be performed using either a multi-bucket model (`soilDomains = "buckets"`), the common single domain model (`soilDomains = "single"`) or the dual permeability model (`soilDomains = "dual"`), which follows with some modifications the MACRO model (Jarvis et al., 1991; Larsbo et al., 2005).

In the **multi-bucket model**, the water coming from rainfall events - i.e. sum of net rainfall (Pr_{net}) and runoff (Ro) - is divided between infiltration into the soil (If) and runoff (Ru), following the models explained in 5.6. Rainfall infiltrating into the soil is divided among soil layers, in a crude attempt to emulate the effect of preferential flows (but without accounting for the increase in infiltration caused by macropores). The amount of water reaching each soil layer is modelled using an extinction function that depends on macroporosity (see function `hydrology_infiltrationRepartition`). Next, the model pools the infiltrated water with input/outputs from plant extraction (or lateral processes). After that, water is added to (or subtracted from) each layer and if content is above field capacity the excess percolates to the layer below. If there

is still an excess for the bottom layer, the model will progressively fill upper layers (eventually generating saturation excess if the first layer becomes saturated). Every day if some layers are over field capacity, the model simulates deep drainage up to an amount defined by the saturated conductivity of the bottom layer.

In the **single-domain model** (panel A in Fig. 5.6), the water coming from rainfall events - i.e. sum of net rainfall (Pr_{net}) and run-on (Ro) - is divided between infiltration into the soil (If) and runoff (Ru), following the models explained in 5.6. Rainfall infiltrating into the soil is divided among soil layers, in a crude attempt to emulate the effect of preferential flows (but without accounting for the increase in infiltration caused by macropores). The amount of water reaching each soil layer is modelled using an extinction function that depends on macroporosity (see function `hydrology_infiltrationRepartition`). In contrast to water from rainfall events, water from snow melt (Sm) is assumed to infiltrate slowly and completely into the first layer. Water flows within the soil according to the Richards equation, assuming an homogenous porous media and where fluxes are driven by both gravitational and matric potentials. Root water uptake/redistribution is taken into account as source/sinks from/to the soil layers along with bare soil evaporation. Water leaving the bottom layer (as well as departures from the water balance) is assumed to drain beyond the soil limits (i.e. deep drainage (Dd)). Saturation excess in the topmost layer leads to an increase in runoff (Ru) and a corresponding decrease in infiltration (If).

In the **dual-permeability model** (panel B in Fig. 5.6), total porosity is partitioned into two separate flow regions (matrix and macropores), each with its own degree of saturation, conductivity and flow rate. The water coming from rainfall events - i.e. sum of net rainfall (Pr_{net}) and run-on (Ro) - is divided between infiltration into the soil matrix (If ; following models in 5.6) and water infiltrating into the macropore domain. Water from snow melt (Sm) is assumed to infiltrate slowly and completely into the soil matrix. Fluxes within the soil matrix follow the Richards equation and is driven by both gravity and matric potentials, as in the single-domain model. In contrast, fluxes in the macropore region are gravity-driven only. Saturation of micropores of a given soil matrix layer leads to a water source into the corresponding macropore layer, whereas water uptake by the soil matrix via capillarity from the corresponding macropore layer follows a diffusion equation. Root water uptake/redistribution is taken into account as source/sinks in the soil matrix along with bare soil evaporation. Deep drainage (Dd) can occur from both the soil matrix or the macropore domain. Finally, saturation excess in the macropore domain leads to accumulation of water in the above layer and, eventually, to surface runoff (Ru).




Figure 5.6: Schematic representation of the (A) single domain and (B) dual-permeability models of flows into, out of and within the soil. Symbols are defined in the main text.

Function `hydrology_soilWaterBalance()` performs infiltration and vertical water movement flows, given a soil, net rainfall, snowmelt and source/sink vector for each layer. Source/sink terms are defined as follows for the first layer ($i = 1$):

$$S_1 = -Es - Tr_{herb,1} - Ex_{cohorts,1} \quad (5.23)$$

and for $1 < i \leq N$:

$$S_i = -Tr_{herb,i} - Ex_{cohorts,i} \quad (5.24)$$

where Es is bare soil evaporation, $Tr_{herb,i}$ is herbaceous uptake from layer i and $Ex_{cohorts}$ is plant cohort uptake/redistribution from/to layer i . In the single-domain and dual-permeability models, all right-hand terms are converted from *mm/day* to instantaneous rates (*m/s*) for proper units in S . By default, **24 time steps** of 1 hour are used for vertical water simulation of each day, but the time step is modified if required to close the water balance of the matrix or macropore domains. Function `hydrology_soilWaterBalance()` updates soil moisture and returns daily fluxes corresponding to infiltration, runoff and deep drainage.

The following sections detail the formulation and numerical solving of soil flows, both under the single-domain and the dual-permeability models.

5.7.2 Darcian flow and Richards equation

Water flows in the soil **from high to low potential** as described by Darcy's law. In the soil matrix, the **total potential** is used, i.e. the sum of the **matric potential** (Ψ_s , here expressed in m , see 2.3.3) and vertical **depth** z (also in m). The vertical depth z is taken as positive in the upward direction, so that $z = 0$ is the ground surface. In the vertical dimension the **flux** Q , i.e. rate of water movement (in ms^{-1}), according to Darcy's law is:

$$Q = -K(\theta) \cdot \frac{\delta(\Psi + z)}{\delta z} = -K(\theta) \cdot \frac{\delta \Psi}{\delta z} - K(\theta) \quad (5.25)$$

where $K(\theta)$ is the hydraulic conductivity (in ms^{-1}) that describes the ease with which water can move through the porous media space, and depends on soil moisture (θ). In the macropore domain, only the vertical depth (i.e. gravity potential) is considered, which leads to a **unit gradient**, and hence the flux Q is simply:

$$Q = -K_{ma}(\theta_{ma}) \quad (5.26)$$

where K_{ma} is the macropore hydraulic conductivity (in ms^{-1}), depending on the macropore water content θ_{ma} (in $m^3 \cdot m^{-3}$).

Consider a volume of soil with horizontal area $\Delta x \Delta y$, thickness Δz and with non-rock proportion $\lambda = 1 - P_{rocks}$. The mass flux entering the parcel from above is $\rho_w Q_{in} \Delta x \Delta y$ and the flux out of it is $\rho_w Q_{out} \Delta x \Delta y$. Conservation requires that the difference between fluxes and source/sinks equals the rate of change in water storage. The mass of water in the soil volume is $\rho_w \theta \Delta x \Delta y \Delta z \lambda$ so the change in water over a time interval Δt is:

$$\rho_w \frac{\Delta \theta}{\Delta t} \Delta x \Delta y \Delta z \lambda = -\rho_w (Q_{in} - Q_{out}) \Delta x \Delta y \quad (5.27)$$

and gives the continuity equation:

$$\frac{\Delta \theta}{\Delta t} \lambda = -\frac{(Q_{in} - Q_{out})}{\Delta z} \quad (5.28)$$

which expressed in the notation of calculus is:

$$\frac{\delta \theta}{\delta t} \lambda = -\frac{\delta Q}{\delta z} \quad (5.29)$$

Substituting Darcy's law for Q , and accounting for (lateral) sources/sinks S , gives the Richards equation:

$$\frac{\delta \theta}{\delta t} \lambda = \frac{\delta}{\delta z} \left[K(\theta) \frac{\delta \Psi}{\delta z} + K(\theta) \right] \pm S = K(\theta) \frac{\delta^2 \Psi}{\delta^2 z} + \frac{\delta K}{\delta z} \frac{\delta \Psi}{\delta z} \frac{\delta K}{\delta z} \pm S \quad (5.30)$$

The former is known as the *mixed-form* Richards equation, because it includes θ on the left hand and the vertical gradient in matric potential Ψ on the right hand. The *head-based* or Ψ -based form expresses the left hand term using Ψ as dependent variable:

$$C(\Psi) \frac{\delta \Psi}{\delta t} \lambda = K(\Psi) \frac{\delta^2 \Psi}{\delta z^2} + \frac{\delta K}{\delta z} \frac{\delta \Psi}{\delta z} + \frac{\delta K}{\delta z} \pm S \quad (5.31)$$

where $C(\Psi) = \delta\theta/\delta\Psi$ is known as the **specific moisture capacity**, in m^{-1} . In the macropore domain, the use of the former equation is problematic; a major reason being the lack of information concerning the relationship between Ψ and θ close to saturation. If we assume capillarity to be negligible in the macropore domain, then $\delta\Psi/\delta z = 0$ and the corresponding governing equation becomes:

$$\frac{\delta\theta_{ma}}{\delta t} \lambda = \frac{\delta K_{ma}}{\delta z} \pm S \quad (5.32)$$

where θ_{ma} and K_{ma} are the soil moisture and soil conductivity, respectively, in the macropore domain.

5.7.3 Hydraulic properties in the single-domain model

In single-domain simulations, the Van Genuchten (1980) model is used to represent the water retention curve (see section 2.3.2):

$$\theta(\Psi) = \theta_{res} + \frac{\theta_{sat} - \theta_{res}}{[1 + |\alpha \cdot \Psi|^n]^m} \quad (5.33)$$

where $m = 1 - 1/n$ is a shape parameter. It can be re-arranged into:

$$S_e = \frac{\theta(\Psi) - \theta_{res}}{\theta_{sat} - \theta_{res}} = (1 + |\alpha \cdot \Psi|^n)^{-m} \quad (5.34)$$

where S_e is the degree of saturation. The specific moisture capacity, $C(\Psi)$, is the derivative $\delta\theta/\delta\Psi$, i.e.:

$$C(\Psi) = \alpha \cdot m \cdot n \cdot (\theta_{sat} - \theta_{res}) \cdot \frac{|\alpha \cdot \Psi|^{n-1}}{(1 + |\alpha \cdot \Psi|^n)^{m+1}} \quad (5.35)$$

The hydraulic conductivity $K(\Psi)$ for a given water potential Ψ is determined in the Van Genuchten model as a function of S_e using:

$$K(\Psi) = K_{sat} \cdot S_e^l \cdot [1 - (1 - S_e^{1/m})^m]^2 \quad (5.36)$$

where K_{sat} is the saturated hydraulic conductivity and $l = 0.5$ is the tortuosity factor.

5.7.4 Hydraulic properties in the dual-permeability model

In MACRO (Jarvis et al., 1991; Larsbo et al., 2005), a “cut and join” approach is used to define the hydraulic functions. A water potential $\Psi_b = -0.1 \text{ m} = -0.98 \text{ kPa}$ partitions the total porosity into matrix pores and macropores, while the corresponding water content, θ_b and hydraulic conductivity K_b represent the saturated state of the soil matrix.

Matrix domain

Soil water retention in the matrix is calculated using a modified form of Van Genuchten’s (1980) equation:

$$S_{mi} = \frac{\theta_{mi}(\Psi) - \theta_{res}}{\theta_{sat}^{fict} - \theta_{res}} = (1 + |\alpha \cdot \Psi|^n)^{-m} \quad (5.37)$$

where $\theta_{sat}^{fict} = \theta_{sat} - P_{macro}$ is a fictitious saturated content obtained subtracting the proportion of macropores. θ_b is estimated as $\theta_{mi}(\Psi_b)$, whereas K_b is estimated with van Genuchten's K equation but using $S_{mi}(\Psi_b)$ instead of S_e . The capacitance in the matrix domain (C_{mi}) is estimated using the same equation as the single domain model, but replacing θ_{sat} by θ_{sat}^{fict} . Finally, the following form is used to describe the unsaturated hydraulic conductivity function in the matrix, using K_b as the matching point:

$$K_{mi} = K_b \cdot \left(\frac{S_{mi}}{S_{mi}(\Psi_b)} \right)^l \cdot \left[\frac{1 - (1 - S_{mi}^{1/m})^m}{1 - (1 - S_{mi}(\Psi_b)^{1/m})^m} \right]^2 \quad (5.38)$$

where $l = 0.5$ is the tortuosity factor. For water potentials less negative than Ψ_b the hydraulic conductivity is set to K_b .

Macropore domain

In the macropore domain, the water potential is not defined and flow equations are expressed in θ -based form. The degree of saturation in the macropores (S_{ma}) is related to soil moisture in the macropores (θ_{ma}) via:

$$S_{ma} = \frac{\theta_{ma}}{e_{ma}} \quad (5.39)$$

where $e_{ma} = \theta_{sat} - \theta_b$. The hydraulic conductivity function in the macropores (K_{ma}) is given as a simple power law of the degree of saturation:

$$K_{ma} = (K_{sat} - K_b) \cdot S_{ma}^{n_{ma}} \quad (5.40)$$

where $n_{ma} = 2.23$ is a kinematic exponent reflecting macropore size distribution and tortuosity. Unlike in MACRO (Jarvis et al., 1991), the fact that macroporosity and macropore maximum hydraulic conductivity may not be constant but vary due to swelling and shrinkage is not taken into account.

Lateral flows

In the beginning, or at any given step, if $\theta_{mi} > \theta_b$, the excess of water is transferred from the matrix to the macropore domain, i.e. it will be used as a source to increase θ_{ma} . However, the macropores have a finite storage capacity for water, C_{ma} , equal to:

$$C_{ma} = (1 - S_{ma}) \cdot (\theta_{sat} - \theta_b) \quad (5.41)$$

Hence, the flow to the macropore domain is limited by C_{ma} . The reverse flow, i.e. the water uptake by the unsaturated matrix is treated as a first-order approximation to a diffusion equation, assuming a rectangular-slab geometry for the aggregates:

$$U_w = \left(\frac{G_w \cdot D_w \cdot \gamma_w}{d^2} \right) \cdot (\theta_b - \theta_{mi}) \quad (5.42)$$

where d (in m) is an effective diffusion path length, D_w (in $m^2 s^{-1}$) is an effective water diffusivity that depends on S_{ma} , $G_w = 3$ is a geometry factor and $\gamma_w = 0.1$ is a scaling factor. See details in (Larsbo et al., 2005). Water uptake by the unsaturated matrix is treated as a source for the matrix domain and as a sink for the macropore domain.

5.7.5 Finite approximation to flows in the soil matrix

Water movement in the soil matrix was taken from chapter 8 of Bonan (2019), and much of the present text has been summarized and adapted from this source, with modifications for the exchange with macropore moisture. The finite difference approximation for the Richards equation represents the soil as a network of N discrete nodal points that vary in space and time. The network is presented in Fig. 5.7.




Figure 5.7: Schematic representation of network of soil nodes

For reasons of numerical stability, Richards equation is solved using an implicit time discretization in which the spatial derivatives $\delta K/\delta z$, $\delta \Psi/\delta z$ and $\delta^2 \Psi/\delta^2 z$ are written numerically using a central difference approximation at time $n + 1$, and the time derivative uses a backward difference approximation at $n + 1$.

Soil layer i has a thickness Δz_i . Water content θ_i , matric potential Ψ_i and hydraulic conductivity K_i are defined at the center of the layer at depth z_i and are uniform over the layer. Here K_i values have been corrected by λ_i , to account for the fact that the amount of stones in the soil reduce the conductivity, which is expressed per soil area unit (so, in fact, all terms in the mass balance equations are multiplied by λ_i , except the source/sinks). Water inputs/outputs from other processes (i.e. infiltration, snow melt, bare soil evaporation, herbaceous layer transpiration and plant uptake/redistribution) are represented as a source/sink term S_i for each node i . If simulations are done in the dual-permeability model, S_i can include positive or negative terms due to saturation excess flow into the macropore domain or capillary imbibition from the macropore domain.

For a given intermediate layer $1 < i < N$, the outward flux Q_i between layers i and $i + 1$ is defined using:

$$Q_i = -\frac{K_{i+1/2}}{\Delta z_{i+1/2}}(\Psi_i - \Psi_{i+1}) - K_{i+1/2} \quad (5.43)$$

where $z_{i+1/2} = z_i - z_{i+1}$ and $K_{i+1/2} = 0.5(K_i + K_{i+1})$ replaces the vertically varying hydraulic conductivity within the two layers with an effective conductivity for an equivalent homogeneous medium. The inward flux Q_{i-1} between layers $i - 1$ and i is defined similarly:

$$Q_{i-1} = -\frac{K_{i-1/2}}{\Delta z_{i-1/2}}(\Psi_{i-1} - \Psi_i) - K_{i-1/2} \quad (5.44)$$

where $z_{i-1/2} = z_{i-1} - z_i$ and $K_{i-1/2} = 0.5(K_{i-1} + K_i)$. If we apply the fluxes to the water balance equation, with fluxes and capacitance expressed for time $n + 1$, and considering the S_i source/sink term (in ms^{-1}), we obtain the following Ψ -based form:

$$\frac{\Delta z_i}{\Delta t} C_i^{n+1}(\Psi_i^{n+1} - \Psi_i^n) \lambda_i = \frac{K_{i-1/2}^{n+1}}{\Delta z_{i-1/2}}(\Psi_{i-1}^{n+1} - \Psi_i^{n+1}) - \frac{K_{i+1/2}^{n+1}}{\Delta z_{i+1/2}}(\Psi_i^{n+1} - \Psi_{i+1}^{n+1}) + K_{i-1/2}^{n+1} - K_{i+1/2}^{n+1} + S_i^{n+1} \quad (5.45)$$

For the first layer ($i = 1$) the boundary condition is defined so that there is no inward flux (but a source/sink term is added), which yields the following Ψ -based form:

$$\frac{\Delta z_1}{\Delta t} C_1^{n+1}(\Psi_1^{n+1} - \Psi_1^n) \lambda_1 = -\frac{K_{1+1/2}^{n+1}}{\Delta z_{1+1/2}}(\Psi_1^{n+1} - \Psi_2^{n+1}) - K_{1+1/2}^{n+1} + S_1^{n+1} \quad (5.46)$$

Finally, for the last layer ($i = N$) a free drainage condition is specified as boundary condition, where $Q_N = K_N$, which yields the following Ψ -based form:

$$\frac{\Delta z_N}{\Delta t} C_N^{n+1}(\Psi_N^{n+1} - \Psi_N^n) \lambda_N = \frac{K_{N-1/2}^{n+1}}{\Delta z_{N-1/2}}(\Psi_{N-1}^{n+1} - \Psi_N^{n+1}) + K_{N-1/2}^{n+1} - K_N^{n+1} + S_N^{n+1} \quad (5.47)$$

Alternatively, if a water table depth is supplied and is equal or shallower than soil depth, the model can simulate capillarity rise by changing the lower boundary condition to that of a constant saturated layer (i.e. $\Psi_{N+1} = 0$).

$$\frac{\Delta z_i}{\Delta t} C_N^{n+1}(\Psi_N^{n+1} - \Psi_N^n) \lambda_i = \frac{K_{N-1/2}^{n+1}}{\Delta z_{N-1/2}}(\Psi_{N-1}^{n+1} - \Psi_N^{n+1}) - \frac{K_{N+1/2}^{n+1}}{\Delta z_{N+1/2}}(\Psi_N^{n+1} - 0) + K_{N-1/2}^{n+1} - K_{N+1/2}^{n+1} + S_N^{n+1} \quad (5.48)$$

where $K_{N+1/2} = 0.5(K_N + K_{sat})$. If the water table is shallower than a given layer, it is assumed to be saturated and if not the model simulates water filling until saturation is reached. Gravitational fluxes are set to zero for layers completely below the water table, and in this case deep drainage is evaluated using the water draining from the unsaturated layers.

The numerical form of the Ψ -based Richards equation for a soil with N layers is a tridiagonal system of N equations with N unknown values Ψ at time $n+1$. An additional complexity is that Ψ at $n+1$ also appears through K and C . The solution requires linearizing with respect to Ψ through various methods. Here the **predictor-corrector** method is employed, a two-step solution that solves the equation twice (Haverkamp et al., 1977). The predictor step uses explicit linearization to solve for Ψ over one-half of a full time step ($\Delta t/2$) at time $n+1/2$ with K and C from time n . The resulting values for Ψ are used to evaluate K and C at $n+1/2$. The corrector step uses these to obtain Ψ over the full time step (Δt) at $n+1$. An implicit solution is used for the predictor step and the Crank-Nicolson method is used for the corrector step.

5.7.6 Finite approximation to flows in the macropore domain

An implicit solution is used to discretize the local water balance in the macropores of a layer i , using:

$$\frac{\theta_{ma,i}^{n+1} - \theta_{ma,i}^n}{\Delta t} \cdot \lambda_i = \frac{K_{ma,i-1}^{n+1} - K_{ma,i}^{n+1}}{\Delta z_i} + S_i^{n+1} \quad (5.49)$$

since θ_{ma} is related to S_{ma} , we can re-express the previous equation into:

$$S_{ma,i}^{n+1} = S_{ma,i}^n + \frac{\Delta t}{\lambda_i \cdot e_{ma,i}} \cdot \left(\frac{K_{ma,i-1}^{n+1} - K_{ma,i}^{n+1}}{\Delta z_i} + S_i^{n+1} \right) \quad (5.50)$$

In this equation, $K_{ma,i-1}^{n+1}$ represents the conductivity of the layer above for the time step $n+1$, whereas the $K_{ma,i}^{n+1}$ is the conductivity of the current layer for the same time step, hence depending on $S_{ma,i}^{n+1}$. The local water balance is solved for each layer, in turn, from the soil surface downward, assuming a zero flow from the surface $K_{ma,0}^{n+1} = 0$ (infiltration into macropores is given by S_1^{n+1}). Then for any given layer i , $K_{ma,i-1}^{n+1}$ is already defined and a solution for $S_{ma,i}^{n+1}$ can be found using a bisection method (i.e. internal halving). After solving for all layers, if the capacity of macropores is exceeded in any given layer, the excess is added to the macropore storage of the layer above. Excess of the topmost layer results in runoff via saturation excess.

Like in the matrix domain, if the water table is shallower than a given layer, it is assumed to be saturated and if not the model simulates macropore water filling until saturation is reached. Gravitational fluxes are set to zero for layers completely below the water table, and in this case macropore drainage is evaluated using the water draining from the unsaturated layers.

Chapter 6

Plant transpiration, photosynthesis and drought stress

This chapter describes how daily plant cohort transpiration ($Tr_{cohorts}$) is modelled in the basic water balance model, completing the flows needed to evaluate eq. (3.6). The model uses the approach of BILJOU (Granier et al., 1999) to estimate plant cohort maximum transpiration, and then divides this value among the contribution of plant cohorts and estimate actual transpiration depending on soil moisture and cohort functional parameters, which determine plant drought. The chapter also describes how is daily photosynthesis estimated, a process that in this model does not influence transpiration but is necessary for growth calculations (chapter 15). Finally, we include here the description of the estimation of plant water status, the effect of drought stress and how plants recover from drought legacies.

6.1 Plant transpiration and photosynthesis

6.1.1 Maximum transpiration

Potential evapotranspiration (PET , in mm) is assumed to represent open water evaporation potential. Maximum canopy transpiration $Tr_{cohorts,max}$ depends on both PET and the amount of transpiring surface, i.e. the current cumulative LAI of the vegetation, LAI_{act} . To estimate $Tr_{cohorts,max}$ the model uses the empirical equation of Granier et al. (1999), where $Tr_{cohorts,max}/PET$ is a function of LAI_{act} :

$$\frac{Tr_{cohorts,max}}{PET} = -0.006 \cdot (LAI_{act})^2 + 0.134 \cdot LAI_{act} + 0.036 \quad (6.1)$$

This equation has already been adopted for Mediterranean biomes (Fyllas and Troumbis, 2009; Ruffault et al., 2013).

Since medfate's **ver. 2.1**, empirical coefficients can be species-specific and Granier et al. (1999) empirical equation has been generalized to:

$$\frac{Tr_{cohorts,max}}{PET} = T_{max,LAI} \cdot LAI_{act} + T_{max,sqLAI} \cdot (LAI_{act})^2 \quad (6.2)$$

where $T_{max,LAI}$ and $T_{max,sqLAI}$ are species-specific parameters (if missing, they are given default values 0.134 and -0.006, respectively).

The maximum transpiration for a given plant cohort i is calculated using the portion of Tr_{max} defined by f_i the fraction of total absorbed SWR that is due to cohort i (see section 4.2):

$$Tr_{max,i} = Tr_{cohorts,max} \cdot \frac{f_i^{0.75}}{\sum_j f_j^{0.75}} \quad (6.3)$$




Figure 6.1: Empirical relationship between $Tr_{cohorts,max}/PET$ and LAI_{act}

Before **ver 2.5.0** $Tr_{max,i}$ was linearly related to f_i , but an exponent 0.75 is now used to decrease *ad hoc* the proportion of transpiration corresponding to cohorts absorbing high SWR fractions with respect to those absorbing low amounts of SWR, which has been found to better match the transpiration predicted using the advanced transpiration model. Note that the actual amount of radiation energy absorbed by plants is much more difficult to determine, involving diffuse and direct radiation, as well as net long-wave radiation (see chapter 9).

6.1.2 Actual plant transpiration and plant water potential

Actual plant transpiration depends on soil moisture and is calculated for each plant cohort and each soil layer separately. $Tr_{i,s}$ (in mm) represents the transpiration made by cohort i from layer s . Actual plant transpiration from a given layer is regulated by soil moisture and the resistance to water flow through the plant. For each plant cohort i and soil layer s , the model first estimates the a whole-plant relative water transpiration, $K_{i,s}$, which varies between 0 and 1 depending on $\Psi_{extract,i}$, the potential at which transpiration is 50% of maximum, and Ψ_s , the water potential in layer s . This relationship is modelled using a Weibull function (see function `hydraulics_psi2K()`):

$$K_{i,s} = K_i(\Psi_s) = \exp \left\{ \ln(0.5) \cdot \left[\frac{\Psi_s}{\Psi_{extract,i}} \right]^{c_{extract,i}} \right\} \quad (6.4)$$

where $c_{extract,i}$ is an exponent that modulates the steepness of the decrease in relative conductance when soil potential becomes negative and $\ln(0.5)$ is used to ensure that $K_i(\Psi_{extract,i}) = 0.5$ (Fig. 6.2).

If cavitation has occurred in previous steps then the capacity of the plant to transport water is impaired. Hence $K_{i,s}$ from eq. (6.4) cannot be larger than the one-complement of the degree of conductance loss in the stem ($PLC_{stem,i}$), so that:

$$K_{i,s}^{PLC} = \min\{K_{i,s}, 1.0 - PLC_{stem,i}\} \quad (6.5)$$

where $PLC_{stem,i}$ is the proportion of stem conductance lost for cohort i (see 6.2.2).




Figure 6.2: Whole-plant relative water conductance functions for different water potential values ($c_{extract} = 3$ in all cases)

Until **ver 2.5.0**, actual transpiration of plant cohort i from a given soil layer s , $Tr_{i,s}$, was defined as the product of (Mouillot et al., 2001): the maximum transpiration of the plant cohort ($Tr_{\max,i}$), the relative whole-plant transpiration ($K_{i,s}^{PLC}$) corresponding to the species and water potential in layer s , and the proportion of plant fine roots in layer s , $FRP_{i,s}$:

$$Tr_{i,s} = Tr_{\max,i} \cdot K_{i,s}^{PLC} \cdot FRP_{i,s} \quad (6.6)$$

Since **ver 2.5.0**, this equation has been slightly modified and is now:

$$Tr_{i,s} = Tr_{\max,i} \cdot K_{i,s}^{PLC} \cdot \frac{K_{unsat,s}^{0.5} \cdot FRP_{i,s}}{\sum_r K_{unsat,r}^{0.5} \cdot FRP_{i,r}} \quad (6.7)$$

where $K_{unsat,s}$ is the unsaturated soil conductivity of layer s . This modification was introduced to allow the plant to increase the amount of water drawn from layers with higher levels of moisture in seasons where the usual water pools are depleted. This formulation decreases the differences in soil moisture between soil layers, which in the advanced model is achieved thanks to hydraulic redistribution process.

The total amount of water transpired by plant cohort i , Tr_i (in mm) is the sum of $Tr_{i,s}$ values over all soil layers:

$$Tr_i = \sum_s Tr_{i,s} \quad (6.8)$$

Analogously, the total amount of water transpired by plant cohorts, $Tr_{cohorts}$ (in mm), is the sum of Tr_i values over all plant cohorts:

$$Tr_{cohorts} = \sum_i Tr_i = \sum_s \sum_i Tr_{i,s} \quad (6.9)$$

Assuming no water limitations (i.e. $K_{i,s} = 1$), we have that $Tr_{cohorts} = Tr_{cohorts,\max}$ and $Tr_i = Tr_{\max,i}$. Cohort transpiration and stand transpiration will be lower than maximum values if soil water potential in

any layer is negative enough to cause a significant reduction in whole-plant conductance. At the plant level, the transpiration of a given plant cohort will be lower than that of others if:

- The cohort is under the shade (it reduces f_i and hence $Tr_{\max,i}$)
- The cohort has a lower amount of leaf area (it reduces f_i and hence $Tr_{\max,i}$)
- The soil layers exploited by the cohort have more negative water potentials (it reduces $K_{i,s}$ and $K_{i,s}^{PLC}$).

6.1.3 Plant water potential

The basic water balance model does not allow estimating a water potential drop from soil to the leaf. Moreover, in a multilayered soil it is difficult to know what would be the water potential of the plant. Despite these limitations, a gross surrogate of ‘plant’ water potential ($\Psi_{plant,i}$; in MPa) may be obtained averaging whole-plant relative conductance values across soil layers and inverting the whole-plant relative transpiration function:

$$\Psi_{plant,i} = K_i^{-1} \left(\sum_s K_{i,s} \cdot FRP_{i,s} \right) \quad (6.10)$$

where K_i^{-1} is the inverse of the relative whole-plant transpiration eq. (6.4). The plant water potential is used to determine plant stress levels and cavitation, as explained in 6.2.1 and 6.2.2.

Importantly, changes in $\Psi_{plant,i}$ induce changes in plant water content (see section 10.3), which may induce a release of water in the transpiration stream or an absorption of water from it, which leads to the cohort transpiration, Tr_i , being different from its water extraction from the soil, Ex_i .

6.1.4 Minimum transpiration

The description of actual transpiration in the previous section incorporates the assumption that plants can regulate water losses up to zero transpiration, neglecting water leaks through the leaf cuticle and/or incomplete stomatal closure (Duursma et al., 2018). To avoid this assumption we can estimate minimum transpiration for a given plant cohort, $Tr_{\min,i}$ (in mm), and prevent $Tr_i < Tr_{\min,i}$.

Minimum transpiration flow per leaf area $E_{\min}$ in $mol H_2O \cdot s^{-1} \cdot m^{-2}$ can be calculated if we know (1) the minimum leaf conductance to water vapor ($g_{sw,\min}$; in $mol H_2O \cdot s^{-1} \cdot m^{-2}$); (2) the leaf vapor pressure deficit (VPD_{leaf} ; in kPa); and (3) atmospheric pressure (P_{atm} ; also in kPa):

$$E_{\min} = g_{sw,\min} \cdot \frac{VPD_{leaf}}{P_{atm}} \quad (6.11)$$

In turn, the vapor pressure deficit in the leaf (VPD_{leaf}) is calculated as:

$$VPD_{leaf} = e_{leaf} - e_{air} \quad (6.12)$$

Where e_{air} is the water vapor pressure of the air surrounding the leaf (but beyond the leaf boundary layer) and e_{leaf} is the vapor pressure inside the leaf. While e_{air} can be derived from the weather input, e_{leaf} should be calculated from leaf temperature (T_{leaf}) and leaf water potential (Ψ_{leaf}):

$$e_{leaf}(T_{leaf}, \Psi_{leaf}) = e_{sat}(T_{leaf}) \cdot \exp \left\{ \frac{2.17 \cdot \Psi_{leaf}}{T_{leaf} + 273.15} \right\} \quad (6.13)$$

where $e_{sat}(T)$ is a function giving the saturated vapor pressure for temperature T (see utility functions of the **meteoland** reference manual). Given the simplicity of the basic water balance model, here we make the assumption that $\Psi_{leaf} = \Psi_{plant}$ (from the previous time step). Furthermore, we assume that leaf temperature is coupled with the temperature of the surrounding air. Given the non-linearity of e_{leaf} we average the vapor pressure values provided by $e_{leaf}(T_{\min}, \Psi_{plant})$ and $e_{leaf}(T_{\max}, \Psi_{plant})$.

Once the leaf level $E_{\min}$ is known for a give cohort i , we can scale this instantaneous rate to daily transpiration (in mm) at the cohort level using:

$$Tr_{\min,i} = E_{\min,i} \cdot 24 \cdot 3600 \cdot LAI_{act,i} \cdot 0.018 \quad (6.14)$$

6.1.5 Transpiration with water pools

Considering water pools (i.e. setting `rhizosphereOverlap = "partial"` or `rhizosphereOverlap = "total"` in the control parameters) involves partitioning the stand area into fractions corresponding to the abundance of each plant cohort. More specifically, the model defines as many plant water pools as plant cohorts, with proportions defined by their LAI values. For example, the fraction corresponding to plant cohort j would be:

$$f_{pool,j} = \frac{LAI_{live,j}}{LAI_{live}} \quad (6.15)$$

Overall soil moisture is then related to moisture of plant water pools by the following equation, that is fulfilled for each soil layer s :

$$W_s = \sum_j f_{pool,j} \cdot W_{j,s} \quad (6.16)$$

where $W_{j,s}$ is the soil moisture (relative to field capacity) of layer s in the water pool of cohort j , i.e. under the area of the stand attributed to cohort j .

When transpiration occurs and water pools are considered, the model assumes that the rhizosphere of each plant cohort occupies its own water pool, but may enter into the water pools under other plant cohorts if the overlap is partial. In other words, the root systems of different cohorts may overlap horizontally. A given plant cohort i will have its roots in layer s partitioned among different water pools. Let us assume that we know $fr_{i,s,j}$, the (horizontal) proportion of fine roots of cohort i in layer s of the water pool j (see section 2.4.3.5). These proportions sum one when summed across water pools:

$$\sum_j fr_{i,s,j} = 1 \quad \forall i, s \quad (6.17)$$

The proportion of fine roots in a given water pool, with respect to the whole root system, will then be equal to:

$$FRP_{i,s,j} = fr_{i,s,j} \cdot FRP_{i,s} \quad (6.18)$$

Estimating transpiration with water pools is analogous to estimating transpiration without them, but replacing the role of $FRP_{i,s}$ by $FRP_{i,s,j}$, and $K_{i,s}$ by $K_{i,s,j}$, in eq. (6.7), which leads to the estimation of $Tr_{i,s,j}$. Analogous replacements have to be made when estimating the plant water potential in eq. (6.10).

Finally, note that the amount of water extracted from the soil layer s of given water pool j (i.e. from $W_{j,s}$) will not be $Tr_{i,s,j}$ (whose units are referred to the overall stand) but $Tr_{i,s,j}/f_{pool,j}$.

6.1.6 Transpiration of hemi-parasitic plants

When mistletoes are considered in water balance simulations, they are included in radiation extinction calculations. The maximum transpiration of mistletoe in cohort i is ($Tr_{mistletoe,i,max}$):

$$Tr_{mistletoe,i,max} = PET \cdot (T_{max,LAI} \cdot LAI_{act} + T_{max,sqLAI} \cdot (LAI_{act})^2) \cdot \frac{f_i^{0.75}}{\sum_j f_j^{0.75}} \cdot \frac{LAI_{mistletoe,i}}{LAI_{act,i} + LAI_{mistletoe,i}} \quad (6.19)$$

Maximum transpiration is then reduced to actual transpiration (i.e. $Tr_{mistletoe,i}$) according to the water potential of the plant cohort ($\Psi_{plant,i}$), following a sigmoid function with parameters $\Psi_{50,gs,bald,mistletoe}$ and $slope_{gs,bald,mistletoe}$ (defined in control parameters). Soil water extraction corresponding to mistletoe transpiration is added to extraction caused by the corresponding plant cohort.

6.1.7 Plant photosynthesis

Because it is useful for growth, and for compatibility with the ‘Sperry’ transpiration mode (chapter 7), the water balance model using ‘Granier’ transpiration mode also calculates gross assimilated carbon. Gross

photosynthesis for a given plant cohort i (in $gC \cdot m^{-2}$) is estimated as a function of transpiration:

$$A_{g,i} = Tr_i \cdot WUE_{\max,i} \cdot (L_i^{PAR})^{WUE_{PAR,i}} \cdot (1 - e^{WUE_{CO2,i} \cdot C_{air}}) \cdot VPD^{WUE_{VPD,i}} \quad (6.20)$$

where Tr_i is the transpiration of plant cohort i , $WUE_{\max,i}$ is the maximum water use efficiency of the cohort (in $gC \cdot mm^{-1}$) - a species-specific parameter where WUE assumed to be estimated under conditions of $VPD = 1kPa$, no atmospheric CO_2 limitations and maximum light availability -; L_i^{PAR} is the proportion of photosynthetically-active radiation available at the mid-crown of cohort i , according to the Beer-Lambert extinction model; C_{air} is the atmospheric CO_2 concentration (in ppm); VPD is the vapor pressure deficit (in kPa); $WUE_{CO2,i}$ and $WUE_{VPD,i}$ are species-specific coefficients regulating the influence of CO_2 concentration and VPD on WUE, respectively; and $WUE_{PAR,i}$ is a species-specific exponent indicating how relative WUE decreases with lower light levels.

Different effects are taken into account in the estimation of gross photosynthesis (eq. (6.20)). The dependency of photosynthesis on drought stress is represented in the Tr_i term, whereas the dependency on CO_2 concentration is given by the negative exponential term and the progressive reduction of photosynthesis under higher VPD (assuming a reduction of stomatal conductance but the same water flow) is modelled using negative $WUE_{VPD,i}$ exponents. Finally, the reduction of photosynthesis due to low light availability (for cohorts in the understory) is represented by the reduction of relative WUE given by low values of L_i^{PAR} :




Figure 6.3: Relationship between the percentage of PAR available to a cohort and the water use efficiency (left) or the water use efficiency relative to the maximum value observed in the stand for the same species (right). WUE values were estimated using the advanced water balance model and a non-linear regression was fitted.

6.2 Plant drought stress and cavitation

6.2.1 Daily drought stress

Similarly to Mouillot et al. (2002), daily drought stress of a given plant cohort i , DDS_i , is defined as the one-complement of relative whole-plant transpiration (see eq. (6.4)) estimated from its plant water potential ($\Psi_{plant,i}$):

$$DDS_i = \phi_i \cdot (1 - K_i(\Psi_{plant,i})) \quad (6.21)$$

Leaf-phenological status is included (only in winter-deciduous or winter semi-deciduous plants) to avoid drought stress during leafless periods. Daily drought stress values can be later used to define other drought stress indices for larger temporal scales.

6.2.2 Cavitation and hydraulic recovery

The water balance model is normally run assuming that although soil drought may reduce transpiration, embolized xylem conduits are automatically refilled when soil moisture recovers (in other words, cavitation is reversible and `stemCavitationRecovery` = "total"). It is possible to simulate irreversible cavitation by setting `stemCavitationRecovery` = "none", `stemCavitationRecovery` = "rate" or `stemCavitationRecovery` = "annual" in the control parameters. Any of these options cause the model to keep track of the maximum value of cavitation so far experienced using the proportion of lost conductance for leaves and the stem the plant cohort i ($PLC_{leaf,i}$ and $PLC_{stem,i}$), which are modelled from Ψ_{plant} using Weibull functions:

$$PLC_{leaf,i} = \max \left\{ PLC_{leaf,i}, 1 - \exp \left\{ \left[\frac{\Psi_{plant,i}}{d_{leaf}} \right]^{c_{leaf}} \right\} \right\} \quad (6.22)$$

$$PLC_{stem,i} = \max \left\{ PLC_{stem,i}, 1 - \exp \left\{ \left[\frac{\Psi_{plant,i}}{d_{stem}} \right]^{c_{stem}} \right\} \right\} \quad (6.23)$$

For simulations of less than one year one can use `stemCavitationRecovery` = "none" to keep track of the maximum cavitation. However, for simulations of several years, it is normally advisable to allow recovery. If `stemCavitationRecovery` = "annual", $PLC_{leaf,i}$ and $PLC_{stem,i}$ values are set to zero at the beginning of each year, assuming that embolized plants overcome the conductance loss by creating new xylem tissue. Finally, if `stemCavitationRecovery` = "rate" the model simulates stem refilling at daily steps as a function of symplasmic water potential. First, a daily recovery rate (r_{refill} ; in $cm^2 \cdot m^{-2} \cdot day^{-1}$) is estimated as a function of $\Psi_{plant,i}$:

$$r_{refill}(\Psi_{plant,i}) = r_{max,refill} \cdot \max\{0.0, (\Psi_{plant,i} + 1.5)/1.5\} \quad (6.24)$$

Where $r_{max,refill}$ is the control parameter `cavitationRecoveryMaximumRate` indicating a maximum refill rate. The right part of the equation normalizes the water potential, so that $r_{refill} = r_{refill,max}$ if $\Psi_{plant,i} = 0$ and $r_{refill} = 0$ if $\Psi_{plant,i} \leq -1.5MPa$. The proportion of conductance lost in stems and leaves is then updated using:

$$PLC_{leaf,i} = \max\{0.0, PLC_{leaf,i} - (r_{refill}/H_v)\} \quad (6.25)$$

$$PLC_{stem,i} = \max\{0.0, PLC_{stem,i} - (r_{refill}/H_v)\} \quad (6.26)$$

where H_v is the Huber value (sapwood area to leaf area ratio) in units of $cm^2 \cdot m^{-2}$.

6.2.3 Cavitation-induced defoliation and recovery

By default, the water-balance model assumes that high levels of leaf hydraulic impairment will lead to defoliation (although this process can be deactivated if `cavitationInducedDefoliation` = FALSE). When cavitation-induced defoliation is considered, the model assumes that crowns are made of a population of leaves whose hydraulic vulnerability follows a Gaussian distribution centered on the input water potential corresponding to 50% loss of conductance (which is dictated by c_{leaf} and d_{leaf}) and with a known coefficient of variation set to 10 % (by default), a modelling strategy developed by Hervé Cochard. Leaves are hydraulically disconnected and die when their embolism rate exceeds a critical value set to 88 % of percent loss of conductance, by default (see defoliation control parameters and function `hydraulics_proportionDefoliationWeibull()`). Ultimately, the assumed variability of hydraulic vulnerability and the current plant water potential lead to a proportion of cavitation-induced crown defoliation.

In the water-balance model, the proportion of crown defoliation leads to a corresponding decrease LAI_{act} , and recovery of foliage after drought will follow the conduit refilling process associated to decrease in PLC_{leaf} .

In the growth model, the proportion of crown defoliation is used to determine leaf senescence and the recovery of foliage is simulated via explicit leaf growth (see chapter 17).

Part III

Advanced water balance modelling

Chapter 7

Advanced water balance model

This chapter provides an overview of the advanced water and energy balance model available in **medfate**. It shares some characteristics with the basic water balance model described in previous chapters. Like the previous one, the model is run using function `spwb()`, for a set of days, or function `spwb_day()`, for a single day. Compared to the basic model, the advanced water balance model adds more detail in some processes (radiation transfer, plant hydraulics and photosynthesis) and incorporates other processes that were missing (canopy/soil energy balances and plant water balance). Importantly, two alternative approaches are possible regarding *plant hydraulics* and *stomatal conductance* sub-models:

- If `transpirationMode = "Sperry"`, the sub-models of plant hydraulics and stomatal conductance follow Sperry et al. (2017) and were described in detail in De Cáceres et al. (2021).
- If `transpirationMode = "Sureau"`, the sub-models of plant hydraulics and stomatal conductance follow SurEau-ECOS and were described in Ruffault et al. (2022).

This chapter presents an overview of the advanced water balance model and is structured as in chapter 3. Process formulations shared with the basic water balance model were presented in previous chapters and will not be repeated. The chapters following this one provide a detailed description of those processes that are represented when using the advanced model only. Whenever necessary, we will detail the difference between the two sub-models of plant hydraulics and stomatal conductance corresponding to De Cáceres et al. (2021) and Ruffault et al. (2022). We recommend readers to familiarize themselves with basic water balance modelling processes before reading this and the subsequent chapters. In addition, users of this model should have enough background on the processes implemented to fully understand and exploit model outputs. Since the scope of this book is focused on the model description, interested readers should address relevant eco-physiological bibliography complement the material exposed here.

7.1 About Sureau

SurEau is a comprehensive plant hydraulic model that already knows different versions. The first and most detailed version was developed in C by Hervé Cochard (Sureau.c, Cochard et al. (2021)), which follows a preliminary version coded in Excell and in R that has been used to understand the stomatal functioning of plants around the world (Martin-StPaul et al., 2017).

In Sureau.c, the plant is relatively well detailed (large roots, trunk, branches, leaves, buds and even flowers). The numerical resolution of the calculations is done according to an explicit scheme, where the elementary transfers of water between each compartment must be calculated at very small time-steps to avoid numerical instabilities known as the Courant–Friedrichs–Lewy condition (for a tree time step must be around 0.001s). This complexity is an advantage for exploring questions about the physiological functioning of the tree. However, the computational demand is also a limit for large-scale application of the model and for integration in dynamic vegetation or land surface models.

Researchers from INRAE-URFM (François Pimont, Julien Ruffault, Nicolas Martin-StPaul at Avignon) and INRAE-PIAF (Hervé Cochard, at Clermont-Ferrand) therefore joined forces to develop a new version, coded in R native, where the numerical scheme for solving water transport calculations was improved (SurEau-Ecos, Ruffault et al. (2022)). They implemented two alternative resolution schemes (the so called semi-implicit and implicit) allowing to solves the water transport accounting for capacitance and cavitation at much larger time steps (of the order of 10 to 30 minutes). In this version the authors also used a tree simplified to only two organs: leaves and stems (and therefore four compartments with an apoplasm and a symplasm for each organ) to facilitate parameterization. This advance has been decisive for **medfate** because it made the model compatible in terms of plant structure and and computational demand. In **medfate** we refer to **Sureau** when this model is used for the computation of transpiration and plant water related processes.

7.2 Design principles

The advanced model performs soil/plant water balance and soil/canopy energy balance for a given forest stand and for the period corresponding to input weather data. Some components of the water balance (rainfall interception, soil evaporation, snow melt, etc) are calculated on a daily step basis whereas energy balances and plant transpiration are determined at subdaily steps (typically 1h steps). The model considers only the vertical spatial dimension of the stand, and not the horizontal distribution of plants within it. Still, the forest stand is divided into: (a) canopy vertical layers; (b) living plant cohorts of different species (and maybe a collective herbaceous layer). Like the basic model, the soil is discretized into a number of soil layers, and the soil water balance, including water movement in the bulk soil, is addressed in daily substeps.

Hydrological processes are the same as in chapter 3. Thus, the model includes water interception loss (Gash et al., 1995; Liu, 2001), plant transpiration, evaporation from soil (Ritchie, 1972), transpiration of the herbaceous layer, infiltration into the soil matrix (Boughton, 1989; Green and Ampt, 1911) and vertical water movement within the soil, the latter following either a single-domain model (Bonan, 2019) or a dual-permeability model (Jarvis et al., 1991; Larsbo et al., 2005), where total porosity is partitioned into two separate flow regions (matrix and macropores), each with its own degree of saturation, conductivity and flux rate.

A great deal of the complexity of the advanced model concerns the estimation of plant cohort transpiration and photosynthesis. The model simulates stomatal regulation of gas exchange separately for *sunlit* and *shade* leaves at sub-daily steps, involving detailed calculations of plant hydraulics, leaf energy balance and photosynthesis. Compared to the basic water balance model of chapter 3, the higher level of complexity of the advanced model allows a more precise estimation of carbon and water flows, as well as a more nuanced evaluation of plant water status, at the expense of a more complex parametrization.

To calculate radiation balance and incident PAR at the leaf level, the leaf area of each plant cohort is divided into vertical layers according to its crown limits. Photosynthesis and transpiration of sunlit and shade leaves are treated separately (De Pury and Farquhar, 1997). Given an input direct and diffuse short-wave irradiance at the top of the canopy, the amount of light absorbed by sunlit leaves, shade leaves and the soil follows the radiation transfer model of Anten and Bastiaans (2016). The scheme for long-wave radiation balance between the atmosphere, the different canopy layers and the soil follows the SHAW model (Flerchinger et al., 2009).

Energy balances are conducted at the *stand* scale and the *leaf* scale:

- At the *stand* scale, the model keeps track of temperature variation within the canopy (i.e. canopy energy balance) and in the uppermost soil surface (i.e. soil energy balance) as the result of energy exchanges between them and with the atmosphere. Heat conduction also occurs among soil layers. Energy balance can be performed assuming a single canopy layer, with soil and canopy energy balance equations being similar to those of Best et al. (2011) for JULES. Alternatively, energy balance can be conducted for each vertical layer following Bonan et al. (2014). In this case, the model allows considering within-canopy gradients in temperature, water vapor pressure, and CO_2 as well as the turbulent transport of heat, water vapor and CO_2 between canopy layers (Ma and Liu, 2019), determined using a canopy turbulence model by Katul et al. (2004).

- At the *leaf* scale, radiation inputs include shortwave radiation from the atmosphere absorbed by the leaf and the net balance of long-wave radiation. Leaf temperature is determined assuming that the temperature of the surrounding air is that of the canopy (or canopy layer). After determining stomatal regulation, the model up-scales the transpiration flux to the canopy (or canopy layer) scale and the corresponding latent heat is used to complete the calculation of the energy balance at the canopy (or canopy layer) scale. Latent heat fluxes from evaporation from the soil and evaporation of intercepted rainfall are also included in the canopy/soil energy balances.

Plant hydraulics and stomatal regulation follow different designs depending on the sub-model employed:

- If `transpirationMode = "Sperry"`, plant hydraulics follows the steady-state approach of Sperry et al. (1998) model, and stomatal regulation follows the profit maximization approach of Sperry et al. (2017).
- If `transpirationMode = "Sureau"`, plant water compartments are always included, and water flows follow Darcy's law, as described in Ruffault et al. (2022). In this case, stomatal regulation follows a Baldocchi-type or Jarvis-type model.

Like in the basic water model, the advanced model assumes by default that soil moisture under all plant cohorts is the same (i.e. vertical soil layers are shared among cohorts). However, variations in soil moisture beneath plant cohorts (and, implicitly, horizontal variation of soil moisture) can be simulated. In this case, hydrological processes are replicated for the fraction of soil corresponding to each cohort. Transpiration of each plant cohort depends on the soil moisture beneath itself and, depending on root overlap, on the soil moisture beneath other plant cohorts.

Optionally, the model can simulate additional transpiration caused by hemi-parasitic plants (mistletoe) attached to specific plant cohorts, as done in the basic water model. This causes water uptake from the corresponding soil water pools to be higher than cohort transpiration. The model does not try to simulate mistletoe hydraulics, because the focus is on estimating the effect of mistletoe transpiration on the host plant cohort. Mistletoe hydraulic conductance (or water potential drop) is not explicitly represented. Instead, the water potential driving transpiration in mistletoe is assumed to be *equal to the leaf water potential of the host plant*. In other words, it is assumed that the drop in water potential in mistletoe branches and leaves is comparable to the drop from the mistletoe branch insertion point to the host leaf (but see different water potential values in Sancho-Knapik et al. (2026)). Another limitation is that mistletoe transpiration flow is not extracted from plant tissues directly, as in Zweifel et al. (2012), but added to the water uptake from the soil layers (or soil water pools) where the plant is connected. Therefore, increased stress effects on the host plant are better represented when soil water pools are considered.

7.3 State variables

The following are state variables in the model under all simulations:

- Cumulative degree days (a) to budburst, (b) to complete unfolding or (c) to senescence ($S_{eco,d}$, $S_{unf,d}$ or $S_{sen,d}$; all in $^{\circ}C$), are tracked by the model to determine leaf phenological status (see 4.1).
- Daily soil moisture content dynamics on each layer s are tracked using $W_s = \theta_s(\Psi_s)/\theta_{fc,s}$, the *proportion of soil moisture in relation to field capacity*, where moisture at field capacity, $\theta_{fc,s}$, is assumed to correspond to $\Psi_{fc} = -0.033$ MPa. Note that W_s values larger than one are possible if the soil is between field capacity and saturation.
- The air temperature within the canopy (T_{can} ; $^{\circ}C$) and the temperature of each soil layer s ($T_{soil,s}$; $^{\circ}C$) are tracked for every subdaily step.
- Leaf area index (LAI_i) of each plant cohort i will change during simulations according to phenology if the species is winter-deciduous.

Additional state variables depend on the activation of specific control flags (see subsection 7.6.3):

- If `transpirationMode = "Sureau"`, the model also tracks $\Psi_{apo,stem,i}$, the water potential of the stem apoplastic compartment, as well as the water potential of stem and leaf symplastic compartments,

- $\Psi_{symp,stem,i}$ and $\Psi_{symp,leaf,i}$ respectively.
- If *stem cavitation is not completely reversible*, the model tracks $PLC_{stem,i}$ and $PLC_{leaf,i}$, the proportion of conductivity lost the stem and leaves, respectively, of plants of cohort i .
- If *plant water pools are considered*, the model also tracks $W_{i,s}$, the proportion of soil moisture in relation to field capacity for layer s within the areal fraction of stand covered by cohort i .
- If *multi-layer canopy energy balance is activated*, the model tracks for each vertical layer $j \in \{1, l\}$:
 - Air temperature, $T_{air,j}$
 - Water vapor pressure, $e_{air,j}$
 - CO_2 concentration, $C_{air,j}$.
- If *cavitation-induced defoliation is considered* (i.e. `cavitationInducedDefoliation = TRUE`), then leaf area index (LAI_i) will become a state variable for all plant cohorts, including evergreen species (see 14.5).

7.4 Water and energy balances

7.4.1 Water balance

The water balance equations for the soil and snowpack of the advanced model are essentially the same as introduced in section 3.3, with some minor modifications. Although the advanced water balance model includes hydraulic redistribution, this does not change the total amount of water in the overall soil, so the water balance equation is very similar to that of the basic water balance model (eq.(3.6)), but where plant cohort water uptake ($Ex_{cohorts}$) is used instead of transpiration ($Tr_{cohorts}$).

$$\Delta V_{soil} = (If + Lw + Cr) - (Dd + Se + Es + Tr_{herb} + Ex_{cohorts}) \quad (7.1)$$

where all the other terms were defined in 3.3. Soil water extraction ($Ex_{cohorts}$) and plant transpiration ($Tr_{cohorts}$) will be equal if plant capacitance effects or transpiration by hemi-parasitic (mistletoe) plants ($Tr_{mistletoe}$) are not considered. More generally, the amount of water stored in the plant tissues changes according to:

$$\Delta V_{plant} = Ex_{cohorts} - Tr_{cohorts} - Tr_{mistletoe} \quad (7.2)$$

Variations of the water balance can occur depending on specific control flags, as explained in section 3.3. Unlike the basic water balance, in this model water balance components contributing to latent heat of vaporization (plant transpiration, evaporation from soil surface and rainfall intercepted by canopy surfaces) or latent heat of fusion (snow melt) are included in the energy balance equations of the canopy and soil, therefore coupling water and energy balances.

7.4.2 Single-layer canopy energy balance

For the sake of performing energy balance, the canopy can be considered a single layer that exchanges energy with the soil and the atmosphere (Best et al., 2011). First, it absorbs shortwave radiation from the atmosphere ($K_{abs,can}$) and long-wave radiation resulting from net radiation balance ($L_{net,can}$). Other energy fluxes considered are convective exchanges between the canopy and atmosphere ($H_{can,atm}$) and between the canopy and the soil ($H_{can,soil}$). Energy is absorbed from the canopy through latent heat of vaporization ($LE_{vap,can}$) produced via transpiration (Tr) and evaporation of water intercepted by the canopy (In); or via latent heat of fusion due to snow melt ($LE_{fus,snow}$). Combining all energy flows (all of them in $W \cdot m^{-2}$), the instantaneous energy balance equation for the canopy is:

$$TC_{can} \cdot \frac{\delta T_{can}}{\delta t} = K_{abs,can} + L_{net,can} - LE_{vap,can} - LE_{fus,snow} - H_{can,atm} - H_{can,soil} \quad (7.3)$$

where TC_{can} is the canopy thermal capacitance (in $J \cdot m^{-2} \cdot K^{-1}$).

7.4.3 Multiple-layer canopy energy balance

Alternatively, the model allows performing an energy balance for each vertical layer in the canopy, including its heat storage capacity (Ma and Liu, 2019). The energy balance equation for canopy layer j is:

$$TC_j \cdot \frac{\delta T_{air,j}}{\delta t} = K_{abs,j} + L_{net,j} - LE_{vap,j} + H_j \quad (7.4)$$

where $K_{abs,j}$ is the short-wave radiation absorbed by leaves in the layer, $L_{net,j}$ is the net long-wave radiation balance of the layer, $LE_{vap,j}$ is the latent heat flux and H_j is the sensible heat flux, which includes heat coming from leaves or soil surfaces as well as turbulent heat transport from adjacent layers. Finally, TC_j is the thermal capacitance (in $J \cdot m^{-2} \cdot K^{-1}$) of canopy layer j . Latent heat of fusion ($LE_{fus,snow}$) needs to be subtracted in the right-hand side of the equation from the bottom canopy layer.

7.4.4 Soil energy balance

Like the canopy, the uppermost soil layer absorbs short-wave radiation from the atmosphere ($K_{abs,soil}$) and has a net long-wave radiation balance ($L_{net,soil}$). The uppermost soil layer also exchanges heat with the canopy ($H_{can,soil}$) and releases energy in the form of latent heat of vaporisation due to evaporation from the soil surface ($LE_{vap,soil}$). Finally, the soil layer exchanges thermal energy by conduction with the layer below ($G_{1,2}$). The energy balance equation for the uppermost soil layer is:

$$TC_{soil,1} \cdot \frac{\delta T_{soil,1}}{\delta t} = K_{abs,soil} + L_{net,soil} + H_{can,soil} - LE_{vap,soil} - G_{1,2} \quad (7.5)$$

where $TC_{soil,1}$ is the thermal capacitance of soil layer 1 (the uppermost layer).

Instantaneous soil temperature changes on each soil layer depend on the balance between upward and downward energy flows. The downward energy flow for the uppermost soil layer depends on its energy exchanges with the canopy and the atmosphere, as shown in eq. (7.5), whereas the remaining energy flows are generated by heat conduction and depend on the temperature gradients between soil layers. The energy balance for a soil layer $s > 1$ is:

$$TC_{soil,s} \cdot \frac{\delta T_{soil,s}}{\delta t} = G_{s-1,s} - G_{s,s+1} \quad (7.6)$$

where $G_{s-1,s}$ is the energy inflow from layers above, $G_{s,s+1}$ is the energy outflow towards layers below and $TC_{soil,s}$ is the thermal capacitance of soil layer s (in $J \cdot m^{-2} \cdot K^{-1}$).

Unlike Best et al. (2011), the model considers the snow pack and the uppermost soil layer as a single layer in terms of energy exchange, but treats this layer differently depending on whether snow is present or not. The snow pack absorbs shortwave radiation (section 5.2) and snow melt is included as latent heat in the energy balance, but the snow layer does not exchange long-wave radiation with the soil nor the atmosphere. $K_{abs,soil}$ is set to zero when snow pack is present (because shortwave radiation is absorbed by snow), but the soil layer still exchanges long-wave radiation and convective energy with the canopy and the atmosphere. In addition, remember that bare soil evaporation does not occur when the snow pack is present (section 5.4).

7.5 Process scheduling

For every day to be simulated, the model performs the following steps:

1. Update leaf area values according to the phenology of species (section 4.1).
2. Determine rainfall interception loss (In) and net rainfall (Pr_{net}) (section 5.3). The amount of water reaching the soil surface from the rainfall event is the sum of net rainfall (Pr_{net}) and surface runoff (Ro), if given. If snow dynamics are included, increase snow pack from snow precipitation (Ps) and decrease it following snow melt (Sm) (section 5.2).
3. Determine bare soil evaporation (Es), if snow is not present (section 5.4), and herbaceous layer transpiration (Tr_{herb} ; section 5.5) if required.

4. Determine sub-daily air temperature and direct/diffuse irradiance variations (chapter 8).
5. Determine the proportion of sunlit/shade leaves (section 9.1.2), the shortwave radiation absorbed by plants (sections 9.1.3) and the soil (section 9.1.4) at sub-daily steps.
6. Set up plant hydraulics (chapter 10). If `transpirationMode = "Sperry"`, this implies building the water supply function of each plant cohort from the current soil moisture levels, whereas if `transpirationMode = "Sureau"` the state of the hydraulic network from the previous day is recovered.
7. Determine instantaneous plant cohort transpiration, photosynthesis rates and close soil/canopy energy balance at sub-daily steps (chapters 12 and 13). This involves the following actions for each sub-daily step:
 - a. Determine long-wave radiation exchange between soil, canopy layers and atmosphere (section 9.2).
 - b. If `transpirationMode = "Sperry"`, for each plant cohort:
 - i. Calculate leaf energy balance and photosynthesis (for shade and sunlit leaves of each plant cohort) corresponding to each transpiration value of the supply function (section 11.1). If plant capacitance effects are considered this concerns the stem-to-leaf hydraulic supply function.
 - ii. Determine stomatal conductance, transpiration and photosynthesis on shade and sunlit leaves of each plant cohort according to Sperry's (2017) profit maximization strategy (section 12.1.3).
 - iii. Update leaf and stem PLC values.
 - iv. If necessary, determine mistletoe transpiration ($Tr_{mistletoe}$) and the corresponding additional soil water uptake (section 12.4).
 - c. If `transpirationMode = "Sureau"`, for each plant cohort leaf energy balance, stomatal and cuticular conductances, transpirational flows, photosynthesis and plant hydraulics are computed iteratively in small temporal sub-steps (e.g. 10 min):
 - i. Perform energy balance of sunlit and shade leaves (section 11.1), according to the energy inputs of the current subdaily time step and the transpiration flux of the previous small sub-step.
 - ii. Calculate shade-leaf, sunlit-leaf and stem cuticular conductances and cuticular transpiration according to current temperature (12.2.2).
 - iii. Calculate stomatal regulation and stomatal transpiration for sunlit and shade leaves (12.2.3).
 - iv. Calculate instantaneous gross and net photosynthesis rates for sunlit and shade leaves.
 - v. Update water potentials, water fluxes (including soil water uptake) and hydraulic conductances across the plant hydraulic network.
 - vi. If necessary, determine mistletoe transpiration ($Tr_{mistletoe}$) and the corresponding additional soil water uptake (section 12.4).
 - d. Scale transpiration ($Tr_{cohorts}$), soil water extraction ($Ex_{cohorts}$) and photosynthesis to the plant cohort level (section 12.3).
 - e. Complete energy balance of the canopy and the soil, after translating plant transpiration, evaporation from bare soil, snow melt and interception loss to latent heat and calculating convective heat exchange for both the canopy and the soil. Alternatively, perform energy balance for individual canopy layers, while including turbulent heat exchange and turbulent transport of water vapor and CO_2 between layers (chapter 13).
8. Determine day-level photosynthesis and transpiration from instantaneous rates and update soil moisture (section 14.1). Determine drought stress for each plant cohort (section 14.3).
9. Determine rainfall infiltration (section 5.6) and vertical water movement within the soil, while considering soil layer source/sinks from infiltration, bare soil evaporation and herbaceous transpiration and plant cohort water uptake/redistribution (section 5.7). This updates soil moisture and lower boundary condition of water movement corresponds to deep drainage (Dd).
10. If fire hazard calculations are requested, estimate live and dead fuel moisture and potential fire behaviour (chapter 26).

Steps 1-3 are the same as described in the basic model, and steps 9-10 here correspond to steps 5-6 in the basic model. The set of steps corresponds to simulations if plant water pools *are not* considered, so that only the water balance of the overall soil matters. If plant water pools *are* considered, steps 1, 2 and 10 are

the same, because they apply to the whole forest stand (note that at present infiltration does not depend on antecedent conditions). However, steps 3 is performed for each water pool separately because it depends on antecedent soil moisture. Steps 6-8 also involve defining plant cohort water uptake from different plant water pools depending on the proportion of fine roots of each plant cohort in each pool (see 6.1.5). Soil water flows from these three steps are then averaged at the stand level. After that, Step 9 is performed on the soil corresponding to each pool, and the resulting drainage estimates are also averaged. Finally, the soil moisture of the overall soil is estimated by averaging the soil moisture of water pools.

7.6 Inputs and outputs

7.6.1 Soil, vegetation and meteorology

Soil

Soil input requirements are the same as for the basic water balance model and were fully described in section 2.3. Genuchten (1980) water retention curves are forced for this model, because rhizosphere conductance is modelled using van Genuchten hydraulic equations (see subsection 10.2).

Vegetation

Vegetation input requirements were fully described in section 2.4. Trees, shrubs and herbs do not need to be characterized with different variables in soil water balance calculations, as long as all of them have a LAI value. In most cases, users only need to estimate the leaf area index corresponding to live leaves, i.e. LAI_{live} , because normally at the starting point all leaves are expanded (i.e. $LAI_{act} = LAI_{live}$) and one can assume no dead leaves in the canopy (i.e., $LAI_{dead} = 0$). Vegetation characteristics stay constant during simulations using function `spwb()`, although the actual expanded leaf area (LAI_{act}) and dead leaf area may vary if the species is winter deciduous. For each plant cohort with hemi-parasitic (mistletoe) plants living on crowns, a corresponding value of mistletoe leaf area index ($LAI_{mistletoe}$) can be defined.

Meteorology

The minimum weather variables required to run the model are min/max temperatures (T_{min} and T_{max}), min/max relative humidity (RH_{min} and RH_{max}), precipitation (P) and solar radiation (Rad). Wind speed (u) is also needed, but the user may use missing values if not available (a default value will be used in this case). Wind speed is assumed to have been measured at a specific height above the canopy (by default at 2 m). Atmospheric CO_2 concentration (C_{atm}) may also be specified, but if missing a default constant value is assumed, which is taken from the control parameters. Definitions and units of these variables were given in section 2.5.

7.6.2 Plant functional parameters

The advanced water balance model requires many functional parameters to be specified for plant cohorts. A first set of parameters refers to leaf phenology (`paramsPhenology`):

Symbol	Units	R	Description
		PhenologyType	Phenology type ("evergreen", "winter-deciduous", or "winter-semideciduous").
		SenescencePeriod	Senescence for evergreen plants ("whole-year", "spring" or "autumn").
		GrowthPeriod	Period of primary growth for evergreen plants ("whole-year" or "spring").
		GrowthDeterminacy	Growth determinacy ("determined", "indetermined" or "intermediate"), only relevant for the growth model (see 15.1.3).
<i>LD</i>	years	LeafDuration	Average duration of leaves (in years).
<i>BFD</i>	days	BudFormationDays	Number of days required to complete bud formation (only relevant for the growth model (see 15.1.3).

Symbol	Units	R	Description
S_{eco}^*	$^{\circ}\text{C}$	Sgdd	Cumulative degree days required for leaf budburst (see section 4.1.3).
S_{unf}^*	$^{\circ}\text{C}$	Ugdd	Cumulative degree-days required to complete leaf unfolding after budburst has occurred (see section 4.1.3).
T_{eco}	$^{\circ}\text{C}$	Tbgdd	Base temperature for the calculation of degree days to leaf budburst and unfolding (see section 4.1.3).
S_{sen}^*	$^{\circ}\text{C}$	Ssen	Degree days corresponding to leaf senescence (see section 4.1.4).
F_{sen}^*	$^{\circ}\text{C}$	Fsen	Cumulative degree days corresponding to complete leaf senescence (see section 4.1.4).
Ph_{sen}	hours	Phsen	Photoperiod corresponding to start counting senescence degree-days (see section 4.1.4).
T_{sen}	$^{\circ}\text{C}$	Tbsen	Base temperature for the calculation of degree days to leaf senescence (see section 4.1.4).

A second set of functional parameters relate to light extinction and water interception (**paramsInterception**):

Symbol	Units	R	Description
λ_{leaf}	degrees	LeafAngle	The (average) angle between the leaf plane and the horizontal plane (i.e. leaf zenith angle)
k_{PAR}	(unitless)	kPAR	Extinction coefficient for diffuse PAR
α_{SWR}	[0-1]	alphaSWR	Short-wave radiation leaf absorbance coefficient
γ_{SWR}	[0-1]	gammaSWR	Short-wave radiation leaf reflectance (albedo)
s_{water}	$mm\ H_2O \cdot LAI^{-1}$	g	Crown water storage capacity (i.e. depth of water that can be retained by leaves and branches) per LAI unit

A third set includes parameters related to plant anatomic and morphological attributes (**paramsAnatomy**):

Symbol	Units	R param	Description
H_{med}	cm	Hmed	Median plant height
$1/H_v$	$m^2 \cdot m^{-2}$	A12As	Ratio of leaf area to sapwood area
LW	cm	LeafWidth	Leaf width
SLA	$m^2 \cdot kg^{-1}$	SLA	Specific leaf area
ρ_{leaf}	$g \cdot cm^{-3}$	LeafDensity	Leaf tissue density
ρ_{wood}	$g \cdot cm^{-3}$	WoodDensity	Wood tissue density
$\rho_{fineroot}$	$g \cdot cm^{-3}$	FineRootDensity	Fine root tissue density
SRL	$cm \cdot g^{-1}$	SRL	Specific root length
RLD	$cm \cdot cm^{-3}$	RLD	Fine root length density (i.e. density of root length per soil volume)
$r_{6.35}$		r635	Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm

A fourth set of parameters are related to transpiration and photosynthesis (**paramsTranspiration**):

Symbol	Units	R param	Description
g_{swmin}	$mol\ H_2O \cdot s^{-1} \cdot m^{-2}$	Gwmin	Minimum stomatal conductance to water vapour
g_{swmax}	$mol\ H_2O \cdot s^{-1} \cdot m^{-2}$	Gwmax	Maximum stomatal conductance to water vapour

Symbol	Units	R param	Description
$J_{max,298}$	$\mu\text{mol } e \cdot m^{-2} \cdot s^{-1}$	Jmax298	Maximum rate of electron transport at 298K
$V_{max,298}$	$\mu\text{mol } CO_2 \cdot s^{-1} \cdot m^{-2}$	Vmax298	Rubisco's maximum carboxylation rate at 298K
$K_{stem,max,ref}$	$kg \cdot s^{-1} \cdot m^{-1} \cdot MPa^{-1}$	Kmax_stemxylem	Maximum stem sapwood reference conductivity per leaf area unit
$K_{root,max,ref}$	$kg \cdot s^{-1} \cdot m^{-1} \cdot MPa^{-1}$	Kmax_rootxylem	Maximum root sapwood reference conductivity per leaf area unit
$k_{leaf,max}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	VCleaf_kmax	Maximum leaf conductance (per leaf area unit)
$k_{stem,max}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	VCstem_kmax	Maximum stem conductance (per leaf area unit)
$k_{root,max,s}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	VCroot_kmax	Maximum root conductance (per leaf area unit) for each soil layer
$k_{rhizo,max,s}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	VGrhizo_kmax	Maximum hydraulic conductance of the rhizosphere for each soil layer
c_{leaf}, d_{leaf}	(unitless), MPa	VCleaf_c, VCleaf_d	Parameters of the vulnerability curve for leaves (Weibull)
$\Psi_{50,leaf}, slope_{leaf}$	MPa, %/MPa	VCleaf_P50, VCleaf_slope	Parameters of the vulnerability curve for leaves (Sigmoid)
c_{stem}, d_{stem}	(unitless), MPa	VCstem_c, VCstem_d	Parameters of the vulnerability curve for stem xylem (Weibull)
$\Psi_{50,stem}, slope_{stem}$	MPa, %/MPa	VCstem_P50, VCstem_slope	Parameters of the vulnerability curve for stem xylem (Sigmoid)
c_{root}, d_{root}	(unitless), MPa	VCroot_c, VCroot_d	Parameters of the vulnerability curve for root xylem (Weibull)
$\Psi_{50,root}, slope_{root}$	MPa, %/MPa	VCroot_P50, VCroot_slope	Parameters of the vulnerability curve for root xylem (Sigmoid)
$T_{gsw,optim,jarv}$	°C	Gsw_Toptim_Jarvis	Temperature corresponding to maximal stomatal conductance (Jarvis stomatal sub-model)
$T_{gsw,sens,jarv}$		Gsw_Tsens_Jarvis	Stomatal sensitivity to temperature (Jarvis stomatal sub-model)
$m_{wue,bald}$		Gsw_AC_slope_Baldocchi	Slope of the Gsw vs Ac/Cs relationship (Baldocchi stomatal sub-model, see 12.2.3)
$\Psi_{50,gsb,bald}$	MPa	Gsw_P50_Baldocchi	Water potential causing 50% reduction in stomatal conductance (Sureau sub-model)
$slope_{gsb,bald}$	%/MPa	Gsw_slope_Baldocchi	Rate of decrease in stomatal conductance at $\Psi_{50,gsb,bald}$ (Baldocchi stomatal sub-model)

It is important to mention that stem- and root-level hydraulic conductances (i.e. $k_{stem,max}$ and $k_{root,max}$) are not given as inputs by the user, but are the result from scaling tissue-level conductivity parameters (i.e. $K_{stem,max,ref}$ and $K_{root,max,ref}$) according to plant size (see A.4.1 and A.4.3 for details). Similarly, parameters of Weibull vulnerability curves are estimated from the input in form of Ψ_{50} , Ψ_{12} and Ψ_{88} available in SpParams. The *slope* of the Sigmoid vulnerability curves can also be estimated from the same values, or provided by the user directly as a column in SpParams.

A fifth (final) set of parameters are related to water storage and water relations in plant tissues (**paramsWaterStorage**):

Symbol	Units	R	Description
$LFMC_{\max}$	%	maxFMC	Maximum live fuel moisture content, corresponding to fine fuels (< 6.35 mm twigs and leaves).
ϵ_{leaf}	MPa	LeafEPS	Modulus of elasticity of leaves
ϵ_{stem}	MPa	StemEPS	Modulus of elasticity of symplastic xylem tissue
$\pi_{0,leaf}$	MPa	LeafPI0	Osmotic potential at full turgor of leaves
$\pi_{0,stem}$	MPa	StemPI0	Osmotic potential at full turgor of symplastic xylem tissue
$f_{apo,leaf}$	[0-1]	LeafAF	Apoplastic fraction in leaf tissues
$f_{apo,stem}$	[0-1]	StemAF	Apoplastic fraction in stem tissues
V_{leaf}	$l \cdot m^{-2}$	Vleaf	Leaf water capacity per leaf area unit
$V_{sapwood}$	$l \cdot m^{-2}$	Vsapwood	Sapwood water capacity per leaf area unit

Vegetation functional attributes are normally filled for each cohort by function **spwbInput()** from species identity. However, different parameters can be specified for different cohorts of the same species if desired (see section 2.4.4).

7.6.3 Control parameters

Control parameters modulate the overall behaviour of water balance simulations (see section 2.6). There are three **key control parameters**:

1. **transpirationMode** defines the transpiration model. If **transpirationMode** = "Sperry" or **transpirationMode** = "Sureau" then the advanced water balance model is run with the corresponding plant hydraulics and stomatal regulation sub-models. In advanced water balance models, the water retention curve model is forced to be that of Van Genuchten, i.e. **soilFunctions** = "VG" (see section 2.3.2).
2. **multiLayerBalance** is a boolean flag to indicate that multi-layer energy balance is desired (by default **multiLayerBalance** = FALSE).

Other relevant control parameters relevant for the advanced water balance model are:

- **ndailysteps** [= 24]: Number of steps into which each day is divided for determination of soil water balance, stomatal conductance, transpiration and photosynthesis (24 equals 1-hour intervals).
- **nsubsteps_canopy** [= 3600]: Defines the number of substeps for multi-layer energy balance calculations. By default, substeps are 1 sec if **ndailysteps** = 24.
- **soilDomains** [= "buckets"]: Either "buckets" (for multi-bucket model), "single" (for single-domain Richards model) and "dual" (for dual-permeability model). See 5.7.
- **max_nsubsteps_soil** [= 300]: Maximum number of substeps for soil water balance solving.
- **defaultWindSpeed** [= 2.5]: Default value for wind speed (in $m \cdot s^{-1}$) when this is missing (only used for leaf fall, see section 4.1).
- **defaultCO2** [= 386]: Default atmospheric (abovecanopy) CO_2 concentration (in $\mu mol \cdot CO_2 \cdot mol^{-1}$ = ppm). This value will be used whenever CO_2 concentration is not specified in the weather input.
- **defaultRainfallIntensityPerMonth** [= c(1.5, 1.5, 1.5, 1.5, 1.5, 1.5, 5.6, 5.6, 5.6, 5.6, 5.6, 1.5)]: A vector of twelve values indicating the rainfall intensity (mm/h) per month. By default synoptic storms (1.5 mm/h) are assumed between December and June, and convective storms (5.6 mm/h) are assumed between July and November (Miralles et al., 2010).
- **snowpack** [= TRUE]: Whether dynamics of snow pack are included (see section 5.2).
- **leafPhenology** [= TRUE]: Whether leaf phenology is simulated for winter-deciduous species (see section 4.1). If FALSE then all species are assumed to be evergreen.
- **bareSoilEvaporation** [= TRUE]: Whether evaporation from the soil surface is simulated (see section 5.4).

- `interceptionMode` [= "Gash1995"]: Infiltration model, either "Gash1995" or "Liu2001"; see section 5.3.
- `infiltrationMode` [= "GreenAmpt1911"]: Infiltration model, either "GreenAmpt1911" or "Boughton1989"; see section 5.6.
- `sapFluidityVariation` [= TRUE]: Flag to indicate that temperature affects sap fluidity (see 10.2.3).
- `windMeasurementHeight` [= 200]: Distance (in cm) above the top of the canopy corresponding to the measurement of wind speed.
- `boundaryLayerSize` [= 2000]: Size of the canopy boundary layer (in cm). The boundary layer causes the model to define vertical layers above the top of the canopy. The parameter is only relevant when multi-layer canopy energy balance is performed.
- `gCrown0` [= 0.150]: Reference canopy conductance (in $\text{mol} \cdot \text{s}^{-1} \cdot \text{m}^{-2}$).
- `unlimitedSoilWater` [=FALSE]: Boolean flag to indicate the simulation of plant transpiration assuming that soil water is always at field capacity.
- `interceptionMode` [= "Gash1995"]: Infiltration model, either "Gash1995" or "Liu2001".
- `infiltrationMode` [= "Green-Ampt"]: Infiltration model, either "Green-Ampt" or "Boughton".
- `rhizosphereOverlap` [= "partial"]: A string indicating the degree of rhizosphere spatial overlap between plant cohorts:
 - "none" - no overlap (independent water pools).
 - "partial" - partial overlap determined by soil moisture.
 - "total" - total overlap (plants extract from common soil pools).
- `thermalCapacityLAI` [= 1000000]: Canopy thermal capacitance per LAI unit (see section 13.1.3).
- `verticalLayerSize` [= 100]: The size of vertical layers (in cm) for leaf area distribution and light extinction calculations (see section 2.4.3.3).
- `segmentedXylemVulnerability` [=TRUE]: If FALSE leaf and root vulnerability curves will be equal to those of stem.
- `leafCavitationRecovery` / `stemCavitationRecovery` [= "rate"]: A string indicating how to model recovery of xylem conductance (see section 14.4). Allowed values are:
 - "none" - no recovery.
 - "annual" - complete recovery every first day of the year.
 - "total" - instantaneous complete recovery.
 - "rate" - following a rate of new leaf or sapwood formation.
- `cavitationRecoveryMaximumRate` [= 0.05]: Maximum rate of conductance recovery (see section 14.4).
- `lfmcComponent` [= "fine"]: Plant component used to estimate LFMC, either "leaf" or "fine" (for fine fuel) (see section 10.3.4).
- `mistletoeParams`: A list with the following elements:
 - `kPAR` [= 0.5]: Mistletoe light extinction coefficient.
 - `g` [= 0.8]: Mistletoe water storage capacity per LAI unit.
 - `Gs_P50_Baldocchi` [= -2.5]: Water potential corresponding to 50% reduction of mistletoe stomatal conductance (i.e. $\Psi_{50_{gs,bald,mistletoe}}$).
 - `Gs_slope_Baldocchi` [= 30]: Rate of decrease in mistletoe stomatal conductance at `Gs_P50` (i.e. $slope_{gs,bald,mistletoe}$).
 - `Gsw_AC_slope_Baldocchi` [=8]: Slope of the `Gsw` vs `Ac/Cs` relationship ($m_{wue,bald,mistletoe}$ of the Baldocchi model).
 - `Vmax298` [= 80]: Maximum Rubisco carboxylation rate at 298K.
 - `Jmax298` [= 120]: Maximum rate of electron transport at 298K.
 - `LeafWidth` [=1]: Mistletoe leaf width (cm).
- `defoliationParams`: A list with the following elements:
 - `cavitationInducedDefoliation` [= TRUE]: Whether leaf hydraulic impairment induces defoliation.
 - `criticalLeafPLC` [= 0.88]: Level of PLC (a proportion) corresponding to 50 % crown defoliation.
 - `cvLeafP50` [= 10.0]: Coefficient of variation (in percent) of leaf P50 within crown leaves.

A set of parameters is only relevant for `transpirationMode = "Sperry"`:

- `leafCavitationEffects / stemCavitationEffects [= FALSE/TRUE]`: A flag indicating whether cavitation effects on conductance of leaves and stem are applied.
- `numericParams`: A list with parameters for numerical approximation routines:
 - `maxNsteps [= 400]` - Maximum number of steps in supply function.
 - `ntrial [= 200]` - Number of iteration trials when finding root of equation system.
 - `psiTol [= 0.0001]` - Tolerance value for water potential.
 - `ETol [= 0.0001]` - Tolerance value for flow.

whereas another set is only relevant for `transpirationMode = "Sureau"`:

- `plantCapacitance [= TRUE]`: Whether the effect of (symplasmic or apoplastic) plant water compartments is considered in simulations.
- `leafCuticularTranspiration [= TRUE]`: Whether the effect of leaf cuticular transpiration is considered in simulations.
- `stemCuticularTranspiration [= FALSE]`: Whether the effect of stem cuticular transpiration is considered in simulations.
- `C_SApoInit [= 2.0e-5]`: Maximum capacitance of the stem apoplasm ($\text{mmol} \cdot \text{m}^{-2}$).
- `C_LApoInit [= 1.0e-5]`: Maximum capacitance of the leaf apoplasm ($\text{mmol} \cdot \text{m}^{-2}$).
- `k_SSym [= 0.26]`: Conductance from stem apoplasm to stem symplasm ($\text{mmol} \cdot \text{s}^{-1} \cdot \text{m}^{-2}$).
- `k_LSym [= 2.0]`: Conductance from leaf apoplasm to leaf symplasm ($\text{mmol} \cdot \text{s}^{-1} \cdot \text{m}^{-2}$).
- `gs_NightFrac [= 0.05]`: Stomatal conductance at night as fraction of maximum stomatal conductance ([0-1]).
- `stomatalSubmodel [= "Baldocchi"]`: Stomatal regulation sub-model, either “Jarvis” or “Baldocchi”.
- `JarvisPAR [= 0.003]`: Parameter regulating the response of stomatal conductance to light (PAR) in the Jarvis model.
- `fTRBToLeaf [= 0.8]`: Fraction of surface of bark exposed to air per leaf area.

Finally, a number of control parameters are used to modify how hydraulic parameters are estimated when calling function `spwbInput()`:

- `taper [= TRUE]`: Whether taper of xylem conduits is accounted for when estimating aboveground stem conductance from xylem conductivity.
- `fracRootResistance [= 0.40]`: Fraction of plant total resistance (leaf+stem+root) that corresponds to root system.
- `averageFracRhizosphereResistance [= 0.15]`: Fraction to total continuum (stem+root+rhizosphere) resistance that corresponds to rhizosphere (averaged across soil water potential values).

7.6.4 Model output

7.6.4.1 Daily output

Function `spwb()` with `transpirationMode = "Sperry"` or `transpirationMode = "Sureau"` returns a list of a class with the same name. The first four elements of this list (i.e., `latitude`, `topography`, `weather` and `spwbInput`) are simply copies of model inputs. The next element is `spwbOutput`, which contains the state of the input object at the end of the simulation (this can be used to perform further simulations starting with current values of state variables). The remaining elements are:

Element	Description
WaterBalance	Climatic input and water balance flows (i.e. net precipitation, infiltration, runoff, transpiration...). All of them in $\text{mm} = \text{l} \cdot \text{m}^{-2}$.
EnergyBalance	Energy balance components (instantaneous values) for the soil and the canopy
Temperature	Minimum and maximum temperature values for topsoil, canopy and atmosphere. Only returned if <code>temperatureResults = TRUE</code> .

Element	Description
Soil	Soil variables for each soil layer: Moisture relative to field capacity (W_s), water potential (Ψ_s) and volumetric water content (V_s). Only returned if <code>soilResults = TRUE</code> .
Stand	Stand-level variables, such as LAI_{act} , LAI_{dead} , the water retention capacity of the canopy (S_{canopy}) or the fraction of light reaching the ground (L_{ground}^{PAR} and L_{ground}^{SWR}). Only returned if <code>standResults = TRUE</code> .
Plants	Plant-level variables, such as $LAI_{act,i}$, plant transpiration, plant water potential, etc. Only returned if <code>plantResults = TRUE</code> .
SunlitLeaves	Leaf-level output variables for sunlit leaves. Only returned if <code>leafResults = TRUE</code> .
ShadeLeaves	Leaf-level output variables for shade leaves. Only returned if <code>leafResults = TRUE</code> .
FireHazard	Fire hazard variables (fuel moisture, rate of spread, fire potentials, etc). Only returned if <code>fireHazardResults = TRUE</code> .

Compared to the output of the basic water balance 3.5.4, new output elements are produced. Many results are given as sub-elements of **Plants**, each of them a data frame with dates in rows and plant cohorts in columns:

Element	Symbol	Units	Description
LAI	LAI_{act}	$m^2 \cdot m^{-2}$	Actual leaf area index (expanded).
LAIlive	LAI_{live}	$m^2 \cdot m^{-2}$	Leaf area index (live) without accounting for phenology or drought-induced defoliation.
Phenology			Daily phenological status for each plant cohort (bud formation, bud dormancy, leaf elongation, leaf senescence), codified into an integer value.
AbsorbedSWR		MJ	Shortwave radiation absorbed per ground area.
AbsorbedLWR		MJ	Longwave radiation absorbed per ground area.
Transpiration	Tr	mm	Transpiration per ground area.
GrossPhotosynthesis		$gC \cdot m^{-2}$	Gross photosynthesis per ground area.
NetPhotosynthesis		$gC \cdot m^{-2}$	Net photosynthesis per ground area.
dEdP	$dE/d\Psi_{leaf}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	Derivative of the (aboveground) supply function.
PlantWaterBalance		mm	Internal daily plant water balance (balance of soil extraction and transpiration).
LeafPsiMin	Ψ_{md}	MPa	Minimum (midday) ‘average leaf’ water potential.
LeafPsiMax	Ψ_{pd}	MPa	Maximum (predawn) ‘average leaf’ water potential.
StemPsi	Ψ_{stem}	MPa	Minimum stem water potential.
RootPsi	$\Psi_{rootcrown}$	MPa	Minimum root-crown water potential.
RhizoPsi	$\Psi_{rootsurf,s}$	MPa	Minimum root surface water potential for each soil layer.
LeafPLC	PLC_{leaf}	MPa	Proportion of leaf conductance loss.
StemPLC	PLC_{stem}	MPa	Proportion of stem conductance loss.
LeafRWC	RWC_{leaf}	%	Mean leaf relative water content.
StemRWC	RWC_{stem}	%	Mean stem relative water content.
LFMC	$LFMC$	%	Live fuel moisture content (as percent of dry weight), corresponding to fine fuels (< 6.35 mm twigs and leaves).
PlantStress	DDS	[0-1]	Drought stress level suffered by each plant cohort (relative whole-plant conductance).

The following table describes the child elements of **SunlitLeaves** and **ShadeLeaves**, each of them a data frame with dates in rows and plant cohorts in columns:

Element	Symbol	Units	Description
LeafPsiMin	$\Psi_{md,i}^{sunlit}$, $\Psi_{md,i}^{shade}$	MPa	Minimum (midday) (sunlit or shade) leaf water potential.
LeafPsiMax	$\Psi_{pd,i}^{sunlit}$, $\Psi_{pd,i}^{shade}$	MPa	Maximum (predawn) (sunlit or shade) leaf water potential.
TempMin	$T_{leaf,min}^{sunlit}$, $T_{leaf,min}^{shade}$	°C	Minimum leaf temperature.
TempMax	$T_{leaf,max}^{sunlit}$, $T_{leaf,max}^{shade}$	°C	Maximum leaf temperature.
GSWMin	$g_{w,i}^{sunlit}$, $g_{w,i}^{shade}$	$mol \cdot s^{-1} \cdot m^{-2}$	Minimum stomatal conductance to water vapour.
GSWMax	$g_{w,i}^{sunlit}$, $g_{w,i}^{shade}$	$mol \cdot s^{-1} \cdot m^{-2}$	Maximum stomatal conductance to water vapour.

The output of simulations can be inspected using `plot`, `shinyplot` and `summary` functions specific to `spwb` objects (examples are given in the corresponding package *vignette*).

7.6.4.2 Sub-daily output

Calling function `spwb_day()` with the advanced water balance model results in sub-daily results for the simulated date. When running the advanced water balance model with function `spwb()` one can request sub-daily results to be included with the function output. This is done via setting `subdailyResults = TRUE` in the control parameter list. In that case, the output list of `spwb()` will include an element `subdaily`, whose sub-elements correspond to the results of `spwb_day()` for each date simulated. Refer to the documentation of functions `spwb_day()` and `transp_transpirationSperry()` or `transp_transpirationSureau()` for details of sub-daily output.

7.7 Applications

The advanced water balance model requires larger parameterization efforts and computational capabilities for large-scale applications. These features make it appropriate for eco-hydrological applications that require high predictive capacity of water flows and water status at the plant or forest stand levels.

The advanced water balance model (Sperry) has been used for the prediction of pre-dawn water potential, in applications focusing on the semi-mechanistic prediction of live fuel moisture content (Balaguer-Romano et al., 2022, 2023). Focusing on plant drought stress, Balaguer-Romano et al. (2025) used advanced water balance model (Sureau) to examine potential hypotheses regarding drought sensitivity of secondary-growth forests. Similarly, Vilà-Vilardell et al. (2023) employed the advanced water balance model (Sperry) to assess the effect of thinning and prescribed burning on drought stress of *Pinus nigra* stands. In a paper employing two different process-based models, Saponaro et al. (2024) studied the combined effect of climatic changes and forest management on carbon and water fluxes in beech forests. Finally, Cabon et al. (2024) used advanced model simulations to obtain estimates of photosynthesis and leaf water potential, and evaluate the relative importance of source and sink limitations to tree growth.

Chapter 8

Sub-daily atmospheric temperature and radiation variation

After estimating leaf phenology (chapter 4) and simulating the main hydrological processes (chapter 5), the advanced water balance model (chapter 7) starts simulating *subdaily* processes, which include energy balances and photosynthesis/transpiration. In order to properly simulate these processes it is important to consider sub-daily variations in temperature and radiation. Here we detail how sub-daily estimates of above-canopy air temperature and atmospheric incoming radiation are derived from daily input values.

8.1 Above-canopy air temperature

Above-canopy air temperature (T_{atm} , in $^{\circ}C$) diurnal variations are modeled assuming a *sinusoidal pattern* with $T_{atm} = T_{min}$ at sunrise and $T_{atm} = (T_{min} + T_{max})/2$ at sunset. Air temperature varies linearly between sunset and sunrise (McMurtrie et al., 1990). Sunrise and sunset hours are determined from latitude and sun declination (see section 4.2 of the reference manual for package **meteoland**).

8.2 Incoming diffuse and direct short-wave radiation

Daily global radiation (Rad , in $MJ \cdot m^{-2} \cdot d^{-1}$) is assumed to include both direct and diffuse short-wave radiation (SWR). Using latitude information and whether is a rainy day, this quantity is partitioned into instantaneous direct and diffuse SWR and PAR for different daily substeps. Values of instantaneous direct and diffuse SWR and PAR above the canopy (I_{beam} and I_{dif} , in $W \cdot m^{-2}$) are calculated using the methods described in Spitters et al. (1986), which involve comparing daily global radiation with daily potential radiation. All these calculations are performed using routines in package **meteoland** (see details in section 4.6 of the reference manual for this package).

8.3 Incoming long-wave radiation

Once the above-canopy air temperature for a given time step is known, instantaneous long-wave radiation (LWR) coming from the atmosphere (L_{atm} , in $W \cdot m^{-2}$) can be calculated following Campbell and Norman (1998):

$$L_{atm} = \epsilon_a \cdot \sigma \cdot (T_{atm} + 273.16)^{4.0} \quad (8.1)$$

where T_{atm} is air temperature, $\sigma = 5.67 \cdot 10^{-8.0} W \cdot K^{-4} \cdot m^{-2}$ is the Stephan-Boltzmann constant and ϵ_a is the emmissivity of the atmosphere, calculated using:

$$\epsilon_a = (1 - 0.84 \cdot c) \cdot \epsilon_{ac} + 0.84 \cdot c \quad (8.2)$$

$$\epsilon_{ac} = 1.72 \cdot \left(\frac{e_{atm}}{T_{atm} + 273.16} \right)^{1/7} \quad (8.3)$$

where e_{atm} is the average daily water vapor pressure (in kPa; see 2.5) and c is the proportion of clouds ($c = 1$ in rainy days and $c = 0$ otherwise).

Chapter 9

Radiation transfer

Radiation transfer is not the same throughout the day, but changes with solar hour. This is represented using β , the solar elevation angle (see section 4.3 of the reference manual for package **meteoland**).

In the previous chapter we detailed sub-daily variation in temperature, direct radiation and diffuse short-wave radiation. Here we deal with the absorption of diffuse and direct shortwave radiation through the canopy as well as the long-wave radiation balance for canopy layers and the soil layer. These are necessary to estimate the different components of the canopy and soil energy balances (see eqs. (7.3), (7.4) and (7.5)) as well as to determine leaf energy balance, transpiration and photosynthesis (chapters 11 and 12). Short-wave radiation transfer can be determined for all sub-daily steps once the variation of atmospheric inputs are known. In contrast, the long-wave radiation (LWR) balance for any surface depends on its temperature and therefore, net LWR for canopy and soil layers need to be determined at each sub-daily step after closing the energy balance of the previous step (see chapter 13).

The vertical distribution of leaves in the stand is a key element of radiation transfer. As explained in section 2.4.3.3, the canopy is divided into vertical layers (whose size is determined by the control parameter **verticalLayerSize**), and the expanded and dead leaf area index of each cohort within each layer is determined. Let l be the number of canopy layers, c the number of plant cohorts, and $LAI_{all,i,j} = LAI_{act,i,j} + LAI_{dead,i,j}$ be the leaf area index of cohort i in layer j .

9.1 Short-wave radiation

In section 8.2 we explained how the model estimates instantaneous and direct/diffuse SWR and PAR at the top of the canopy. Given an input direct and diffuse short-wave irradiance at the top of the canopy, the amount of light absorbed by sunlit leaves, shade leaves and the soil follows the model of Anten and Bastiaans (2016), which we detail in the following subsections.

9.1.1 Direct and diffuse irradiance across the canopy

The average irradiance reaching the top of each canopy layer j is calculated separately for direct beam and diffuse radiation (the same equations are valid for SWR or PAR, but with different coefficients):

$$I_{beam,j} = (1 - \gamma_i) \cdot I_{beam} \cdot \exp \left[\sum_{h=j+1}^l \sum_i^c -k_{b,i} \cdot \alpha_i^{0.5} \cdot LAI_{all,i,h} \right] \quad (9.1)$$

$$I_{dif,j} = (1 - \gamma_i) \cdot I_{dif} \cdot \exp \left[\sum_{h=j+1}^l \sum_i^c -k_{d,i} \cdot \alpha_i^{0.5} \cdot LAI_{all,i,h} \right] \quad (9.2)$$

where I_{beam} and I_{dif} are the direct and diffuse irradiance at the top of the canopy (in $W \cdot m^{-2}$), γ_i is the leaf reflectance ($\gamma_{SWR,i}$ is an input parameter, whereas $\gamma_{PAR,i} = 0.8 \cdot \gamma_{SWR,i}$), $k_{b,i}$ is the extinction coefficient

of cohort i for direct light, $k_{d,i}$ is the extinction coefficient of cohort i for diffuse light (i.e. $k_{PAR,i}$ or $k_{SWR,i}$) and α_i is the absorbance coefficient ($\alpha_{SWR,i}$ is an input parameter, whereas $\alpha_{PAR,i} = \alpha_{SWR,i} \cdot 1.35$).

Plant leaves are not disposed horizontally but have an inclination (i.e. a leaf angle λ_{leaf}) which may change from one species to the other and determines radiation extinction along with solar elevation. The extinction coefficient for direct light, k_b , depends on the leaf angle (λ_{leaf}) and the solar elevation angle (β), as described in Sellers (1985). First λ_{leaf} is used to determine χ_{leaf} , a parameter describing the departure of leaf angle distribution from the spherical distribution (which assumes all angles as equally probable):

$$\chi_{leaf}(\lambda_{leaf}) = 2.0 \cdot \cos(\lambda_{leaf}) - 1.0 \quad (9.3)$$

χ_{leaf} equals 0 for +1 for horizontal leaves (i.e. for $\lambda_{leaf} = 0$ degrees), 0 for random leaves (i.e. for $\lambda_{leaf} = 60$ degrees) and -1 for vertical leaves (i.e. for $\lambda_{leaf} = 90$ degrees). χ_{leaf} and β are then used to estimate G_b the relative projected area of leaves in the direction of the solar beam:

$$G_b(\lambda_{leaf}, \beta) = \phi_1 + \phi_2 \cdot \sin(\beta); \quad (9.4)$$

where $\phi_1 = 0.5 - 0.633 \cdot \chi_{leaf} - 0.33 \cdot \chi_{leaf}^2$ and $\phi_2 = 0.877 \cdot (1 - 2 \cdot \phi_1)$ for $-0.4 \leq \chi_{leaf} \leq 0.6$. Finally, k_b is estimated as:

$$k_b = \frac{G_b(\lambda_{leaf}, \beta)}{\sin(\beta)} \quad (9.5)$$

The calculations can be illustrated with function `light_directionalExtinctionCoefficient()`:

```
solarElevation <- 40 # in degrees
meanLeafAngle <- 60 # in degrees
sdLeafAngle <- 20

## Parameters of the beta distribution
beta <- light_leafAngleBetaParameters(meanLeafAngle*(pi/180), sdLeafAngle*(pi/180))
beta

##           p           q
## 2.333333 1.166667

## Direct beam extinction coefficient for solar elevation
kb <- light_directionalExtinctionCoefficient(beta["p"], beta["q"], solarElevation*(pi/180))
kb

## [1] 0.7696812
```

9.1.2 Sunlit and shade leaves

It is generally accepted that sunlit and shade leaves need to be treated separately for photosynthesis calculations (De Pury and Farquhar, 1997). This separation is necessary because photosynthesis of shade leaves has an essentially linear response to irradiance, while photosynthesis of leaves in sunflecks is often light saturated and independent of irradiance. The proportion of sunlit leaves, i.e. leaves in a canopy layer j that the direct light beams (sunflecks) reach is:

$$f_{SL,j} = \exp \left(\sum_{k>j}^l \sum_i^c -k_{b,i} \cdot LAI_{all,i,k} \right) \cdot \exp \left(\sum_i^c -k_{b,j} \cdot 0.5 \cdot LAI_{all,i,j} \right) \quad (9.6)$$

where it is assumed that only half the leaf area of layer j contributes self-shading. Note that since k_b depends on the time of the day, also $f_{SL,j}$ may be different depending on the solar hour. From fraction $f_{SL,j}$ we can

derive the (expanded) leaf area of each layer that is affected by direct light beams (i.e. the amount of sunlit and shade leaves):

$$LAI_{sunlit,i,j} = f_{SL,j} \cdot LAI_{act,i,j} \quad (9.7)$$

$$LAI_{shade,i,j} = (1 - f_{SL,j}) \cdot LAI_{act,i,j} \quad (9.8)$$

As an example we will consider a canopy of one species of $LAI = 2$, divided into ten layers with constant leaf density:

```
LAI = 2
nlayer = 10
LAIlayerlive = matrix(rep(LAI/nlayer,nlayer),nlayer,1)
LAIlayermax = matrix(rep(LAI/nlayer,nlayer),nlayer,1)
LAIlayerdead = matrix(0,nlayer,1)
LAIlayermist = matrix(0,nlayer,1)
```

We will also define the coefficients of diffuse extinction, leaf absorbance and leaf reflection for PAR and SWR:

```
alpha_PAR <- 0.9
alpha_SWR <- 0.7
gamma_PAR <- 0.04
gamma_SWR <- 0.05
clumping <- 0.75
```

These definitions, and that of direct extinction coefficient, lead to a percentage of the above-canopy irradiance reaching each layer (Anten and Bastiaans, 2016). Extinction of direct radiation also defines the proportion of leaves of each layer that are affected by sunflecks (i.e. the proportion of sunlit leaves). Both outcomes are illustrated in the figure below:

9.1.3 Short-wave radiation absorbed by canopy elements

The amount of absorbed diffuse radiation per leaf area unit of cohort i within a given canopy layer j is calculated as:

$$\Phi_{dif,i,j} = I_{dif,j} \cdot k_{d,i} \cdot \alpha_i^{0.5} \exp \left[\sum_h^c -k_{d,h} \cdot \alpha_i^{0.5} \cdot 0.5 \cdot LAI_{all,h,j} \right] \quad (9.9)$$

The amount of absorbed scattered beam radiation per leaf area unit of cohort i within a given canopy layer j is calculated as:

$$\Phi_{sca,i,j} = I_{b,j} \cdot k_{b,j} \cdot (A - B) \quad (9.10)$$

$$A = \alpha_i^{0.5} \cdot \exp \left(\sum_h^c -k_{b,i} \cdot \alpha_i \cdot 0.5 \cdot LAI_{all,h,i} \right) \quad (9.11)$$

$$B = \frac{\alpha_i}{(1 - \gamma_i)} \cdot \exp \left(\sum_h^c -k_{b,i} \cdot 0.5 \cdot LAI_{all,h,i} \right) \quad (9.12)$$

Finally, the direct radiation absorbed by a unit of sunlit leaf area of cohort i in a canopy layer j does not depend on the position of the canopy layer and is:

$$\Phi_{dir,i,j} = I_{beam} \cdot \alpha_i \cdot k_{b,i} \quad (9.13)$$

Once these components are known, the amount of light absorbed by sunlit/shaded foliage of cohort i in layer j per leaf area unit ($\Phi_{sunlit,i,j}$ and $\Phi_{shade,i,j}$, respectively) is:

$$\Phi_{sunlit,i,j} = \Phi_{dif,i,j} + \Phi_{sca,i,j} + \Phi_{dir,i,j} \quad (9.14)$$

$$\Phi_{shade,i,j} = \Phi_{dif,i,j} + \Phi_{sca,i,j} \quad (9.15)$$




Figure 9.1: The left panel shows irradiance per layer of diffuse and direct PAR and SWR assuming a LAI of 2 equally distributed among layers (see function `light_layerIrradianceFraction()`). The right panel shows the corresponding proportions of sunlit and shade leaves in each layer (see function `light_layerSunlitFraction()`).

Let us show how all this works in an example. Regarding incoming light, we assume the following direct and diffuse irradiance at the top of the canopy:

```
solarElevation = 0.67
SWR_direct = 1100
SWR_diffuse = 300
PAR_direct = 550
PAR_diffuse = 150
```

Solar elevation is the angle between the sun and the horizon (i.e. the complement of the zenith angle). Under these conditions, and for the same canopy used for fig. 9.1, the amount of PAR and SWR absorbed per unit of leaf area at each canopy layer is (Anten and Bastiaans, 2016):

```
ZF <- c(0.178, 0.514, 0.308) ##Standard overcast sky
Zangles <- c(15.0, 45.0, 75.0);
KDIFF <- matrix(NA, 3, 1);
for(k in 1:3){
  KDIFF[k,1] <- light_directionalExtinctionCoefficient(beta["p"], beta["q"], Zangles[k]*(pi/180.0));
}
KDIFF
```

```
##           [,1]
## [1,] 1.9888914
## [2,] 0.6926432
## [3,] 0.4853480
```




Figure 9.2: PAR (left) and SWR (right) absorbed per unit of sunlit/shade leaf area at each canopy layer ($I_{sunlit,i,j}$ and $I_{shade,i,j}$, respectively; see function `light_cohortSunlitShadeAbsorbedRadiation()`).

The SWR absorbed per ground area unit by sunlit and shade foliage of cohort i in canopy layer j is:

$$K_{abs,sunlit,i,j} = \Phi_{sunlit,i,j} \cdot LAI_{sunlit,i,j} \quad (9.16)$$

$$K_{abs,shade,i,j} = \Phi_{shade,i,j} \cdot LAI_{shade,i,j} \quad (9.17)$$

The previous quantities can be aggregated across cohorts or layers. The SWR absorbed per ground area unit by cohort i ($K_{abs,i}$) is found using:

$$K_{abs,i} = \sum_j^l K_{abs,sunlit,i,j} + K_{abs,shade,i,j} = K_{abs,sunlit,i} + K_{abs,shade,i} \quad (9.18)$$

where $K_{abs,sunlit,i}$ and $K_{abs,shade,i}$ are the SWR absorbed per ground area unit by sunlit and shade foliage, respectively, of cohort i . Similarly, the SWR absorbed by canopy layer j ($K_{abs,j}$) is found using:

$$K_{abs,j} = \sum_i^c K_{abs,sunlit,i,j} + K_{abs,shade,i,j} \quad (9.19)$$

and the SWR absorbed by the whole canopy ($K_{abs,ca}$) is:

$$K_{abs,ca} = \sum_j^l K_{abs,j} = \sum_i^c K_{abs,i} = \sum_i^c \sum_j^l K_{abs,sunlit,i,j} + K_{abs,shade,i,j} \quad (9.20)$$

9.1.4 Short-wave radiation absorbed by the soil

The instantaneous shortwave radiation reaching the soil is calculated separately for direct beam and diffuse radiation:

$$I_{beam,soil} = I_{beam} \cdot \exp \left[\sum_{h=j+1}^l \sum_i^c -k_{b,i} \cdot \alpha_i^{0.5} \cdot LAI_{all,i,h} \right] \quad (9.21)$$

$$I_{dif,soil} = I_{dif} \cdot \exp \left[\sum_{h=j+1}^l \sum_i^c -k_{d,i} \cdot LAI_{all,i,h} \right] \quad (9.22)$$

where I_{beam} and I_{dif} are the direct and diffuse irradiance at the top of the canopy, $k_{b,i}$ is the extinction coefficient of cohort i for direct light and $k_{d,i}$ is the extinction coefficient of cohort i for diffuse SWR. From these, the SWR absorbed by the soil ($K_{abs,soil}$) is found by:

$$K_{abs,soil} = (1 - \gamma_{SWR,soil}) \cdot (I_{beam,soil} + I_{dif,soil}) \quad (9.23)$$

where $\gamma_{SWR,soil} = 0.10$ is the SWR reflectance (10% albedo) of the soil.

9.2 Long-wave radiation

Long-wave radiation (LWR) transfer within the canopy is based on the SHAW model (Flerchinger et al., 2009).

9.2.1 Long-wave radiation balance by canopy layers

Assume canopy layers are ordered from $j = 1$ (in contact with the soil) to $j = l$ (in contact with the atmosphere above the boundary layer). The procedure first calculates downward LWR below each canopy layer, $L_{down,j}$, from $j = l$ to $j = 1$. The downward LWR below layer j is calculated as:

$$L_{down,j} = \tau_j \cdot L_{down,j+1} + (1 - \tau_j) \cdot \epsilon_{can} \cdot \sigma \cdot (T_{air,j} + 273.16)^4 \quad (9.24)$$

where $L_{down,j+1}$ is the downward LWR of the layer above (L_{atm} in the case of $j = l$), $T_{air,j}$ is the air temperature of layer j , τ_j is the diffuse transmissivity of layer j , $\sigma = 5.67 \cdot 10^{-8} W \cdot K^{-4} \cdot m^{-2}$ is the Stephan-Boltzmann constant and $\epsilon_{can} = 0.97$ is the canopy emissivity. Diffuse transmissivity values are defined by cohort i and layer j using:

$$\tau_{i,j} = \exp(-k_{LWR} \cdot LAI_{all,i,j}) \quad (9.25)$$

where $k_{LWR} = 0.7815$ is the extinction coefficient for LWR. The transmissivity of layer j is simply $\tau_j = \sum_i \tau_{i,j}$. Once we know $L_{down,1}$ we can calculate the upwards LWR from the soil ($L_{up,s}$) using:

$$L_{up,soil} = (1 - \epsilon_{soil}) \cdot L_{down,1} + \epsilon_{soil} \cdot \sigma \cdot (T_{soil,1} + 273.16)^4 \quad (9.26)$$

where $T_{soil,1}$ is the temperature of the topmost soil layer and $\epsilon_{soil} = 0.97$ is the soil emissivity. Now we calculate upward LWR from $j = 1$ to $j = J$:

$$L_{up,j} = \tau_j \cdot L_{up,j-1} + (1 - \tau_j) \cdot \epsilon_{can} \cdot \sigma \cdot (T_{air,j} + 273.16)^4 \quad (9.27)$$

where $L_{up,j-1}$ is the downward LWR of the layer below ($L_{up,soil}$ in the case of $j = 1$). Once we have downward and upward LWR fluxes, we can estimate the net LWR absorbed within canopy layer j using:

$$L_{net,j} = \epsilon_{can} \cdot (1 - \tau_j) \cdot (L_{down,j} + L_{up,j} - \sigma \cdot (T_{air,j} + 273.16)^4) \quad (9.28)$$

9.2.2 Net long-wave radiation of the whole canopy and the soil

Net LWR can be aggregated at the scale of the whole canopy. The net LWR of the whole canopy ($L_{net,can}$) is simply the sum across layers of net LWR:

$$L_{net,can} = \sum_j^l L_{net,j} \quad (9.29)$$

whereas the net LWR of the soil ($L_{net,soil}$) is:

$$L_{net,soil} = \epsilon_{soil} \cdot (L_{down,1} - \sigma \cdot (T_{soil,1} + 273.16)^4) \quad (9.30)$$

9.2.3 Long-wave radiation balance by plant cohorts

The net LWR per leaf area unit absorbed within canopy layer j can also be decomposed among plant cohorts. The net LWR of cohort i in layer j is found using the relative proportion of light absorbed by the cohort (i.e. the complement of its diffuse transmissivity):

$$L_{net,i,j} = L_{net,j} \cdot \frac{1 - \tau_{i,j}}{\sum_h 1 - \tau_{h,j}} \quad (9.31)$$

from which we can estimate $L_{net,sunlit,i}$ and $L_{net,shade,i}$, the average net LWR per leaf area unit for sunlit and shade leaves of cohort i , respectively:

$$L_{net,sunlit,i} = \frac{\sum_j^l L_{net,i,j} \cdot LAI_{sunlit,i,j}}{\sum_j^l LAI_{sunlit,i,j}} \quad L_{net,shade,i} = \frac{\sum_j^l L_{net,i,j} \cdot LAI_{shade,i,j}}{\sum_j^l LAI_{shade,i,j}} \quad (9.32)$$

Chapter 10

Plant hydraulics

We devote this chapter to describe the background necessary to understand how plant hydraulics is implemented in the advanced water balance model of **medfate**. Section 10.1 introduces the hydraulic networks from which to choose, depending on the sub-model of plant hydraulics employed (i.e. either `transpirationMode = "Sperry"` or `transpirationMode = "Sureau"`). After that, in sections 10.2 and 10.3 we present the building blocks of models of plant hydraulics, i.e. how to represent in a model the water content of plant tissues and the water flow through vascular elements. In section 10.4 we describe in detail the plant hydraulic sub-model presented in De Cáceres et al. (2021) (without water compartments), and in 10.5 we describe the sub-model of plant hydraulics that was presented as SurEau-ECOS in Ruffault et al. (2022) (see 7.1).

10.1 Hydraulic architecture

The hydraulic architecture of a model describes the set of water compartments and resistances to water flow that are explicitly considered, often implying a simplification and discretization of real plant hydraulic pathways. The advanced water balance model of **medfate** implements different hydraulic architectures depending on the transpiration sub-model:

- If `transpirationMode = "Sperry"`, the hydraulic network is described using $(S \times 2 + 1 + 1)$ resistance elements, with soil being represented in S different soil layers. For each soil layer there is a rhizosphere element in series with a root xylem element. The S soil layers are in parallel up to the root crown. From there there is one stem xylem segment and a final leaf xylem segment in series (Fig. 10.1). When considered (via control parameter `stemCavitationEffects`), stem cavitation affects the resistance between the root crown ($\Psi_{rootcrown}$) and the stem (Ψ_{stem}). Similarly, when considered (via control parameter `leafCavitationEffects`) leaf cavitation affects the resistance between the stem (Ψ_{stem}) and the leaf (Ψ_{leaf})
- If `transpirationMode = "Sureau"`, the soil–plant system in SurEau-ECOS (Ruffault et al., 2022) is discretized into three soil layers and two plant compartments (Fig. 10.2): a leaf and a stem. Each of the two plant organs contains an apoplasm and a symplasm. The stem apoplasm and symplasm include water volumes of all non-leaf compartments, i.e., trunk, root and branches. Symplasmic capacitances (i.e. C_{LSym} and C_{SSym}) mostly buffer water fluxes during well-watered conditions, whereas apoplasm capacitances (i.e. C_{LApo} and C_{SApo}) come into play when cavitation occurs. Water can leave the plant via three flows, through stomata (E_{stom}), leaf cuticles ($E_{cuti,leaf}$) and stem cuticles ($E_{cuti,stem}$). The architecture is completed with nodes for the fine root surface of each soil layer ($\Psi_{rootsurf}$) and for the root crown ($\Psi_{rootcrown}$), but these elements do not correspond to water compartments. Stem cavitation affects the conductances of the pathways between root surfaces ($\Psi_{rootsurf}$) and the root crown ($\Psi_{rootcrown}$), as well as the conductance between the root crown and the stem apoplasm ($\Psi_{stem,apo}$). Leaf cavitation affects the conductance between the stem apoplasm ($\Psi_{stem,apo}$) and the




Figure 10.1: Schematic representation of hydraulics in a whole-plant network

leaf apoplasts ($\Psi_{leaf,apo}$).

The two architectures share as common elements the nodes corresponding to bulk soil (Ψ_{soil}) and root surface ($\Psi_{rootsurf}$) for each soil layer, as well as the water potential at the root crown ($\Psi_{rootcrown}$). Since the architecture of `transpirationModel = "Sperry"` considers plant xylem elements only, the stem water potential (Ψ_{stem}) can be considered analogous to the apoplastic stem water potential of Sureau-ECOS ($\Psi_{stem,apo}$). Similarly its leaf water potential (Ψ_{leaf}) can be considered equivalent to the apoplastic stem water potential of Sureau-ECOS ($\Psi_{leaf,apo}$). Symplasmic compartments of Sureau-ECOS do not have equivalents in the Sperry architecture, but the water potential of symplasmic compartments can be assumed to be equal to those of their corresponding apoplastic compartments. Important differences between the two architectures include the additional consideration of (i) plant capacitance effects, (ii) cavitation flows and (iii) cuticular transpiration flows that are only available in Sureau-ECOS (Fig. 10.1). The two models also differ in how stomatal regulation, water flows and water potentials are solved. These differences are described in the following sub-sections as well as in chapter 12.

10.2 Vulnerability curves

Vulnerability curves form the basis of hydraulic calculations. Let us use k to denote hydraulic conductance, i.e. instantaneous flow rate per leaf surface unit and per pressure drop (in $mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$). Each element of the hydraulic network has a vulnerability curve $k(\Psi)$ that starts at maximum hydraulic conductance ($k_{max} = k(0)$) and declines as water pressure (Ψ) becomes more negative.

10.2.1 Xylem vulnerability curves

Xylem vulnerability curves are modelled using different functions, depending on the transpiration sub-model.




Figure 10.2: Schematic representation of hydraulics in a network with water storage compartments according to the design of SurEau-ECOS

10.2.1.1 Weibull vulnerability curves

When `transpirationMode = "Sperry"`, xylem tissues are assigned a two-parameter Weibull function as the vulnerability curve $k(\Psi)$:

$$k(\Psi) = k_{max} \cdot e^{-((\Psi/d)^c)} \quad (10.1)$$

where k_{max} is the maximum hydraulic conductance, and c and d are species-specific and tissue-specific parameters. Note that parameter d is the water potential (in MPa) at which $k(\Psi)/k_{max} = e^{-1} = 0.367$, i.e. the water potential at which hydraulic conductance is 36.7% of its maximum value. Parameter c controls the shape of the vulnerability curve (*exponential* shape with no threshold has $c \leq 1$; and *sigmoidal* shape with threshold occurs when $c > 1$).

The concept of vulnerability curve can be used to specify the relationship between pressure and conductance in any portion of the flow path. For example, we define the following parameter values for a stem xylem ($k_{stem,max}$ and parameters c_{stem} and d_{stem} of the vulnerability curve):

```
kstemmax = 5.0 # mmol·m-2·s-1·MPa-1
stemc = 3
stemd = -3.0 # MPa
```

For root xylem ($k_{root,max}$), we may assume a higher conductance (i.e. higher efficiency) but also higher vulnerability to cavitation (defined by parameters c_{root} and d_{root}):

```
krootmax = 6.6 # mmol·m-2·s-1·MPa-1
rootc = 2
rootd = -2.5 #MPa
```

For leaf xylem $k_{leaf}(\Psi)$, we may specify higher conductance than in stems but also higher vulnerability:

```
kleafmax = 10
leafc = 2
leafd = -2
```

With these parameter values, the vulnerability curves for root, stem and leaf are (see `hydraulics_xylemConductance()` and `hydraulics_vulnerabilityCurvePlot()`):

The dash-dot line between 0 and $\Psi_{cav} = -2.5$ MPa indicates the modification of the stem xylem vulnerability curve when cavitation has occurred (i.e., previous embolism limits the maximum conductance value), as indicated in Sperry et al. (2017). The corresponding proportion of stem conductance loss (PLC_{stem}) can be found using the stem vulnerability curve. Similarly, PLC_{leaf} can limit conductance in leaves. Although root xylem are more vulnerable to the formation of emboli for a given potential, it is generally accepted that the less negative potentials of root xylem compared to the stem lead to cavitation occurring more often in the stem. The constrain created by cavitation has an effect on the calculation of the flow rates and derived quantities (see below).

10.2.1.2 Sigmoid vulnerability curves

When `transpirationMode = "Sureau"`, xylem tissues are assigned a two-parameter sigmoid function as the vulnerability curve $k(\Psi)$:

$$k(\Psi) = \frac{k_{max}}{1 + e^{(slope/25) \cdot (\Psi - \Psi_{50})}} \quad (10.2)$$

where Ψ_{50} is the water potential corresponding to 50% of conductance, and *slope* is the slope of the curve in percent at that point. The two parameters need to be specified for each plant segment under consideration, analogously to the Weibull functions.




Figure 10.3: Example vulnerability curves corresponding to the parameters defined above for stem, root and leaf segments.

10.2.2 Rhizosphere vulnerability curve

The rhizosphere conductance function $k_{rhizo}(\Psi)$ is modelled as a van Genuchten function (Genuchten, 1980) (this choice is the reason why the water retention curve also needs to be modelled with van Genuchten):

$$k_{rhizo}(\Psi) = k_{rhizo,max} \cdot v^{(n-1)/(2 \cdot n)} \cdot ((1 - v)^{(n-1)/n} - 1)^2 \quad (10.3)$$

$$v = [(\alpha \Psi)^n + 1]^{-1} \quad (10.4)$$

where $k_{rhizo,max}$ is the maximum rhizosphere conductance, and n and α are texture-specific parameters (Leij et al., 1996; Carsel and Parrish, 1988). These are automatically set by function `soil()` when initializing soil objects (see section 2.3.3), but we can use function `soil_vanGenuchtenParamsCarsel()` to derive them from texture types:

```
textures = c("Sandy loam", "Silt loam", "Clay")
#Textural parameters
#Sandy clay loam
p1 = soil_vanGenuchtenParamsCarsel(textures[1])
p1
```

```
##      alpha      n theta_res theta_sat      Ks
## 764.983    1.890    0.065    0.410 69513.524
```

```
alpha1 = p1[1]
n1 = p1[2]
#Silt loam
p2 = soil_vanGenuchtenParamsCarsel(textures[2])
p2
```

```
##      alpha      n theta_res theta_sat      Ks
## 203.9955    1.4100    0.0670    0.4500 7077.1687
```

```
alpha2 = p2[1]
n2 = p2[2]
#Silty clay
p3 = soil_vanGenuchtenParamsCarsel(textures[3])
p3
```

```
##      alpha      n theta_res theta_sat      Ks
## 81.59819    1.09000    0.06800    0.38000 3145.40832
```

```
alpha3 = p3[1]
n3 = p3[2]
```

We can estimate maximum rhizosphere conductance values assuming that they account for an average percentage of the resistance (e.g. 15%) across the continuum (see functions `hydraulics_averageRhizosphereResistancePer` and `hydraulics_findRhizosphereMaximumConductance()`):

```
percentResistance = 15
#Sandy clay loam
krmax1 = hydraulics_findRhizosphereMaximumConductance(percentResistance,
                                                       n1, alpha1, krootmax, rootc, rootd, kstemmax, stemc, stemd,
                                                       kleafmax, leafc, leafd, -2, 0)
krmax1
```

```
## [1] 1.54602e+14
```

```
#Silt loam
krmax2 = hydraulics_findRhizosphereMaximumConductance(percentResistance,
                                                       n2, alpha2, krootmax, rootc, rootd, kstemmax, stemc, stemd,
```

```

kleafmax, leafc, leafd, -2, 0)
krmax2

## [1] 1667139299
#Silty clay
krmax3 =hydraulics_findRhizosphereMaximumConductance(percentResistance,
n3,alpha3, krootmax, rootc,rootd, kstemmax, stemc, stemd,
kleafmax, leafc, leafd, -2, 0)
krmax3

## [1] 30061324

```

With these parameters, the resulting $k_{rhizo}(\Psi)$ functions can be displayed using the function `hydraulics_vanGenuchtenCondu`




Figure 10.4: Example rhizosphere vulnerability curves (i.e. Van Genuchten functions) for three different soil textures.

10.2.3 Xylem sap viscosity

The ease of water flow (i.e. hydraulic conductance or resistance) is dependent on water viscosity, which fluctuates as a function of temperature and dissolved substances. Xylem sap can be considered to have negligible amounts of solutes, but temperature is still a relevant factor of viscosity. The dynamic viscosity of water V_w varies with temperature (relative to its viscosity at 20°C) according to the Vogel equation:

$$V_w(T) = \exp \left[-3.7188 + \frac{578.919}{-137.546 + T} \right] \quad (10.5)$$

where T is the temperature of the medium in Kelvin. Temperature of segments (roots, stem, leaves) is used to correct maximum conductances simply as follows:

$$k_{max,T} = \frac{k_{max}}{V_w(T)} \quad (10.6)$$

where k_{max} is the maximum conductance describing the vulnerability curve, assumed to be determined at 20°C.

10.3 Water content of plant tissues

Following Martin-StPaul et al. (2017), we consider two kind of tissues in (leaf and stem) plant segments (Tyree and Yang, 1990). The first are conduits (tracheids or vessels), which will release water due to cavitation and may be refilled with water from adjacent living tissue or upstream segments. The second source of water is formed by more elastic living cells (i.e. parenchyma) and can potentially be a large source of water. This source can be described using the relative water content of a symplasmic tissue.

10.3.1 Relative water content of symplasmic tissues

A pressure-volume curve of a tissue relates its water potential against its *relative water content* (RWC ; $kg\ H_2O \cdot kg^{-1}\ H_2O$ at saturation). Pressure-volume theory is usually applied to leaf tissues (Bartlett et al., 2012), but it can also be applied to other tissues such as sapwood or cambium cells.

For living cells, the relationship between Ψ and RWC of the symplasmic fraction (RWC_{sym}) is achieved by separating Ψ into osmotic (solute) potential (Ψ_S) and the turgor potential (Ψ_P):

$$\Psi = \Psi_S + \Psi_P \quad (10.7)$$

The relationship for Ψ_P is:

$$\Psi_P = -\pi_0 - \epsilon \cdot (1.0 - RWC_{sym}) \quad (10.8)$$


where π_0 (MPa) is the osmotic potential at full turgor (i.e. when $RWC_{sym} = 1$), and ϵ is the modulus of elasticity (i.e. the slope of the relationship). Assuming constant solute content, the relationship for Ψ_S is:

$$\Psi_S = \frac{-\pi_0}{RWC_{sym}} \quad (10.9)$$

When $\Psi \leq \Psi_{tlp}$, the water potential at turgor loss point, then $\Psi_P = 0$ and $\Psi = \Psi_S$. If $\Psi > \Psi_{tlp}$ then the two components are needed. The water potential at turgor loss point (Ψ_{tlp}) can be found by (Bartlett et al., 2012):

$$\Psi_{tlp} = \frac{\pi_0 \cdot \epsilon}{\pi_0 + \epsilon} \quad (10.10)$$

As an example, the following figure draws the pressure-volume curve for a tissue with $\epsilon = 12$ and $\pi_0 = -3.0$ MPa:




To calculate RWC_{sym} from the water potential of a tissue, the previous equations need to be combined and, after isolating RWC_{sym} , a quadratic relationship is obtained (Martin-StPaul et al., 2017).

10.3.2 Relative water content of apoplastic tissues

In **medfate** it is assumed that apoplastic water fraction of an organ corresponds to the water in its xylem conduits. Xylem conduits consist of inelastic cells that have very small changes of water volume in relation to changes in water potential. However, xylem conduits release their water to the transpiration stream following the formation of emboli. As in Hölttä et al. (2009), we equate the relative water content of the apoplastic reservoir of a segment (leaves or stem) to the proportion of conductance lost (relative to the maximum conductance) for a given water potential (see (10.1)):

$$RWC_{apo}(\Psi_{apo}) = \frac{k(\Psi_{apo})}{k_{max}} = e^{-((\Psi_{apo}/d)^c)} \quad (10.11)$$

Hence we assume that the relationship between Ψ_{apo} and the relative water content follows the same function as the relationship between Ψ_{apo} and the proportion of conductance loss (i.e. the hydraulic vulnerability curve, see ?(vulnerabilitycurves)).



10.3.3 Average relative water content of a compartment

The average relative water content in a given segment or organ (RWC) can be obtained by calculating RWC_{apo} and RWC_{sym} followed by assuming a constant apoplastic fraction f_{apo} :

$$RWC = RWC_{apo} \cdot f_{apo} + RWC_{sym} \cdot (1 - f_{apo}) \quad (10.12)$$

Normally the apoplastic fraction is large ($\sim 80\%$) in the stem and small ($\sim 15\%$) in leaves. Hence, the relative water content in stems will be mostly dictated by the level of cavitation, whereas that of leaves will mostly follow leaf symplastic water potential and the symplastic pressure-volume curve.

10.3.4 Live fuel moisture content

Given an average relative water content (RWC) of a plant organ, its live fuel moisture content ($LFMC$ in $gH_2O \cdot g^{-1}$ of dry tissue) can be calculated using:

$$LFMC = RWC \cdot \Theta \cdot \frac{\rho_{H_2O}}{\rho} = RWC \cdot LFMC_{max} \quad (10.13)$$

where Θ is the tissue porosity (cm^3 of water per cm^3 of tissue), ρ is the density of the tissue and ρ_{H_2O} is the density of water. In practice estimates of maximum fuel moisture content $LFMC_{max}$ are easier to

obtain than porosity estimates, so it is more straightforward to estimate $LFMC$ as the product of RWC and $LFMC_{max}$.

In medfate, $LFMC$ estimates are calculated differently depending on the value of the control variable `lfmcComponent`. If `lfmcComponent` = "leaf", $LFMC$ estimates correspond to the leaf segment, i.e.:

$$LFMC_{leaf} = RWC_{leaf} \cdot LFMC_{max} \quad (10.14)$$

where RWC_{leaf} is the relative water content of in leaves (already averaging symplasmic and apoplasmic tissues). In contrast, if `lfmcComponent` = "fine", $LFMC$ estimates correspond to fine fuels (i.e. leaves and twigs of < 6.5 mm) and are estimated from stem and leaf water content:

$$LFMC_{fine} = (RWC_{leaf} \cdot (1.0/r_{6.35}) + RWC_{stem} \cdot (1.0 - (1.0/r_{6.35}))) \cdot LFMC_{max} \quad (10.15)$$

where $r_{6.35}$ is the ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm and RWC_{stem} is the relative water content of in stem (already averaging symplasmic and apoplasmic tissues).

10.4 Sperry's sub-model of plant hydraulics

The supply-loss theory of plant hydraulics, presented by Sperry and Love (2015) and used in Sperry et al. (2017), uses the physics of flow through soil and xylem to quantify how steady-state canopy water supply declines with drought and ceases by hydraulic failure. The theory builds on the hydraulic model of Sperry et al. (1998) and can be applied to different segmentations of the soil-plant continuum.

The supply function describes the steady-state rate of water supply (i.e. flow) for transpiration (E) as a function of water potential drop. The steady-state flow rate E_i through any i element of the continuum is related to the flow-induced drop in pressure across that element ($\Delta\Psi_i = \Psi_{down} - \Psi_{up}$) by the integral transform of the element's vulnerability curve $k_i(\Psi)$ (Sperry and Love, 2015):

$$E_i = \int_{\Psi_{up}}^{\Psi_{down}} k_i(\Psi) d\Psi \quad (10.16)$$

where Ψ_{up} and Ψ_{down} are the upstream and downstream water potential values, respectively. The integral transform assumes infinite discretization of the flow path.

The supply function can be defined for individual elements of the continuum or for the whole soil-plant continuum using different hydraulic networks. Supply functions are used in the Sperry's sub-model to determine photosynthesis, stomatal conductance and transpiration (see chapters 11 and 12).

In the following subsections we illustrate the supply function for different cases.

10.4.1 Supply function for a single xylem element

In the case of a single xylem element, the supply function describes the steady-state flow rate as a function of pressure at the stem top (Ψ_{canopy}). It can be calculated by numerical integration or approximated using an *incomplete gamma function*. The shape of the supply function starting at different root crown water potential values ($\Psi_{rootcrown}$) is (see function `hydraulics_EXylem()`):

The supply function of a *rhizosphere* element relates the steady-state flow rate to the pressure inside the roots (Ψ_{root}). It is calculated by numerical integration of the van Genuchten function, for which we use the analytical approximation of Van Lier et al. (2009) (see function `hydraulics_EVanGenuchten()`). In the figure below, we draw examples of the supply function for the rhizosphere. The nearly vertical lines indicate that for many values of E_i the corresponding drop in water potential through the rhizosphere will be negligible. Only for increasingly negative soil water potential values the decrease in water potential through the rhizosphere becomes relevant. Both in the case of a xylem element or a rhizosphere element the derivative $dE_i/d\Psi$ of the supply function is equal to the corresponding vulnerability curve.




Figure 10.5: Supply function of a single xylem element starting at different root crown water potential values. Left pane shows the uncavitated supply functions and right pane shows the supply functions that are obtained in the case of a cavitated xylem (i.e. without refilling), assuming that the minimum water potential experienced so far was -2.5 MPa. Note the linear part of the flow rate between $\Psi_{rootcrown}$ and this limit.




Figure 10.6: Supply functions of the rhizosphere starting at the four different values of bulk soil pressure (Ψ_{soil}) and for the same three texture types used for vulnerability curves.

10.4.2 Supply function of two elements in series

Let us describe the soil-plant continuum is represented using *two* elements in series (rhizosphere + stem xylem). In this case, the supply function has to be calculated by using the previous supply functions sequentially. The E_i is identical for each element and equal to the canopy E . Since Ψ_{soil} is known, one first inverts the supply function of the rhizosphere to find Ψ_{root} (see function `hydraulics_E2psiVanGenuchten()`) and then inverts the supply function of the xylem to find Ψ_{canopy} (see function `hydraulics_E2psiXylem()`). The two operations can be summarized in a single supply function describing the potential rate of water supply for transpiration (E) as function of the canopy xylem pressure (Ψ_{canopy}), starting from different bulk soil (Ψ_{soil}) values (see function `hydraulics_supplyFunctionTwoElements()`):




Figure 10.7: Example of two-element supply functions describing the potential rate of water supply for transpiration (E) as function of the canopy xylem pressure (Ψ_{canopy}), starting from different bulk soil (Ψ_{soil}) values and for different soil textures. Left/right panel shows uncavitated/cavitated supply functions.

The supply function for the whole continuum contains much information. The Ψ intercept at $E = 0$ represents the (predawn) canopy sap pressure which integrates the rooted soil moisture profile. As E increments from zero, the disproportionately greater drop in Ψ_{canopy} results from the loss of conductance. As the soil dries the differences in flow due to soil texture become more apparent.

The derivative of the whole continuum supply function, $dE/d\Psi$, is not equal to either of the vulnerability curves and it has to be obtained numerically. The derivative functions corresponding to the supply functions shown in the previous figure are:

The derivative $dE/d\Psi_{canopy}$ is the conductance if the entire continuum was exposed to Ψ_{canopy} (Sperry and Love, 2015). It corresponds to the local loss of hydraulic conductance at the downstream end of the flow path. It falls towards zero for asymptotic critical values (E_{crit}). For a cavitated system $dE/d\Psi_{canopy}$ can be rather flat, in accordance with the close to linear part of the supply function.




Figure 10.8: Derivatives of two-element supply functions corresponding to figure 10.7. Left/right panel shows the derivatives of uncavitated/cavitated supply functions.

10.4.3 Supply function of three elements in series

If the soil-plant continuum is represented using *three* elements in series (rhizosphere + stem xylem + leaf), the resulting overall conductance and resistance fractions (under wet conditions) are:

```
rstemmin = 1/kstemmax
rleafmin = 1/kleafmax

#Percentages of minimum resistance
rvec = c(rstemmin,rleafmin)
100*rvec/sum(rvec)

## [1] 66.66667 33.33333

#Maximum overall conductance
1/sum(rvec)

## [1] 3.333333
```

As before, the supply function has to be calculated *sequentially*, knowing that E_i is identical through each element. Since Ψ_{soil} is known, one first inverts the supply function of the rhizosphere to find Ψ_{root} and then inverts the supply function of the xylem to find Ψ_{stem} . Finally, one inverts the supply function of the leaf element to find Ψ_{leaf} . As before, the three operations can be summarized in a single supply function describing the potential rate of water supply for transpiration (E) as function of the leaf pressure (Ψ_{leaf}), starting from different bulk soil (Ψ_{soil}) values (see function `hydraulics_supplyFunctionThreeElements()`):




Figure 10.9: Example of three-element supply functions describing the potential rate of water supply for transpiration (E) as function of the leaf pressure (Ψ_{leaf}), starting from different bulk soil (Ψ_{soil}) values and for different soil textures. Left/right panel shows uncavitated/cavitated supply functions.

Note that overall conductance and the maximum flow of the supply function are smaller in this case than in the representation using two elements in series, because we added a new resistance (leaves). While the rhizosphere component only adds a significant resistance when the soil dries, considering the leaf segment (or a root xylem segment) substantially increases the overall resistance of the continuum. Higher vulnerability of leaves also makes the curve to saturate for less negative soil water potentials. The derivative functions corresponding to the supply functions shown in the previous figure are (note the highest value being equal to the overall maximum conductance):




Figure 10.10: Derivatives of three-element supply functions corresponding to figure 10.9. Left/right panel shows the derivatives of uncavitated/cavitated supply functions.

10.4.4 Supply function of a root system

So far we considered supply functions of elements in series, but resistance elements will be in parallel if the soil is represented using $S > 1$ different layers. For each soil layer there is a rhizosphere element in series with a root xylem element. The S soil layers are in parallel up to the root crown.

Network of S rhizosphere components and root layers in parallel there are $S + 1$ unknown pressures: the S root surface pressures ($\Psi_{rootsurf,1}, \dots, \Psi_{rootsurf,S}$) and the root crown pressure at the downstream junction for all root components ($\Psi_{rootcrown}$). The unknown pressures are solved, for each specified total flow value E , using multidimensional Newton-Raphson on a set of equations for steady-state flow (Sperry et al., 2016):

$$E_{rhizo,s} - E_{root,s} = 0 \quad (10.17)$$

$$\sum_{s=1}^S E_{root,s} - E = 0 \quad (10.18)$$

where $E_{rhizo,s}$ and $E_{root,s}$ are steady-state supply flows calculated using the integrals of either van Genuchten

or Weibull function as vulnerability curves, respectively. In the case of rhizosphere elements, $\Psi_{up} = \Psi_{soil,s}$ and in the case of root elements $\Psi_{up} = \Psi_{rootsurf,s}$. Solving the steady-state equations also provides values for flow across each of the parallel paths $E_{rhizo,s} = E_{root,s}$.




Figure 10.11: Schematic representation of hydraulics in a root network

As an example, we start by defining the water potential of three soil layers corresponding to four situations (analogously with the soil water potentials defined above):

```
psiSoilLayers1 = c(-0.3,-0.2,-0.1)
psiSoilLayers2 = c(-1.3,-1.2,-1.1)
psiSoilLayers3 = c(-2.3,-2.2,-2.1)
psiSoilLayers4 = c(-3.3,-3.2,-3.1)
```

In a network of several soil layers, one has to divide the total rhizosphere and root xylem conductances among layers. Let layer widths (d_s) be:

```
d = c(300,700,3000) #Soil layer widths in mm
```

Now let FRP_1 , FRP_2 and FRP_3 be the proportion of fine root biomass in each soil layer (see section 2.4.3.4), which can be calculated using:

```
Z50 = 200 #Parameter of LDR root distribution
Z95 = 1200 #Parameter of LDR root distribution
FRP = root_ldrDistribution(Z50, Z95, NA, d)
FRP
```

```
##           [,1]      [,2]      [,3]
## [1,] 0.6652935 0.2749944 0.05971209
```

In the case of the rhizosphere conductances, we can simply define them (for each soil texture type) as:

```
krhizomaxvec1 = krmx1*FRP
krhizomaxvec2 = krmx2*FRP
krhizomaxvec3 = krmx3*FRP
```

To divide maximum root xylem conductance among soil layers we need weights inversely proportional to the length of transport distances (Sperry et al., 2016). Vertical transport lengths can be calculated from soil depths and radial spread can be calculated assuming cylinders with volume that has to be in accordance with the xylem root conductance (TO BE DESCRIBED). The whole process can be done using functions `root_coarseRootSoilVolumeFromConductance()` and `root_coarseRootLengthsFromVolume()`:

```
rfc = c(20,50,70)
Vol = root_coarseRootSoilVolumeFromConductance(1.0, 2500,krootmax,
                                                FRP,d, rfc)
lengths = root_coarseRootLengthsFromVolume(Vol, FRP, d, rfc)
lengths
```

```
## [1] 3621.093 2768.105 4513.795
```

```
weights = lengths/sum(lengths)
```

Transport weights are quite different than the fine root biomass proportions. This is because radial lengths are largest for the first (top) layer and vertical lengths are largest for the third (bottom) layer. The root xylem conductances are (in this case they do not depend on soil texture):

```
krootmaxvec = krootmax*weights
krootmaxvec
```

```
## [1] 2.191987 1.675640 2.732373
```

Having all these maximum conductances, we can now build the supply functions for each soil texture and starting from the different soil water potential configurations (see function `hydraulics_supplyFunctionBelowground()`):

The derivative of $dE/d\Psi_{rootcrown}$ for the supply function of the root system is again obtained numerically.

Solving the previous system of equations provides the water potentials in different points of the root system. Here we plot them for the results of silt loam texture and the first and last soil potential vectors defined above:

Note that when soil is not dry (first situation) pressure drop in the rhizosphere is negligible, but not the pressure drop in the root xylem. For drier soils rhizosphere becomes more relevant.

We can also plot the flow rates across each of the parallel paths (again corresponding to the results of silt loam texture and for the four soil potential vectors):

Note that the contribution of each soil layer depends on the soil conditions and the total amount of flow. For a low total flow rate some layers may have negative flows if their potential is lower than others, which in a dynamic context will cause *hydraulic redistribution* of water among soil layers.

10.4.5 Supply function of the soil-plant continuum

As mentioned above, **medfate** uses a hydraulic network of $(S \times 2 + 2 + 1)$ resistance elements to represent the soil-plant continuum, with soil being represented in S different layers. As before, the S soil layers are in parallel up to the root crown and each soil layer requires at least a rhizosphere and a root segment. From the root crown there are two stem xylem elements in series and a final leaf element (Fig. 10.1).

To build the supply function for the whole hydraulic network, we proceed by calculating water potentials in the network for each value of flow. For any given E value we start by calculating flows and potentials within the root system. After that, the water potential at the mid of the stem ($\Psi_{stem,1}$) is obtained using the inverse of the stem supply function and setting $\Psi_{up} = \Psi_{rootcrown}$, and the water potential at the upper end of the stem, $\Psi_{stem,2}$, is obtained similarly but setting $\Psi_{up} = \Psi_{stem,1}$. In both cases the maximum conductance for




Figure 10.12: Example of supply function for a root system (left) and its derivative (right) under different soil textures and starting from different soil water potential vectors.




Figure 10.13: Water potentials inside fine roots and at the root crown, for overall flow rates corresponding




Figure 10.14: Flow rates inside roots corresponding to each soil layer, for overall flow rates corresponding to fig. 10.12.

segments equal to $2 \cdot k_{stem,max,i}$. Leaf water potential (Ψ_{leaf}) is finally obtained using the inverse of the leaf supply function and setting $\Psi_{up} = \Psi_{stem,2}$ and assuming a steady-state flow E . The whole supply function $E(\Psi_{leaf})$ is obtained repeating these operations from $E = 0$ to a critical value E_{crit} .

The following figure shows network supply functions for each soil texture and starting from the different soil water potential configurations (see function `hydraulics_supplyFunctionNetwork()`):




Figure 10.15: Examples of supply functions for a hydraulic network describing the potential rate of water supply for transpiration (E) as function of the leaf pressure (Ψ_{leaf}), starting from different bulk soil (Ψ_{soil}) values and for different soil textures. Left/right panel shows uncavitated/cavitated supply functions.

As with previous representations of the soil-plant continuum, the derivative of $dE/d\Psi_{leaf}$ for the network topology is obtained numerically:

As with the root system, we can know the water potentials in different points of the continuum. Here we plot them for the results of silt loam texture and the first and last soil potential vectors defined above:

A supply function of the soil-plant hydraulic network defines a set steady-state flows and corresponding water potentials given fixed values of soil moisture. Hence, supply functions need to be updated whenever soil moisture is updated, i.e. once a day in the water balance model (see subsection 7.5). This frequency should be enough in most cases, since soil water potentials change more slowly than plant water potentials.

10.5 Sureau's sub-model of plant hydraulics

10.5.1 Constitutive equations

Water dynamics of the SPA system in SurEau-Ecos (Ruffault et al., 2022) are governed by partial differential equations for water mass conservation.

The water balance of each of the four plant compartments (symplasm and apoplasm for leaf and stem, respectively) is determined according to a generic equation. For a given plant compartment i , the water




Figure 10.16: Derivatives of supply functions corresponding to figure 10.15. Left/right panel shows the derivatives of uncavitated/cavitated supply functions.

balance differential equation is:

$$C_i \cdot \frac{d\Psi_i}{dt} + \sum_j k_{ij} \cdot (\Psi_i - \Psi_j) - S = 0 \quad (10.19)$$

and solved at each time step. In the above equation Ψ_i and Ψ_j would be the water potential of compartments i and j , respectively (i.e. $\Psi_{leaf, symp}$, $\Psi_{leaf, apo}$, $\Psi_{stem, symp}$ or $\Psi_{stem, apo}$; see Fig. 10.2); C_i is the capacitance associated to the compartment i and defined generically as:

$$C_i = \frac{Q_i}{\Psi_i} \quad (10.20)$$

where Q_i is the water volume content of the compartment; k_{ij} is the conductance between the two compartments and S represents an outflow of the target compartment. Flows in/out of compartments are stem cuticular transpiration ($E_{cuti, stem}$), leaf cuticular transpiration ($E_{cuti, leaf}$), leaf stomatal transpiration (E_{stom}), stem cavitation flux ($F_{cav, stem}$) and leaf cavitation flux ($F_{cav, leaf}$).

10.5.2 Plant and soil conductances

Overall plant conductance k_{plant} is defined as the inverse of the sum of resistances across the hydraulic network Fig. 10.2:

$$k_{plant} = \left(\frac{1}{\sum_{s=1}^S k_{root, s}} + \frac{1}{k_{stem, apo}} + \frac{1}{k_{leaf, apo}} + \frac{1}{k_{leaf, symp}} \right)^{-1} \quad (10.21)$$

Whenever needed, for each soil layer s , the conductance from the root surface to the root crown is updated using the current level of PLC_{stem} :

$$k_{root, s} = k_{root, max, s} \cdot (1.0 - PLC_{stem}) \quad (10.22)$$




Figure 10.17: Water potential in network elements corresponding to overall flow rates of figure 10.15. Left/right panel shows the derivatives of uncavitated/cavitated supply functions.

whereas the conductance from the root crown to the stem apoplasm is updated similarly:

$$k_{stem,apo} = k_{stem,apo,max} \cdot (1.0 - PLC_{stem}) \quad (10.23)$$

and the conductance from the stem apoplasm to the leaf apoplasm is updated using the current level of PLC_{leaf} :

$$k_{leaf,apo} = k_{leaf,apo,max} \cdot (1.0 - PLC_{leaf}) \quad (10.24)$$

Unlike plant conductances, which are updated at short time steps, the conductance in the bulk soil for each soil layer ($k_{soil,s}$) is updated once per day, from $\Psi_{soil,s}$ and using Van Genuchten equations.

10.5.3 Plant capacitances

Symplasmic reservoirs behave as variable plant capacitances related to the pressure volume curve, which corresponds to the water quantity (Q) changes in symplasmic cells, i.e.:

$$dQ/dt = C \cdot (d\Psi/dt) \quad (10.25)$$

from which we get an expression to update C :

$$C = \frac{dQ}{d\Psi} = Q^{sat} \cdot \frac{dRWC}{d\Psi} \quad (10.26)$$

where Q^{sat} is the water volume of the compartment at saturation. From the former general equation, the capacitance of the leaf symplasm (C_{LSym} ; in $mmol \cdot MPa^{-1} \cdot m^{-2}$) is updated using the derivative of the leaf symplasmic pressure-volume curve (RWC_{sym}^{leaf}) with respect to $\Psi_{leaf,sym}$ (Ruffault et al., 2022):

$$C_{LSym} = Q_{LSym}^{sat} \cdot \frac{dRWC_{sym}^{leaf}(\Psi_{leaf,sym})}{d\Psi_{leaf,sym}} \quad (10.27)$$

and the capacitance of the stem symplasm (C_{SSym} ; in $mmol \cdot MPa^{-1} \cdot m^{-2}$) is updated analogously:

$$C_{SSym} = Q_{SSym}^{sat} \cdot \frac{dRWC_{sym}^{stem}(\Psi_{stem,sym})}{d\Psi_{stem,sym}} \quad (10.28)$$

Equations for derivatives of the relative water content are given in Ruffault et al. (2022).

10.5.4 Cavitation flows

SurEau-Ecos also considers the capacitance effect of xylem cavitation (Hölttä et al., 2009), i.e., the water released to the streamflow when cavitation occurs (Ruffault et al., 2022). The non-cavitated part of the xylem receives a water flux from the cavitated part and is then transferred to adjacent compartments. The amount of water corresponding to a new cavitation event (F_{cav}) is derived from the quantity of water in the apoplasm at saturation (Q_{apo}^{sat}) and the temporal variations in PLC as follows:

$$F_{cav} = Q^{sat} \cdot \max\left(\frac{dPLC}{dt}, 0\right) \quad (10.29)$$

This flux is linearized in temporal variations in Ψ in order to express this flux in the form of a Darcy's law. For that purpose an equivalent conductance k_{cav} is introduced as follows:

$$F_{cav} = Q^{sat} \cdot \max\left(\frac{dPLC}{d\Psi} \cdot \frac{d\Psi}{dt}, 0\right) \quad (10.30)$$

$$\approx k_{cav} \cdot \max(\Psi_{cav} - \Psi, 0) \quad (10.31)$$

where Ψ_{cav} is the minimum value of Ψ reached for the apoplastic compartment, $dPLC/d\Psi$ is the derivative of PLC with respect to Ψ and k_{cav} is:

$$k_{cav} = -\frac{Q^{sat} \cdot (dPLC/d\Psi)}{dt} \quad (10.32)$$

The equation for $dPLC/d\Psi$ is given in Ruffault et al. (2022).

10.5.5 Transpiration flows

Stem cuticular transpiration ($E_{cuti,stem}$), leaf cuticular transpiration ($E_{cuti,leaf}$) and leaf stomatal transpiration (E_{stom}) are described in section 12.2.1.

10.5.6 Numerical scheme

Three numerical schemes were presented and compared in Ruffault et al. (2022) to solve the system of differential equations, but only the semi-implicit solving approach has been implemented in **medfate**.

Chapter 11

Leaf/canopy photosynthesis

In chapter 10 we introduced plant hydraulics under Sperry and SurEau models. In this chapter we present the building blocks for the calculation of leaf and canopy photosynthesis. How transpiration and photosynthesis values are determined in the model will be explained in chapter 12.

11.1 Leaf energy balance, gas exchange and photosynthesis

Let us assume that leaf transpiration rate E and leaf water potential Ψ_{leaf} are known. If we also know air temperature, water vapor pressure and the light conditions in which leaves are, we can calculate leaf photosynthesis rate. In a nutshell, E is first used to calculate leaf temperature from an evaluation of the leaf energy balance. The diffusive conductances of the leaf to H_2O and CO_2 are obtained from water supply and leaf water vapor deficit. Finally, the gross/net assimilation rate is then obtained from the diffusive conductance and a modelled curve between assimilation and leaf internal CO_2 concentration. The following subsections detail each of these steps separately.

11.1.1 Leaf temperature

Leaf temperature (T_{leaf} ; in $^{\circ}C$) can be calculated for any given flow rate E using an energy balance equation (Campbell and Norman, 1998):

$$T_{leaf} = T_{air} + \frac{\Phi_{SWR,leaf} + L_{net,leaf} - \lambda_v(T_{air}) \cdot E_{two-sided}}{C_p \cdot (g_r + g_{Ha})} \quad (11.1)$$

where $\Phi_{SWR,leaf}$ (in $W \cdot m^{-2}$) is the instantaneous shortwave radiation absorbed per leaf area unit, $L_{net,leaf}$ is the instantaneous net long-wave radiation per leaf area unit, $E_{two-sided}$ is the flow rate E converted to $mol \cdot s^{-1} \cdot m^{-2}$ per two-sided leaf area basis, ϵ is longwave radiation emissivity (0.97), σ is the Stephan-Boltzman constant, T_{air} is the temperature of the air surrounding the leaf (in $^{\circ}C$), $C_p = 29.3 J \cdot mol^{-1} \cdot ^{\circ}C^{-1}$ is the specific heat capacity of dry air at constant pressure and $\lambda_v(T_{air})$ is the latent heat of vaporization (in $J \cdot mol^{-1}$) corresponding to T_{air} (see utility functions in the **meteoland** reference manual). If canopy energy balance is performed assuming a single canopy layer T_{air} corresponds to T_{can} , the whole-canopy air temperature resulting from the energy balance of the previous step (see section 13.1.3). If a multi-layer canopy energy balance is used, T_{air} corresponds to $T_{air,j}$, the air temperature of the canopy layer j that contains the leaf. Finally, g_r and g_{Ha} are the radiative and heat conductance values (in $mol \cdot m^{-2} \cdot s^{-1}$), respectively (Campbell and Norman, 1998):

$$g_r = \frac{4 \cdot \epsilon \cdot \sigma \cdot (T_{air} + 273.15)^3}{C_p} \quad (11.2)$$

$$g_{Ha} = 0.189 \cdot (u_{leaf}/d)^{0.5} \quad (11.3)$$

where u_{leaf} is the leaf-level wind speed (in $m \cdot s^{-1}$) and d is 0.72 times the leaf width (species parameter **LeafWidth** in cm). As before, u_{leaf} corresponds to u_j , the wind-speed at the canopy layer j where the leaf is located (see 25.2).

11.1.2 Leaf vapor pressure deficit

Vapor pressure deficit in the leaf (VPD_{leaf} , in kPa) is the driving force for leaf transpiration. It is calculated as the difference between leaf and surrounding air vapor pressures:

$$VPD_{leaf} = e_{leaf} - e_{air} \quad (11.4)$$

Where e_{air} is the water vapor pressure of the air surrounding the leaf (but beyond the leaf boundary layer) and e_{leaf} is the leaf vapor pressure, which can be calculated from leaf temperature (T_{leaf}) and leaf water potential (Ψ_{leaf}):

$$e_{leaf} = e_{sat}(T_{leaf}) \cdot \exp \left\{ \frac{2.17 \cdot \Psi_{leaf}}{T_{leaf} + 273.15} \right\} \quad (11.5)$$

where $e_{sat}(T)$ is a function giving the saturated vapor pressure for temperature T (see utility functions of the **meteoland** reference manual). If a single-canopy energy balance is considered, the water vapor pressure of the air surrounding the leaf is taken as the average atmospheric water vapor pressure of the day, i.e. $e_{air} = e_{atm}$. This assumes a well-coupled canopy in terms of water vapor. When a multi-layer canopy energy balance is simulated, e_{air} will correspond to the water vapor pressure $e_{air,j}$ of the canopy layer j where the leaf is located.

11.1.3 Leaf gas exchange

Diffusive conductance to water vapor (g_w ; in $mol H_2O \cdot s^{-1} \cdot m^{-2}$) and to carbon dioxide (g_c ; in $mol CO_2 \cdot s^{-1} \cdot m^{-2}$) between the leaf mesophyll and the air surrounding the leaf are assumed to be related by a single constant:

$$g_c = g_w / 1.6 \quad (11.6)$$

Moreover, if transpiration flow E (here expressed in $mol H_2O \cdot s^{-1} \cdot m^{-2}$) and VPD_{leaf} are known, then we can use Fick's law to estimate the diffusive conductance g_w :

$$g_w = E \cdot \frac{P_{atm}}{VPD_{leaf}} \quad (11.7)$$

Otherwise, g_w or g_c can be estimated by its components. At this point it is important to distinguish that in the Sperry sub-model leaf water is assumed to flow through stomata only, whereas in SurEau sub-model cuticular transpiration is also possible.

Gas exchange through stomata

In the Sperry sub-model, g_w is the result of combining leaf stomatal conductance and the conductance of the leaf boundary layer:

$$g_w = (g_{sw}^{-1} + g_{bound}^{-1})^{-1} \quad (11.8)$$

where g_{sw} is the stomatal conductance and g_{bound} is the leaf boundary conductance (all in $mol H_2O \cdot s^{-1} \cdot m^{-2}$). If we estimate g_{bound} using:

$$g_{bound} = 0.397 \cdot (u_{leaf}/d)^{0.5} \quad (11.9)$$

then the only unknown to determine g_w (or g_c) is stomatal conductance g_{sw} . Sperry's sub-model makes use of eq. (11.7) to estimate g_w from E . Then, stomatal conductance is determined using eq. (11.8), i.e.:

$$g_{sw} = (g_w^{-1} - g_{bound}^{-1})^{-1} \quad (11.10)$$

Ensuring here that $g_{sw} \geq 0.0$. If transpiration flow E is not known, one needs to find alternative ways to determine g_{sw} , which has lead to the development of algorithms that simultaneously determine g_{sw} , E and photosynthesis (see chapter 12).

Gas exchange through the leaf cuticle

In the SurEau sub-model, the diffusive conductance regulating gas exchange through the cuticle g_{wc} is the result of combining cuticular conductance and the conductance of the leaf boundary layer:

$$g_{wc} = (g_{cuti}^{-1} + g_{bound}^{-1})^{-1} \quad (11.11)$$

which means that the overall diffusive conductance is, instead of eq. (11.8):

$$g_w = ((g_{sw} + g_{cuti})^{-1} + g_{bound}^{-1})^{-1} \quad (11.12)$$

11.1.4 Leaf photosynthesis

Rubisco-limited photosynthesis rate A_c (in $\mu\text{mol CO}_2 \cdot \text{s}^{-1} \cdot \text{m}^{-2}$) is modelled using (Collatz et al., 1991; Medlyn et al., 2002):

$$A_c = \frac{V_{max} \cdot (C_i - \Gamma^*)}{C_i + K_c \cdot (1 + O_a/K_o)} \quad (11.13)$$

where V_{max} is Rubisco's maximum carboxylation rate (in $\mu\text{mol CO}_2 \cdot \text{s}^{-1} \cdot \text{m}^{-2}$), C_i is the internal carbon dioxide concentration (in $\mu\text{mol} \cdot \text{mol}^{-1}$), Γ^* is the compensation point (in $\mu\text{mol} \cdot \text{mol}^{-1}$), K_c (in $\mu\text{mol} \cdot \text{mol}^{-1}$) and K_o (in $\text{mmol} \cdot \text{mol}^{-1}$) are Michaelis-Menten constants for carboxylation and oxygenation, respectively, and O_a is the atmospheric oxygen concentration (i.e. $209 \text{ mmol} \cdot \text{mol}^{-1}$). Γ^* , K_c and K_o depend on leaf temperature (T_{leaf} , in $^{\circ}\text{C}$) (Bernacchi et al., 2001):

$$\Gamma^* = 42.75 \cdot e^{\frac{37830 \cdot (T_{leaf} - 25)}{298 \cdot R \cdot (T_{leaf} - 273)}} \quad (11.14)$$

$$K_c = 404.9 \cdot e^{\frac{79430 \cdot (T_{leaf} - 25)}{298 \cdot R \cdot (T_{leaf} - 273)}} \quad (11.15)$$

$$K_o = 278.4 \cdot e^{\frac{36380 \cdot (T_{leaf} - 25)}{298 \cdot R \cdot (T_{leaf} - 273)}} \quad (11.16)$$

Electron transport-limited photosynthesis A_e (in $\mu\text{mol CO}_2 \cdot \text{s}^{-1} \cdot \text{m}^{-2}$) was obtained from Medlyn et al. (2002):

$$A_e = \frac{J}{4} \cdot \frac{C_i - \Gamma^*}{C_i + 2 \cdot \Gamma^*} \quad (11.17)$$

$$J = \frac{(\alpha \cdot Q_{PAR,leaf} + J_{max}) - \sqrt{(\alpha \cdot Q_{PAR,leaf} + J_{max})^2 - 4.0 \cdot c \cdot \alpha \cdot Q_{PAR,leaf} \cdot J_{max}}}{2 \cdot c} \quad (11.18)$$

where $\alpha = 0.3 \text{ mole} \cdot \text{mol photon}^{-1}$ is the quantum yield of electron transport, $Q_{PAR,leaf}$ is the absorbed PAR photon flux density per leaf area ($\mu\text{mol photon} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$), which is calculated from absorbed PAR per leaf area ($\Phi_{PAR,leaf}$; in $\text{W} \cdot \text{m}^{-2}$):

$$Q_{PAR,leaf} = \Phi_{PAR,leaf} \cdot 546 \cdot 0.836 \cdot 10^{-2} \quad (11.19)$$

where 546 is the wavelength in nm . J_{max} and J are the maximum and actual rate of electron transport (both in $\mu\text{mol e} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) and $c = 0.9$ defines the curvature of the light-response curve. The gross assimilation rate A at a given C_i is the minimum of A_e and A_c . To obtain a smooth A -vs- C_i curve we used (Collatz et al., 1991):

$$A = \frac{(A_c + A_e) - \sqrt{(A_c + A_e)^2 - 4.0 \cdot c' \cdot A_e \cdot A_c}}{2 \cdot c'} \quad (11.20)$$

where $c' = 0.98$ is a curvature factor. The temperature dependence of J_{max} and V_{max} relative to 25°C (i.e., relative to input parameters $J_{max,298}$ and $V_{max,298}$) is modelled using Leuning (2002) (his eq. 1 with parameters from his Table 2). The internal CO_2 concentration, C_i , needs to be known to calculate A using the previous equations. Sperry et al. (2016) use a second equation for A which uses g_c , the diffusive conductance to CO_2 (see eq. (11.6)):

$$A = g_c \cdot (C_{air} - C_i) \quad (11.21)$$

where C_{air} is the CO_2 concentration (in $\mu mol \cdot mol^{-1}$) in the air surrounding the leaf (beyond the leaf boundary layer). If a single-layer canopy energy balance is used then $C_{air} = C_{atm}$, the atmospheric CO_2 concentration (see weather input and parameter `Catm` in function `defaultControl()`), whereas if a multi-layer canopy energy balance is used C_{air} corresponds to $C_{air,j}$, the CO_2 concentration in the canopy layer j where the leaf occurs, analogously to T_{air} and e_{air} . Combining the two equations for A and finding the root of the resulting equation using Newton-Raphson method allows determining C_i and therefore A . Finally, leaf net photosynthesis (A_n ; i.e. accounting for autotrophic respiration) is calculated as:

$$A_n = A - 0.015 \cdot V_{max} \quad (11.22)$$

11.2 Crown/canopy photosynthesis

In the previous subsection we showed how to calculate photosynthesis at the leaf level, but gross or net assimilation needs to be up-scaled to the crown or canopy levels, while accounting for the variation of leaf photosynthetic conditions across canopies. In this section, we briefly describe different alternatives for canopy/crown photosynthesis and justify our modelling choice.

11.2.1 Multi-layer canopy photosynthesis model

Ideally, crown or canopy photosynthesis requires repeating the calculations of leaf temperature, leaf VPD, leaf gas conductance and photosynthesis for every leaf. Multi-layer canopy models, however, divide the canopy into vertical layers and determine photosynthesis for each one (and each plant species or plant cohort, if required). Multi-layer models can account for the fact that environmental variables usually vary across the crown, including direct/diffuse absorbed radiation, air temperature, wind speed, water vapor pressure and CO_2 concentration. The amount of SWR and PAR absorbed per unit of leaf area at each canopy layer was shown in fig. 9.2.

Not only environmental factors, but leaves themselves may be different across canopy layers. Importantly, it is generally accepted that sunlit and shade leaves need to be treated separately (De Pury and Farquhar, 1997). Separating the two kinds of leaves acknowledges that they operate at different parts of the light-saturation curve. Following De Pury and Farquhar (1997), we further assume that maximum carboxylation and electron transport rates are highest for leaves at the top of the canopy and there is an exponential decrease from there towards the bottom, where maximum rates are 50% of those at the top:

$$V_{max,298,i,j} = V_{max,298,i} \cdot \exp(-0.713 \cdot \sum_{h>i} LAI_{act,h,j} / LAI_{act,i}) \quad (11.23)$$

$$J_{max,298,i,j} = J_{max,298,i} \cdot \exp(-0.713 \cdot \sum_{h>i} LAI_{act,h,j} / LAI_{act,i}) \quad (11.24)$$

where $LAI_{act,i,j}$ is the LAI value of the plant cohort i at a given canopy layer j and $LAI_{act,i}$ is the expanded LAI of the plant cohort. The following figure illustrate this decrease for the single-species canopy example of section 9.1.3:

In a multi-layer canopy photosynthesis model, gross and net photosynthesis values (i.e. A and A_n) are determined for sunlit and shade leaves of each cohort in each canopy layer. Then, sunlit and shade photosynthesis values should be averaged across the crown for each plant cohort. Assuming that Ψ_{leaf} is equal for all leaves across the crown, the function $A(\Psi_{leaf})$ would be obtained for each plant cohort.

11.2.2 Big-leaf canopy photosynthesis model

Multi-layer canopy photosynthesis models allow evaluating leaf conditions, stomatal conductance and photosynthesis for different points of the canopy. However, this comes at high computational cost. For this reason, many models implement what is called the *big-leaf approximation*. Assuming that wind speed, temperature, water vapor pressure and CO_2 concentration are similar for all leaves and that the distribution of photosynthetic capacity between leaves is in proportion to the profile of absorbed irradiance then the equation describing leaf photosynthesis will also represent canopy photosynthesis (Sellers et al. 1992).




Figure 11.1: Decrease of Rubisco maximum carboxylation rate across the canopy

11.2.3 Sun-shade canopy photosynthesis model

An alternative between multi-layer and big-leaf canopy photosynthesis models is to collapse variation of photosynthetic conditions into two leaf classes: sunlit and shade leaves. While big-leaf canopy models are known to be unaccurate under some situations, sun-shade canopy models (De Pury and Farquhar, 1997) provide estimates that are close to multiple layer models (Hikosaka et al., 2016). The sun-shade canopy photosynthesis model was adopted here. Assuming that wind speed, temperature, water vapor pressure and and CO_2 concentration are similar for all leaves, sun-shade models involve the following steps:

- a. Aggregate the leaf area of sunlit/shade leaves across layers:

$$LAI_{sunlit,i} = \sum_{j=1}^l LAI_{sunlit,i,j} \quad (11.25)$$

$$LAI_{shade,i} = \sum_{j=1}^l LAI_{shade,i,j} \quad (11.26)$$

where $LAI_{sunlit,i,j}$ and $LAI_{shade,i,j}$ are the leaf area index of sunlit and shade leaves for cohort i in canopy layer j , from eq. (9.8).

- b. Average the SWR/PAR absorbed by leaves of each kind across layers. The average light absorbed by sunlit/shaded foliage of cohort i per ground area unit is found using:

$$\Phi_{abs,sunlit,i} = \frac{\sum_{j=1}^l K_{abs,sunlit,i,j}}{LAI_{sunlit,i}} \quad (11.27)$$

$$\Phi_{abs,shade,i} = \frac{\sum_{j=1}^l K_{abs,shade,i,j}}{LAI_{shade,i}} \quad (11.28)$$

where $K_{abs,sunlit,i,j}$ and $K_{abs,shade,i,j}$ are the light absorbed per ground area unit by sunlit/shade leaves of cohort i at layer j (see section 9.1.3). Analogous equations were already given for the net long-wave radiation balance of sunlit leaves ($L_{net,sunlit,i}$) and shade leaves ($L_{net,shade,i}$) in section 9.2.3.

- c. Average the maximum carboxylation (respectively, electron transport) rates across layers, again

separating sunlit and shade leaves:

$$V_{max,sunlit,298,i} = \frac{\sum_{j=1}^l V_{max,298,i,j} \cdot LAI_{sunlit,i,j}}{LAI_{sunlit,i}} \quad (11.29)$$

$$V_{max,shade,298,i} = \frac{\sum_{j=1}^l V_{max,298,i,j} \cdot LAI_{shade,i,j}}{LAI_{shade,i}} \quad (11.30)$$

- d. Use $V_{max,sunlit,298,i}$ as $V_{max,298}$ in Leuning (2002) to obtain V_{max} for eq. (11.13); $\Phi_{SWR,sunlit,i}$ as $\Phi_{SWR,leaf}$ and $L_{net,sunlit,i}$ as $L_{net,leaf}$ in eq. (11.1); and $\Phi_{PAR,sunlit,i}$ as $\Phi_{PAR,leaf}$ in eq. (11.19) to estimate sunlit leaf photosynthesis, which can be up-scaled to the crown level multiplying by $LAI_{sunlit,i}$. The same would be done for shade leaves. In a sun-shade canopy model one then calls the photosynthesis function twice (i.e. once for shade leaves and once for sunlit leaves) for each plant cohort i .

In the sun-shade photosynthesis model, the question arises on how to determine layer j for sunlit or shade leaves of a given cohort. The choice is done by calculating the height corresponding to the mass center of sunlit leaves or shade leaves. The canopy layer j where this mass center height is contained is chosen as the layer from which environmental conditions will be taken. For any given plant cohort, sunlit leaves will take their environmental conditions from layers above (or equal to) those corresponding to shade leaves.

11.2.4 Within-canopy variation in environmental conditions

The environmental conditions surrounding leaves depend on the choice of canopy representation:

- If a single-layer canopy energy balance is used, CO_2 concentration and vapor pressure are assumed equal to the atmosphere (i.e., $e_{air} = e_{atm}$ and $C_{air} = C_{atm}$), whereas air temperature is that of the canopy (i.e., $T_{air} = T_{can}$), which assumes a well-coupled canopy. However, leaf-level wind speed (u_{leaf}) can still be different for different canopy layers, which will lead to different leaf temperature and transpiration due to the influence of wind on boundary layer conductance for water vapor and heat.
- If a multi-layer canopy energy balance is used, all four environmental variables can differ between canopy layers (i.e. $T_{air,j}$, $e_{air,j}$, $C_{air,j}$ and u_j).

Chapter 12

Stomatal regulation, transpiration and photosynthesis

Plants must open their stomata to acquire CO_2 and perform photosynthesis, but doing so promotes water loss. This trade-off has resulted in a tight coordination between capacity to supply and transpire water (hydraulic conductance and diffusive conductance to water vapor) and the maximum capacity for photosynthesis (carboxylation rate and electron-transport rate). For modelling purposes, this carbon-for-water trade-off means that hydraulics, stomatal conductance, transpiration and photosynthesis need to be estimated simultaneously. In chapters 10 and 11 we introduced plant hydraulics and photosynthesis, respectively, but we did not explain how are actual transpiration and photosynthesis values determined. This depends on the sub-model chosen and is the subject of the present chapter. The following two sections detail how stomatal regulation and instantaneous transpiration, soil water uptake and photosynthesis rates are determined under Sperry and SurEau sub-models. The final section of the chapter is devoted to scaling instantaneous rates to the subdaily time-step.

12.1 Stomatal regulation, transpiration and photosynthesis under Sperry's sub-model

Sperry et al. (2017) proposed estimating stomatal conductance from the instantaneous maximization of profit, defined as the difference between photosynthesis gain and hydraulic cost (both normalized for comparability). First, water supply and photosynthesis functions are defined. Stomatal regulation, transpiration and photosynthesis are then estimated separately for sunlit and shade leaves, according to profit maximization. Finally, a leaf-average instantaneous transpiration rate is determined depending on sunlit/shade transpiration rates and their leaf area contributions. This final step also determines the water potentials along the plant hydraulic network. The following sections provide details for all these steps.

12.1.1 Water supply function

In chapter 10 we introduced the necessary concepts of steady-state plant hydraulics to define the *hydraulic supply function for the soil-plant continuum*. The supply function specifies the flow rate, as per leaf area, for values of leaf water potential and given a soil moisture status (Sperry et al., 1998; Sperry and Love, 2015; Sperry et al., 2016). The supply function is calculated once per day for each plant cohort. Here we assume that hydraulic conductance k is in $mmol H_2O \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$ of leaf area, transpiration rate E_{leaf} in $mmol H_2O \cdot s^{-1} \cdot m^{-2}$ of leaf area and leaf water potential Ψ_{leaf} is in MPa.

12.1.2 From water supply to sunlit/shade photosynthesis functions

According to what we learned in section 11.1, for each pair of E_{leaf} and Ψ_{leaf} values, we have a corresponding leaf temperature (T_{leaf} ; in °C), leaf-to-air vapor pressure deficit (VPD_{leaf} ; in kPa), leaf water vapor conductance (g_{sw} ; in $\text{mol } H_2O \cdot s^{-1} \cdot m^{-2}$) and, finally the leaf gross and net (i.e. after accounting for autotrophic respiration) photosynthesis assimilation rates (A_g and A_n ; both in $\mu\text{mol } CO_2 \cdot s^{-1} \cdot m^{-2}$). In short, the supply function generates a **photosynthesis function**. Since the model deals with canopies and not single leaves, however, different parts of the crowns of plant cohorts may be in different canopy positions. Calculating photosynthesis at the canopy level requires dividing the canopy into vertical layers, differentiating between *sunlit* and *shade* leaves and determining **photosynthesis functions for sunlit and shade leaves separately**, as explained in section 11.2.

In this section we describe leaf photosynthesis functions, which define leaf gas exchange, photosynthesis and energy balance for values of leaf water potential. This requires knowing wind, water vapor and temperature conditions of the air surrounding leaves (section 13.1.3), the short-wave radiation absorbed by leaf elements (section 9.1.3) and the leaf long-wave radiation balance (section 9.2).

Leaf temperature

By inspecting eq. (11.1), we can conclude that transpiration flow decreases leaf temperature, whereas radiation increases it and wind speed makes it more similar to the temperature of the surrounding air. The following figures illustrate the effect of varying wind speed and flow rate on T_{leaf} for two contrasted leaf widths (see function `biophysics_leafTemperature`):




Figure 12.1: Values of T_{leaf} for two leaf widths and varying values of wind speed and flow rate, calculated for 24°C air temperature and $740 \text{ W} \cdot m^{-2}$ instantaneous absorbed radiation (including SWR and LWR).

Let us now fix wind speed at the leaf level to $u_{leaf} = 2 \text{ m/s}$. The application of the leaf energy balance equation (11.1) to the $E(\Psi_{leaf})$ curves corresponding to the complete hydraulic network (10.4.5) yields the following $T_{leaf}(\Psi_{leaf})$ curves:

Leaf vapor deficit




Figure 12.2: Examples of leaf temperature functions for a hydraulic network, corresponding to fig. 10.15 and for different soil textures. Left/right panel shows values for uncavitated/cavitated supply functions.

Since e_{leaf} decreases when leaf temperature decreases in eq. (11.5), increasing transpiration decreases leaf VPD as a result of decreasing leaf temperature. To illustrate this effect, let us assume the following values of relative humidity, yielding a $e_{air} = e_{atm} = 1.91 \text{ kPa}$:

```
RHmin = 60
RHmax = 75
VPatm = meteoland::utils_averageDailyVP(Tmin, Tmax, RHmin, RHmax)
VPatm
```

```
## [1] 1.912181
```

The application of eqs. (11.4) and (11.5) to the $T_{leaf}(\Psi_{leaf})$ curves of fig. 12.2 yields the following $VPD_{leaf}(\Psi_{leaf})$ curves:

Note that the VPD decreasing curves do not start at the same VPD_{leaf} value despite corresponding to the same T_{leaf} value, because of the effect of Ψ_{leaf} on e_{leaf} in eq. (11.5).

Stomatal conductance

The application of equations for g_w , g_{bound} and g_{sw} to the $VPD_{leaf}(\Psi_{leaf})$ curves yields the following stomatal conductance $g_{sw}(\Psi_{leaf})$ curves:

In the previous figure we have indicated the thresholds of $g_{sw,min}$ and $g_{sw,max}$, the species-specific minimum and maximum water vapour conductances (i.e. conductances when stomata are fully closed and fully open, respectively; see parameters `Gswmin` and `Gswmax` in `SpParamsMED`). g_{sw} cannot exceed $g_{sw,max}$ so that some flow rates may not be possible (see stomatal regulation below). However, $g_{sw,max}$ should quickly become non-limiting as soil dries (i.e. reducing E) or VPD_{leaf} increases (Sperry et al., 2017).

Leaf photosynthesis




Figure 12.3: Examples of leaf vapour pressure deficit (VPD_{leaf}) functions for a hydraulic network, corresponding to fig. 10.15 and for different soil textures. Left/right panel shows values for uncavitated/cavitated supply functions.




Figure 12.4: Examples of stomatal conductance to water vapor (g_{sw}) functions for a hydraulic network, corresponding to fig. 10.15 and for different soil textures. Left/right panel shows values for uncavitated/cavitated supply functions. Minimum and maximum conductance values ($g_{sw,min} = 0.0045$ and $g_{sw,max} = 0.3$) are indicated using dashed lines.

Thus, after defining PAR photon flux density, canopy air CO_2 concentration and maximum rate parameters:

```
Q = 2000
Cair = 386
Vmax298 = 100
Jmax298 = 1.67*Vmax298
```

one can obtain the following $A(\Psi_{leaf})$ curves from $T_{leaf}(\Psi_{leaf})$ and $g_{sw}(\Psi_{leaf})$:




Figure 12.5: Examples of gross photosynthesis (A) functions for a hydraulic network, corresponding to fig. 10.15 and for different soil textures. Left/right panel shows values for uncavitated/cavitated supply functions.


Comparison of big-leaf, sun-shade and multi-canopy photosynthesis models

The figure below provides the canopy photosynthesis functions obtained using the multi-layer canopy photosynthesis model (top), a sunshade canopy photosynthesis model (center) or a big-leaf photosynthesis model (bottom). These were generated using functions `photo_multilayerPhotosynthesisFunction()`, `photo_sunshadePhotosynthesisFunction()` and `photo_leafPhotosynthesisFunction()`, respectively, and assuming homogeneous wind, temperature and water vapor pressure through the canopy. Thus, only absorbed radiation varied across layers and leaf types. Note the coincidence between the multi-layer and the sun-shade models.

12.1.3 Leaf stomatal regulation by profit maximization

Wolf et al. (2016) proposed the **carbon maximization criterion**, which states that at each instant in time the stomata regulate canopy gas exchange and pressure to achieve the maximum profit, which is the maximum difference between photosynthetic gains and costs, the latter associated with hydraulic vulnerability attained with low water potentials. Such approach has been shown to be supported by data from global forest biomes (Anderegg et al., 2018). Building on this approach, Sperry et al. (2017) presented a profit maximization function where hydraulic costs of opening the stomata are compared against photosynthetic gains. Details of their formulation are given in this section. Stomatal regulation is performed in **medfate** separately for sunlit and shade leaves.

Cost and gain functions



The hydraulic supply function is used to derive a transpirational *cost function* $\theta_1(\Psi_{leaf})$ that reflects the increasing damage from cavitation and the greater difficulty of moving water along the continuum (Sperry et al., 2016):

$$\theta(\Psi_{leaf}) = \frac{k_{c,max} - k_c(\Psi_{leaf})}{k_{c,max} - k_{crit}} \quad (12.1)$$

where $k_c(\Psi_{leaf}) = dE/d\Psi(\Psi)$ is the slope of the supply function corresponding to (leaf) water potential Ψ_{leaf} , $k_{c,max}$ is the maximum slope of the supply function (which occurs when $E_{leaf} = 0$), i.e. the maximum whole-plant conductance for the current soil moisture conditions, and $k_{crit} = k_c(\Psi_{crit})$ is the slope of the supply function at $E_{leaf} = E_{crit}$ the critical flow beyond which hydraulic failure occurs.

The following figures illustrate the $\theta(\Psi_{leaf})$ curves corresponding to the supply functions:




Figure 12.7: Cost functions (i.e. $\theta(\Psi_{leaf})$) obtained for a hydraulic network, corresponding to fig. 10.15 and different soil textures and soil water potentials. Left/right panels show values for uncavitated/cavitated supply functions, respectively.

The normalized photosynthetic *gain function* $\beta(\Psi_{leaf})$ reflects the actual assimilation rate with respect to the maximum (Sperry et al., 2016):

$$\beta(\Psi_{leaf}) = \frac{A(\Psi_{leaf})}{A_{max}} \quad (12.2)$$

where A_{max} is the instantaneous maximum (gross) assimilation rate estimated over the full Ψ_{leaf} range (including values that imply a stomatal conductance larger than the maximum).

The following figures illustrate the $\beta(\Psi_{leaf})$ curves corresponding to the supply and assimilation functions:

Profit maximization at the leaf level

Sperry et al. (2017) suggested that stomatal regulation can be effectively estimated by determining the maximum of the *profit function* ($Profit(\Psi_{leaf})$):

$$Profit(\Psi_{leaf}) = \beta(\Psi_{leaf}) - \theta(\Psi_{leaf}) \quad (12.3)$$




Figure 12.8: Gain function ($\beta(\Psi_{leaf})$) obtained for a hydraulic network, corresponding to fig. 10.15 and different soil textures and soil water potentials. Left/right panels show values for uncavitated/cavitated supply functions, respectively.

The maximization is achieved when the slopes of the gain and cost functions are equal:

$$\frac{\delta\beta(\Psi_{leaf})}{\delta\Psi_{leaf}} = \frac{\delta\theta(\Psi_{leaf})}{\delta\Psi_{leaf}} \quad (12.4)$$

Instantaneous profit maximization assumes a ‘use it or lose it’ reality with regards to available soil water. Because the gain function accelerates more quickly from zero and reaches 1 sooner than the cost function, their maximum difference occurs at intermediate Ψ_{leaf} values. Once Ψ_{leaf} that maximizes profit is determined, the values of the remaining variables are also determined. At this point, it may happen that $g_{sw}(\Psi_{leaf})$ is lower than the minimum (i.e. cuticular) water vapor conductance ($g_{sw,min}$) or larger than the maximum water vapor conductance ($g_{sw,max}$). These thresholds need to be taken into account when determining the maximum of the profit function.

The following figures illustrate the $Profit(\Psi_{leaf})$ curves corresponding to the previous cost and gain curves:

Squares in the previous figures indicate the maximum profit points in each situation. The drier the soil, the closer is the maximum profit Ψ_{leaf} to soil water potential as one would expect intuitively (i.e. a smaller drop in water potential along the hydraulic pathway). Note that when the soil is very dry the squares are to the right of the true maximum. This is because the `transp_profitMaximization()` function takes into account the minimum and maximum stomatal conductance and, in this case, does not allow optimum stomatal conductances below the minimum (cuticular) value.

Note that θ , β and $Profit$ functions can be expressed as a function of stomatal conductance, instead of leaf water potential. This allows visualizing more clearly the effect of $g_{sw,min}$ and $g_{sw,max}$ thresholds on the maximum profit optimization strategy, as illustrated in the following figures.




Figure 12.9: Profit functions (i.e. $Profit(\Psi_{leaf})$) obtained for a hydraulic network, corresponding to fig. 10.15 and different soil textures and soil water potentials. Left/right panels show values for uncavitated/cavitated supply functions, respectively.




Figure 12.10: Profit function as a function of stomatal conductance, corresponding to fig. 12.9 and different soil textures and soil water potentials. Left/right panels show values for uncavitated/cavitated supply functions, respectively.

12.1.4 Plant-level transpiration rates and plant water potentials

In the previous section, we considered stomatal regulation at the level of sunlit or shade leaves only. At the plant cohort level, the gain function could be build from a crown photosynthesis function $A(\Psi_{leaf})$, as shown in section 11.2. However, applying the profit maximization approach for a single crown photosynthesis function would imply the assumption that the same stomatal aperture occurs in all leaves of the plant cohort, independently of whether they are in shade or sunlit. A more realistic approach is to determine stomatal regulation by profit maximization for sunlit and shade leaves separately. The gain function and profit maximization calculations conducted for each leaf type yield instantaneous leaf water potentials $\Psi_{leaf,sunlit}$ and $\Psi_{leaf,shade}$ and instantaneous flow values E_{shade} and E_{sunlit} from the supply functions. The corresponding photosynthesis functions allow determining values for leaf temperatures $T_{leaf,sunlit}$ and $T_{leaf,shade}$, vapor pressure deficits $VPD_{leaf,sunlit}$ and $VPD_{leaf,shade}$, stomatal conductance $g_{sw,sunlit}$ and $g_{sw,shade}$ and net photosynthesis rates $A_{n,sunlit}$ and $A_{n,shade}$. This is a lot of useful information at the leaf level for each plant cohort, but we also need transpiration and photosynthesis values at the plant cohort-level.

The average instantaneous flow rate ($\hat{E}$, in $mmol H_2O \cdot s^{-1} \cdot m^{-2}$) per leaf area unit of a plant cohort is the weighed average:

$$\hat{E} = \frac{E_{shade} \cdot LAI_{sunlit} + E_{sunlit} \cdot LAI_{shade}}{LAI_{act}} \quad (12.5)$$

where LAI_{sunlit} and LAI_{shade} are the cohorts LAI values for sunlit and shade leaves, from eq. (9.8), and LAI_{act} is the leaf area index of the plant cohort. The model then uses the hydraulic supply function to find the transpiration rate E numerically closest to $\hat{E}$ (remember that the supply function is build in discrete steps). Finding the E value numerically closest to $\hat{E}$ determines the E_{plant} , the instantaneous transpiration flow, and also leads to setting values for current time step of several other variables, such as water potentials (Ψ_{leaf} , Ψ_{stem} , $\Psi_{rootcrown}$, ...), the slope of the supply function ($dE/d\Psi$), and instantaneous soil water uptake rates (U_s).

12.2 Stomatal regulation, transpiration and photosynthesis under Sureau's sub-model

As mentioned in 7.5, with SurEau's sub-model, leaf energy balance, stomatal and cuticular conductances, transpirational flows, photosynthesis and plant hydraulics are computed iteratively in small temporal sub-steps (e.g. 10 min). The following sections detail how leaf energy balance, conductances, transpiration and photosynthesis are estimated, whereas plant hydraulics were described in 10.5.

12.2.1 Transpiration flows

In Sureau-ECOS (Ruffault et al., 2022), plants lose water through stomatal transpiration (E_{stom}), cuticular transpiration of the leaf ($E_{leaf,cuti}$) and cuticular transpiration of the stem ($E_{stem,cuti}$), see Fig. 10.2. Therefore, the total plant transpiration E_{plant} is decomposed as:

$$E_{plant} = E_{leaf} + E_{stem,cuti} = E_{stom} + E_{leaf,cuti} + E_{stem,cuti} \quad (12.6)$$

Since there are sunlit and shade, leaves, both E_{stom} and $E_{leaf,cuti}$ are computed as weighted averages of sunlit and shade leaf values:

$$E_{stom} = \frac{E_{stom,sunlit} \cdot LAI_{sunlit} + E_{stom,shade} \cdot LAI_{shade}}{LAI_{act}} \quad (12.7)$$

$$E_{leaf,cuti} = \frac{E_{leaf,cuti,sunlit} \cdot LAI_{sunlit} + E_{leaf,cuti,shade} \cdot LAI_{shade}}{LAI_{act}} \quad (12.8)$$

In turn, $E_{stom,sunlit}$, $E_{stom,shade}$, $E_{leaf,cuti,sunlit}$ and $E_{leaf,cuti,shade}$ values (all in $mmol H_2O \cdot s^{-1} \cdot m^{-2}$) are the result of multiplying the water vapor gradient by a compound conductance formed by either stomatal or cuticular conductances (which will likely be different for sunlit and shade leaves) and the leaf boundary layer conductance, i.e. (we omit the sunlit/shade superscripts):

$$E_{stom} = (g_{sw}^{-1} + g_{bound}^{-1})^{-1} \cdot \frac{VPD_{leaf}}{P_{atm}} \quad (12.9)$$

$$E_{leaf,cuti} = (g_{leaf,cuti}^{-1} + g_{bound}^{-1})^{-1} \cdot \frac{VPD_{leaf}}{P_{atm}} \quad (12.10)$$

where VPD_{leaf} (in MPa) is the vapor pressure deficit of the leaf (11.1.2), P_{atm} is the atmospheric pressure, g_{sw} is the stomatal conductance, $g_{leaf,cuti}$ is the cuticular conductance of the leaf and g_{bound} is the conductance of the leaf boundary layer (here all in $mmol H_2O \cdot s^{-1} \cdot m^{-2}$). The overall transpiration rate of the leaf, E_{leaf} has the following expression in the Fick's law:

$$E_{leaf} = g_w \cdot \frac{P_{atm}}{VPD_{leaf}} = ((g_{sw} + g_{cuti})^{-1} + g_{bound}^{-1})^{-1} \cdot \frac{P_{atm}}{VPD_{leaf}} \quad (12.11)$$

Note that in Ruffault et al. (2022) a crown boundary conductance was also considered, but within the design of **medfate** it is not needed, since VPD_{leaf} represents the gradient in vapor pressure between the leaf mesophyll and the canopy (or canopy layer) air.

Finally, $E_{stem,cuti}$ is calculated as follows:

$$E_{stem,cuti} = (g_{stem,cuti}^{-1} + g_{bound}^{-1})^{-1} \cdot \frac{VPD_{air}}{P_{atm}} \quad (12.12)$$

where VPD_{air} is the vapor pressure deficit in the canopy air, i.e.:

$$VPD_{air} = e_{sat}(T_{air}) - e_{air} \quad (12.13)$$

12.2.2 Cuticular conductances

Leaf cuticular conductances are not only species-specific, but can also change with temperature, according to changes in the permeability of lipid layers in the leaf epidermis. $g_{leaf,cuti}$ is a function of leaf temperature (T_{leaf}), which is based on a single or double Q10 equation depending on whether leaf temperature is above or below the transition phase temperature (T_{phase}). If $T_{leaf} \leq T_{phase}$ we have:

$$g_{leaf,cuti} = g_{leaf,cuti,20} \cdot Q_{10a}^{\frac{T_{leaf}-20}{10}} \quad (12.14)$$

whereas if $T_{leaf} > T_{phase}$ we have:

$$g_{leaf,cuti} = g_{leaf,cuti,20} \cdot Q_{10a}^{\frac{T_{leaf}-20}{10}} \cdot Q_{10b}^{\frac{T_{leaf}-T_{phase}}{10}} \quad (12.15)$$

Sunlit and shade leaves will have different $g_{leaf,cuti}$ because of their different temperature. Unlike $g_{leaf,cuti}$, $g_{stem,cuti}$ is assumed non temperature dependent.

12.2.3 Stomatal conductance

g_{sw} is the stomatal conductance taking into account the dependence of stomata on light and temperature, as well as leaf water status, i.e:

$$g_{sw} = \lambda(\Psi_{leaf,sym}) \cdot g_{sw,light,temp} \quad (12.16)$$

where $g_{sw,light,temp}$ is the stomatal conductance value without water stress, and λ is a regulation factor that varies between 0 and 1 to represent stomatal closure according to leaf symplasmic water potential ($\Psi_{leaf,sym}$) using a sigmoid-type function:

$$\lambda(\Psi_{leaf,sym}) = 1 - \left(1 + e^{(slope_{gs,bald}/25) \cdot (\Psi_{leaf,sym} - \Psi_{50_{gs,bald}})}\right)^{-1} \quad (12.17)$$

depending on the potential at 50 % of stomatal closure ($\Psi_{50_{gs}}$) and a shape parameter ($slope_{gs}$) describing the rate of decrease in stomatal conductance per unit of water potential drop. With the current implementation, λ will be the same for sunlit and shade leaves (i.e. there is only one symplasmic water potential for both kind of leaves) but $g_{sw,light,temp}$ will differ because of the different light and temperature conditions.

Stomatal regulation according to light and temperature can be performed following either a Jarvis (1976) or Baldocchi (1994), as dictated by the control parameter `stomatalSubmodel` (by default set to Baldocchi-type).

Jarvis-type stomatal regulation

In the Jarvis-type approach (Jarvis, 1976), stomatal conductance according to light and temperature ($g_{sw,light,temp}$) is defined as:

$$g_{sw,light,temp} = g_{sw,night,temp} + (g_{sw,max,temp} - g_{sw,night,temp}) \cdot (1 - e^{-Q_{PAR,leaf} \cdot c_{PAR}}) \quad (12.18)$$

where $g_{sw,max,temp}$ and $g_{sw,night,temp}$ are the maximum stomatal conductance and nocturnal stomatal conductance after accounting for temperature effects, $Q_{PAR,leaf}$ is the absorbed PAR photon flux per leaf area ((11.19)) and c_{PAR} is an empirical coefficient. Temperature effects on maximum stomatal conductance are modelled using:

$$g_{sw,max,temp} = g_{sw,max} \cdot \left[1 + \left(\frac{T_{leaf} - T_{gs,optim,jarv}}{T_{gs,sens,jarv}}\right)^2\right]^{-1} \quad (12.19)$$

where T_{leaf} is the current leaf temperature, $T_{gs,optim,jarv}$ is the optimal temperature and $T_{gs,sens,jarv}$ is an empirical coefficient. Temperature effects on nocturnal stomatal conductance are modelled analogously.

Baldocchi-type stomatal regulation

Baldocchi (1994) presented an analytical solution of the coupled leaf photosynthesis and stomatal conductance models. Baldocchi's approach uses Farquhar's model to determine photosynthesis, and the stomatal conductance equation is:

$$g_{sw,light,temp} = g_{sw,0} + m_{wue,bald} \cdot \frac{A_n}{C_s} \quad (12.20)$$

where $m_{wue,bald}$ is a dimensionless slope representing the water use efficiency, A_n is net photosynthesis as determined by Farquhar's model (which accounts for light and temperature effects), $g_{sw,0}$ is the zero intercept when $A_n = 0$, here taken as $g_{sw,0} = g_{sw,night}$ and C_s is the CO_2 concentration inside stomata, i.e.:

$$C_s = C_{air} - (A_n / g_{bc}) \quad (12.21)$$

where C_{air} is the inside the canopy CO_2 concentration and g_{bc} is the boundary layer for CO_2 :

$$g_{bc} = g_{bound} / 1.6 \quad (12.22)$$

Note that A_n is only calculated here to find the analytical solution to $g_{sw,light,temp}$. It does not represent actual photosynthesis, which has to be calculated taking into account leaf water status (i.e. $\lambda(\Psi_{leaf,sym})$). This is detailed in the next subsection.

12.2.4 Photosynthesis

Instantaneous gross and net photosynthesis rates for sunlit and shade leaves (i.e., $A_{g,shade}$, $A_{g,sunlit}$, $A_{n,shade}$, $A_{n,sunlit}$) are calculated using Farquhar's model (11.1.4) and assuming that g_c , the leaf diffusive conductance for CO_2 is influenced by leaf boundary (g_{bound}), stomatal (g_{sw}) and leaf cuticular ($g_{leaf,cuti}$) conductances, as explained in 11.1.3. Hence, non-zero gross assimilation can still occur with stomata completely closed.

12.3 Scaling of transpiration, soil water uptake and photosynthesis rates

The amount of transpiration of a plant cohort in a given time step t ($Tr_{cohorts,t}$, in $l \cdot m^{-2}$ of ground area, i.e. mm) is:

$$Tr_{cohorts,t} = E_{plant} \cdot LAI_{act} \cdot f_{dry} \cdot 10^{-3} \cdot 0.01802 \cdot \Delta t_{step} \quad (12.23)$$

where E_{plant} is the instantaneous cohort-level transpiration rate ($mmol H_2O \cdot s^{-1} \cdot m^{-2}$ of leaf area), 0.01802 is the molar weight (in $kg = l$) of water and $\Delta t_{step} = 86400/n_t$ the size of the time step in seconds, being n_t the number of time steps. f_{dry} is the fraction of the canopy that is assumed dry, which is necessary to avoid overestimating total evapo-transpiration. In non-rainy days $f_{dry} = 1$ whereas in rainy days it is derived from comparing the daily evaporation due to rainfall interception (In ; in mm) with Penman's potential evapo-transpiration (PET ; in mm):

$$f_{dry} = 1.0 - \frac{In}{PET} \quad (12.24)$$

Instantaneous water uptake rates from a given soil layer s (U_s ; in $mmol H_2O \cdot s^{-1} \cdot m^{-2}$ of leaf area) are scaled to the time step as done for transpiration:

$$Ex_{cohorts,s,t} = U_s \cdot LAI_{act} \cdot f_{dry} \cdot 10^{-3} \cdot 0.01802 \cdot \Delta t_{step} \quad (12.25)$$

Under the Sperry model, the sum of soil extraction equals transpiration, i.e.:

$$Tr_{cohorts,t} = Ex_{cohorts,t} = \sum_s Ex_{cohorts,s,t} \quad (12.26)$$

In contrast, under the SurEau model, soil extraction may not be equal to transpiration, leading to a change in plant water content (see (7.2)).

Instantaneous net photosynthesis per leaf area of sunlit and shade leaves in a given time step t (i.e. $A_{n,sunlit,t}$ and $A_{n,shade,t}$) are also aggregated into $A_{n,t}$, the net photosynthesis of the plant cohort for the time step t , in $g C \cdot m^{-2}$ of ground area:

$$A_{n,t} = (A_{n,sunlit,t} \cdot LAI_{sunlit} + A_{n,shade,t} \cdot LAI_{shade}) \cdot 10^{-6} \cdot 12.01017 \cdot \Delta t_{step} \quad (12.27)$$

Scaling of gross photosynthesis rates is done analogously.

12.4 Stomatal regulation and transpiration of hemi-parasitic plants

Stomatal regulation and transpiration of mistletoe is performed separately for sunlit and shade leaves. Leaf temperature (i.e. energy balance) is first estimated, using the absorbed short-wave radiation and net long-wave radiation per leaf area unit of the corresponding sunlit or shade leaves of the host cohort, but employing the mistletoe transpiration rate of the previous subdaily time step and its leaf width (see 11.1). Next, the corresponding vapor pressure deficit of sunlit and shade mistletoe leaves (i.e. $VPD_{leaf,mistletoe}$) are estimated as described in 11.1.2. Stomatal regulation follows an approach similar to that described for the SurEau

submodel (see 12.2.1). Stomatal conductance (and photosynthesis) is first estimated for current temperature and light conditions assuming no water limitation (i.e. $g_{sw,light,temp,mistletoe}$), using the approach of Baldocchi (1994):

$$g_{sw,light,temp,mistletoe} = g_{sw,0,mistletoe} + m_{wue,bald,mistletoe} \cdot \frac{A_{n,mistletoe}}{C_{s,mistletoe}} \quad (12.28)$$

where $m_{wue,bald,mistletoe}$ is a dimensionless slope representing the water use efficiency, $A_{n,mistletoe}$ is net mistletoe photosynthesis as determined by Farquhar's model (which accounts for light and temperature effects on mistletoe photosynthetic capacity parameters), $g_{sw,0,mistletoe}$ is the zero intercept when $A_{n,mistletoe} = 0$, here taken as $g_{sw,0,mistletoe} = g_{sw,night}$ and $C_{s,mistletoe}$ is the CO_2 concentration inside mistletoe stomata, i.e.:

$$C_{s,mistletoe} = C_{air} - (A_{n,mistletoe}/g_{bc}) \quad (12.29)$$

where C_{air} is the inside the canopy CO_2 concentration and g_{bc} is the boundary layer for CO_2 :

$$g_{bc} = g_{bound}/1.6 \quad (12.30)$$

The stomatal conductance of mistletoe taking into account the dependence of leaf water status (i.e. $g_{sw,mistletoe}$) is:

$$g_{sw,mistletoe} = \lambda_{mistletoe}(\Psi_{leaf,sym}) \cdot g_{sw,light,temp,mistletoe} \quad (12.31)$$

where $\lambda_{mistletoe}$ is a regulation factor that varies between 0 and 1 to represent stomatal closure according to the host leaf symplasmic water potential ($\Psi_{leaf,sym}$) using a sigmoid-type function:

$$\lambda_{mistletoe}(\Psi_{leaf,sym}) = 1 - \left(1 + e^{(slope_{gs,bald,mistletoe}/25) \cdot (\Psi_{leaf,sym} - \Psi_{50_{gs,bald,mistletoe}})}\right)^{-1} \quad (12.32)$$

with $\Psi_{50_{gs,bald,mistletoe}}$ and $slope_{gs,bald,mistletoe}$ being parameters of the stomatal vulnerability curve described as control parameters.

Mistletoe transpiration flow $E_{mistletoe}$ is estimated (for sunlit or shade leaves) using Fick's law:

$$E_{mistletoe} = g_{w,mistletoe} \cdot \frac{VPD_{leaf,mistletoe}}{P_{atm}} \quad (12.33)$$

where $VPD_{leaf,mistletoe}$ is the vapor pressure deficit of the mistletoe leaf and $g_{w,mistletoe}$ is the diffusive conductance incorporating both mistletoe stomatal conductance and boundary layer conductance.

Mistletoe transpiration effects are not modelled directly on the plant host, but rather as additional soil water uptake from the soil layers (or pools) where the host is connected, which feedbacks on the water status of the host at later time steps. Under the Sperry transpiration submodel, mistletoe transpiration is added to the average instantaneous flow rate $\hat{E}$ and the hydraulic supply function of the host is used to find the transpiration rate E numerically closest to $\hat{E}$ (remember that the supply function is build in discrete steps). Finding the E value numerically closest to $\hat{E}$ allows determining the total (i.e. host + mistletoe) instantaneous soil water uptake rates (U_s) from each soil layer s . Under the Sureau transpiration submodel, mistletoe transpiration is calculated at small time steps and it is added to soil water uptake following the same soil layer uptake proportions determined for the direct uptake of the host plant cohort.

Chapter 13

Closing energy balances

In chapter 9 we described the details of radiation energy transfer. Transpiration is an important latent heat component of the energy balance, and the details of how plant transpiration is determined were given in chapters 10, 11 and 12. This chapter explains how to calculate the remaining components of the canopy energy balance, while distinguishing between assuming a single-layer canopy or considering a multiple-layer canopy.

13.1 Single-layer canopy

The first part of the section describes the calculation of latent heat and convective energy exchanges between the soil, canopy and atmosphere layers. These are the remaining energy components that are needed to close the energy balances of the canopy (eq. (7.3)) and soil layers (eqs. (7.5) and (7.6)). Temperature changes applied to the canopy and soil layers are also described.

13.1.1 Latent heat

Canopy exchanges of latent heat include latent heat exchanged from plant transpiration and evaporation of rain intercepted by the canopy. In the single-layer canopy energy balance, vaporisation latent heat exchange for the whole canopy ($LE_{vap,can}$; in $W \cdot m^{-2}$) is calculated as:

$$LE_{vap,can} = \lambda_v(T_{can}) \cdot \left(\frac{In_t + \sum_{i=1}^c Tr_{i,t}}{\Delta t_{step}} \right) \quad (13.1)$$

where $\lambda_v(T_{can})$ is the latent heat of vaporization at temperature T_{can} (in $J \cdot kg^{-1}$), $Tr_{i,t}$ is the amount (in $mm = kg \cdot m^{-2}$) of transpiration for cohort i during time step t (12.23), and In_t is the amount of intercepted water evaporated from the canopy for time step t (total daily interception In is divided into timesteps according to the proportion of shortwave radiation absorbed by the canopy at each time step).

Fusion latent heat due to snow melting ($LE_{fus,soil}$; in $W \cdot m^{-2}$) is estimated using:

$$LE_{fus,snow} = \frac{\lambda_f \cdot Sm_t}{\Delta t_{step}} \quad (13.2)$$

where $\lambda_f = 0.33355 J \cdot kg^{-1}$ is the latent heat of fusion. The latent heat exchange between the soil and the atmosphere comes as vaporisation latent heat from water evaporated from the soil surface ($LE_{vap,soil}$; in $W \cdot m^{-2}$). It is estimated from water flow in $mm = kg \cdot m^{-2}$ using:

$$LE_{vap,soil} = \frac{\lambda_v(T_{soil}) \cdot Es_t}{\Delta t_{step}} \quad (13.3)$$

where $\lambda_v(T_{soil})$ is the latent heat of vaporization at temperature T_{soil} .

As before, daily evaporation from bare soil (Es) and snow melt water equivalent (Sm) are divided into time steps according to the proportion of shortwave radiation absorbed by the soil layer each step.

13.1.2 Convective energy

Convective energy fluxes between atmosphere and the canopy ($H_{can,atm}$) and between the canopy and the soil ($H_{can,soil}$) are determined as follows:

$$H_{can,atm} = \frac{\rho_{atm} \cdot c_p}{r_{can,atm}} \cdot (T_{can} - T_{atm}) \quad (13.4)$$

$$H_{can,soil} = \frac{\rho_{can} \cdot c_p}{r_{can,soil}} \cdot (T_{can} - T_{soil,1}) \quad (13.5)$$

where ρ_{atm} and ρ_{can} are the air density above-canopy and inside-canopy, respectively, calculated from the corresponding temperatures (see utility functions in meteoland reference manual); $c_p = 1013.86 \text{ J} \cdot \text{kg}^{-1} \cdot \text{C}^{-1}$ is the specific heat capacity of the air. $r_{can,atm}$ and $r_{can,soil}$ are the atmosphere-canopy and canopy-soil aerodynamic resistances (in $\text{s} \cdot \text{m}^{-1}$). These, in turn, are calculated using the FAO56 calculation procedure and canopy height, total LAI and above-canopy and below-canopy wind speeds. Wind speed below the canopy is calculated as explained in 25.2 assuming a height of 2 m.

13.1.3 Canopy temperature changes

After evaluating the canopy energy balance equation (7.3) one has to translate energy balance into temperature variation of the canopy. Rearranging the same equation and expressing it in discrete time steps we have:

$$\Delta T_{can} = \Delta t_{step} \cdot \frac{K_{abs,can} + L_{net,can} - LE_{can} - H_{can,atm} - H_{can,soil}}{TC_{can}} \quad (13.6)$$

where TC_{can} is the canopy thermal capacitance (in $\text{J} \cdot \text{m}^{-2} \cdot \text{K}^{-1}$) and $\Delta t_{step} = 86400/n_t$ is the size of the time step in seconds. Canopy thermal capacitance depends on the leaf area index of the stand:

$$TC_{can} = TC_{LAI} \cdot \frac{0.8 \cdot LAI_{live} + 1.2 \cdot (LAI_{act} + LAI_{dead})}{2} \quad (13.7)$$

where TC_{LAI} is the thermal capacitance per LAI unit, which is specified by the control parameter `thermalCapacityLAI`. By using both the maximum leaf area of the stand, LAI_{live} , and its current live/dead leaf area ($LAI_{act} + LAI_{dead}$) it is assumed that part of thermal capacitance corresponds to stems and branches, so that capacitance does not drop to zero for deciduous canopies.

13.1.4 Soil temperature changes

Analogously to the canopy, the change in temperature for a given soil layer s in the current time step is given by rearranging eq. (7.6):

$$\Delta T_{soil,s} = \Delta t_{step} \cdot \frac{G_{s-1,s} - G_{s,s+1}}{C_{soil,s}} \quad (13.8)$$

where the energy balance on each soil layer s depends on the balance between energy coming from above ($G_{s-1,s}$) and energy going to below ($G_{s,s+1}$) and $C_{soil,s}$, the thermal capacitance of soil layer s . Energy inflow to the first (uppermost) layer (i.e. $G_{0,1}$) is the result of energy exchanges between the soil layer and the canopy and atmosphere, i.e. from eq. (7.5):

$$G_{0,1} = K_{soil} + L_{net,soil} + H_{can,soil} - LE_{soil} \quad (13.9)$$

Heat conduction between layers s and $s+1$ (i.e. $G_{s,s+1}$) depend on the soil temperature gradient (see function `soil_temperatureGradient()`):

$$G_{s,s+1} = \lambda_{soil,s} \cdot \frac{\delta T_{soil,s}}{\delta z} = \lambda_{soil,s} \cdot \frac{T_{soil,s} - T_{soil,s+1}}{(Z_{s-1} - Z_{s+1})/2} \quad (13.10)$$

where Z_{s-1} and Z_{s+1} are expressed in m and $\lambda_{soil,s}$ is the *thermal conductivity* of layer s , calculated from soil moisture and texture following Dharssi et al. (2009) (see function `soil_thermalconductivity()`). The gradient in the lowermost layer is calculated assuming a temperature of the earth (at 10 m) of 15.5 Celsius.

Finally, $C_{soil,s}$ the thermal capacitance of soil layer s is calculated as:

$$C_{soil,s} = VTC_{soil,s} \cdot d_s \quad (13.11)$$

where d_s is the soil width of layer s (expressed in m) and VTC_s is the *volumetric thermal capacity* of soil layer s (in $J \cdot m^{-3} \cdot K^{-1}$), calculated from soil moisture and texture following a simplification of Cox et al. (1999) (see function `soil_thermalcapacity()`).

13.2 Multi-layer canopy

A multi-layer canopy energy balance requires the calculation of sensible heat and latent heat exchanges between the air in each canopy layer and the leaves it contains. In addition, one has to consider heat exchanges between each layer and the ones below/above, derived from turbulent flow.

13.2.1 Latent heat of vaporization

The latent heat of vaporization for a given canopy layer j ($LE_{vap,j}$; in $W \cdot m^{-2}$) includes transpiration and evaporation of intercepted rain, and is estimated analogously to the single-canopy case (eq. (13.1)):

$$LE_{vap,j} = \lambda_v(T_{air,j}) \cdot \left(\frac{In_{j,t}}{\Delta t_{step}} + E_{j,t} \right) \quad (13.12)$$

where $\lambda_v(T_{air,j})$ is the latent heat of vaporization at temperature $T_{air,j}$ (in $J \cdot kg^{-1}$), $In_{j,t}$ is the intercepted water that is evaporated from the canopy layer, determined by dividing In_t among canopy layers according to the fraction of total LAI in each one:

$$In_{j,t} = In_t \cdot \frac{\sum_{i=1}^c LAI_{act,i,j}}{LAI_{act}} \quad (13.13)$$

and $E_{j,t}$ is the instantaneous transpiration rate in layer j (in $kg H_2O \cdot m^{-2} \cdot s^{-1}$), given by:

$$E_{j,t} = 10^{-3} \cdot 0.01802 \cdot \sum_{i=1}^c E_{sunlit,i,t} \cdot LAI_{sunlit,i,j} + E_{shade,i,t} \cdot LAI_{shade,i,j} \quad (13.14)$$

where $E_{sunlit,i,t}$ and $E_{shade,i,t}$ are the instantaneous transpiration rates (in $mmol H_2O \cdot m^{-2} \cdot s^{-1}$) of sunlit and shade leaves, respectively, of cohort i during time step t ; and $LAI_{sunlit,i,j}$ and $LAI_{shade,i,j}$ are the leaf area index of sunlit and shade leaves, respectively, of cohort i in layer j (9.8).

Latent heat derived from soil water evaporation and snow melting ($LE_{vap,soil}$ and $LE_{fus,snow}$; in $W \cdot m^{-2}$) are estimated as in the single-canopy case, using eqs. (13.3) and (13.2).

13.2.2 Sensible heat between leaves and the canopy air space

The sensible heat exchanged between the air of canopy layer j and the leaves within it ($H_{leaf,j}$; $W \cdot m^{-2}$) is obtained using:

$$H_{leaf,j} = \sum_{i=1}^c LAI_{sunlit,i,j} \cdot H_{sunlit,i,j} + LAI_{shade,i,j} \cdot H_{shade,i,j} \quad (13.15)$$

where $LAI_{sunlit,i,j}$ and $LAI_{shade,i,j}$ are the leaf area index of sunlit and shade leaves of plant cohort i in layer j , respectively (see eq. (9.8)). $H_{sunlit,i,j}$ and $H_{shade,i,j}$ are the sensible heat exchange between layer j and sunlit or shade leaves of cohort i (Ma and Liu, 2019):

$$H_{sunlit,i,j} = -2 \cdot C_p \cdot (T_{leaf,sunlit,i} - T_{air,j}) \cdot g_{Ha,i,j} \quad (13.16)$$

$$H_{shade,i,j} = -2 \cdot C_p \cdot (T_{leaf,shade,i} - T_{air,j}) \cdot g_{Ha,i,j} \quad (13.17)$$

where $T_{leaf,sunlit,i}$ and $T_{leaf,shade,i}$ are, respectively, the sunlit and shade leaf temperatures for cohort i (determined using eq. (11.1)); $T_{air,j}$ is the current air temperature in canopy layer j , $C_p = 29.37152 J \cdot mol^{-1} \cdot C^{-1}$ is the specific heat capacity of dry air at constant pressure; and $g_{Ha,i,j}$ is the leaf boundary layer conductance for heat ($mol \cdot m^{-2} \cdot s^{-1}$) for leaves of cohort i and wind speed in layer j (see eq. (11.3)).

13.2.3 Turbulent heat exchange

Turbulent heat exchanges occur between the soil and the bottom canopy layer (i.e. $j = 1$):

$$H_{1,soil} = \frac{\rho_1 \cdot c_p}{r_{1,soil}} \cdot (T_{air,1} - T_{soil,1}) \quad (13.18)$$

where $c_p = 1013.86 J \cdot kg^{-1} \cdot C^{-1}$ is the specific heat capacity of the air, ρ_1 is the air density at $j = 1$, $T_{air,1}$ is the air temperature of layer 1, $T_{soil,1}$ is the temperature of the top soil layer and $r_{1,soil}$ is the aerodynamic resistance between the soil and canopy layer 1 (in $s \cdot m^{-1}$). The bottom layer also have turbulent heat exchanges with the layer above, determined by the canopy turbulence model. The turbulent heat flux gradient for layer $j = 1$ is:

$$(\delta F_H / \delta z)_1 = \frac{-c_p \cdot \rho_1 \cdot (T_{air,2} - T_{air,1}) \cdot \bar{u} \bar{w}_1}{\Delta z \cdot (\delta u / \delta z)_1} - H_{1,soil} \quad (13.19)$$

where Δz is the size of vertical layers (in m); and $(\delta u / \delta z)_1$ and $\bar{u} \bar{w}_1$ are the wind speed gradient and Reynolds stress at $j = 1$.

For all intermediate layers (i.e. $1 < j < l$) we have turbulent exchanges with the neighboring layers $j - 1$ and $j + 1$, which results in the following turbulent heat flux gradient:

$$(\delta F_H / \delta z)_j = \frac{-c_p \cdot \rho_j \cdot (T_{air,j+1} - T_{air,j-1}) \cdot \bar{u} \bar{w}_j}{2 \cdot \Delta z \cdot (\delta u / \delta z)_j} \quad (13.20)$$

And, finally, the topmost layer (i.e. $j = l$) has turbulent heat exchanges with the layer below $l - 1$ and the atmosphere, so that:

$$(\delta F_H / \delta z)_l = \frac{-c_p \cdot \rho_l \cdot (T_{atm} - T_{air,l-1}) \cdot \bar{u} \bar{w}_l}{2 \cdot \Delta z \cdot (\delta u / \delta z)_l} \quad (13.21)$$

13.2.4 Canopy and soil temperature changes

Evaluation of the canopy layer energy balance equation (7.4) is done taking into account that sensible heat exchange for a given layer j includes heat exchange with leaves (i.e. $H_{leaf,j}$) and the turbulent heat flux gradient (i.e. $(\delta F_H / \delta z)_j$), so that the variation of air temperature in layer j for a given time interval $\Delta t_{substep}$ is:

$$\Delta T_{air,j} = \Delta t_{substep} \cdot \frac{K_{abs,j} + L_{net,j} - LE_j + H_{leaf,j} + (\delta F_H / \delta z)_j}{TC_j} \quad (13.22)$$

where TC_j is the thermal capacitance of canopy layer j (in $J \cdot m^{-2} \cdot K^{-1}$), found analogously to TC_{can} (eq. (13.7)):

$$TC_j = TC_{LAI} \cdot \frac{0.8 \cdot LAI_{live,j} + 1.2 \cdot (LAI_{act,j} + LAI_{dead,j})}{2} \quad (13.23)$$

$\Delta t_{substep}$ is the time step for closing the energy balance, which has to be smaller than Δt_{step} to avoid numerical instabilities due to turbulent flow.

When simulating a multiple-layer energy balance, we have to replace $H_{can,soil}$ by $H_{1,soil}$ in the energy inflow of soil layer 1: (13.9):

$$G_{0,1} = K_{soil} + L_{net,soil} + H_{1,soil} - LE_{vap,soil} \quad (13.24)$$

Soil temperature changes are estimated in the same way as for single-canopy energy balance, except that a smaller temporal step is used:

$$\Delta T_{soil,s} = \Delta t_{substep} \cdot \frac{G_{s-1,s} - G_{s,s+1}}{C_{soil,s}} \quad (13.25)$$

13.2.5 Within-canopy changes in water vapor and CO_2

Multi-layer canopy energy balance also entails the possibility of considering gradients of scalars such as water vapor or CO_2 . Analogously to the turbulent heat exchange, wind turbulence leads to changes in concentration of water vapor or CO_2 within canopy layers. In addition, plant physiology also alters the concentration of these gases through transpiration in photosynthesis. To model temporal changes in these scalars for canopy layers, we first have to estimate CO_2 concentration in $mg \cdot m^{-3}$ and H_2O concentration in $kg \cdot m^{-3}$ from $C_{air,j}$ (ppm) and $e_{air,j}$ (kPa) using:

$$[CO_2]_j = 0.409 \cdot 44.01 \cdot C_{air,j} \quad (13.26)$$

$$[H_2O]_j = 0.622 \cdot \frac{e_{air,j} \cdot \rho_j}{P_{atm}} \quad (13.27)$$

The one-dimensional scalar flux balance for an homogeneous turbulent flow can be described by the following conservation equations (Lai et al., 2000):

$$(\delta[CO_2]/\delta t)_j = (\delta F_{[CO_2]}/\delta z)_j + S_{[CO_2],j} \quad (13.28)$$

$$(\delta[H_2O]/\delta t)_j = (\delta F_{[H_2O]}/\delta z)_j + S_{[H_2O],j} \quad (13.29)$$

where $(\delta F_{[CO_2]}/\delta z)_j$ and $(\delta F_{[H_2O]}/\delta z)_j$ are the gradients of vertical flux of CO_2 and H_2O in layer j , respectively; and $S_{[H_2O],j}$ and $S_{[CO_2],j}$ are the source strengths for $[H_2O]$ (in $kg H_2O \cdot m^{-3} \cdot s^{-1}$) and $[CO_2]$ (in $mg CO_2 \cdot m^{-3} \cdot s^{-1}$) in layer j , respectively, which derive from transpiration (source of water) and photosynthesis (sink of CO_2):

$$S_{[H_2O],j} = \frac{(In_{j,t}/\Delta t) + E_{j,t}}{\Delta z} \quad (13.30)$$

$$S_{[CO_2],j} = -\frac{A_{n,j,t}}{\Delta z} \quad (13.31)$$

where $E_{j,t}$ and $A_{n,j,t}$ are the instantaneous transpiration and photosynthesis rates in layer j . The former was already defined in eq. (13.14), whereas the latter ($A_{n,j,t}$; in $mg CO_2 \cdot m^{-2} \cdot s^{-1}$) is defined analogously:

$$A_{n,j,t} = 10^{-3} \cdot 44.01 \cdot \sum_{i=1}^c A_{n,sunlit,i,t} \cdot LAI_{sunlit,i,j} + A_{n,shade,i,t} \cdot LAI_{shade,i,j} \quad (13.32)$$

where $A_{n,sunlit,i,t}$ and $A_{n,shade,i,t}$ are the instantaneous net photosynthesis rates (in $\mu mol CO_2 \cdot m^{-2} \cdot s^{-1}$) of sunlit and shade leaves, respectively, of cohort i during time step t ; and $LAI_{sunlit,i,j}$ and $LAI_{shade,i,j}$ are the leaf area index of sunlit and shade leaves, respectively of cohort i in layer j (9.8). For the special case of the lowest canopy layer, water vapor needs to include the evaporation from soil, i.e.:

$$S_{[H_2O],1} = \frac{((In_{1,t} + Es_t)/\Delta t) + E_{1,t}}{\Delta z} \quad (13.33)$$

For any layer $1 < j < l$ the gradients of vertical scalar fluxes are found using:

$$(\delta F_{[H_2O]}/\delta z)_j = -\frac{c_p \cdot \rho_j \cdot ([H_2O]_{j+1} - [H_2O]_{j-1}) \cdot \bar{u}\bar{w}_j}{2 \cdot \Delta z \cdot (\delta u/\delta z)_j} \quad (13.34)$$

$$(\delta F_{[CO_2]}/\delta z)_j = -\frac{c_p \cdot \rho_j \cdot ([CO_2]_{j+1} - [CO_2]_{j-1}) \cdot \bar{u}\bar{w}_j}{2 \cdot \Delta z \cdot (\delta u/\delta z)_j} \quad (13.35)$$

whereas for the bottom layer ($j = 1$) we have:

$$(\delta F_{[H_2O]}/\delta z)_1 = -\frac{c_p \cdot \rho_1 \cdot ([H_2O]_2 - [H_2O]_1) \cdot \bar{u}\bar{w}_1}{\Delta z \cdot (\delta u/\delta z)_1} \quad (13.36)$$

$$(\delta F_{[CO_2]}/\delta z)_1 = -\frac{c_p \cdot \rho_1 \cdot ([CO_2]_2 - [CO_2]_1) \cdot \bar{u}\bar{w}_1}{\Delta z \cdot (\delta u/\delta z)_1} \quad (13.37)$$

and for the top layer ($j = l$) we have:

$$(\delta F_{[H_2O]}/\delta z)_l = - \frac{c_p \cdot \rho_l \cdot ([H_2O]_{atm} - [H_2O]_{l-1}) \cdot \bar{u}\bar{w}_l}{2 \cdot \Delta z \cdot (\delta u/\delta z)_l} \quad (13.38)$$

$$(\delta F_{[CO_2]}/\delta z)_l = - \frac{c_p \cdot \rho_j \cdot ([CO_2]_{atm} - [CO_2]_{l-1}) \cdot \bar{u}\bar{w}_l}{2 \cdot \Delta z \cdot (\delta u/\delta z)_l} \quad (13.39)$$

Equation (13.29) is solved via discretization both in terms of space (vertical layers) and time, the latter using $\Delta t_{substep}$ as for the energy balance.

Chapter 14

Day-level results

In the preceeding chapters we detailed how the model performs canopy and soil energy balances for subdaily time steps and how transpiration and photosynthesis values are determined for any given substep. This chapter indicates how these are aggregated at the daily scale and how other day-level model outputs are calculated.

14.1 Photosynthesis and transpiration

Cohort's transpiration Tr_t (eq. (12.23)) are added across subdaily steps to yield daily transpiration (Tr , in $mm\ H_2O$):

$$Tr = \sum_{t=1}^{n_t} Tr_t \quad (14.1)$$

An the same for water extraction $Ex_{s,t}$ for each soil layer s (eq.(12.25)):

$$Ex_s = \sum_{t=1}^{n_t} Ex_{s,t} \quad (14.2)$$

Ex_s are subtracted from the water content of the corresponding soil layer, closing the soil water balance of the day (eq. (7.1)).

Daily values of gross and net carbon assimilation for plant cohorts are obtained similarly. $A_{g,t}$ and $A_{n,t}$ (eq. (12.27)) are added across subdaily steps to obtain the daily gross and net assimilation values at the cohort level (in $gC \cdot m^{-2}$).

14.2 Plant water potentials and relative water contents

Because the model determines optimum transpiration for every subdaily time step, this leads to a daily sequence of leaf water potential ($\Psi_{leaf,t}$), stem water potential ($\Psi_{stem,t}$), root crown water potential ($\Psi_{rootcrown,t}$) and root surface water potential ($\Psi_{rootcrown,s,t}$) values (for soil layer s in the last case). The model defines the following daily water potentials:

- *Pre-dawn leaf water potential* (Ψ_{pd}): the maximum of $\Psi_{leaf,t}$ values.
- *Pre-dawn shade leaf water potential* ($\Psi_{pd,shade}$): the maximum of $\Psi_{leaf,shade,t}$ values.
- *Pre-dawn sunlit leaf water potential* ($\Psi_{pd,sunlit}$): the maximum of $\Psi_{leaf,sunlit,t}$ values.
- *Mid-day leaf water potential* (Ψ_{md}): the minimum of $\Psi_{leaf,t}$ values.
- *Mid-day shade leaf water potential* ($\Psi_{md,shade}$): the minimum of $\Psi_{leaf,shade,t}$ values.
- *Mid-day sunlit leaf water potential* ($\Psi_{md,sunlit}$): the minimum of $\Psi_{leaf,sunlit,t}$ values.
- *Stem water potential* (Ψ_{stem}): the minimum of $\Psi_{stem,t}$ values.

- *Root-crown water potential* ($\Psi_{rootcrown}$): the minimum of $\Psi_{rootcrown,t}$ values.
- *Root surface water potentials* ($\Psi_{rootsurf,s}$): the minimum of $\Psi_{rootsurf,s,t}$ values for each soil layer s .

Analogously, relative water content of stems and leaves is known for every subdaily time step, which results in a daily sequence of leaf relative water content ($RWC_{leaf,t}$) and stem relative water content ($RWC_{stem,t}$). These are summarized at the daily level:

- *Leaf relative water content* (RWC_{leaf}): the mean $RWC_{leaf,t}$ values.
- *Stem relative water content* (RWC_{stem}): the mean of $RWC_{stem,t}$ values.

Finally, the daily sequence of slopes of the supply function ($dE/d\Psi_t$) is also averaged at the daily level:

- *Slope of the supply function* ($dE/d\Psi$): the mean $dE/d\Psi_t$ values.

14.3 Plant drought stress

In order to have an estimate of daily drought stress for the plant cohort, the model uses the complement of relative whole-plant hydraulic conductance. In turn, the relative conductance is determined by dividing the current whole-plant hydraulic conductance by its maximum value:

$$DDS = \phi \cdot \left(1.0 - \frac{k_{plant}}{k_{maxplant}} \right) \quad (14.3)$$

where ϕ is the leaf phenological status, k_{plant} is the current whole-plant hydraulic conductance and $k_{maxplant}$ is the maximum whole-plant conductance. Leaf-phenological status is included (only in winter-deciduous or winter semi-deciduous plants) to avoid drought stress during leafless periods. Since k_{plant} varies through the day, average daily values are used for the determination of drought stress. Furthermore, remember that in the Sperry transpiration model we have that the whole-plant conductance is the slope of the vulnerability curve:

$$k_{plant} = dE/d\Psi \quad (14.4)$$

In the Sureau sub-model, k_{plant} is estimated from the resistances across the hydraulic network (see 10.5.2).

14.4 Cavitation and conduit refilling

Like with the basic water balance model, the advanced water balance model is normally run assuming that although soil drought may reduce transpiration. Embolized xylem conduits may be automatically refilled when soil moisture recovers if `stemCavitationRecovery` = "total". Automatic refilling is always assumed for root segments, but not for leaves and stems. There are three other options to deal with leaf and stem xylem cavitation besides automatic refilling. Any of them cause the model to keep track of the proportion of lost conductance in the leaf and stem xylem of the plant cohort (PLC_{leaf} and PLC_{stem}) at successive time steps:

$$PLC_{leaf,t} = \max\{PLC_{leaf,t-1}, 1 - k_{leaf}(\Psi_{leaf})/k_{max,leaf}\} \quad (14.5)$$

$$PLC_{stem,t} = \max\{PLC_{stem,t-1}, 1 - k_{stem}(\Psi_{stem})/k_{max,stem}\} \quad (14.6)$$

In the Sperry sub-model, PLC values are evaluated every subdaily (hourly) time step, whereas in the Sureau sub-model it PLC values are updated more frequently.

For simulations of less than one year one can use `stemCavitationRecovery` = "none" to keep track of the maximum drought stress. However, for simulations of several years, it is normally advisable to allow recovery. If `stemCavitationRecovery` = "annual", PLC_{leaf} and PLC_{stem} values are set to zero at the beginning of each year, assuming that embolized plants overcome the conductance loss by creating new xylem tissue. Finally, if `stemCavitationRecovery` = "rate" the model simulates stem refilling at daily steps as a function

of symplasmic water potential. First, a daily recovery rate (r_{refill} ; in $cm^2 \cdot m^{-2} \cdot day^{-1}$) is estimated as a function of $\Psi_{symp,stem}$:

$$r_{refill}(\Psi_{symp,stem}) = r_{max,refill} \cdot \max\{0.0, (\Psi_{symp,stem} + 1.5)/1.5\} \quad (14.7)$$

Where $r_{max,refill}$ is the control parameter `cavitationRecoveryMaximumRate` indicating a maximum refill rate. The right part of the equation normalizes the water potential, so that $r_{refill} = r_{refill,max}$ if $\Psi_{symp,stem} = 0$ and $r_{refill} = 0$ if $\Psi_{symp,stem} \leq -1.5MPa$. The proportion of conductance lost in leaves and stem are then updated using:

$$PLC_{leaf} = \max\{0.0, PLC_{leaf} - (r_{refill}/H_v)\} \quad (14.8)$$

$$PLC_{stem} = \max\{0.0, PLC_{stem} - (r_{refill}/H_v)\} \quad (14.9)$$

where H_v is the Huber value in units of $cm^2 \cdot m^{-2}$.

14.5 Cavitation-induced defoliation and recovery

By default, the water-balance model assumes that high levels of leaf hydraulic impairment will lead to leaf hydraulic disconnection and defoliation (although this process can be deactivated if `cavitationInducedDefoliation = FALSE`). When cavitation-induced defoliation is considered, the model assumes that crowns are made of a population of leaves whose hydraulic vulnerability follows a Gaussian distribution centered on the water potential corresponding to 50% loss of conductance (i.e. $\Psi_{50,leaf}$, or the corresponding value dictated by c_{leaf} and d_{leaf}) and with a known coefficient of variation set to 10 % (by default), a modelling strategy developed by Hervé Cochard. The $slope_{leaf}$ parameter of the vulnerability curve, or the corresponding c_{leaf} exponent in the case of a Weibull function, is considered constant. Leaves are hydraulically disconnected and die when their embolism rate exceeds a critical value set to 88 % of percent loss of conductance, by default (see defoliation control parameters as well as functions `hydraulics_proportionDefoliationWeibull()` and `hydraulics_proportionDefoliationSigmoid()`). Ultimately, the assumed variability of hydraulic vulnerability and the current leaf water potential lead to a proportion of cavitation-induced crown defoliation.

In the water-balance model, the proportion of crown defoliation leads to a corresponding decrease LAI_{act} , and recovery of foliage after drought will follow the conduit refilling process associated to decrease in PLC_{leaf} . In the growth model, the proportion of crown defoliation is used to determine leaf senescence and the recovery of foliage is simulated via explicit leaf growth (see chapter 17).

Part IV

Forest growth modelling

Chapter 15

Forest growth model

This chapter provides an overview of a forest growth model, which builds on the previous two models and allows simulating carbon and water balance as well as plant growth and mortality on a daily step basis for a given forest stand and for the period corresponding to input weather data. The initial version of the model was described in De Cáceres et al. (2023), but several modifications have occurred since then. The model is run using function `growth()`, for a set of days, or function `growth_day()`, for a single day.

The forest growth model can be run in two different levels of complexity, depending on the water balance submodel employed (basic and advanced, corresponding to chapters 3 and 7). In this and the following chapters, we indicate those design and formulation elements where the forest growth model building on the basic water balance differs from that building on the advanced one.

The information provided in the current chapter should be enough to understand what the growth model does, but the reading previous chapters and the following ones provide a more detailed understanding on how the different processes are designed and formulated.

15.1 Design principles

15.1.1 Overview

Like in the preceding models, the physical structure of the stand is represented in one (vertical) dimension (i.e. the horizontal coordinates of plants are not explicit). The model uses *cohort-based* representation of living plants, meaning that similar plant individuals are represented using a single entity with average characteristics (e.g. tree height and diameter at breast height) and a density variable is used to scale from (an average) **individual level** to the **cohort level**. Additionally, the model keeps track of carbon pools for **dead standing plants** (i.e. snags), **litter** (aboveground or belowground) and **soil organic matter**, represented at cohort-level (for snags), species-level (for litter) or aggregated level (for soil organic matter), respectively.

Processes in the forest stand are modeled either at the cohort level (water balance, photosynthesis, or plant mortality) or at the individual level (carbon balance, tissue growth and senescence). Energy balance, water balance and plant photosynthesis processes follow the design of the previous models and will not be repeated here (see chapters 3 and 7).

At the *individual level*, the growth model tries to reproduce the phenological, physiological and physical processes that modulate live and dead leaf area changes, sapwood area changes and fine root area changes of an *average individual* of the plant cohort. Formation of new sapwood leads to diameter and height increases, so that physiological processes involve **plant structural changes**. Since the model does not implement all processes and drivers (such as nutrient availability) potentially affecting growth, user input parameters are

used to empirically constrain growth rates and plant size, i.e. maximum relative sapwood area growth rates, maximum plant height and diameter-height ratios.

At the *plant cohort level*, the forest growth model implements **plant mortality** for woody cohorts, which reduces cohort density, arising from different processes. A baseline mortality rate is incremented either because of exhaustion of carbon reserves (i.e. plant starvation) or dehydration of plant tissues (i.e. plant desiccation). Additionally, fire behavior and effects (i.e. fire severity) can be simulated. Fires can partially or totally burn crown leaves and buds. Fire heat can also induce necrosis in the cambium, leading to full above-ground mortality.

Finally, the model also simulates the **decomposition** of organic matter produced by senescence and mortality processes (root exudation, litter fall and plant mortality). Plant litter (i.e. fine roots, leaves, branches, etc.) is allocated into different structural carbon pools depending on the species that produced it. Decomposition of organic matter in those structural pools leads to fluxes towards surface and soil carbon matter pools for further decomposition. Decomposition processes involve heterotrophic respiration, which releases CO_2 back to the atmosphere.

15.1.2 Plant carbon pools and carbon flows

Different *plant carbon compartments (organs)* are considered separately: **leaves**, **twigs** (branchlets), **sapwood** (including stem, branches and coarse roots) and **fine roots**. Twigs are kept separated from the rest of woody organs because their biomass dynamics is assumed to be coupled with leaf biomass. Each compartment has one or more *carbon pools* where carbon is in different form. Carbon pools are differentiated between **structural** (live organs), **dead** (dead organs) and **labile**. In turn, labile carbon involves **metabolic** and **storage** (Richardson et al., 2013; Dietze et al., 2014), with a conversion rate regulated by the need to maintain, as much as possible, an equilibrium in the amount of metabolic carbon to sustain cell metabolism and for osmotic purposes. Similar designs of two non-structural carbon compartments have been implemented in other forest ecosystem models (Cho et al., 2022).

Details of the carbon pools in different compartments are as follows (Fig. 15.1):

- **Structural carbon** corresponds to the carbon forming cell membranes, cell walls and the cytosolic machinery of *living plant organs*. The amount of structural carbon (together with metabolic carbon) in a given compartment is used to determine *maintenance respiration* costs, and the structural carbon per tissue unit determines *growth respiration* costs. Structural carbon occurs in *leaves*, *twigs*, *sapwood* and *fine roots* compartments, which have different maintenance respiration costs per dry biomass. Additionally, only live (parenchymatic) sapwood/twig cells, accounting for between 5 and 40% of volume (Plavcová and Jansen, 2015), contribute to estimate sapwood/twig maintenance respiration costs. The fraction of embolized conduits is also discounted from sapwood/twig respiratory biomass. Structural carbon is transformed into dead carbon or exported to litter via tissue senescence. Meristematic tissues (buds) are not represented as structural carbon (although a state variable is used to determine the availability of living meristems).
- **Dead carbon** corresponds to dead organs that are still standing as part of living plants (as opposed to being litter or snags). This includes the mass of *dead leaves* and *dead twigs* corresponding to dead leaf area that has not yet fallen to become litter, as well as the mass of *heartwood*, in the case of tree cohorts.
- **Metabolic carbon** is a labile carbon form assumed to correspond to **sugars** (e.g., glucose or fructose) used to sustain cell functioning (i.e. provide carbon for maintenance respiration). Metabolic carbon occurs in leaves and sapwood compartments only. *Leaf metabolic carbon* is generated from photosynthesis or leaf starch degradation and used in different functions, including leaf maintenance respiration and export to sapwood via phloem transport. Sugar concentration in the phloem is assumed to be (passively) in equilibrium with leaf metabolic sugar concentration. *Sapwood metabolic carbon* comes from phloem transport, or sapwood starch degradation, and is used to directly sustain maintenance respiration of sapwood and fine roots.
- **Storage carbon** is a labile carbon form assumed to loosely correspond to **starch**. Like metabolic carbon, storage carbon occurs in leaves and sapwood compartments only. *Leaf storage carbon* is

intended to represent short-term storage (mainly starch within chloroplasts) and its inclusion in the model follows the need to buffer daily variations in leaf metabolic carbon as well as in the corresponding leaf osmotic potential. *Sapwood storage carbon* in parenchymatic tissues of branches, stems and coarse roots conforms a much larger carbon pool and is intended to serve as carbon source for growth and long-term (seasonal and inter-annual) storage (Plavcová and Jansen, 2015).




Figure 15.1: Plant labile (metabolic and storage), structural and dead carbon pools and processes determining carbon flows between them (indicated using continuous arrows). Decomposition pools are indicated in gray, and flows towards the atmosphere or decomposition pools are indicated using dashed arrows. Translocation of reserves during senescence is not represented.

Different processes contribute to the carbon balance of plant's structural and labile compartments:

- Carbon balance of **labile carbon pools** is performed by considering **photosynthesis, maintenance respiration, growth costs, sugar-starch conversion, root exudation and phloem transport**. Photosynthesis estimates come from the water balance submodel. Respiration is modelled following the *growth-maintenance* paradigm, but we acknowledge that such approach has limitations and that others have been proposed such as assuming a constant *carbon use efficiency* or the *process/residual approach* (Cannell and Thornley, 2000; Gifford, 2003). Maintenance respiration demand of leaves and sapwood depends on both structural and metabolic carbon, whereas only structural carbon counts for fine root maintenance respiration because they lack labile carbon pools. Maintenance respiration of leaves and fine roots can be species-specific rates, but can also be estimated from species-specific values of nitrogen concentration (Reich et al., 2008). In contrast, the species-specific maintenance respiration per sapwood unit mass (in fact, only parenchymatic tissues contribute to sapwood respiration) is not estimated from nitrogen concentration and should be calibrated. In all three cases, however, temperature dependencies are included in the actual respiration rates (Tjoelker et al., 2001). Growth costs include the carbon cost of building new structures as well as the respiration expenses of growth. Sugar-starch conversion is designed to maintain an equilibrium in the concentration of metabolic sugar. Any surplus in sapwood storage carbon over storage capacity is assumed to be diverted towards root exudation. Hence, root exudation is not modeled as a process actively competing for metabolic carbon, but as a consequence

of plant C storage capacity being surpassed (Prescott et al., 2020).

- Biomass balance in **structural pools** and **dead pools** comes from the interplay between **growth** (i.e. new tissue formation) and **senescence** (see 15.1.4). Structural growth rates are lower than growth costs for labile pools, since growth costs also include growth respiration expenses. During leaf senescence it is assumed that non-structural carbon sources (i.e. leaf labile carbon pools) are recycled and relocated to sapwood storage. Analogously, when sapwood is converted into heartwood, labile carbon in the protoplasm of parenchyma cells is assumed to be re-absorbed by neighboring living cells as storage (Spicer, 2005).

Further design details and formulation regarding the estimation of carbon pool sizes and carbon balance components are given in chapter 16.

15.1.3 Primary growth phenology

Whereas in the water balance models expanded leaf area corresponded to the portion of live leaf area that was unfolded according to the phenological status, in the growth model leaf area changes result from explicit simulation of growth. However, the leaf phenology submodel still controls the duration of phenophases corresponding to leaf elongation and leaf senescence (see 4.1), as well as persistent bud formation and dormancy (see Fig. 15.2).

Primary growth phenology is defined by combining leaf phenology types (i.e. **PhenologyType**), seasonality of primary growth (i.e. **GrowthPeriod**) and the degree of determinacy of primary growth (i.e. **GrowthDeterminacy**), which describes how primary growth in the current year occurs either (Baumgarten et al., 2026): (i) from elongating organs preformed the preceding year and contained within persistent buds: *determined growth*; (ii) from continuous meristematic activity forming new shoots *indetermined growth*; or (iii) from both strategies: *intermediate growth*. The following five combinations are common (see Fig. 15.2):

- Evergreen / Whole-year growth / Indetermined growth*: Evergreen plants with whole-year indetermined growth are assumed to shed old leaves at the same time as new leaves appear. Leaf programmed senescence and leaf formation can occur continuously throughout the year, and leaf growth occurs without bud phase. In this growth phenology the variation of leaf area arises as the interplay between leaf senescence and growth.
- Evergreen / Spring growth / Determinate growth*: Evergreen plants with determined, spring growth are assumed to form persistent buds before winter and concentrate budburst, leaf elongation and leaf senescence during the same period in spring or early summer. The amount of leaf area to be elongated is decided during the persistent bud formation period of the preceding year. Once preformed leaf elongation is over, new buds are formed for the new year to come.
- Evergreen / Spring growth / Intermediate growth*: Evergreen plants with spring growth and intermediate growth determinacy have a mixed growth strategy between the former two. Like plants with determinate spring growth, persistent buds are formed the preceding year, and a first set of leaves is flushed out in spring. Additionally, the plant can continue forming new leaves after these initial ones have finished elongation, in the same way as evergreen plants with indeterminate growth (until there is a decrease in photoperiod). From that moment on, persistent buds are formed for the next year.
- Winter (semi-)deciduous / Spring growth / Determined growth*: In winter deciduous and semi-deciduous plants with determined spring growth leaf bud formation occurs at the end of the leaf elongation. Buds remain under dormancy until next year, when budburst and elongation occurs. Winter deciduous species shed their all their leaves at the time of leaf senescence in autumn, whereas semi-deciduous species keep dead leaves in the crown during winter until budburst of new leaves.
- Winter (semi-)deciduous / Spring growth / Intermediate growth*: Winter (semi-)deciduous species with spring intermediate growth are the analogue of intermediate evergreen species but with winter (semi-)deciduous phenology. Budburst and elongation of preformed leaves can be followed by new leaf formation during the growth season. Like in intermediate evergreen growth plants, bud formation is set to occur between the decrease of photoperiod and the start of dormancy.

Legend

- Persistent bud formation
- Bud dormancy
- Budburst and leaf elongation
- Neogrowth (free growth)




Figure 15.2: Schematic representation of processes and timings for different primary growth phenology types

Although phenology parameters allow some flexibility in growth phenology (e.g., to define the period of preformed leaf elongation or the duration of bud formation after cessation of leaf growth), by design leaf formation cannot occur after photoperiod has decreased beyond some threshold ($Ph_d > Ph_{sen}$) and bud formation cannot extend beyond the point of programmed winter dormancy (i.e. $S_{sen,d} \geq S_{sen}^*$) (see 4.1.4).

15.1.4 Growth and senescence processes

The forest growth model allows simulating the variation of living tissues via the following growth and senescence processes (see fig. 15.3):

- i. **Bud formation:** Organogenesis during persistent bud formation does not consume any carbon, and is scheduled to last a number of days (see parameter `BudFormationDays`).
- ii. **Leaf/twig growth:** Leaf area increase occurs during *leaf elongation* periods (after budburst) or during *indetermined (free) growth*. Unlike bud formation, leaf elongation or indetermined (free) leaf growth does consume carbon. Moreover, leaf growth includes an additional carbon cost in *twig growth*, according to the biomass ratio between both.
- iii. **Leaf/twig senescence and leaf fall:** Leaf/twig senescence (and the corresponding increases in dead leaf area) can occur in the model due to *leaf aging* in evergreen species, due to *programmed leaf senescence* in deciduous species, or due to *leaf dessication*. Leaf dessication is assumed to occur because cavitation in tracheids or vessels leads to a hydraulic disconnection of leaves, which inhibits replacing the water lost. To avoid sudden defoliation of whole crowns, the model assumes a variation in the hydraulic vulnerability of the branches supporting leaves. Leaf senescence increases the leaf area index corresponding to dead leaves, thereby increasing the leaf and twig dead carbon pools. *Leaf fall* is the process that transforms dead leaf area into leaf and twig litter, which triggers decomposition processes (see fig. 15.1). In addition, fires can induce *foliage burning*, which also leads to a decrease in dead and live leaf area (but where non-structural carbon reallocation is not possible) and produces carbon emissions due to leaf and twig combustion.
- iv. **Branch senescence and recovery:** Branch biomass is not explicitly modeled. Instead, the model tracks a state variable representing the percentage of viable crown buds with respect to a fully developed crown, which implicitly represents the availability of live branches. Cavitation-induced *bud dessication* processes lead to a decrease in the amount of crown buds, which then limits primary growth. In turn, *primary growth* (i.e. leaf growth) is assumed to lead to the progressive recovery in the percentage of viable crown buds, again implicitly assuming the growth of new branches.
- v. **Sapwood growth:** Sapwood formation corresponds mostly to secondary growth (but primary growth occurs at the same time due to diameter-height relationships, see 15.1.7). The model assumes that the hormonal signals controlling cambium division have been mainly synthesized in leaves. Auxins are known to be primarily generated in the plant's apical meristems, young leaves, and flower buds; from where they are transported throughout the plant via the phloem (Bajguz and Piotrowska-Niczyporuk, 2023). More specifically, we assume that the levels of secondary growth hormones (auxines) depend linearly on the *current amount of foliage relative to the expected maximum foliage of the same plant without competition*. This entails that: (a) leaf phenological status of deciduous species will affect sapwood growth rates; (b) for a given plant size (i.e. cambium surface) the amount of foliage will be higher or lower depending on the shading created by taller plants and, hence, understory plants or densely-crowded plants will have lower sapwood growth rates than plants that are isolated or overtop the other plants; (c) defoliation events (e.g. from drought impacts) will have an effect on sapwood growth rates, as observed empirically (Javoy et al., 2025).
- vi. **Sapwood senescence:** Losses in sapwood area (i.e. conversion to heartwood) occur via tissue *aging* (i.e. assuming a daily rate of conversion). Since xylem tracheids and vessels are already dead cells, sapwood senescence refers to the death of parenchymatic tissue. Similarly to 3-PG (Landsberg and Waring, 1997) we assume that the relative turnover rate of sapwood is smallest for young plants, and it increases up to a maximum value. However, recent reviews indicate that heartwood formation should be viewed as an active developmental program during which a conductive but vulnerable sapwood is transformed in a non-conductive but durable heartwood (Spicer, 2005). Sapwood senescence also occurs whenever the sapwood area exceeds the amount required to sustain the current leaf area target. Note that stem *cavitation* reduces the area of sapwood that is effectively functional for water transport,

but does not lead to sapwood senescence *per se*, unless mortality occurs. Sapwood senescence produces heartwood biomass in trees, which does not decompose as long as the trees are alive. In contrast, sapwood senescence leads to increase the mass of standing dead branches in shrubs (see fig. 15.1).

- vii. **Fine root growth:** Fine root growth can in principle occur any time of the year (i.e. the model does not explicitly consider root phenology), but since fine root biomass targets depend on leaf area, seasonal fine root area patterns may follow phenological changes in leaf area.
- viii. **Fine root senescence:** Fine root senescence occurs via *aging* with a turnover rate depending on soil temperature.

15.1.5 Source and sink limitations to growth

The growth rates leaf/twig, sapwood and fine root compartments are primarily determined by parameters specifying *maximum daily growth rates* (an exception to this rule is leaf preformation within buds). Actual growth rates are determined by modulating maximum growth rates by taking into account both **source** (i.e. carbon availability) and **sink** (i.e. biophysical constraints on plant tissue expansion) limitations to growth (Fatichi et al., 2014; Guillemot et al., 2015; Körner, 2015) (see fig. 15.3).

Carbon availability limitations to growth apply to leaf and fine root growth because storage carbon levels need to be at more than 10% of their capacity. In contrast, sapwood formation is only allowed if storage carbon levels are above a given species-specific threshold which are normally much higher (e.g. 50%) than required for leaf or fine root growth. Thus, when carbon levels are relatively low, sapwood growth rates fall to zero, whereas replacement of leaves and fine roots continues to sustain metabolism. This design was chosen to stop sapwood growth much before emptying storage carbon reserves, so that maintenance respiration can be sustained longer whenever photosynthesis is low, as in shadow-tolerant species.

Temperature and turgor sink limitations to growth are implemented following Cabon et al. (2020a) and Cabon et al. (2020b). These authors developed their approach for tracheid production and enlargement only, whereas we apply it to different kinds of tissues. Bud formation and leaf elongation of preformed leaves can both be reduced due to turgor limitations during the corresponding organogenesis or expansion process, but these processes do not take into account temperature effects. In contrast, the rate of indetermined (free) leaf area growth is affected by both temperature and turgor conditions. Note that in determinate growth organogenesis and bud formation is temporally separated from elongation, the turgor conditions during different periods (spring or the preceding summer/autumn) will affect the final amount of leaves produced, which can lead to complex leaf area patterns. In the case of fine roots, differences in soil water potential between layers can lead to different degrees of turgor limitation for the growth of fine roots, resulting in acclimation of fine root distribution to soil moisture dynamics.

15.1.6 Growth coordination

Leaf area target limits the maximum amount of leaves formed during leaf organogenesis and this target is determined by plant size (i.e. tree diameter) and stand structure (i.e. competition for light) via allometric equations of foliar biomass (see fig. 15.3). This limits tissue growth to the replacement of aging tissues, unless there are changes in forest structure, i.e. competition is released or the target tree increases in diameter. Sapwood growth targets are determined from functional relationships (e.g. target Huber value or whole-plant conductance) with leaf area targets (see below). Since sapwood formation leads to tree diameter increases, and therefore to forest structural changes, maximum sapwood growth rate indirectly determine the increase in leaf area and fine root targets, hence influencing the whole-plant maximum growth rate.

Two options are possible regarding the functional coordination between structural carbon of leaves and sapwood (see fig. 15.3):

1. *Stable Huber value:* In this strategy, growth of leaf area and sapwood area need are coordinated so that the ratio of sapwood area to leaf area (i.e. Huber value) is in dynamic equilibrium around its initial value. To this aim, sapwood senescence is scheduled whenever the current Huber value is larger than the target (i.e. there is an excess of sapwood area). This approach is similar to the LPG model (Sitch et al., 2003), where different turnover rates are applied for different tissues, but then the model tries to




Figure 15.3: Design of leaf, fine root and sapwood growth and senescence drivers.

satisfy the pipe model (Shinozaki et al., 1964) by allocating C where it is more limiting. Note that in this strategy the whole-plant conductance will progressively decrease with the increase in plant height, because of the increase in the length of stem xylem pathway.

2. *Stable whole-plant conductance*: Ratios of sapwood area to leaf area (i.e. Huber values) can vary within species, due to environmental conditions (Mencuccini and Grace, 1995). In this allocation strategy, growth of leaf area, sapwood area and height needs to be coordinated so that whole-plant conductance (i.e. from roots to leaves) is in dynamic equilibrium around its initial value. Sapwood senescence is scheduled whenever the current whole-plant conductance is above the target value. In this strategy, the Huber value will tend to increase with the increase in plant height, thereby increasing the sapwood respiratory demands.

When running the growth model along with the basic water balance submodel, a stable Huber value is the only strategy available and fine root area growth also follows the need to achieve a stable ratio between leaf area and fine root area. In contrast, the two strategies are possible when simulating growth along with the advanced water balance. In this case, fine root area growth is coordinated with the formation of the other tissues via the need to ensure an *average resistance fraction in the rhizosphere*, which is used to define the fine root area target and, hence, the allocation to fine roots.

15.1.7 Update of structural variables

Tree diameter increment follows the creation of sapwood area, whereas sapwood area decreases do not affect diameter, as they imply conversion to heartwood. Tree height increments are estimated from diameter increments following diameter-height ratios that are modulated by the proportion of light available to the plant cohort, as frequently done in forest gap models (Lindner et al., 1997). Tree crown ratios are updated using static allometric relationships, whereas shrub height and cover are estimated from new leaf area values by inverting static leaf biomass allometries (see chapter 23).

15.1.8 Plant mortality

Mortality pathways

Dynamic vegetation models implement mortality processes in very different ways, from purely-empirical to process-based (Hawkes, 2000; Keane et al., 2001; Bugmann et al., 2019). The fact that proximal causes of mortality are complex, co-occurring, interrelated and variable has prompted the question of whether a mechanistic approach will improve prediction ability (Adams et al., 2013). In medfate, (above-ground parts of) woody plants can die following a combination of empirical and process-based approaches:

1. Trees with $DBH < 7.5$ cm (i.e. saplings) are assumed to suffer mortality due to a self-thinning process of unspecified cause. Specifically, the model represents the self-thinning process occurring between small trees (e.g. 1 cm DBH) resulting from seed establishment (or resprouting) and trees achieving the size (e.g. 7.5 cm) corresponding to inclusion as individual in forest inventories (e.g. ingrowth size). Although implemented here, this self-thinning process is most relevant for simulations of mid- and long-term forest dynamics (see 19).
2. All woody cohorts are assumed to die at a basal rate due to processes not included in the model (e.g. biotic attacks). This rate is empirically determined and can be a constant value or, in the case of trees, can depend on the degree of competition (i.e. basal area). This latter option is important to ensure a limit in stand basal area, which may not occur as a result of mechanistic mortality processes.
3. Mortality rates can be higher than the basal rates whenever physiological thresholds presumed to lead to plant mortality are surpassed (Hawkes, 2000). The model allows plants to die from either *starvation* (if metabolic carbon is exhausted) or *dessication* (tissue dehydration) (McDowell et al., 2008; McDowell, 2011). Allowing for starvation is important to simulate death due to shading by other plant cohorts, whereas plant dessication is important to represent situations of low soil water availability.

With respect to intra-cohort variability, the model can be run assuming that all individuals within plant cohorts are functionally equal, so that mortality applies to the entire plant cohort. Alternatively, mortality can be applied to a fraction of the plant cohort, in proportion to the mortality likelihood, which implicitly acknowledges intra-cohort differences. As common in other models, mortality process can be simulated in a deterministic or stochastic fashion.

Physiological underpinnings of drought-related mortality

Multiple mechanisms determine plant survival or mortality under drought (McDowell et al., 2008, 2022). However, hydraulic failure is widely recognized as one of the key mechanisms of drought-induced mortality in woody plants. Lethal water potential have been defined as a critical threshold beyond which trees are unable to recover from drought, even after rewatering. Experimental work by Urli et al. (2013) determined that irreversible damage in angiosperm saplings occurred only after 88% losses in stem hydraulic conductance. Similarly, Hammond et al. (2019) found probability of mortality to strongly increase beyond 80% of loss in hydraulic conductivity in saplings a gymnosperm species. In a multi-species synthesis, Adams et al. (2017) assessed found that all studied species showed at least 60% or higher loss of xylem hydraulic conductivity, whereas evidence for carbon starvation was more common for gymnosperms. While loss of hydraulic function seems ubiquitous in drought-related mortality, not all researchers agree that hydraulic failure be the primary cause of mortality. Körner (2019) argued that hydraulic failure often correlates with a critical degree of tree dehydration and that the disruption of capillary continuum from soil to the root because of soil dehydration is more important because it forces trees to drop their leaves and rely on their own water reserves. Trees will die when the symplast surpasses its dehydration tolerance, and Körner (2019) considers xylem embolism as a byproduct rather than the cause of critical tree dehydration under severe drought, so that mechanistic models should focus on the dehydration process. Other authors have stressed the importance of the interplay between water reserves and water loss rates through cuticle after stomatal closure (Blackman et al., 2016; Martin-StPaul et al., 2017). Regardless of causality, loss of hydraulic function could still be a good indicator of mortality likelihood. However, recently Martinez-Vilalta et al. (2019) suggested use the relative water content as an indicator of mortality risk, as it integrates several aspects related to hydraulics, stomatal responses and carbon economy during drought.

Starvation thresholds

We use the amount of metabolic sugar in the stem as indicator of carbon starvation. Martínez-Vilalta et al. (2016) found that plants can have very distinct amounts of non-structural carbon concentration but minimum seasonal values are around 46% of maximum values. Accordingly, we define the indicator of starvation in relative terms (see control variable `mortalityRelativeSugarThreshold`, set to 30% by default) with respect to the stem equilibrium sugar concentration.

Dessication thresholds

Following Martínez-Vilalta et al. (2019), we primarily use the relative water content (RWC) in the stem symplastic tissues as the stress variable increasing dessication likelihood under drought. Whereas the stem symplastic pressure-volume curve is always used to determine RWC from the stem water potential, in the basic water balance model the latter is assumed to be equal to the estimated plant water potential (i.e. $\Psi_{sym,stem} = \Psi_{plant}$; see 6.2.1). Moreover, in the advanced water balance model there is no distinction between stem apoplastic and symplastic (i.e. parenchymatic) water potentials (i.e. $\Psi_{sym,stem} = \Psi_{apo,stem}$) unless plant water compartments are explicitly simulated.

Using stem relative water content as indicator of dessication has the inconvenient, however, that the parameters $\pi_{0,stem}$ and ϵ_{stem} specifying the stem pressure-volume curve are often not known, so that estimates have to come from its covariation with other traits (here wood density, see @ref{inputationPV}). The uncertainty coming from this parameterization issue can be avoided if one combines the use of RWC with the loss of xylem hydraulic conductivity, since parameters of the hydraulic vulnerability curve are much more frequently available. For this reason, we use the **maximum of stem relative water content and relative xylem hydraulic conductance** as the drought stress variable to determine dessication.

The critical threshold for dessication is specified in the control variable `mortalityRWCThreshold`, set to 40% by default. Note that most studies focus on leaf RWC, for which Kursar et al. (2009) indicate ranges between 16% and 61%, and Körner (2019) advocates for values around 50%.

Mortality effects

Mortality processes lead to the estimation of the number of dead individuals during the current day (N_{dead}). The model then determines what fraction of these individuals will resprout from the root collar (although resprouts are actually created by `fordyn()` at the end of the year). The fine root and coarse root biomass of non-resprouting individuals are added to the corresponding litter pools. Leaf area of all dead trees is added to dead leaf area (which effectively transfers leaf biomass and twig biomass to dead carbon pools). The aboveground wood biomass (i.e. heartwood and sapwood) of all dead trees is stored in cumulative variables ($D_{smallbranches}$ and $D_{largewood}$) for future snag creation.

Details of the implementation of plant mortality are given in section 17.4.

15.1.9 Fire occurrence, severity and emissions

Fire occurrence and behavior

The occurrence of fires is represented in the growth model in a forcing variable that describes the daily probability that a wildfire strikes the stand. Whenever a fire occurs, the model first simulates fire behavior according to the conditions of the stand (see chapter 26) and then fire severity and carbon emissions are evaluated.

Fire severity

The degree to which vegetation is impacted by fire depends on the heat fluxes incident on plant tissues and organs, which is an outcome of various fire behavior variables (such as fireline intensity and residence time) and the position of plant organs in the stand, and the physiological responses triggered thereafter. Fire effects on plants can be classified as two types: first- and second-order effects (Bär et al., 2019; Hood et al., 2018). First-order effects comprise the immediate impacts of heat transfer on plant tissues. Nonlethal first-order heat injuries can trigger second-order effects, such as physiological limitations in carbon and water relations or increased susceptibility to insect attacks and pathogenic infections. The growth model currently deals with first-order effects only. Moreover, the model focuses on effects in above-ground tissues (leaves/twigs,

buds and cambium), disregarding fire effects on below-ground structures such as seed banks, or below-ground meristems, which are also important for the resilience of ecosystems (Stephan et al., 2010).

Fire-induced plant injury and mortality models can be grouped into two categories: (1) statistical models that involve regression equations relating injury and mortality to observable indicators (Peris-Llopis et al., 2024); and (2) process-based models that explicitly simulate the underlying thermal and biophysical processes occurring when a plant is heated (Butler and Dickinson, 2010; Hood et al., 2018). The design of first-order fire effects on vegetation follows, with some modifications, the process-based model proposed by Michaletz and Johnson (2008) for surface fires, and it is outlined in fig. 15.4. The surface fire behavior variables required for severity evaluation are the residence time and fireline intensity. With them, the model estimates the air temperature gradient of the plume, which is then compared to the distribution of foliage for the determination of the proportion of the crown foliage and crown buds that die due to heat accumulation. In addition, the model determines the effects on the cambium by estimating the radial necrosis depth and comparing it with bark thickness. If enough heat reaches the cambium, the tree is considered girdled, which leads to above-ground mortality. Otherwise, the degree of crown bud necrosis will determine whether the plant can resprout from the crown (and rebuild the leaf area lost) or not.

Crown fires will be considered to impact the forest stand whenever surface fireline intensity is larger than the critical value considered necessary for crowing (Scott and Reinhardt, 2002). Crown fires are considered to burn for all plant cohorts, 100% of crown foliage and, most likely, 100% of crown buds (although very fast crown fires could spare buds if sufficiently protected, (Michaletz and Johnson, 2008)). When crown fires do not occur at the stand level, torching may still occur for particular plant cohorts depending on its crown base height, with the same effects.

If 100% of crown buds have died but the cambium is alive and the species has dormant buds beneath the bark, then the plant has suffered crown kill but it can resprout epicormically without losing height (Hood et al., 2018). If the cambium is dead, however, all above-ground plant organs will be considered dead (i.e. topkill; Hood et al. (2018)). At this point, the plant may still exhibit basal resprouting from the root crown or below-ground meristems, but is dealt in the forest dynamics model (see chapter 19).

While the severity sub-model described above is based on heat transfer and accumulation in tissues, causing necrosis, it is important to note that a growing number of studies increasingly have been documenting fire impacts on xylem hydraulic function, both via increased vapor pressure deficit (causing cavitation) and via structural damage in the xylem, affecting its efficiency and vulnerability (Bär et al., 2019; Hood et al., 2018). These first-order effects are not considered here.

Fire emissions

Whenever foliage burning occurs, the corresponding dry biomass of leaves and twigs is added to the carbon emissions. In addition, for cohorts where above-ground mortality occurs 20% of tree wood biomass and 100% of shrub wood biomass are assumed to be lost via combustion and are added to fire carbon emissions (see fig. 15.4). The remaining 80% of aboveground tree wood biomass contributes to dead snag biomass. Belowground biomass assumed to die (unless resprouting is scheduled) but does not contribute to emissions. Fire emissions include combustion of litter compartments (only leaves, twigs and small branches) and snag compartments (only small branches).

Details of the implementation of fire severity and emissions are given in section 17.5.

15.1.10 Organic matter decomposition

The design of carbon decomposition processes is a modification of the CENTURY model, and its daily variant DAYCENT (Parton et al., 1988, 1993, 1998), as described in Bonan et al. (2019). It is summarized in fig. 15.5 below.

The decomposition model simulates a number of chemically distinct organic matter pools with varying decay rates. The CENTURY/DAYCENT models partition plant litter into metabolic material that readily decomposes and structural material that is more resistant to decomposition. Leaf litter enters the surface pools, and root litter enters the belowground pools. The model additionally represents twig, small-branch




Figure 15.4: Design of fire effects on foliage, buds and cambium.

and large wood debris (on the surface) as well as coarse root debris (belowground). Only leaf litter fall and fineroot litter fall generate metabolic material, whereas the other types of litter (large wood, small branches, twigs and coarse roots) generate only structural material. Litter pools include two important differences with respect to the description of DAYCENT provided in Bonan et al. (2019). First, structural litter pools are species-specific (yellow boxes in fig. 15.5), which allows decomposition rates to be dependent on the species identity of the plant that generated the litter, thus accounting for potential differences in lignin content. Second, the model also considers aboveground standing dead wood (i.e. snags), which originate from plant mortality (or aboveground dead structures in resprouting cohorts) and will normally decompose at slower rates than downed deadwood (magenta gray boxes in fig. 15.5). While the growth model deals with decomposition of these pools, dead branch fall and snag fall processes, which lead to transfer of carbon from standing pools to downed litter pools, are dealt with in the forest dynamics (i.e. `fordyn()`) model (see 19.1.4).

In CENTURY/DAYCENT, soil organic matter is represented by *active*, *slow* and *passive* carbon pools, that represent material with increasingly longer turnover time. These pools have a surface and belowground component, except the passive pool, which only occurs belowground (light gray boxes in fig. 15.5). Decomposition process in a given pool transfers carbon to other pools, while releasing carbon to the atmosphere due to heterotrophic respiration. This conceptualization results in a system of first-order linear differential equations. Each pool has a specific base decomposition rate, which is modified for abiotic factors. Abiotic factors (temperature, pH and moisture) in the surface and belowground carbon pools are those corresponding to the first soil layer, whereas abiotic factors for standing dead pools correspond to air temperature and relative humidity. Processes of the nitrogen cycle (i.e. immobilization, mineralization and plant uptake) are not simulated at present.

15.2 State variables

15.2.1 State variables related to live plant cohorts

True state variables

Many *cohort-level state variables* are inherited by the corresponding water balance model and depend on its level of complexity (see 3.2 and 7.3). Several variables were treated as fixed parameters in the water balance models and become variable here:

- Leaf area index of live leaves, LAI_{live} ($m^2 \cdot m^{-2}$).
- Leaf area index expected assuming no light competition from larger plants LAI_{NC} ($m^2 \cdot m^{-2}$).
- Actual expanded leaf area, LAI_{act} ($m^2 \cdot m^{-2}$).
- Dead leaf area, LAI_{dead} ($m^2 \cdot m^{-2}$).
- Percent cover of shrubs, $Cover$ (%).
- Density of living individuals in the plant cohort, N ($ind \cdot ha^{-1}$).
- Density of dead individuals in the plant cohort, N_{dead} ($ind \cdot ha^{-1}$).
- The proportion of fine roots, FRP_s , in each soil layer s ([0-1]).
- Cumulative mass of small dead branches (diameter between 0.635 cm and 7.5 cm; 10h and 100h fuels) corresponding to cohort dead individuals $D_{smallbranches}$ ($gC \cdot m^{-2}$).
- Cumulative mass of large dead wood (diameters > 7.5 cm; 1000h and 10000h fuels) corresponding to cohort dead individuals $D_{largewood}$ ($gC \cdot m^{-2}$).

Variables $D_{smallbranches}$ and $D_{largewood}$ are not to be confounded with state variables corresponding to decomposing snags (see below). In growth simulations $D_{smallbranches}$ and $D_{largewood}$ are used to track mass of dead wood only. In forest dynamic simulations, the values of $D_{smallbranches}$ and $D_{largewood}$ at the end of the simulated year are used by function `fordyn()` to create new snag cohorts to start the decomposition process the year after.

Individual-level state variables are needed in growth simulations:

- Plant height, H (cm).
- Diameter at breast height for trees, DBH (cm).




Figure 15.5: Design of dead carbon pools and decomposition processes. Boxes are carbon pools (some of them being cohort-specific or species-specific). Dashed arrows indicate processes originating surface or belowground litter. Continuous arrows indicate carbon transfer from one pool to another, and most of them include release of carbon to the atmosphere via heterotrophic respiration.

- Crown ratio, CR ([0-1]).
- Actual leaf area, LA_{act} (m^2).
- Target leaf area, LA_{target} (m^2).
- Dead leaf area LA_{dead} (m^2).
- Preformed leaf area in persistent buds, $LA_{preformed}$ (m^2).
- Leaf area programmed to enter senescence, $LA_{progsen}$ (m^2).
- Sapwood area, SA (cm^2).
- Target sapwood area, SA_{target} (cm^2).
- The percentage of the crown with active buds, $PCAB$ (%).
- Fine root biomass, $B_{fineroot,s}$ ($g\ dry$), and the target fine root biomass, $B_{fineroot,target,s}$ ($g\ dry$), in each soil layer s .
- Sugar concentration in leaves, SS_{leaf} ($mol\ gluc \cdot L^{-1}$).
- Starch concentration in leaves, ST_{leaf} ($mol\ gluc \cdot L^{-1}$).
- Sugar concentration in sapwood (stem, branches and coarse roots), $SS_{sapwood}$ ($mol\ gluc \cdot L^{-1}$).
- Starch concentration in sapwood (stem, branches and coarse roots), $ST_{sapwood}$ ($mol\ gluc \cdot L^{-1}$).

Note that LA_{act} is the same state variable as LAI_{act} , but expressed at the individual level. The same happens between LA_{live} (a derived variable) and LAI_{live} ; and between LA_{dead} and LAI_{dead} . These pairs of state variables are related by cohort density (N).

Derived variables

Other individual-level variables can be derived from the true state variables and, hence, hence do not need to be treated as state variables themselves but are often used in equations instead of the true state variables:

- Live leaf area, LA_{live} , from LA_{target} and the percentage of crown active buds ($PCAB$).
- Leaf biomass, B_{leaf} ($g\ dry$), from LA_{act} .
- Twig biomass, B_{twig} ($g\ dry$), from LA_{act} .
- Dead leaf mass, D_{leaf} ($g\ dry$), from LA_{dead} .
- Dead twig (diameter < 0.635 cm) mass, D_{twig} ($g\ dry$), from LA_{dead} .
- The radial length of coarse roots in each soil layer s , $L_{radial,s}$ (mm), from FRP_s .
- Sapwood biomass, $B_{sapwood}$ ($g\ dry$), from SA , H and FRP_s and $L_{radial,s}$.
- Heartwood mass, for trees, $D_{heartwood}$ ($g\ dry$), from SA , DBH , H and FRP_s and $L_{radial,s}$.
- Leaf storage volume ($V_{storage,leaf}$; in L) from B_{leaf} .
- Sapwood storage volume ($V_{storage,sapwood}$; in L) from SA , H and FRP_s and $L_{radial,s}$.
- Fine root area (FRA_s ; in m^2) in each soil layer s , from $B_{fineroot,s}$.

Some belowground variables (such as the radial length of coarse roots) are only considered dynamic when simulating growth with advance water balance.

15.2.2 State variables related to carbon decomposition

Dead standing wood cohorts

Cohorts of dead standing wood (i.e. snag cohorts) are identified by combinations of species, DBH of the original tree, height and dead age (i.e. the years passed since mortality occurred). Those properties match the element `snagData` of forest objects (see 2.4.2.4) and are modified by the model of forest dynamics, whereas the growth model deals with the daily decomposition process. In the growth model, the following state variables are tracked for cohorts of dead standing wood:

- Mass of small dead branches, with diameters comprised between 0.635 cm and 7.5 cm (i.e. 10h and 100h fuels), $SN_{smallbranches}$ ($g\ C \cdot m^{-2}$).
- Mass of large dead wood, with diameters > 7.5 cm (i.e. 1000h and 10000h fuels), $SN_{deadwood}$ ($g\ C \cdot m^{-2}$).

Structural litter pools

Structural litter pools are **species-specific**. For each species, the following pools are tracked, matching the element `litterData` of forest objects (see 2.4.2.4):

- Leaf structural litter, LI_{leaves} ($gC \cdot m^{-2}$).
- Twig (diameter < 0.635 cm; 1h fuels) structural litter, LI_{twigs} ($gC \cdot m^{-2}$).
- Small branches (diameter between 0.635 cm and 7.5 cm; 10h and 100h fuels) structural litter, $LI_{smallbranches}$ ($gC \cdot m^{-2}$).
- Large wood (diameter > 7.5 cm; 1000h and 10000h fuels) structural litter, $LI_{largewood}$ ($gC \cdot m^{-2}$).
- Coarse roots structural litter, $LI_{coarseroots}$ ($gC \cdot m^{-2}$).
- Fine roots structural litter, $LI_{fineroots}$ ($gC \cdot m^{-2}$).

Soil carbon pools

The decomposition model keeps track of the carbon amount in different organic matter pools which come from the CENTURY model and match the element `SOCData` of forest objects (see 2.4.2.5):

- Surface metabolic litter, $LI_{surface,metabolic}$ ($gC \cdot m^{-2}$).
- Surface active decomposition pool, $SOC_{surface,active}$ ($gC \cdot m^{-2}$).
- Surface slow decomposition pool, $SOC_{surface,slow}$ ($gC \cdot m^{-2}$).
- Soil metabolic litter, $LI_{soil,metabolic}$ ($gC \cdot m^{-2}$).
- Soil active decomposition pool, $SOC_{surface,active}$ ($gC \cdot m^{-2}$).
- Soil slow decomposition pool, $SOC_{surface,slow}$ ($gC \cdot m^{-2}$).
- Soil passive decomposition pool, $SOC_{surface,passive}$ ($gC \cdot m^{-2}$).

15.3 Mass and energy balances

15.3.1 Water and energy balance

Since the growth model builds on soil and plant water balance models, the water and energy balances explained in sections 3.3 or 7.4 (depending on `transpirationMode`) also hold for this model.

15.3.2 Labile carbon balance

The model performs daily carbon balance of *labile carbon* in leaves and sapwood (stem, branches and coarse roots) for each plant cohort. In each of these two compartments carbon balance is performed for the two labile carbon forms, dissolved sugars (i.e. *metabolic carbon*) and starch (i.e. *storage carbon*). In the following all carbon flows (i.e. the components of right hand side of equations) are assumed to be in units of $g\text{ gluc.}$. At the leaf level, the carbon balance equations for dissolved sugars and starch are:

$$\Delta SS_{leaf} \cdot V_{storage,leaf} = A - MR_{leaf} - F_{phloem} - SC_{leaf} - TS_{leaf} \quad (15.1)$$

$$\Delta ST_{leaf} \cdot V_{storage,leaf} = SC_{leaf} - TT_{leaf} \quad (15.2)$$

where A is (gross) photosynthesis, MR_{leaf} is leaf maintenance respiration, F_{phloem} is the sugar flow via phloem towards the stem and roots, SC_{leaf} is the conversion from leaf sugar to leaf starch, TS_{leaf} and TT_{leaf} are the translocation of leaf sugars and starch into sapwood storage, respectively, during leaf senescence (see eq. (15.6) below) or if leaf storage capacity is surpassed.

The corresponding carbon balance equations for sapwood dissolved sugars and starch are:

$$\Delta SS_{sapwood} \cdot V_{storage,sapwood} = F_{phloem} - MR_{twig} - MR_{sapwood} - MR_{fineroot} \quad (15.3)$$

$$-SC_{sapwood} - TS_{sapwood} \quad (15.4)$$

$$\Delta ST_{sapwood} \cdot V_{storage,sapwood} = SC_{sapwood} + TS_{leaf} + TT_{leaf} + TS_{sapwood} \quad (15.5)$$

$$-GC_{sapwood} - GC_{leaf} - GC_{twig} - GC_{fineroot} - RE \quad (15.6)$$

where F_{phloem} is the sugar inflow via phloem transport, MR_{twig} is the maintenance respiration of twigs, $MR_{sapwood}$ is the sum of maintenance respiration of branches, stem and coarse roots, $MR_{fineroot}$ is the maintenance respiration of fine roots; GC_{leaf} , GC_{twig} , $GC_{sapwood}$ and $GC_{fineroot}$ are the carbon costs of

forming new leaves, twigs, sapwood or fine roots (respectively), $SC_{sapwood}$ is the conversion of sapwood sugar to sapwood starch, $TS_{sapwood}$ is the translocation of sapwood sugars into starch during leaf senescence and RE is the root exudation. If we define S_{labile} as the total amount of labile carbon for an individual plant (in $g\ gluc$):

$$S_{labile} = (SS_{leaf} + ST_{leaf}) \cdot V_{storage,leaf} + (SS_{sapwood} + ST_{sapwood}) \cdot V_{storage,sapwood} \quad (15.7)$$

then labile carbon balance at the plant level reduces to:

$$\Delta S_{labile} = A - MR - GC - RE \quad (15.8)$$

where $MR = MR_{leaf} + MR_{twig} + MR_{sapwood} + MR_{fineroot}$ is plant-level maintenance respiration and $GC = GC_{leaf} + GC_{twig} + GC_{sapwood} + GC_{fineroot}$ is plant-level growth costs.

15.3.3 Plant biomass balance

In addition to changes in labile carbon pools, the model also implements changes in the size of structural and dead carbon compartments, as a result of the interplay between growth, senescence and foliage burning processes, as well as cohort-level biomass balance once mortality processes are taken into account.

Live biomass balance For the balance of structural carbon pools, we can define the following structural equations (all variables in $g\ dry$):

$$\Delta B_{leaves} = B_{leaves,growth} - B_{leaves,senescence} - B_{leaves,burnt} \quad (15.9)$$

$$\Delta B_{twigs} = B_{twigs,growth} - B_{twigs,senescence} - B_{twigs,burnt} \quad (15.10)$$

$$\Delta B_{sapwood} = B_{sapwood,growth} - B_{sapwood,senescence} \quad (15.11)$$

$$\Delta B_{fineroot} = B_{fineroot,growth} - B_{fineroot,senescence} \quad (15.12)$$

where $B_{leaves,growth}$, $B_{twigs,growth}$, $B_{sapwood,growth}$ and $B_{fineroot,growth}$ are the biomass increase due to growth for leaves, twigs, sapwood and fineroots, respectively; $B_{leaves,senescence}$, $B_{twigs,senescence}$, $B_{sapwood,senescence}$ and $B_{fineroot,senescence}$ are the biomass loss due to senescence for leaves, twigs, sapwood and fineroots, respectively; $B_{leaves,burnt}$ and $B_{twigs,burnt}$ are the biomass burnt of leaves and twigs.

The following equation defines the biomass balance for (live) structural pools (in $g\ dry$):

$$\Delta B_{structure} = \Delta B_{leaves} + \Delta B_{twigs} + \Delta B_{sapwood} + \Delta B_{fineroot} \quad (15.13)$$

Dead mass balance

Senescence processes increase carbon for dead tissue pools, whereas litter fall and foliage burn decrease them. Hence, we can define the following equations (all variables in $g\ dry$):

$$\Delta D_{leaves} = B_{leaves,senescence} - D_{leaves,fall} - D_{leaves,burnt} \quad (15.14)$$

$$\Delta D_{twigs} = B_{twigs,senescence} - D_{twigs,fall} - D_{twigs,burnt} \quad (15.15)$$

$$\Delta D_{heartwood} = B_{sapwood,senescence} \quad (15.16)$$

$$(15.17)$$

Where $D_{leaves,fall}$ and $D_{twigs,fall}$ are the mass of dead leaves and twigs, respectively, that falls into litter compartments; and $D_{leaves,burnt}$ and $D_{twigs,burnt}$ are the mass of dead leaves and twigs, respectively, that are burnt. For trees, sapwood senescence ($B_{sapwood,senescence}$) results in a corresponding increase in heartwood ($D_{heartwood}$). In contrast, for shrubs heartwood is not considered and sapwood senescence ($B_{sapwood,senescence}$) results in dead standing branch (snag) creation.

Total biomass balance

The overall mass balance at the **individual level** is completed if we add the labile biomass balance to the structural and dead tissue mass balances (in $g\ dry$):

$$\Delta B_{plant} = \Delta B_{structure} + \Delta S_{labile} + \Delta D_{leaves} + \Delta D_{twigs} + \Delta D_{heartwood} \quad (15.18)$$

To evaluate biomass balance at the cohort level, we need to take into account that some individuals will die, so that the biomass balance needs to incorporate mortality losses (all variables in $g\ dry \cdot m^{-2}$):

$$\Delta B_{cohort} = \Delta B_{plant} \cdot N_{cohort} - D_{above,dead} - D_{below,dead} \quad (15.19)$$

where N_{cohort} is the initial cohort density (before mortality occurred), $D_{above,dead}$ is the aboveground (leaves, twigs, small branches and large wood) biomass loss due to mortality of individuals and $D_{below,dead}$ is the belowground (coarse roots and fine roots) biomass loss due to mortality of individuals. The latter two are calculated using:

$$D_{above,dead} = (B_{plant,above} + \Delta B_{plant,above}) \cdot N_{dead} \quad (15.20)$$

$$D_{below,dead} = (B_{plant,below} + \Delta B_{plant,belowground}) \cdot N_{dead,nonresp} \quad (15.21)$$

where $B_{plant,above}$ is the initial plant aboveground biomass (leaves, twigs, small branches and large wood), $B_{plant,below}$ is the initial plant belowground biomass (coarse roots and fine roots); N_{dead} is the density of dead individuals and $N_{dead,nonresp}$ is the density of individuals that are dead and will not resprout next year (resprouting is handled by the forest dynamic model). In turn, $D_{above,dead}$ is divided between standing wood ($D_{wood,non-burnt}$) and wood emissions due to wildfire combustion ($D_{wood,burnt}$):

$$D_{above,dead} = D_{wood,non-burnt} + D_{wood,burnt} \quad (15.22)$$

Note that $D_{above,dead}$ should include dead biomass of leaves and twigs, that is not shown here for simplicity.

15.3.4 Mass balance of decomposition pools

Let ΔSN , ΔLI and ΔSOC be the change in the amount of carbon in snag, litter and soil organic carbon pools ($g\ C \cdot m^{-2}$). Now let ΔDEC be the sum of mass changes in all pools taken together:

$$\Delta DEC = \Delta SN + \Delta LI + \Delta SOC \quad (15.23)$$

ΔDEC is equal to litter and root exudation inputs minus heterotrophic respiration and snag/litter combustion:

$$\Delta DEC = \sum_i (LI_{cohort,i} + RE_i) - SN_{burnt} - LI_{burnt} - HR \quad (15.24)$$

where RE_i is root exudation for cohort i , SN_{burnt} is the mass of snags (small branches) burnt, LI_{burnt} is the mass of litter burnt and HR is heterotrophic respiration, all expressed in $g\ C \cdot m^{-2}$ units. Finally, $LI_{cohort,i}$ is the sum of litter inputs from cohort i , i.e.:

$$LI_{cohort,i} = D_{leaves,fall,i} + D_{twigs,fall,i} + B_{fineroot,senescence,i} + D_{below,dead,i} \quad (15.25)$$

where $D_{leaves,fall,i}$ and $D_{twigs,fall,i}$ is the mass of fallen leaves, respectively, $B_{fineroot,senescence,i}$ is the mass of dead fine roots due to senescence (all after multiplication by cohort density, $N_{cohort,i}$, and carbon content), $D_{below,dead,i}$ is the mass of coarse and fine roots entering decomposition after belowground mortality. The litter inputs corresponding to small branches and large wood are not included here because they are created by the model of forest dynamics as a result of snag fall.

15.3.5 Fire combustion

Fire effects led to carbon emissions (FC ; $g\ C \cdot m^{-2}$) corresponding the combustion of live and dead biomass from plant cohorts, snags and litter:

$$FC = \sum_i B_{burnt,i} + SN_{burnt} + LI_{burnt} \quad (15.26)$$

where SN_{burnt} is the mass of snags burnt (small branches), LI_{burnt} is the mass of litter burnt (only leaves, twigs and small branches) and $B_{burnt,i}$ is the biomass burnt for cohort i :

$$B_{burnt,i} = B_{leaves,burnt,i} + B_{twigs,burnt,i} + D_{leaves,burnt,i} + D_{twigs,burnt,i} + D_{wood,burnt,i} \quad (15.27)$$

where $B_{leaves,burnt,i}$ and $B_{twigs,burnt,i}$ are the mass of live leaves and twigs, respectively, that are burnt; $D_{leaves,burnt,i}$ and $D_{twigs,burnt,i}$ are the mass of dead leaves and twigs, respectively, that are burnt; $D_{wood,burnt,i}$ is the mass of burnt wood for cohort i .

15.3.6 Stand-level carbon balance

The model estimates several carbon fluxes at the stand level, all expressed in units of $gC \cdot m^{-2}$:

- **Gross primary production (GPP)** is the sum of gross photosynthesis (A_g ; in $gC \cdot m^{-2}$) across plant cohorts:

$$GPP = \sum_i A_{g,i} \quad (15.28)$$

where $A_{g,i}$ is the (cohort-level) gross photosynthesis for cohort i .

- **Maintenance respiration (MR)** is the sum across plant cohorts of the carbon emitted to the atmosphere as a result of respiration required for the maintenance of living tissues (leaves, fine roots and sapwood parenchyma):

$$MR = \frac{m_{carbon} \cdot 6}{m_{gluc}} \cdot \sum_i MR_i \cdot (N_i/10000) \quad (15.29)$$

where MR_i is plant-level maintenance respiration for cohort i .

- **Synthesis respiration (SR)** is the sum across plant cohorts of the carbon returned to the atmosphere as a result of the process of synthesizing new structures:

$$SR = \frac{m_{carbon} \cdot 6}{m_{gluc}} \cdot \sum_i SR_i \cdot (N_i/10000) \quad (15.30)$$

where SR_i is plant-level synthesis respiration for cohort i . Note that this is less than the total carbon used for growth, as it does not include the carbon allocated to the new structures.

- **Net primary production (NPP)** is gross primary production minus the maintenance respiration and synthesis respiration terms:

$$NPP = GPP - MR - SR \quad (15.31)$$

Note that net primary production is in some models equal to the carbon allocated to new plant structures (i.e. growth), but in medfate NPP can be used to replenish labile carbon pools.

- **Heterotrophic respiration (HR)** is the amount of carbon returned to the atmosphere as a result of respiration occurred during decomposition processes.
- **Fire combustion (FC)** is the amount of carbon returned to the atmosphere as a result of combustion of plant tissues.
- **Net ecosystem production (NEP)** is the net carbon source of the forest ecosystem, i.e. the net primary production minus heterotrophic respiration and fire combustion:

$$NEP = \Delta C_{stand} = NPP - HR - FC \quad (15.32)$$

NEP expresses the overall carbon balance of the forest stand (ΔC_{stand}), i.e.:

$$\Delta C_{stand} = \Delta SN + \Delta LI + \Delta SOC + \sum_i (\Delta B_{cohort,i} + D_{snags,i}) \quad (15.33)$$

where $D_{snags,i}$ stands for standing dead wood from cohort i that that will become snag cohorts (managed by the model of forest dynamics) and includes non-burnt aboveground mortality ($D_{wood,nonburnt,i}$) and shrub sapwood senescence ($B_{sapwood,senescence,i}$):

$$D_{snags,i} = D_{wood,nonburnt,i} + B_{sapwood,senescence,i} \quad (15.34)$$

15.4 Process scheduling

Growth simulations are conducted in **daily** time steps. For every day to be simulated, the model performs the following blocks of steps:

A. Water and energy balance

Water and energy balance calculations are performed by calling the basic or the advanced water balance model (3.4).

B. Individual-level carbon balance, growth and senescence

The model performs the following steps for each plant cohort:

1. Determine the amount of live structural biomass (in leaves, sapwood and fine root compartments), storage volume (in leaves and sapwood), storage capacity (16.1) and the construction costs per unit tissue (area or dry weight).
2. If $LA_\phi > 0$, estimate gross photosynthesis per individual (A) from cohort-level photosynthesis (A_g).
3. Estimate maintenance respiration for leaves (MR_{leaf}), twigs (MR_{twig}), sapwood ($MR_{sapwood}$) and fine roots ($MR_{fineroot}$) (16.2).
4. If leaf growth is permitted, evaluate leaf temperature and turgor sink limitations, leaf and twig growth costs (GC_{leaf} and GC_{twig} , respectively), leaf and twig synthesis respiration (SR_{leaf} and SR_{twig} , respectively) and leaf area increase (17.1.3).
5. If sapwood growth is permitted, evaluate sapwood temperature and turgor sink limitations, sapwood growth costs ($GC_{sapwood}$), sapwood synthesis respiration ($SR_{sapwood}$) and sapwood area increase (17.1.4).
6. If fineroot biomass is lower than the target biomass, evaluate fineroot temperature and turgor sink limitations, fineroot growth costs ($GC_{fineroot}$), fineroot synthesis respiration ($SR_{fineroot}$) and fineroot biomass increase (17.1.5).
7. Update changes in sugar and starch compartments due to photosynthesis, maintenance respiration and growth. Apply transfer from leaf metabolic to sapwood metabolic carbon (16.2.4) and conversions between storage and metabolic carbon (16.2.5).
8. Determine leaf senescence due to aging and leaf/bud senescence due to cavitation effects (17.2.1).
9. Determine sapwood senescence due to aging. For shrubs, convert sapwood senescence into mass of standing dead branches (17.2.2).
10. Determine fine root senescence by aging (17.2.3). Divide dead fine root mass between metabolic soil carbon pool and fine root litter pool.
11. Evaluate labile carbon translocation due to leaf and sapwood senescence (TS_{leaf} , TT_{leaf} and $TS_{sapwood}$). If leaf storage carbon is surpassed, add the excess to TT_{leaf} . Add the sum of translocated carbon to the starch sapwood pool.
12. Move any surplus in sapwood storage carbon to root exudation (RE). This completes the daily carbon balance for leaf labile pools (15.2), sapwood labile pools (15.6), and plant-level labile carbon (15.8). The model checks at this point the plant-level labile carbon balance.
13. Use growth and senescence results to update actual leaf area, dead leaf area, sapwood area, fine root biomass and derived traits (17.3.1).
14. Translate growth into structural variables (i.e., plant height, tree DBH, tree crown ratio and shrub cover) and update leaf area targets (17.3.2),
15. Update sapwood area and fine root biomass targets (17.3.3).
16. Update storage volumes and the concentration of labile pools (17.3.4).

Steps 2-7 can be conducted at sub-daily time steps if the advanced model has been used for water balance (see control option `subdailyCarbonBalance`).

C. Fire occurrence and behavior

Then, the model evaluates fire occurrence and, if so, predicts fire behavior (see chapter 26).

D. Mortality and fire severity effects

D1. Determine if a wildfire stroke the forest stand and, if so, fire severity effects (17.5). D2. Determine if plant death by starvation or desiccation occurred (17.4).

E. Decomposition

15.5 Inputs and outputs

15.5.1 Soil, vegetation and meteorology

Soil

Soil input requirements are the same as for the basic water balance model and were fully described in section 2.3. Genuchten (1980) water retention curves are forced for this model, because rhizosphere conductance is modelled using van Genuchten hydraulic equations (see subsection 10.2).

Vegetation

Vegetation input requirements were fully described in section 2.4. Note that vegetation structural characteristics are daily modified by simulations of this model, which includes leaf area index (LAI). Moreover, temporal variation in LAI values is related to temporal variation in structural parameters via allometric relationships. Hence, in this model one should not specify a LAI value as input, but specify initial values for tree size (diameter, height), tree density and shrub dimensions (cover, height). Nevertheless, for each plant cohort with hemi-parasitic (mistletoe) plants living on crowns, a corresponding value of mistletoe leaf area index ($LAI_{mistletoe}$) can be defined, which is assumed static during simulations.

Meteorological input

The minimum weather variables required to run the model are min/max temperatures (T_{min} and T_{max}), min/max relative humidity (RH_{min} and RH_{max}), precipitation (P) and solar radiation (Rad). Wind speed (u) is also needed, but the user may use missing values if not available (a default value will be used in this case). Wind speed is assumed to have been measured at a specific height above the canopy (by default at 2 m). Atmospheric CO_2 concentration (C_{atm}) may also be specified, but if missing a default constant value is assumed, which is taken from the control parameters. If wildfire occurrence and severity is to be simulated, the user has to specify daily fire probability (P_{fire}) in an additional column. Definitions and units of these variables were given in section 2.5.

15.5.2 Plant functional parameters

The growth model requires many functional parameters to be specified for plant cohorts. Some of them depend on whether the basic or advanced water balance is adopted. Here we report functional parameters needed in addition to those necessary for the water balance submodel.

A first set of parameters refers to anatomical features (**paramsAnatomy**), most of them already required by water balance model simulations:

Symbol	Units	R	Description
H_{max}	cm	Hmax	Maximum plant height
H_{med}	cm	Hmed	Median plant height
$1/H_v$	$m^2 \cdot m^{-2}$	A12As	Leaf area to sapwood area ratio
R_{LR}	$m^2 \cdot m^{-2}$	Ar2A1	Fine root area to leaf area ratio
SLA	$m^2 \cdot kg^{-1}$	SLA	Specific leaf area
LW	cm	LeafWidth	Leaf width
ρ_{leaf}	$g \cdot cm^{-3}$	LeafDensity	Leaf tissue density
ρ_{wood}	$g \cdot cm^{-3}$	WoodDensity	Wood tissue density
$\rho_{fineroot}$	$g \cdot cm^{-3}$	FineRootDensity	Fine root tissue density
$f_{conduits}$	[0-1]	conduit2Sapwood	Ratio of sapwood corresponding to xylem conduits (the remaining corresponds to parenchymatic tissues)

Symbol	Units	R	Description
SRL	$cm \cdot g^{-1}$	SRL	Specific root length
RLD	$cm \cdot cm^{-3}$	RLD	Fine root length density (i.e. density of root length per soil volume)
$r_{6.35}$		r635	Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm

A second set of parameters refers to allometric coefficients needed for structural updates (**paramsAllometries**).

Symbol	Units	R param	Description
a_{ash}		Aash	Coefficient relating the square of shrub height with shrub area
a_{bsh}, b_{bsh}		Absh, Bbsh	Coefficients relating crown volume with dry weight of shrub individuals
bt_{sh}		BTsh	Bark thickness in shrubs
$a_{cr}, b_{1cr}, b_{2cr}, b_{3cr},$ c_{1cr}, c_{2cr}		B1cr, B2cr, B3cr, C1cr, C2cr	Coefficients to calculate crown ratio of trees
a_{cw}, b_{cw}		Acw, Bcw	Regression coefficients used to calculate the crown width of trees (as intermediary step to obtain the crown ratio).
a_{bt}, b_{bt}		Abt, Bbt	Regression coefficients used to calculate the bark thickness of trees

A third set of parameters refers to growth parameters (**paramsGrowth**).

Symbol	Units	R param	Description
N_{leaf}	$mg N \cdot g dry^{-1}$	Nleaf	Leaf nitrogen concentration per dry mass
$N_{sapwood}$	$mg N \cdot g dry^{-1}$	Nsapwood	Sapwood nitrogen concentration per dry mass
$N_{fineroot}$	$mg N \cdot g dry^{-1}$	Nfineroot	Fine root nitrogen concentration per dry mass
MRR_{leaf}	$g gluc \cdot g dry^{-1} \cdot day^{-1}$	RERleaf	Leaf respiration rate at 20 °C
$MRR_{sapwood}$	$g gluc \cdot g dry^{-1} \cdot day^{-1}$	RERSapwood	Living sapwood (parenchymatic tissue) respiration rate at 20 °C
$MRR_{fineroot}$	$g gluc \cdot g dry^{-1} \cdot day^{-1}$	RERfineroot	Fine root respiration rate at 20 °C
$RGR_{leaf,max}$	$m^2 \cdot cm^{-2} \cdot day^{-1}$	RGRleafmax	Maximum leaf area daily growth rate, relative to sapwood area
$RGR_{cambium,max}$	$cm^2 \cdot cm^{-1} \cdot day^{-1}$	RGRcambiummax	Maximum daily tree sapwood growth rate relative to cambium perimeter length
$RGR_{sapwood,max}$	$cm^2 \cdot cm^{-2} \cdot day^{-1}$	RGRsapwoodmax	Maximum daily shrub sapwood growth rate relative to sapwood area

Symbol	Units	R param	Description
$RGR_{fineroot,max}$	$g\ dry \cdot g\ dry^{-1} \cdot day^{-1}$	RGRfinerootmax	Maximum daily fine root relative growth rate
RGR_{bud}	$m^2 \cdot m^{-2}$	RGRbud	Bud (potential leaf area) daily growth rate, relative to leaf area growth
CC_{leaf}	$g\ gluc \cdot g\ dry^{-1}$	CCleaf	Leaf construction cost
$CC_{sapwood}$	$g\ gluc \cdot g\ dry^{-1}$	CCsapwood	Sapwood construction cost
$CC_{fineroot}$	$g\ gluc \cdot g\ dry^{-1}$	CCfineroot	Fine root construction cost
$SR_{sapwood}$	day^{-1}	SRsapwood	Daily sapwood senescence rate
$SR_{fineroot}$	day^{-1}	SRfineroot	Daily fine root senescence rate
$RSSG$	[0-1]	RSSG	Minimum relative starch for sapwood growth to occur.
$f_{HD,min}$	$cm \cdot cm^{-1}$	fHDmin	Minimum height-to-diameter ratio
$f_{HD,max}$	$cm \cdot cm^{-1}$	fHDmax	Maximum height-to-diameter ratio
	$g\ C \cdot g^{-1}dry$	WoodC	Wood carbon content per dry weight

A third set of parameters refers to mortality/regeneration parameters (**paramsMortalityRegeneration**).

Symbol	Units	R param	Description
$P_{mort,base}$	[0 – 1]	MortalityBaselineRate	Deterministic proportion or probability specifying the baseline reduction of cohort's density occurring in a year as a result of processes not explicitly modelled
TS_{surv}	[0 – 1]	SurvivalModelStep	Time step in years of the empirical survival model depending on stand basal area (e.g. 10)
$\beta_{surv,0}$	[0 – 1]	SurvivalB0	Intercept of the logistic baseline survival model depending on stand basal area
$\beta_{surv,1}$	[0 – 1]	SurvivalB1	Slope of the logistic baseline survival model depending on stand basal area
$N_{tree,recr}$	$ind \cdot ha^{-1}$	RecrTreeDensity	Density of tree recruits from seeds.
$N_{tree,ingrowth}$	$ind \cdot ha^{-1}$	IngrowthTreeDensity	Density of trees reaching ingrowth DBH.
$DBH_{tree,recr}$	cm	RecrTreeDBH	DBH for tree recruits from seeds or resprouting (e.g. 1 cm).
$DBH_{tree,ingrowth}$	cm	IngrowthTreeDBH	Ingrowth DBH for trees (e.g. 7.5 cm).

Vegetation functional attributes are normally filled for each cohort by function **growthInput()** from species identity. However, different parameters can be specified for different cohorts of the same species if desired (see section 2.4.4).

15.5.3 Control parameters

Control parameters modulate the overall behaviour of growth simulations (see section 2.6). Parameters controlling water balance have been already described in sections 3.5.3 and 7.6.3. Here we only mention those control parameters specific to the **growth()** function.

First, there are flags that regulate the application of processes:

- `allowDessication [=TRUE]`: Boolean flag to indicate that mortality by dessication is allowed.
- `allowStarvation [=TRUE]`: Boolean flag to indicate that mortality by starvation is allowed.
- `sinkLimitation [=TRUE]`: Boolean flag to indicate that temperature and turgor limitations to growth are applied.
- `shrubDynamics [= TRUE]`: Boolean flag to allow the application of demographic processes to shrubs.
- `herbDynamics [= TRUE]`: Boolean flag to allow dynamic herb leaf area as a function of shading due to leaf area of woody cohorts.

An important parameter concerns the allocation strategy:

- `allocationStrategy [= "Plant_kmax"]` defines the allocation strategy (investment in leaf area vs. sapwood area) of the growth model. If `allocationStrategy = "Plant_kmax"` then the target leaf area is set to maintain the input whole-plant maximum conductance. If `allocationStrategy = "A12As"` then the target leaf area is set to maintain the input Huber value.

A number of control parameters regulate carbon balance, growth and senescence processes:

- `phloemConductanceFactor [= 0.2]`: Factor to transform stem xylem conductance to phloem conductance (only for `transpirationMode = "Sperry"`).
- `nonSugarConcentration [=0.25]`: A constant specifying non-sugar (inorganic) solute concentration ($\text{mol} \cdot \text{L}^{-1}$) in cells.
- `equilibriumOsmoticConcentration [= c(leaf = 0.8, sapwood = 0.6)]`: Equilibrium osmotic concentrations ($\text{mol} \cdot \text{L}^{-1}$) for leaf and sapwood cells. The difference between leaf and sapwood values helps maintaining phloem transport. The equilibrium sugar concentration is found using `equilibriumOsmoticConcentration - nonSugarConcentration` and defaults to `[= c(leaf = 0.55, sapwood = 0.35)]`.
- `minimumRelativeStarchForGrowth [= 0.50]`: Default minimum concentration of storage carbon (starch), relative to the maximum storage capacity, for sapwood growth to occur. This default value is used when species-specific values are not specified via `SpParams`.
- `senescenceRates [= c(sapwood = 0.0001261398, fineroot = 0.001897231)]`: Relative senescence rates (day^{-1}) for sapwood and fine roots. Default are equivalent to 4.5% and 50% annual turnover for sapwood and fine roots, respectively. Leaf senescence rates depend on leaf lifespan (i.e. leaf duration) for evergreen species. These are default values used when relative growth rates are not specified via species-specific `SpParams`.
- `constructionCosts [= c(leaf = 1.5, sapwood = 1.47, fineroot = 1.30)]`: Default construction costs, including respiration and structural carbon, per dry weight of new tissue ($g \text{ gluc} \cdot g \text{ dry}^{-1}$). These are default values used when relative growth rates are not specified via species-specific `SpParams`.
- `maximumRelativeGrowthRates [= c(leaf = 0.01, cambium = 0.005, sapwood = 0.002, fineroot = 0.1)]`: Default maximum relative growth rates for leaves ($\text{m}^2 \cdot \text{cm}^{-2} \cdot \text{day}^{-1}$), tree sapwood ($\text{cm}^2 \cdot \text{cm}^{-1} \cdot \text{day}^{-1}$), shrub sapwood ($\text{cm}^2 \cdot \text{cm}^{-2} \cdot \text{day}^{-1}$) and fine roots ($g \text{ dry} \cdot g \text{ dry}^{-1} \cdot \text{day}^{-1}$). These are default values used when relative growth rates are not specified via species-specific `SpParams`.
- `organogenesisEfficiencyModulator [= 1]`: Exponent relating turgor-related relative growth rate with leaf organogenesis efficiency.
- `elongationEfficiencyModulator [= 1]`: Exponent relating turgor-related relative growth rate with leaf elongation efficiency.

A set of parameters control defoliation and crown bud mortality:

- `defoliationParams`: A list with the following elements:
 - `cavitationInducedDefoliation [= TRUE]`: Whether leaf hydraulic impairment induces defoliation and bud senescence.
 - `criticalLeafPLC [= 0.50]`: Level of PLC (a proportion) corresponding to 50 % crown defoliation.
 - `criticalBudPLC [= 0.88]`: Level of PLC (a proportion) corresponding to 50 % crown bud senescence.
 - `cvLeafP50 [= 10.0]`: Coefficient of variation (in percent) of leaf P50 within crown.
 - `cvBudP50 [= 10.0]`: Coefficient of variation (in percent) of bud P50 within crown.

Finally, we have the following options regarding whole-plant mortality:

- **mortalityMode** [= "density/deterministic"]: String describing how mortality is applied. Current accepted values are combinations of "cohort" vs "density" (for whole-cohort mortality vs reduction of stem density) and "deterministic" vs. "stochastic".
- **mortalityBaselineRate** [= 0.0015]: Default deterministic proportion or probability specifying the baseline reduction of cohort's density occurring in a year as a result of processes not explicitly modelled (for mortalityMode = "density/deterministic" or "density/stochastic").
- **mortalityRelativeSugarThreshold** [= 0.4]: Threshold of stem sugar concentration relative to equilibrium values (by default 40%), resulting in an increased mortality rate/probability whenever levels are below.
- **mortalityRWCThreshold** [= 0.4]: Threshold of stem relative water content resulting in increased mortality rate/probability whenever levels are below (by default 40%).

15.5.4 Model output

15.5.4.1 Daily output

Functions **growth()** returns a list of a class with the same name. The first four elements of this list (i.e., **latitude**, **topography**, **weather** and **growthInput**) are simply copies of model inputs. The next element is **growthOutput**, which contains the state of the input object at the end of the simulation (this can be used to perform further simulations starting with current values of state variables). The remaining list elements are:

Element	Description
WaterBalance	Climatic input and water balance flows (i.e. net precipitation, infiltration, runoff, transpiration...). All of them in $mm = L \cdot m^{-2}$.
EnergyBalance	Energy balance components (instantaneous values) for the soil and the canopy
CarbonBalance	Stand-level carbon balance components (see (15.33)), all of them in $gC \cdot m^{-2}$.
BiomassBalance	Stand-level biomass balance components. All of them in $g dry \cdot m^{-2}$.
Temperature	Minimum and maximum temperature values for topsoil, canopy and atmosphere (if transpirationMode = "Sperry")
Soil	Soil variables for each soil layer: Moisture relative to field capacity (W_s), water potential (Ψ_s) and volumetric water content (V_s).
Stand	Stand-level variables, such as LAI_{stand}^{total} , LAI_{stand}^{act} , LAI_{stand}^{live} , LAI_{stand}^{dead} , the water retention capacity of the canopy (S_{canopy}) or the fraction of light reaching the ground (L_{ground}^{PAR} and L_{ground}^{SWR}).
Plants	Plant-level variables, such as LAI_{act} , LAI_{live} , phenological status, plant transpiration, plant water potential, etc.
SunlitLeaves	Leaf-level output variables for sunlit leaves (advanced model).
ShadeLeaves	Leaf-level output variables for shade leaves (advanced model).
LabileCarbonBalance	A list with the daily results of labile carbon balance components and carbon pools.
PlantBiomassBalance	A list with the daily results of biomass balance components for plant cohorts.
PlantStructure	A list with daily results of plant compartment sizes, in terms of area and biomass.
GrowthMortality	A list with daily results of growth rates for leaf, sapwood and fine roots; as well as daily mortality rates.
DecompositionPools	A list with daily values of carbon decomposition pools.

Elements **WaterBalance** to **ShadeLeaves** are equal to the output of function **spwb()** and were described in previous chapters (see 3.5.4 and 7.6.4).

The following table describes the child elements of **LabileCarbonBalance**, each of them a data frame with dates in rows and plant cohorts in columns:

Element	Symbol	Units	Description
GrossPhotosynthesis	A_g/B_i	$g\ gluc \cdot g\ dry^{-1} \cdot day^{-1}$	Daily gross photosynthesis per dry weight of living biomass.
MaintenanceRespiration	M_r/B_i	$g\ gluc \cdot g\ dry^{-1} \cdot day^{-1}$	Daily maintenance respiration (including leaves, sapwood and fine roots) per dry weight of living biomass.
GrowthCosts	G_i/B_i	$g\ gluc \cdot g\ dry^{-1} \cdot day^{-1}$	Daily growth costs (including leaves, sapwood and fine roots) per dry weight of living biomass.
RootExudation	RE_i	$g\ gluc \cdot g\ dry^{-1} \cdot day^{-1}$	Daily root exudation per dry weight of living biomass.
CarbonBalance	$\Delta S_i/B_i$	$g\ gluc \cdot g\ dry^{-1}$	Daily labile carbon balance (eq. (15.8)) per dry weight of living biomass.
SugarLeaf	$SS_{leaf,i}$	$mol\ gluc \cdot L^{-1}$	Sugar concentration in leaves.
StarchLeaf	$ST_{leaf,i}$	$mol\ gluc \cdot L^{-1}$	Starch concentration in leaves.
SugarSapwood	$SS_{sapwood,i}$	$mol\ gluc \cdot L^{-1}$	Sugar concentration in sapwood.
StarchSapwood	$ST_{sapwood,i}$	$mol\ gluc \cdot L^{-1}$	Starch concentration in sapwood.
SugarTransport	F_{phloem}	$mol\ gluc \cdot s^{-1}$	Average instantaneous rate of carbon transferred between leaves and stem compartments via phloem.

The following table describes the child elements of **PlantCarbonBalance**, each of them a data frame with dates in rows and plant cohorts in columns:

Element	Symbol	Units	Description
StructuralBiomassBalance		$g\ dry \cdot m^{-2} \cdot day^{-1}$	Daily structural biomass balance.
LabileBiomassBalance		$g\ dry \cdot m^{-2} \cdot day^{-1}$	Daily labile biomass balance.
PlantBiomassBalance		$g\ dry \cdot m^{-2} \cdot day^{-1}$	Daily plant biomass balance.
MortalityBiomassLoss		$g\ dry \cdot m^{-2} \cdot day^{-1}$	Daily biomass loss due to mortality.
CohortBiomassBalance		$g\ dry \cdot m^{-2} \cdot day^{-1}$	Daily cohort biomass balance (includes plant biomass balance and mortality).

The following table describes the child elements of **PlantStructure**, again each of them a data frame with dates in rows and plant cohorts in columns:

Element	Symbol	Units	Description
LeafBiomass	$B_{leaf,i}$	$g\ dry \cdot ind^{-1}$	Daily leaf structural biomass for an average individual.
SapwoodBiomass	$B_{sapwood,i}$	$g\ dry \cdot ind^{-1}$	Daily sapwood structural biomass for an average individual.
FineRootBiomass	$B_{fineroot,i}$	$g\ dry \cdot ind^{-1}$	Daily fine root biomass for an average individual.
LeafArea	LA_i	m^2	Daily amount of leaf area for an average individual.
CrownFoliageCompleteness		[0-1]	Leaf area expressed as a proportion of the maximum crown foliage for an average individual of each plant cohort.
SapwoodArea	SA_i	cm^2	Daily amount of sapwood area for an average individual.
FineRootArea	FRA_i	m^2	Daily amount of fine root area for an average individual.
HuberValue	H_v	$cm^2 \cdot m^{-2}$	The ratio of sapwood area to (target) leaf area
RootAreaLeafArea		$m^2 \cdot m^{-2}$	The ratio of fine root area to (target) leaf area
DBH	DBH_i	cm	Daily diameter at breast height for an average individual.
Height	H_i	cm	Daily total height for an average individual.

The following table describes the child elements of **GrowthMortality**, again each of them a data frame with dates in rows and plant cohorts in columns:

Element	Symbol	Units	Description
LAgrowth	$\Delta LA_i / SA_i$	$m^2 \cdot cm^{-2} \cdot day^{-1}$	Leaf area growth relative to sapwood area for an average individual.
SAgrowth	$\Delta SA_i / SA_i$	$cm^2 \cdot cm^{-2} \cdot day^{-1}$	Leaf area growth relative to sapwood area for an average individual.
FRAgrowth	$\Delta FRA_i / SA_i$	$m^2 \cdot cm^{-2} \cdot day^{-1}$	Fine root area growth relative to sapwood area for an average individual.
StarvationRate		$ind \cdot ind^{-1} \cdot day^{-1}$	Daily mortality rate from starvation.
DessicationRate		$ind \cdot ind^{-1} \cdot day^{-1}$	Daily mortality rate from dessication.
MortalityRate		$ind \cdot ind^{-1} \cdot day^{-1}$	Daily mortality rate (any cause).

15.5.4.2 Subdaily output

As with function `spwb()`, subdaily output can also be generated with function `growth()`. This is done via setting `subdailyResults = TRUE` in the control parameter list. In that case, the output list of `growth()` will include an element `subdaily`, whose subelements correspond to the results of `growth_day()` for each date simulated. Refer to the documentation of function `growth_day()` for details of sub-daily output.

Chapter 16

Plant carbon pools and components of the labile carbon balance

16.1 Size of plant carbon pools

Biomass of leaves, twigs, sapwood and fine roots is needed in the model to estimate respiratory costs and the size of the C storage pools.

16.1.1 Leaf structural, dead, metabolic and storage biomass

The live leaf area LA_{live} per individual (in m^2) is:

$$LA_{live} = 10000 \cdot LAI_{live} / N \quad (16.1)$$

where N is the density of the cohort and LAI_{live} is the leaf area index. The actual (expanded) leaf area LA_{act} is estimated analogously from LAI_{act} :

$$LA_{act} = 10000 \cdot LAI_{act} / N \quad (16.2)$$

And so is the dead leaf area, LA_{dead} :

$$LA_{dead} = 10000 \cdot LAI_{dead} / N \quad (16.3)$$

Structural leaf biomass per individual (B_{leaf} ; $g \text{ dry} \cdot ind^{-1}$) is the result of dividing leaf area by SLA ($m^2 \cdot kg^{-1}$), the specific leaf area of the species:

$$B_{leaf} = 1000 \cdot LA_{act} / SLA \quad (16.4)$$

where a factor 1000 is used to convert from kg to g. Hence, only expanded leaf area has respiratory cost (i.e. winter resistance buds do not). An analogous equation is used to estimate dead leaf biomass per individual (D_{leaf} ; $g \text{ dry} \cdot ind^{-1}$):

$$D_{leaf} = 1000 \cdot LA_{dead} / SLA \quad (16.5)$$

Leaf storage volume for an individual ($V_{storage,leaf}$; in $L \cdot ind^{-1}$) is:

$$V_{storage,leaf} = LA_{act} \cdot V_{leaf} \quad (16.6)$$

where V_{leaf} is the water storage capacity of leaf tissue per leaf area unit (in $L \cdot m^{-2}$) (see A.4.5). Leaf storage volume is necessary to estimate the biomass of metabolic (sugars) or storage (starch) from the concentration

of these substances in leaves (SS_{leaf} and ST_{leaf} , which are expressed in $mol\ glucose \cdot L^{-1}$), and conversely. For example:

$$B_{sugar,leaf} = SS_{leaf} \cdot V_{storage,leaf} \cdot m_{gluc} \quad (16.7)$$

where m_{gluc} is the molar mass of glucose. To estimate leaf starch storage capacity, the model assumes that up to 10% of leaf tissue volume can be occupied with starch, so that the leaf starch storage capacity (ST_{leaf}^{max} ; in $mol \cdot ind^{-1}$) is:

$$ST_{leaf}^{max} = 0.1 \cdot 1000 \cdot \frac{V_{storage,leaf} \cdot \rho_{starch}}{m_{starch}} \quad (16.8)$$

where ρ_{starch} is the density of starch and m_{starch} is its molar mass.

16.1.2 Twig structural and dead biomass

Structural twig biomass per individual (B_{twig} ; $g\ dry \cdot ind^{-1}$) is related to the biomass of leaves (B_{leaf}), via the $r_{6.35}$ ratio, which is defined as the ratio between the biomass of leaves and twigs over the biomass of leaves:

$$B_{twig} = B_{leaf} / (r_{6.35} - 1) \quad (16.9)$$

An analogous equation is used to estimate dead twig mass per individual (D_{twig} ; $g\ dry \cdot ind^{-1}$) from the mass of dead leaves (D_{leaf}):

$$D_{twig} = D_{leaf} / (r_{6.35} - 1) \quad (16.10)$$

While B_{twig} represents the structural twig biomass, it cannot be used to estimate twig respiration (see living sapwood biomass below). In addition, we assume that embolized parts of twigs do not contribute to twig respiration. Hence, twig living (respiratory) biomass is estimated using:

$$B_{living,twig} = B_{twig} \cdot (1 - PLC_{stem}) \cdot (1 - f_{conduits}) \quad (16.11)$$

16.1.3 Sapwood structural, metabolic and storage biomass

Sapwood volume ($V_{sapwood}$; in $L \cdot ind^{-1}$) is defined as the product of sapwood area and the sum of aboveground height and belowground coarse root length:

$$V_{sapwood} = 1000 \cdot SA \cdot (H + \sum_s FRP_s \cdot L_s) \quad (16.12)$$

where SA is sapwood area, H is plant height, FRP_s is the proportion of fine roots in soil layer s and L_s is the coarse root length in layer s (all lengths expressed here in m). Sapwood structural biomass per individual ($B_{sapwood}$; $g\ dry \cdot ind^{-1}$) represents the sapwood biomass of sapwood of trunks, branches and coarse roots. It is defined as the product of $V_{sapwood}$ and wood density (ρ_{wood} ; $g\ dry \cdot cm^{-3}$):

$$B_{sapwood} = 1000 \cdot V_{sapwood} \cdot \rho_{wood} \quad (16.13)$$

While $B_{sapwood}$ represents the structural sapwood biomass, it cannot be used to estimate sapwood respiration, since only xylem axial or radial parenchymatic rays (and not dead cells like tracheids or vessels) contribute to sapwood respiration. In addition, we assume that embolized parts of sapwood do not contribute to sapwood respiration. Hence, sapwood living (respiratory) biomass is estimated using:

$$B_{living,sapwood} = B_{sapwood} \cdot (1 - PLC_{stem}) \cdot (1 - f_{conduits}) \quad (16.14)$$

where PLC_{stem} is the current proportion of stem conductance lost and $f_{conduits}$ is the fraction of sapwood volume that corresponds to dead conduits, being the complement of the fraction of parenchymatic tissue. The volume available for metabolic or storage carbon within sapwood is limited by both the sapwood fraction that corresponds to parenchyma and wood porosity (i.e. $(1 - (\rho_{wood}/1.54))$). Sapwood storage volume ($V_{storage,sapwood}$; in $L \cdot ind^{-1}$) is:

$$V_{storage,sapwood} = V_{sapwood} \cdot (1 - (\rho_{wood}/1.54)) \cdot (1 - f_{conduits}) \quad (16.15)$$

Analogously to leaf carbon, $V_{storage,sapwood}$ is used to estimate the biomass of metabolic (sugars) or storage (starch) from the concentration of these substances in sapwood ($SS_{sapwood}$ and $ST_{sapwood}$, which are expressed in $mol\ gluc \cdot L^{-1}$), and conversely. For example:

$$B_{sugar,sapwood} = SS_{sapwood} \cdot V_{storage,sapwood} \cdot m_{gluc} \quad (16.16)$$

where m_{gluc} is the molar mass of glucose. To estimate sapwood storage capacity, the model assumes that up to 50% of volume of parenchymatic cells can be occupied with starch, so that the sapwood starch storage capacity ($ST_{sapwood}^{max}$; in $mol \cdot ind^{-1}$) is:

$$ST_{sapwood}^{max} = 0.5 \cdot 1000 \cdot \frac{V_{storage,sapwood} \cdot \rho_{starch}}{m_{starch}} \quad (16.17)$$

16.1.4 Heartwood biomass

Total wood volume (V_{wood} ; in $L \cdot ind^{-1}$) is defined, analogously as sapwood volume, as the product of stem section area and the sum of aboveground height and belowground coarse root length:

$$V_{wood} = 1000 \cdot SSA \cdot (H + \sum_s FRP_s \cdot L_s) \quad (16.18)$$

where SSA is stem section area (in cm^2 ; derived from DBH), H is plant height, FRP_s is the proportion of fine roots in soil layer s and L_s is the coarse root length in layer s (all lengths expressed here in m). Once total wood volume and sapwood volume are known, we can determine heartwood volume as their difference:

$$V_{heartwood} = V_{wood} - V_{sapwood} \quad (16.19)$$

Heartwood biomass per individual ($D_{heartwood}$; $g\ dry \cdot ind^{-1}$) represents the heartwood mass of trunks, branches and coarse roots. Analogously to sapwood biomass, is defined as the product of $V_{heartwood}$ and wood density (ρ_{wood} ; $g\ dry \cdot cm^{-3}$):

$$D_{heartwood} = 1000 \cdot V_{heartwood} \cdot \rho_{wood} \quad (16.20)$$

16.1.5 Fine root area and biomass

The fine root compartment does not have labile carbon (i.e. the model supplies labile carbon for fine roots from sapwood compartment), so that only the structural biomass of fine roots needs to be estimated. If growth is simulated using the *basic* water balance model, the initial area of fine roots (FRA , m^2) for layer s is estimated from leaf area using:

$$FRA = FRP_s \cdot LA_{act} \cdot RLR \quad (16.21)$$

where FRP_s is the proportion of fine roots in layer s and RLR is the species-specific root area to leaf area ratio. Biomass of fine roots per individual for a given layer s ($B_{fineroot,s}$; $g\ dry \cdot ind^{-1}$) is then estimated from FRA_s using:

$$B_{fineroot,s} = \frac{10^4 \cdot FRA_s}{2.0 \cdot \sqrt{\frac{\pi \cdot SRL}{\rho_{fineroot}}}} \quad (16.22)$$

where $\rho_{fineroot}$ is fine root tissue density ($g\ dry \cdot cm^{-3}$) and SRL ($cm \cdot g\ dry^{-1}$) is the specific root length. Fine root biomass per individual ($g\ dry \cdot ind^{-1}$) is simply the sum across layers:

$$B_{fineroot} = \sum_s B_{fineroot,s} \quad (16.23)$$

If growth is simulated using the *advanced* water balance model, the initial fine root length per area in each soil layer (FLA_s ; $m \cdot m^{-2}$) is estimated from rhizosphere conductance, assuming a cylindrical flow geometry in the rhizosphere. The initial fine root biomass for a given layer s ($g\ dry \cdot ind^{-1}$) is then calculated using:

$$B_{fineroot,s} = \frac{10000 \cdot FLA_s}{N \cdot 0.01 \cdot SRL} \quad (16.24)$$

where N is the density of individuals. Finally, initial fine root area in each soil layer (FRA_s) is then calculated from $B_{fineroot}$ by inverting the equation presented above:

$$FRA_s = 10^{-4} \cdot B_{fineroot,s} \cdot 2.0 \cdot \sqrt{\frac{\pi \cdot SRL}{\rho_{fineroot}}} \quad (16.25)$$

16.1.6 Total living biomass

Total living biomass per individual (i.e. not accounting for heartwood or xylem conduits) is:

$$B_{total} = B_{leaf} + B_{living,twig} + B_{living,sapwood} + B_{fineroot} + \\ + B_{sugar,leaf} + B_{starch,leaf} + B_{sugar,sapwood} + B_{starch,sapwood}$$

B_{total} is used to express carbon balance components per dry weight of living biomass in the model output, which allows a better comparison across plant cohorts of different size and species.

16.2 Components of labile carbon balance

Here we provide the details of labile carbon balance calculations, i.e. how the components of eqs. (15.2) and (15.6) are determined.

16.2.1 Gross photosynthesis

The soil water balance submodel provides values of photosynthesis calculated at the plant cohort level, but note that the user output of function `spwb()` refers to net photosynthesis (A_n) after accounting for an estimate of leaf respiration. In the basic water balance model there is no way to make the distinction between gross and net photosynthesis, so we will assume here that $A_g = A_n$ (in $gC \cdot m^{-2}$) as calculated in section 6.1.7. In the advance water balance model the output is also net photosynthesis, but the distinction is possible (see section 11.1). In any case, here we start with a A_g value in units of carbon per ground area unit, i.e. $gC \cdot m^{-2}$, and to enter the labile carbon balance we need scale photosynthesis to units of glucose per individual (A ; in $ggluc \cdot ind^{-1}$). This is done using:

$$A = \frac{10000 \cdot A_g}{N} \cdot \frac{m_{gluc}}{m_{carbon} \cdot 6} \quad (16.26)$$

where N is the density of individuals per hectare, m_{gluc} is the glucose molar mass and m_{carbon} is the carbon molar mass.

16.2.2 Maintenance respiration

Maintenance respiration (in $ggluc \cdot ind^{-1}$) is calculated for each living compartment (leaves, twigs, sapwood, and fine roots) individually (Mouillot et al., 2001), and differs slightly depending on the complexity of the transpiration submodel.

When simulating growth with the basic water balance, the model uses a Q_{10} relationship of baseline respiration with temperature, which means that for every 10 °C change in temperature there is a Q_{10} factor change in respiration:

$$MR_{leaf} = (B_{leaf} + B_{sugar,leaf}) \cdot MRR_{leaf} \cdot Q_{10}^{(T_{mean}-20)/10} \cdot (L^{PAR})^{WUE_{decay}} \quad (16.27)$$

$$MR_{twig} = B_{living,twig} \cdot MRR_{sapwood} \cdot Q_{10}^{(T_{mean}-20)/10} \quad (16.28)$$

$$MR_{sapwood} = (B_{living,sapwood} + B_{sugar,sapwood}) \cdot MRR_{sapwood} \cdot Q_{10}^{(T_{mean}-20)/10} \quad (16.29)$$

$$MR_{fineroot} = B_{fineroot} \cdot MRR_{fineroot} \cdot Q_{10}^{(T_{mean}-20)/10} \cdot (LA_{act}/LA_{live}) \quad (16.30)$$

where T_{mean} is the average daily temperature (in °C). Baseline maintenance respiration rates per dry biomass (MRR_{leaf} , $MRR_{sapwood}$ and $MRR_{fineroot}$ for leaves, parenchymatic sapwood and fine roots, respectively; in $g\text{ gluc} \cdot g\text{ dry}^{-1} \cdot \text{day}^{-1}$) are can be supplied via species-specific parameters and should refer to 20 °C. Leaf and fine root rates can also be estimated from nitrogen concentration, following Reich et al. (2008) (see section A.3.19). Factor Q_{10} has been found to decrease with temperature (Tjoelker et al., 2001). For example, a 10 °C increase in measurement temperature at low measurement temperatures from 5 to 15 °C results in an approximate 2.8 fold increase in respiration rate, whereas an increase in temperature from 25 to 35 °C results in a less than two-fold (1.8) increase in rate. The general temperature relationship proposed by Tjoelker et al. (2001) is used here:

$$Q_{10} = 3.22 - 0.046 \cdot T_{mean} \quad (16.31)$$

Factor $(L^{PAR})^{WUE_{decay}}$ is added to reduce respiration rates in leaves under shade, analogously to the reduction of water use efficiency explained in 6.1.7. Thus, it is assumed that the carbon use efficiency of the leaves is more or less constant. Factor LA_{act}/LA_{live} is added to the maintenance respiration of fine roots to reduce respiration rates during decreased leaf periods, assuming a decrease in respiration rates of fine roots parallel to that of leaves.

When simulating growth with the advanced model, sub-daily leaf maintenance respiration rates are estimated as the difference between gross and net photosynthesis (see 11.1.4). Sapwood and fine root respiration rates are also estimated at sub-daily steps and, hence, sub-daily canopy temperature variation is taken into account (i.e. sub-daily canopy T_{can} values are used instead of the daily temperature mean, T_{mean}).

16.2.3 Growth costs and synthesis respiration

Construction costs per unit dry weight of new tissue (CC_{leaf} , $CC_{sapwood}$ and $CC_{fineroot}$ for leaves, sapwood and fine roots; in $g\text{ gluc} \cdot g\text{ dry}^{-1}$) are specified as control parameters (i.e. they are not species-specific). These unitary costs include both the carbon used in respiration during growth and structural carbon. Given an increase in leaf area ΔLA , leaf carbon growth costs (in $g\text{ gluc} \cdot \text{ind}^{-1}$) are:

$$GC_{leaf} = \Delta LA \cdot CC_{leaf} \cdot 1000/SLA \quad (16.32)$$

whereas the growth costs for the corresponding twigs are:

$$GC_{twig} = \Delta LA \cdot CC_{sapwood} \cdot (r_{6.35} - 1) \cdot 1000/SLA \quad (16.33)$$

Analogously, given an increase in sapwood area ΔSA , sapwood growth costs (in $g\text{ gluc} \cdot \text{ind}^{-1}$) are:

$$GC_{sapwood} = \Delta SA \cdot CC_{sapwood} \cdot (H + \sum_s FRP_s \cdot L_s) \cdot \rho_{wood} \quad (16.34)$$

Finally, and given an increase in fine root biomass $\Delta B_{fineroot}$ has the following growth costs (in $g\text{ gluc} \cdot \text{ind}^{-1}$) are:

$$GC_{fineroot} = \Delta B_{fineroot} \cdot CC_{fineroot} \quad (16.35)$$

Synthesis respiration (in $g\text{ gluc} \cdot \text{ind}^{-1}$) associated to growth is less than growth costs, but can be found easily from growth costs, using construction costs per unit dry mass:

$$SR_{leaf} = GC_{leaf} \cdot \frac{CC_{leaf} - 1}{CC_{leaf}} \quad (16.36)$$

$$SR_{twig} = GC_{twig} \cdot \frac{CC_{sapwood} - 1}{CC_{sapwood}} \quad (16.37)$$

$$SR_{sapwood} = GC_{sapwood} \cdot \frac{CC_{sapwood} - 1}{CC_{sapwood}} \quad (16.38)$$

$$SR_{fineroot} = GC_{fineroot} \cdot \frac{CC_{fineroot} - 1}{CC_{fineroot}} \quad (16.39)$$

The plant-level total synthesis respiration (required for stand-level carbon balance) is $SR = SR_{leaf} + SR_{twig} + SR_{sapwood} + SR_{fineroot}$.

16.2.4 Phloem transport

When growth is simulated using the basic water balance model, phloem transport between leaf and sapwood (F_{phloem}) is simply modelled as the flow needed to make the concentration of metabolic carbon equal in both compartments. When the advanced water balance submodel is used, F_{phloem} is modelled following Hölttä et al. (2017). Specifically, the instantaneous phloem flow per leaf area basis ($mol\ gluc \cdot m^{-2} \cdot s^{-1}$) depends on k_{phloem} , i.e. the phloem conductance per leaf area basis ($L \cdot m^{-2} \cdot MPa^{-1} \cdot s^{-1}$), sap viscosity relative to water and the difference in turgor between the sieve cells of leaf and sapwood compartments. k_{phloem} is estimated as linear function of stem maximum conductance, $k_{stem,max}$:

$$k_{phloem} = f_{phloem} \cdot k_{stem,max} \quad (16.40)$$

where factor f_{phloem} is specified by the control parameter `phloemConductanceFactor`. Sap viscosity is calculated following Först et al. (2002), which takes into account temperature and the average sugar concentration between the two compartments (no phloem flow can occur if temperature is below zero). Turgor in sieve cells of each compartment depends on symplastic water potential and osmotic water potential, with Van't Hoff's equation being used to calculate the osmotic water potential, based on temperature, sugar concentration and the concentration of other solutes (control parameter `nonSugarConcentration`). Note that here leaf or sapwood sugar concentration is modulated to account for the effect of the relative water content of the compartment on osmotic water potential.

16.2.5 Sugar-starch dynamics

When growth is simulated using the *basic* water balance submodel, sugar-starch dynamics (SC_{leaf} and $SC_{sapwood}$) are simply defined as the conversion between metabolic and storage carbon needed to keep the sugar concentration equal to an equilibrium required for metabolic functioning, which is specified by control parameters `equilibriumOsmoticConcentration` (including two values, for leaves and sapwood respectively) and `nonSugarConcentration`. For example, if leaf osmotic equilibrium concentration is $0.8\ mol\ gluc \cdot L^{-1}$ and non-sugar concentration is $0.3\ mol\ gluc \cdot L^{-1}$, the model will simulate sugar-starch dynamics to stabilize sugar concentration in leaves to $0.5\ mol\ gluc \cdot L^{-1}$. A lower equilibrium osmotic concentration is required for sapwood, by default $0.6\ mol\ gluc \cdot L^{-1}$, so that sugar equilibrium concentration is $0.1\ mol\ gluc \cdot L^{-1}$. That lower equilibrium concentrations are required for sapwood than for leaves is necessary to sustain the phloem transport of sugars from leaves towards sapwood.

When growth is simulated using the *advanced* water balance submodel, sugar-starch dynamics are similar to the previous case, in the sense that equilibrium sugar concentrations are sought for leaves and sapwood. However, instantaneous conversion rates are calculated depending on equations regulating starch synthesis (occurring when sugar concentration is higher than the equilibrium concentration) and starch hydrolysis (occurring when sugar concentration is lower than the equilibrium concentration). Synthesis is modelled as a Michaelis-Menten function, whereas hydrolysis rate is simply a linear function of starch concentration. Maximum synthesis rates and hydrolysis linear factors are larger (i.e. faster sugar-starch dynamics) in leaves than sapwood.

16.2.6 Root exudation

Whereas surplus in leaf storage capacity is relocated into sapwood storage, surplus in sapwood storage carbon is diverted towards root exudation ($RE_{sapwood}$). Hence, root exudation is not a process competing for metabolic carbon, but only a consequence of plant C storage capacity being surpassed (Prescott et al., 2020). Root exudation may happen in the model when growth is restricted due to sink limitations but photosynthesis continues and storage carbon levels have attained the storage capacity in the sapwood compartment ($ST_{sapwood} > ST_{sapwood}^{max}$).

Chapter 17

Growth, senescence and mortality

17.1 Growth

17.1.1 Temperature and turgor sink limitations

Sink limitations due to temperature and turgor effects on growth are modelled following Cabon et al. (2020a) and Cabon et al. (2020b). These authors suggested equations to model sink limitations on cambium cell division and tracheid expansion, but we apply the same approach for simulating growth of leaves, sapwood and fine roots. Cell relative expansion rate (r_{cell}) is central to the approach by Cabon et al. and is defined as the relative time derivative of cell volume:

$$r_{cell} = \frac{dV}{Vdt} \quad (17.1)$$

Cabon et al. (2020a) first suggested to model the dependence of r_{cell} on cell turgor using Lockhart's equation:

$$r_{cell}(\Psi, \pi_0) = \xi_{max} \cdot (\Psi - \pi_0 - Y_P) \quad (17.2)$$

where Ψ is the water potential, π_0 is the osmotic water potential at full turgor, Y_P is the turgor yield threshold and ξ_{max} is the maximum cell wall extensibility. Later, Cabon et al. (2020b) suggested to account for both turgor and temperature limitations on r_{cell} using the following expanded equation:

$$r_{cell}(T, \Psi, \pi_0) = \xi_{max} \cdot (\Psi - \pi_0 - Y_P) \cdot \frac{f_{met}(T_K)}{f_{met}(288.15)} \cdot f_{micro}(T, T_{thr}) \quad (17.3)$$

where T is temperature, $f_{met}(T_K)$ is a function modulating the effect of temperature T_K in Kelvin, on metabolic rate, and $f_{micro}(T, T_{thr})$ is a sigmoidal function modulating the effect of temperature on microtubule stability, depending on a temperature threshold T_{thr} . Function $f_{met}(T_K)$ is defined as:

$$f_{met}(T_K) = \frac{T_K \cdot \exp\left\{\frac{\Delta H_A}{R^n \cdot T_K}\right\}}{1 + \exp\left\{\frac{\Delta S_D}{R^n} \cdot \left(1 - \frac{\Delta H_D}{\Delta S_D \cdot T_K}\right)\right\}} \quad (17.4)$$

where R^n is the ideal gas constant, $\Delta H_A = 87500$ the enthalpy of activation and $\Delta H_D = 333000$ and $\Delta S_D = 1090$ the enthalpy and entropy difference (respectively) between the catalytically active and inactive states of the enzymatic system).

In **medfate**, we assume a constant value of $\pi_0 = -1$ in meristematic tissues, so that r_{cell} depends on Ψ and T only. Scaling from the cell to the tissue level is conducted by assuming that maximum cell-level expansion rates correspond to maximum tissue-level relative growth rates. Maximum cell-level expansion rates, $r_{cell,max}$, are assumed to occur at $T = 30^\circ C$ and $\Psi = 0$. Other parameters for eq. (17.3) are set to $\xi_{max} = 0.5$ (which becomes irrelevant), $Y_P = 0.05 MPa$ and $T_{thr} = 5^\circ C$.

17.1.2 Persistent bud formation

Persistent bud formation is a process required for species whose primary growth determinacy is either determined or intermediate (see 15.1.3). The process is scheduled to start at the end of the leaf growth period, normally between summer and autumn, and its duration is controlled by the species-specific parameter BFD ($BudFormationDays$). Bud formation involves leaf organogenesis and budset formation, and sets the leaf area preformed in persistent buds, $LA_{preformed}$ (m^2):

$$LA_{preformed} = 1.1 \cdot LOE \cdot \frac{LA_{live}}{LD} = 1.1 \cdot LOE \cdot \frac{LA_{target} \cdot (PCAB/100)}{LD} \quad (17.5)$$

Whereas LA_{live}/LD sets the target leaf area to overcome the annual fraction of leaves that are expected to undergo senescence, the 1.1 factor increases this target by 10% to account for expected sink limitations to growth. Finally, LOE is the leaf organogenesis efficiency, which is estimated by averaging turgor limitations over the bud formation period:

$$LOE = \frac{\sum_i^{BFD} \left(\frac{r_{cell}(30, \Psi_i)}{r_{cell,max}} \right)^{m_{LOE}}}{BFD} \quad (17.6)$$

Note from the previous equation that only daily water potential limitations (i.e. Ψ_i) to organogenesis are considered (i.e. temperature is assumed to be optimal). Here m_{LOE} is an exponent (control parameter `organogenesisEfficiencyModulator`) modulating the effect of turgor limitations on organogenesis efficiency.

17.1.3 Leaf growth

Leaf area increment ΔLA only occurs when phenological state is *unfolding*, and is defined as the minimum of three values expressing three corresponding constraints:

$$\Delta LA = \min(\Delta LA_{pheno}, \Delta LA_{source}, \Delta LA_{sink}) \quad (17.7)$$

Target leaf area increase

ΔLA_{pheno} is the increase in leaf area that would be expected by leaf growth targets and crown bud availability. In plants with indetermined primary growth, or with intermediate growth after finishing the elongation of preformed leaves (see 15.1.3), organogenesis is not separated from leaf expansion and ΔLA_{pheno} is directly leaf area increment allowed by the live leaf area (LA_{live} ; see 17.3.3), minus the current expanded leaf area (LA_{act}):

$$\Delta LA_{pheno} = \max(LA_{live} - LA_{act}, 0) = \max(LA_{target} \cdot (PCAB/100) - LA_{act}, 0) \quad (17.8)$$

In contrast, in plants with determined primary growth, or during elongation of preformed leaves in plants with intermediate growth, the leaf are target is dictated by preformed leaves within buds. If there is leaf area to be elongated (i.e. $LA_{preformed} > 0.0$) then ΔLA_{target} is simply equal to the preformed (in buds) leaf area that is to be elongated according to the leaf unfolding model (see 4.1.3):

$$\Delta LA_{pheno} = LA_{preformed} \cdot (\phi_t - \phi_{t-1}) \quad (17.9)$$

where ϕ_t and ϕ_{t-1} are the leaf elongation status at the current day and the previous day, respectively. When $LA_{preformed} = 0.0$ then ΔLA_{pheno} is set as detailed above for indetermined primary growth.

Carbon source limitations

ΔLA_{source} represents the leaf area increment expected given carbon source limitations and is calculated:

$$\Delta LA_{source} = \frac{ST_{sapwood} \cdot m_{gluc} \cdot V_{sapwood,leaf}}{1000 \cdot CC_{leaf}/SLA} \quad (17.10)$$

where $ST_{sapwood}$ is the current concentration of sapwood storage carbon (starch), $V_{storage,sapwood}$ is the sapwood storage volume and m_{gluc} is glucose molar mass and the denominator contains the construction costs per leaf area unit, see eq. (16.32).

Sink limitations

ΔLA_{sink} represents the leaf area increment resulting from sink limitations. If leaf growth is occurring via leaf elongation of preformed leaves, only turgor limitations are considered, because temperature effects are already taken into account in the phenological model of leaf elongation:

$$\Delta LA_{sink} = \Delta LA_{pheno} \cdot \left(\frac{r_{cell}(30, \Psi_i)}{r_{cell,max}} \right)^{m_{LEE}} \quad (17.11)$$

where m_{LEE} is an exponent (control parameter `elongationEfficiencyModulator`) modulating the effect of turgor limitations on leaf elongation efficiency.

For indetermined leaf growth, both temperature and turgor limitations are considered; and ΔLA_{sink} is estimated using the maximum leaf tissue growth rate relative to sapwood area ($RGR_{leaf,max}$; in $m^2 \cdot cm^{-2} \cdot day^{-1}$), the percentage of the crown with active buds ($PCAB$) and the relative cell expansion rate given T_{day} and $\Psi_{symp,leaf}$:

$$\Delta LA_{sink} = SA \cdot (PCAB/100) \cdot RGR_{leaf,max} \cdot \frac{r_{cell}(T_{day}, \Psi_{symp,leaf})}{r_{cell,max}} \quad (17.12)$$

where cell relative expansion rate is divided by the maximum relative cell expansion rate $r_{cell,max}$, so that $RGR_{leaf,max}$ is attained when sink conditions are optimal. The final expression of ΔLA_{sink} reduces to a product of the maximum leaf area growth times two factors (for turgor and temperature) bounded between 0 and 1.

Update of crown active buds

If previous drought or fire impacts had reduced the percentage of the crown with active buds ($PCAB$), this percentage is increased along with primary growth (i.e. with leaf growth), until it achieves 100%:

$$PCAB_{t+1} = \min(PCAB_t + RGR_{bud} \cdot \frac{\Delta LA}{LA_{target}}, 100) \quad (17.13)$$

where RGR_{bud} is the growth rate of buds per unit per unit leaf area growth ($m^2 \cdot m^{-2}$).

17.1.4 Sapwood growth

Newly assimilated carbon is preferentially allocated to leaves and fine roots whenever storage levels are low. ΔSA is defined as the minimum of three values expressing targets, source and sink constraints:

$$\Delta SA = \min(\Delta SA_{pheno}, \Delta SA_{source}, \Delta SA_{sink}) \quad (17.14)$$

Sapwood area target

ΔSA_{pheno} is the increase in sapwood area that would be expected comparing the sapwood target with the current functional sapwood area:

$$\Delta SA_{pheno} = \max(SA_{target} - SA \cdot (1.0 - PLC_{stem})) \quad (17.15)$$

where SA_{target} is the target sapwood area dictated by functional coordination (see 17.3.3) and PLC_{stem} is the proportion of stem conductance loss due to cavitation.

Carbon source limitations

ΔSA_{source} represents the sapwood area increment expected given carbon source limitations and is calculated using:

$$\Delta SA_{source} = \frac{\max(ST_{sapwood} - ST_{sapwood,growth}, 0) \cdot m_{gluc} \cdot V_{storage,sapwood}}{CC_{sapwood} \cdot (H + \sum_s FRP_s \cdot L_s) \cdot \rho_{wood}} \quad (17.16)$$

where $ST_{sapwood}$ is the current starch concentration, $ST_{sapwood,growth}$ is the minimum starch concentration required for sapwood growth, $V_{storage,sapwood}$ is the sapwood storage volume, m_{gluc} is glucose molar mass and the denominator contains the construction costs per sapwood area unit, see eq. (16.34). $ST_{sapwood,growth}$ is related to the minimum relative starch concentration for growth ($RSSG$), which is specified via the species-specific parameter `RSSG` or the control parameter `minimumRelativeStarchForGrowth`. This parameter is important because it allows specifying to which extent a given plant species stops growth and saves carbon to ensure survival (e.g. shadow tolerant species), as opposed to a species strongly investing in growth to reach the top of the canopy and have access to high light levels (e.g. light-demanding species).

Sink limitations

Assuming that the maximum relative sapwood growth rate ($RGR_{cambium,max}$ or $RGR_{sapwood,max}$) corresponds to a maximum rate of daily ring area increase, we have that the daily increase in sapwood area according to sink limitations, ΔSA_{sink} , is defined as:

$$\Delta SA_{sink} = \pi \cdot DBH \cdot (PCAC/100) \cdot RGR_{cambium,max} \cdot \frac{r_{cell}(T_{day}, \Psi_{symp,stem})}{r_{cell,max}} \quad (17.17)$$

for trees, and as:

$$\Delta SA_{sink} = SA \cdot (PCAC/100) \cdot RGR_{sapwood,max} \cdot \frac{r_{cell}(T_{day}, \Psi_{symp,stem})}{r_{cell,max}} \quad (17.18)$$

for shrubs. Sapwood growth in trees is proportional to the cambium length (in cm), whereas in shrubs it is proportional to the current sapwood area (in cm^2). Hence, two different maximum relative growth rates are defined for trees ($RGR_{cambium,max}$) and shrubs ($RGR_{sapwood,max}$). In the above equations, $PCAC$ is the percentage of active cambium cells, which is assumed to be controlled by the amount of foliage via the synthesis of growth hormones that then are transported through phloem. In practice, $PCAC$ is estimated as:

$$PCAC = 100 \cdot \frac{LAI_{act}}{LAI_{NC}} \quad (17.19)$$

where LAI_{act} is the currently expanded leaf area index and LAI_{NC} is the leaf area index assuming no light competition from taller plants. Since LAI_{act} depends on the phenological status for some species, the sapwood growth rate will depend on leaf phenology. Moreover, since $LAI_{act} \leq LAI_{live} \leq LAI_{NC}$, maximum growth rates will be achieved by cohorts having fully developed crowns and lacking competition of taller cohorts and low self-competition. Since competition increases with forest stand development, the inclusion of $PCAC$ ensures that sapwood area growth rates will decrease for forest stands of larger basal area.

17.1.5 Fine root growth

Fine root biomass increment is modelled for each soil layer separately, and is defined analogously to leaf area increment:

$$\Delta B_{fineroot} = \min(\Delta B_{fineroot,pheno}, \Delta B_{fineroot,source}, \Delta B_{fineroot,sink}) \quad (17.20)$$

Fineroot biomass target

$\Delta B_{fineroot,pheno}$ is the maximum fine root biomass increment allowed by the biomass target set by the allocation rule, $B_{fineroot,target}$ (see 17.3.3), in comparison with current biomass, $B_{fineroot}$:

$$\Delta B_{fineroot,pheno} = \max(B_{fineroot,target} - B_{fineroot}, 0) \quad (17.21)$$

Carbon source limitations

$\Delta B_{fineroot,source}$ represents the biomass increment expected given the available storage carbon (starch):

$$\Delta B_{fineroot,source} = \frac{ST_{sapwood} \cdot m_{gluc} \cdot V_{storage,sapwood}}{CC_{fineroot}} \quad (17.22)$$

where $ST_{sapwood}$ is the current sapwood concentration of storage carbon, $V_{storage,sapwood}$ is the sapwood storage volume, m_{gluc} is glucose molar mass and $CC_{fineroot}$ is the construction costs per fine root biomass unit.

Sink limitations

$\Delta B_{fineroot,sink}$ represents the biomass increment expected by taking into account maximum tissue growth rate ($RGR_{fineroot,max}$; in $g\ dry^{-1} \cdot day^{-1}$) and the relative cell expansion rate given temperature (T_{day}) and water potential in the rhizosphere ($\Psi_{rhizo,s}$):

$$\Delta B_{fineroot,sink} = B_{fineroot} \cdot RGR_{fineroot,max} \cdot \frac{r_{cell}(T_{day}, \Psi_{rhizo,s})}{r_{cell,max}} \quad (17.23)$$

cell relative expansion rate is divided by the maximum relative cell expansion rate so that $RGR_{fineroot,max}$ is attained when sink conditions are optimal.

17.2 Senescence

17.2.1 Leaf aging and cavitation effects on leaves and buds

Leaf senescence can occur due to two processes: programmed senescence (aging) or hydraulic disconnection related to cavitation. Each day, the model estimates the proportion of leaves that can be lost due to each process, i.e. $p_{aging,leaf}$ and $p_{cavitation,leaf}$, and the maximum of the two values is finally applied to reduce LA_{act} and increase LA_{dead} correspondingly.

Programmed leaf senescence

In evergreen species whose senescence period is "whole-year", the proportion of leaf area that undergoes programmed senescence each day is determined by the species-specific leaf duration parameter (LD):

$$p_{aging,leaf} = \frac{1}{365.25 \cdot LD} \quad (17.24)$$

For the remaining species (i.e. with "autumn" or "spring" senescence), a target leaf area for programmed senescence ($LA_{progsen}$) is set at the beginning of the senescence period and then leaves progressively die during the senescence period. First, a maximum target leaf area for senescence ($LA_{progsen,max}$) is estimated. In the case of winter (semi-)deciduous species $LA_{progsen,max} = LA_{act}$, whereas for evergreens it is set as $LA_{progsen,max} = LA_{act}/LD$. The actual target leaf area for programmed senescence ($LA_{progsen}$) can be lower if the crown is already defoliated:

$$LA_{progsen} = \max(0.0, LA_{progsen,max} - (LA_{live} - LA_{act})) \quad (17.25)$$

This has the advantage of avoiding conducting programmed senescence on crowns that have already suffered from drought-induced defoliation (i.e. where $LA_{act} - LA_{live}$) at the start of the senescence period.

During each day of the leaf senescence period, the proportion of leaf senescence is set as:

$$p_{aging,leaf} = \frac{LA_{progsen} \cdot (\phi_{t-1} - \phi_t)}{LA_{act}} \quad (17.26)$$

for "autumn" senescence, and:

$$p_{aging,leaf} = \frac{LA_{progsen} \cdot (\phi_t - \phi_{t-1})}{LA_{act}} \quad (17.27)$$

for "spring" senescence.

Drought-induced senescence

Leaf senescence can also occur as a consequence of hydraulic disconnection, as described in 6.2.3 and 14.5, for the basic or advanced water balance models, respectively.

If the proportion of leaf conductance loss (i.e. PLC_{leaf}) has increased with respect to the preceeding day, the model determines $LA_{cavitation}$, the maximum leaf area allowed according to the current level of leaf cavitation, by estimating the cavitation-induced defoliation proportion and applying it to the maximum leaf area (LA_{target}). Drought-induced leaf senescence effects are modulated with control parameters `criticalLeafPLC` and `cvLeafP50`. If $LA_{cavitation} < LA_{act}$ then the corresponding proportion $p_{cavitation,leaf}$ is estimated for senescence.

Xylem cavitation not only causes defoliation but also mortality of persistent buds, which has the effect of slowing down post-drought leaf area recovery. The percentage of the crown with active buds ($PCAB$) is reduced whenever the proportion of conductance loss has increased with respect to the preceding day, using the same approach taken for leaves, but using PLC_{stem} as driver and with different thresholds (see control parameters `criticalBudPLC` and `cvBudP50`).

17.2.2 Sapwood senescence

The daily rate of sapwood senescence is specified via the species-specific parameter $SR_{sapwood}$ or, when missing, via a control parameter. Prentice et al. (1993) assumed a constant annual rate of 4% for the conversion from sapwood to heartwood. Similarly, Sitch et al. (2003) assumed a sapwood annual turnover rate of 5% for all biomes. In our model, the proportion of sapwood area that is transformed into heartwood daily is estimated using:

$$p_{aging,sapwood} = \frac{SR_{sapwood}}{1 + 15 \cdot e^{-0.01 \cdot H}} \cdot \frac{\max(T_{day} - 5, 0)}{20} \quad (17.28)$$

where $SR_{sapwood}$ is a species-specific parameter, T_{day} is the average day temperature and 0.01 is a constant causing shorter plants to have slower senescence rates. It is important to mention that, while stem cavitation PLC_{stem} reduces the amount of functional sapwood in with respect to hydraulics (and therefore transpiration and photosynthesis), it does not increase the rate of sapwood senescence. Hence, when xylem embolism occurs air bubbles are formed within vessels but surrounding parenchymatic cells (as well as the storage carbon they contain) are assumed unaffected.

17.2.3 Fine root senescence

Aging is the only process leading to fine root senescence. The daily turnover proportion for fine roots ($SR_{fineroot}$) is assumed to correspond to a temperature of 25 °C, is specified via species-specific parameters or the control parameters `turnoverRates`. Actual turnover proportion for a given soil layer ($p_{aging,fineroot}$) decreases linearly with soil temperature down to zero at 5 °C:

$$p_{aging,fineroot} = SR_{fineroot} \cdot \frac{\max(T_{soil,s} - 5, 0)}{20} \quad (17.29)$$

where $T_{soil,s}$ is the temperature of soil layer s . When $SR_{fineroot}$ is missing from species parameter, a default control value for $SR_{fineroot}$ is taken, which produces an annual 50% turnover of fine roots.

17.3 Update of plant traits, structural variables and allocation targets

17.3.1 Plant functional traits

Firstly, the model updates tissue variables at the individual level, i.e. actual leaf area (LA_{act}), sapwood area (SA) and fine root biomass at each soil layer (FRB_s), from growth and senescence results.

Secondly, actual leaf area index of the plant cohort (i.e. LAI_{act}) is updated from actual leaf area at the individual level (i.e. LA_{act}) inverting eq. (16.2), and the same is done for the dead leaf area index (i.e. LAI_{dead} ; (16.3)). By definition, fine root biomass changes in each soil layer ($B_{fineroot,s}$) lead to updates in the proportion of fine roots in each layer (FRP_s):

$$FRP_s = \frac{B_{fineroot,s}}{\sum_l B_{fineroot,l}} \quad (17.30)$$

Thirdly, several anatomic and physiological parameters are updated after applying changes in the size of leaf, sapwood and fine root compartments, which creates feedbacks to hydraulic and physiological processes simulated in the water balance submodel:

- a. The Huber value (H_v , the sapwood area to leaf area ratio; in $m^2 \cdot m^{-2}$) is updated using:

$$H_v = \frac{SA/10000}{LA_{live}} \quad (17.31)$$

Note that we use LA_{live} (and not LA_{act}) in the denominator to avoid variations in Huber value due to leaf phenology.

- b. The proportion of conductance loss due to cavitation in stem (PLC_{stem}) and leaf (PLC_{leaf}) segments is reduced whenever sapwood area (resp. leaf area) growth occurs:

$$PLC_{stem,t+1} = \min(PLC_{stem,t} - \frac{\Delta SA}{SA}, 0) \quad (17.32)$$

$$PLC_{leaf,t+1} = \min(PLC_{leaf,t} - \frac{\Delta LA}{LA_{act}}, 0) \quad (17.33)$$

- c. When using the advanced model, stem maximum conductance per leaf area unit ($k_{stem,max}$; in $mmol \cdot m^{-2} \cdot s^{-1} \cdot MPa^{-1}$) is determined as a function of species-specific xylem conductivity ($K_{xylem,max}$; in $kg \cdot m^{-1} \cdot s^{-1} \cdot MPa^{-1}$), leaf area, sapwood area and tree height (Christoffersen et al., 2016):

$$k_{stem,max} = \frac{1000}{0.018} \cdot \frac{K_{xylem,max} \cdot H_v}{(H/100)} \cdot \chi_{taper} \quad (17.34)$$

where χ_{taper} is a factor to account for taper of xylem conduit with height (Savage et al., 2010; Christoffersen et al., 2016), 0.018 is the molar weight of water (in $kg \cdot mol^{-1}$). Both an increase in SA or a decrease in LA_{live} (i.e. an increase in H_v) increase $k_{stem,max}$ and, hence, alleviate drought effects (i.e. a lower decrease in water potential across the stem for the same flow). In contrast, an increase in plant height will decrease stem conductance and increase drought stress.

- d. When using the advanced model, changes in stem maximum conductance have cascade effects on root maximum conductance. First, coarse root minimum resistance is defined as a fixed proportion of whole-plant minimum resistance, so an increase in stem maximum conductance will increase whole-plant conductance and coarse root conductance, $k_{root,max}$.
- e. When using the advanced model, rhizosphere maximum conductance per leaf area unit in a given soil layer s ($k_{rhizo,max,s}$; in $mmol \cdot m^{-2} \cdot s^{-1} \cdot MPa^{-1}$) depends on fine root biomass in this layer ($B_{fineroot,s}$) and on leaf area (i.e. LA_{live}). The equations regulating these relationships are modulated by several soil and species parameters, such as soil saturated hydraulic conductance, species-specific root length, root length density and density of fine roots.

17.3.2 Structural variables and leaf area targets

Tree structural variables

In the case of tree cohorts, the new sapwood area (ΔSA , in cm^2) is translated to an increment in DBH (ΔDBH , in cm) following:

$$\Delta DBH = 2 \cdot \sqrt{(DBH/2)^2 + (\Delta SA/\pi)} - DBH \quad (17.35)$$

Furthermore, the model assumes that increments in height are linearly related to increments in diameter through a function f_{HD} (Lindner et al. 1997):

$$\Delta H = f_{HD} \cdot \Delta DBH \quad (17.36)$$

Hence, f_{HD} represents the height increment (in cm) per each cm of diameter increment. It was customary in forest gap models to prevent height from being larger than a species-specific value $H_{\max}$, so that beyond some point trees only grew in size by increasing their diameter. Moreover, light conditions influence growth in height with trees living under the shade of others generally showing larger increases in height than trees living in open conditions. Hence, our formulation for f_{HD} is (Lindner et al., 1997; Rasche et al., 2012):

$$f_{HD} = [f_{HD,\min} \cdot L^{PAR} + f_{HD,\max} \cdot (1 - L^{PAR})] \cdot \left(1 - \frac{H - 137}{H_{\max} - 137}\right) \quad (17.37)$$

where $f_{HD,\min}$ would be the height-diameter ratio for a tree of 137 cm height growing in full light and $f_{HD,\max}$ would be the same ratio for a tree of the same height growing in the shadow, and L^{PAR} is the proportion of photosynthetically active radiation available at mid-crown height (4.2). This formulation seems slightly easier to calibrate than that presented in Rasche et al. (2012). $H_{\max}$ could be dependent on environmental conditions, but we skip this here, because environmental conditions already affect growth rate and carbon balance.

After updating tree diameter (DBH) and tree height (H), the model updates tree crown ratio (CR) by applying allometric relationships that take into account tree size and competition (see details in 23.2.3).

Tree leaf area target

Foliar biomass without tree competition is first estimated using the following allometry (see chapter 23):

$$FB_{tree,NC} = a_{fbt} \cdot DBH^{b_{fbt}} \cdot e^{-0.0001 \cdot N} \quad (17.38)$$

In turn, $FB_{tree,NC}$ is used to estimate leaf area index without competition effects LAI_{NC} :

$$LAI_{NC} = SLA \cdot FB_{tree,NC} \cdot (N/10000) \quad (17.39)$$

Leaf area allocation target (LA_{target}) is estimated by adding to $FB_{tree,NC}$ the competition effect of due to basal area of larger trees (BAL):

$$LA_{target} = SLA \cdot FB_{tree,NC} \cdot e^{c_{fbt} \cdot BAL} \quad (17.40)$$

and from there, live leaf area (LA_{live}) and live leaf area index (LAI_{live}) are also updated after accounting for the percentage of crown active buds ($PCAB$):

$$LA_{live} = LA_{target} \cdot (PCAB/100) \quad (17.41)$$

$$LAI_{live} = LA_{live} \cdot (N/10000) \quad (17.42)$$

Shrub leaf area target and structural variables

Since shrub structural variables are height and cover, shrub growth is done in a way somewhat different from trees. Shrubs are often multi-stemmed (some trees also are), so that increases in sapwood area are not easily related to diameter growth. Since leaf biomass is related to sapwood area, one may model shrub growth assuming an allometric relationship between phytovolume of individual shrub crowns and photosynthetic biomass. This strategy entails that shrubs may grow or shrink in size depending on their C balance, in the same way that tree crowns would become denser or sparser depending on their C balance. Hence, shrubs can be understood as crowns in the floor.

Given an updated sapwood area (SA), the model first estimates the corresponding target and live leaf areas:

$$LA_{live} = \frac{SA}{10000 \cdot H_{v,target}} \quad (17.43)$$

$$LA_{target} = LA_{live} \cdot (100/PCAB) \quad (17.44)$$

Using LA_{live} , then we can estimate the foliar weight per shrub individual, neglecting drought-induced defoliation or leaf phenology (in $kg \cdot ind^{-1}$):

$$W_{leaves} = \frac{LA_{live}}{SLA} \quad (17.45)$$

An allometric relationship relating the biomass of leaves plus small branches and crown phytovolume (PV ; in $m^3 \cdot ind^{-1}$) can be drawn from fuel calculations:

$$W_{leaves+branches} = W_{leaves} \cdot r_{6.35} = a_{bsh} \cdot PV^{b_{bsh}} \quad (17.46)$$

where a_{bsh} and b_{bsh} are allometric relationships and $r_{6.35}$ is a species-specific ratio relating the dry weight of leaves plus small branches to the dry weight of leaves. Inverting this relationship we obtain an expression of shrub crown phytovolume:

$$PV = \left[\frac{W_{leaves} \cdot r_{6.35}}{a_{bsh}} \right]^{1/b_{bsh}} \quad (17.47)$$

Phytovolume is defined as the volume occupied by the shrub individual, i.e.:

$$PV = (A_{sh}/10000) \cdot (H/100) \quad (17.48)$$

where A_{sh} is the area of a single shrub individual (in cm^2). If we use the following quadratic relationship between A_{sh} and H :

$$A_{sh} = a_{ash} \cdot H^{b_{ash}} \quad (17.49)$$

we can calculate shrub height from phytovolume using:

$$H = \left[\frac{10^6 \cdot PV}{a_{ash}} \right]^{1/(1+b_{ash})} \quad (17.50)$$

Finally, the new value for shrub cover (in percent) can be obtained from H and N (in $ind \cdot ha^{-1}$):

$$Cover = 100 \cdot (N/10000) \cdot (A_{sh}/10000) = \frac{N \cdot a_{ash} \cdot H^2}{10^6} \quad (17.51)$$

Note that crown ratio for shrubs is assumed constant in the model. Like for trees, shrub height is limited to a maximum height H_{max} . However, unlike trees, shrubs are not allowed to continue growing once this maximum size is attained. If this happens, the preceeding equations are inversed to limit LA_{live} and LA_{target} .

17.3.3 Sapwood and fineroot targets

Sapwood area target

If the allocation strategy pursues a *constant Huber value* (i.e. `allocationStrategy = "A12As"`), the sapwood area target (SA_{target}) is defined from leaf area target (LA_{target}) according to a constant value $H_{v,target}$:

$$SA_{target} = \frac{H_{v,target} \cdot LA_{live}}{10000} \quad (17.52)$$

With this formulation, $H_v = H_{v,target}$ and sapwood area target will increase/decrease following changes in LA_{live} (hence being affected by decreases in $PCAB$ due to drought-induced cavitation).

When using the basic model, the only allocation strategy is the constant Huber value. However, when using the advanced model one can choose the allocation strategy pursuing a *constant whole-plant conductance* (i.e. `allocationStrategy = "Plant_kmax"`). In this case, the model keeps $k_{plant,max}$ close to an initial value $k_{plant,target}$, and here the sapwood area target is defined as:

$$SA_{target} = \frac{H_{v,target} \cdot LA_{live}}{10000} \cdot \frac{k_{plant,target}}{k_{plant,max}} \quad (17.53)$$

In this strategy, target sapwood area will be higher whenever the current whole-plant conductance is below the target value (i.e. $k_{plant,max} > k_{plant,target}$) and, conversely, if $k_{plant,max} < k_{plant,target}$ then sapwood area target will be reduced. Note that with this formulation $H_v \neq H_{v,target}$, and in general the Huber value will increase with height, to compensate height increases to achieve a constant whole-plant conductance.

Fine root biomass target

When using the basic model, fine root biomass target is set using:

$$B_{fineroot,target} = \frac{10^4 \cdot LA_{target} \cdot RLR}{2.0 \cdot \sqrt{\frac{\pi \cdot SRL}{\rho_{fineroot}}}} \quad (17.54)$$

where RLR is the root area to leaf area ratio, $\rho_{fineroot}$ is fine root tissue density ($g\ dry \cdot cm^{-3}$) and SRL ($cm \cdot g\ dry^{-1}$) is the specific root length.

When using the advanced model, the target of fine root biomass for any given soil layer s ($B_{fineroot,target,s}$) directly follows changes in maximum whole-plant conductance. The average resistance in the rhizosphere is assumed to correspond to a fixed percentage of total soil-plant resistance. Hence, changes in the conductance of leaves, stem or coarse roots will entail a variation in the absolute rhizosphere maximum conductance to be targeted ($k_{rhizo,max,target,s}$), which in turn will determine $B_{fineroot,target,s}$. For example $k_{rhizo,max,target,s}$ will increase as a consequence of sapwood area growth. The model thus first estimate $k_{rhizo,max,target,s}$ for each layer s and then translates $k_{rhizo,max,target,s}$ values to $B_{fineroot,target,s}$ using the relationships based on soil saturated hydraulic conductance, species-specific root length, root length density and density of fine roots mentioned above.

17.3.4 Storage concentrations

Leaf and sapwood carbon storage capacity depend on structural parameters such as sapwood area (SA), leaf area (LA_{act}), height (H) and root distribution. Hence, sugar and starch concentrations need to be updated after tissues and structural variables have been modified.

17.4 Plant mortality

17.4.1 Self-thinning of small trees

As explained in the mortality design section (15.1.8), the model implements a mortality process for young individual trees, meant to represent a self-thinning process occurring between trees of small diameter ($DBH_{tree,recr}$, typically 1 cm) and trees having a diameter corresponding to inclusion as individual in forest inventories ($DBH_{tree,ingrowth}$, typically 7.5 cm). The aim is to ensure that the tree density is progressively reduced until the tree cohort reaches $DBH_{tree,ingrowth}$, where the density should be $N_{tree,ingrowth}$. The following relationship between tree diameter and density is used:

$$N = a_{st} \cdot DBH^{b_{st}} \quad (17.55)$$

where a_{st} and b_{st} are parameters regulating the speed of the self-thinning process. Note that if we know the diameter and density of recruitment (i.e. $DBH_{tree,recr}$ and $N_{tree,recr}$), as well as $N_{tree,ingrowth}$, the density we want to ensure when the tree reaches $DBH_{tree,ingrowth}$, the self-thinning curve is completely determined. Hence, we can estimate b_{st} using:

$$b_{st} = \frac{\log(N_{tree,ingrowth}/N_{tree,recr})}{\log(DBH_{tree,ingrowth}/DBH_{tree,recr})} \quad (17.56)$$

whereas a_{st} can be estimated using:

$$a_{st} = \frac{N_{tree,ingrowth}}{DBH_{tree,ingrowth}^{b_{st}}} \quad (17.57)$$

Once we know the parameters of the self-thinning curve, we can determine the maximum cohort density allowed for any cohort of diameter DBH_i . If the tree cohort has density N_i , the density decrease due to self-thinning mortality can be estimated using:

$$N_{dead,i} = N_i - \min(N_i, a_{st} \cdot DBH_i^{b_{st}}) \quad (17.58)$$

Note that the self-thinning process does not distinguish between tree resprouts or trees recruited from seeds.

17.4.2 Basal mortality rates

Adult trees ($DBH > 7.5$ cm) and shrubs die at a basal rate due to unspecific causes. For shrubs, the basal rate is always a species-specific constant ($P_{mort,base}$). For trees, the basal rate can alternatively depend on the degree of (symmetric) competition, which is measured using the tree basal area BA ($m^2 \cdot ha^{-1}$) of the stand, through a simple survival logistic model:

$$P_{surv,base} = \text{logit}^{-1}(\beta_{surv,0} + \beta_{surv,1} \cdot \sqrt{BA}) = \frac{\exp(\beta_{surv,0} + \beta_{surv,1} \cdot \sqrt{BA})}{1 + \exp(\beta_{surv,0} + \beta_{surv,1} \cdot \sqrt{BA})} \quad (17.59)$$

where $\beta_{surv,0}$ and $\beta_{surv,1}$ are species-specific coefficients obtained by fitting Generalized Linear Models on repeated forest inventory plot data. These models can be developed for arbitrary time steps (normally 5 or 10 years), but the basal mortality rate has to be finally expressed as a daily probability ($P_{base,daily}$).

17.4.3 Mortality due to starvation

The starvation stress indicator is $ST_{relative}^{sapwood}$, the starch concentration in sapwood relative to the equilibrium value:

$$ST_{relative}^{sapwood} = \frac{ST_{sapwood}}{ST_{equilibrium}^{sapwood}} \quad (17.60)$$

where $ST_{sapwood}$ ($mol\ glucose \cdot l^{-1}$) is the current starch concentration in sapwood and $ST_{equilibrium}^{sapwood}$ is the equilibrium concentration (by default $ST_{equilibrium}^{sapwood} = 0.35\ mol\ glucose \cdot l^{-1}$). Assuming that a threshold of relative starch concentration ($ST_{relative}^{thresh}$) corresponds to 50% annual mortality due to starvation, a logistic sigmoidal function is used to estimate the probability of annual starvation from a given $ST_{relative}^{sapwood}$:

$$P_{starv,annual} = 1.0 - \frac{\exp(40 \cdot (ST_{relative}^{sapwood} - ST_{relative}^{thresh}))}{1.0 + \exp(40 \cdot (ST_{relative}^{sapwood} - ST_{relative}^{thresh}))} \quad (17.61)$$

Since the model operates at the daily temporal resolution, $P_{starv,annual}$ is re-expressed as a daily probability using:

$$P_{starv,daily} = 1.0 - \exp(\log(1.0 - P_{starv,annual})/356) \quad (17.62)$$

Assuming a relative threshold $ST_{relative}^{thresh} = 0.4$, the figure below illustrates the shape of the sigmoidal function (top) and the corresponding daily probability function.

```
## Warning: Using `size` aesthetic for lines was deprecated in ggplot2 3.4.0.
## i Please use `linewidth` instead.
## This warning is displayed once per session.
## Call `lifecycle::last_lifecycle_warnings()` to see where this warning was
## generated.
```




Figure 17.1: Annual (top) and daily (bottom) probability of starvation as a function of sapwood starch concentration relative to its equilibrium value.

17.4.4 Mortality due to dessication

As explained in the section 15.1.8, the dessication stress indicator (D_{stem}) is the maximum of relative water content in the stem (RWC_{stem}) and relative hydraulic conductance in the stem, i.e. the complement of PLC_{stem}

$$D_{stem} = (RWC_{stem} \cdot 1.0 - PLC_{stem}) / 2.0 \quad (17.63)$$


Assuming that a threshold of relative water content (RWC_{thresh}) corresponds to 50% annual mortality due to dessication, its probability is estimated analogously to starvation. First, a logistic sigmoidal function is used to estimate the probability of annual dessication from a given D_{stem} value:

$$P_{dessic,annual} = 1.0 - \frac{\exp(40 \cdot (D_{stem} - RWC_{thresh}))}{1.0 + \exp(40 \cdot (D_{stem} - RWC_{thresh}))} \quad (17.64)$$

Since the model operates at the daily temporal resolution, $P_{dessic,annual}$ is re-expressed as a daily probability using:

$$P_{dessic,daily} = 1.0 - \exp(\log(1.0 - P_{dessic,annual}) / 356) \quad (17.65)$$


Assuming a relative threshold $RWC_{thresh} = 0.4$, the figure below illustrates the shape of the sigmoidal




function (top) and the corresponding daily probability function.

Uncertainty of stem pressure-volume parameters


As explained in 15.1.8, using stem relative water content as indicator of dessication has the inconvenient that the parameters $\pi_{0,stem}$ and ϵ_{stem} specifying the pressure-volume curve are often not known, so that estimates have to come from wood density (see section A.3.14). The following figure shows the water potential corresponding to the dessication threshold for species with different wood density:



If the relationship between wood density and the stem pressure-volume curve (A.3.14) does not hold for a particular species the mortality rate may be under or overestimated. Here an example would be *Cistus albidus*, which has a low wood density but high dessication resistance. This is illustrated when we draw a scatter plot between the water potential corresponding to 60% stem PLC (i.e. 40% stem relative hydraulic conductance) against the water potential corresponding to 40% stem RWC:



As explained in 15.1.8, one can avoid the potential uncertainty in dessication mortality by defining the dessication stress indicator (D_{stem}) as the maximum of relative water content in the stem (RWC_{stem}) and relative hydraulic conductance in the stem. When we do so, the following curves are obtained for each of the previous species:



It is obvious in the former figure that using D_{stem} as indicator results in curves that are not continuous in all their range. However, we prefer this inconvenient to the potential overestimation of mortality due to an uncertainty in the pressure-volume curve parameters.

17.4.5 Overall mortality probability

Every day the model determines for woody cohorts the overall probability of mortality ($P_{mort,daily}$) as the maximum of the basal probability, starvation probability and desiccation probability:

$$P_{mort,daily} = \max(P_{base,daily}, P_{starv,daily}, P_{dessic,daily}) \quad (17.66)$$

At this point $P_{mort,daily}$ may be used deterministically (i.e., as a proportion of N to kill) or stochastically (i.e. throwing a random number between 0 and 1 to determine the mortality event).

17.5 Fire severity

Fire effects on plants follow, with some modifications, the model by Michaletz and Johnson (2008). Buoyant plume theory is first used to estimate the vertical plume temperature T_{plume} distribution that will drive heat transfer to the vascular cambium and vegetative organs. Heat transfer theory is then used to calculate

the depth of vascular cambium necrosis and height of crown foliage and crown bud necrosis. Finally, these severity metrics are used to define the fate of the plant cohort, as described in section 15.1.9.

17.5.1 Plume temperature distribution

The surface fire is assumed to occur at the time of the day where temperature is T_{max} is the maximum daily temperature. For a line-source plume in a quiescent atmosphere (no wind), the plume temperature T_{plume} at height z (in m) can be estimated using (Michaletz and Johnson, 2008):

$$T_{plume}(z) = \min \left[900, C_{plume} \cdot \left(\frac{1}{z} \right) \cdot \left(\frac{T_{max} + 273.15}{g} \right)^{1/3} \cdot \left(\frac{I_B}{c_{p,air} \cdot \rho_{air}} \right)^{2/3} + T_{max} \right] \quad (17.67)$$

where $C_{plume} = 2.6$ is the plume proportionality constant, $g = 9.8 \text{ m} \cdot \text{s}^{-2}$ is the gravity constant, I_B is Byram's surface fireline intensity (eq. (26.131); expressed in $\text{kW} \cdot \text{m}^{-1}$), $c_{p,air} = 1.007 \text{ J} \cdot \text{kg}^{-1} \cdot ^\circ\text{C}^{-1}$ is the specific heat capacity of the air and ρ_{air} is the air density at temperature T_{max} . Because this similarity analysis fails at heights comparable with the fireline width, T_{plume} is constrained to a maximum flame temperature of 900°C .

17.5.2 Foliage and crown bud necrosis

The model predicting the distribution of plume temperature and the residence time of surface fires t_R (eq. (26.132)), are used to determine the height of necrosis for plant organs, here foliage or crown buds, depending also on their heat capacitance. The estimate (foliage or bud) necrosis height z_n (in m) is given by (Michaletz and Johnson, 2008):

$$z_n = C_{plume} \cdot \left(\frac{1}{T_{crit} - T_{max}} \right) \cdot \left(\frac{T_{max} + 273.15}{g} \right)^{1/3} \cdot \left(\frac{I_B}{c_{p,air} \cdot \rho_{air}} \right)^{2/3} \quad (17.68)$$

where T_{crit} is the critical temperature of the plume required for organ necrosis, given a residence time t_R , which is estimated using:

$$T_{crit} = \frac{T_n - \theta_T \cdot T_{max}}{1 - \theta_T} \quad (17.69)$$

Here, T_n is the temperature leading to necrosis, assumed to be 60°C and θ_T is defined as the excess temperature ratio, which depends on the organ thermal factor (TF) and t_R :

$$\theta_T = \exp(-TF \cdot t_R) \quad (17.70)$$

The thermal factor TF is normally larger for leaves than buds, given their higher surface area to volume ratio, resulting in lower values of T_{crit} and, therefore, higher values of necrosis height. In the case of leaves, the thermal factor (TF_{leaves}) is estimated from specific leaf area, SLA , using (Michaletz and Johnson, 2006, 2008):

$$TF_{leaves} = SLA \cdot (h_{leaves}/c_{leaves}) \quad (17.71)$$

where $h_{leaves} = 130$ stands for the convection heat transfer coefficient of leaves and $c_{leaves} = 2500$ is the specific heat capacity of leaves. Typical values would be $TF_{leaves} = 0.208$ for $SLA = 4$ and $TF_{leaves} = 0.624$ for $SLA = 12$. Currently, for crown buds a constant $TF_{buds} = 0.130$ is assumed, on the basis of values given in Michaletz and Johnson (2006), but we acknowledge that TF_{buds} should be at least species-specific. Once z_n is defined, for either leaves or crown buds, depending on their TF value, the vertical leaf distribution described in 2.4.3.3 is used to determine the proportion of crown leaves or crown buds with necrosis.

If the case of a crown fires, i.e. if surface fireline intensity is larger than van Wagner's critical intensity; eq. (26.160), then all crown foliage is burned. The same happens if torching occurs for the target plant cohort, which is determined by calculating a cohort-specific critical intensity. Crown bud necrosis is decided in those cases by calculating T_{crit} using the crown fire residence time and comparing it to the flame temperature, i.e. 900°C . Despite these calculations, in practice all crown buds will also normally suffer necrosis under a crown fire.

17.5.3 Cambium necrosis

Vascular cambium necrosis is a one-dimensional transient conduction problem depending the thermal diffusivity of the bark, α_{bark} , and the surface fire residence time t_R . Assuming that the bark surface temperature is equal to the plume temperature at $z = 0.1$, i.e. $T_{plume}(0.1)$, the radial bole necrosis depth, x_n (in m), can be estimated using:

$$x_n = 2 \cdot (\alpha_{bark} \cdot t_R)^{1/2} \cdot \text{erf}^{-1} \left(\frac{T_n - T_{plume}(0.1)}{T_{max} - T_{plume}(0.1)} \right) \quad (17.72)$$

where erf^{-1} is the inverse error function. Bark diffusivity (α_{bark}) is estimated following the equations given in Michaletz and Johnson (2008), which depend on bark's tissue density, moisture content (estimated assuming a fine dead fuel) and temperature. Cambium necrosis is determined by comparing x_n (in m) with x_{ba} (in mm) the bark thickness of the plant, which in case of trees depends on DBH:

$$x_{ba} = a_{bt} \cdot DBH^{b_{bt}} \quad (17.73)$$

and in the case of shrubs x_{ba} is assumed to be equal to a fixed species-specific value bt_{sh} .

Chapter 18

Decomposition and soil biogeochemistry

18.1 Carbon pools, losses and fluxes

Carbon pools and fluxes between them were described in the design section and illustrated in fig. 15.5. The following table provides the terminologic mapping between carbon pools in MEDFATE, which mainly follows Bonan et al. (2019), and the corresponding pools in CENTURY:

MEDFATE	CENTURY (Parton et al.)	CENTURY code
Small branches snags*		
Large wood snags*		
Twig litter*		
Small branch litter	Dead fine branch	wood1c
Large wood litter	Dead large wood	wood2c
Coarse root litter	Dead coarse roots	wood2c
Structural leaf litter	Surface structural	strucc(1)
Metabolic surface	Surface metabolic	metabc(1)
Active surface	Surface microbe	som1c(1)
Slow surface		som2c(1)
Structural fine root litter	Belowground structural	strucc(2)
Metabolic belowground	Belowground metabolic	metabc(2)
Active belowground	Active organic	som1c(2)
Slow belowground	Slow organic	som2c(2)
Passive belowground	Passive	som3c

The first three carbon pools (marked with an asterisk) are those new to MEDFATE, in comparison with Bonan et al. (2019). In CENTURY, only leaf litter and fine root litter are those conforming structural pools, whereas dead wood (either fine branches, large wood or coarse roots) is considered separately.

Each (daily) time step a decomposition rate k_i is estimated for each carbon pool i . This decomposition rate leads to an amount of carbon (ΔC_i) being loss from the carbon pool:

$$\Delta C_i = k_i \cdot \Delta t \quad (18.1)$$

The carbon lost can exit the carbon pool as heterotrophic respiration (HR_i) or it can lead to a carbon flux to another carbon pool j , e.g. $F_{i,j}$. The dynamics of the whole set of pools can be solved using matrix algebra.

The following sections detail how decomposition rates are estimated as well as the estimation of respiration and fluxes between carbon pools.

18.2 Decomposition rates

Each carbon pool i has its own base decomposition rate ($k_{\text{base},i}$). The base decomposition rates of a carbon pool i are reduced by multiplicative functions of environmental factors:

$$k_i = k_{\text{base},i} \cdot f(T) \cdot f(\theta) \cdot f(pH) \quad (18.2)$$

Where $k_{\text{base},i}$ is the base daily decomposition rate for pool i which is multiplied by three modifiers: (T) is the effect of temperature, $f(\theta_{\text{rel}})$ is the effect of moisture relative to saturation, and $f(pH)$ is the effect of soil pH. Additional factors are considered for specific pools. For example, belowground pools include a factor to account for anaerobic conditions. Similarly, structural carbon pools include a factor to account for the effect of lignin percent on decay rates.

18.2.1 Effect of pH

The effect of pH on decomposition, $f(pH)$ is calculated using:

$$f(pH) = b + \frac{c}{\pi} \cdot \tan^{-1}[d \cdot (pH - a) \cdot \pi] \quad (18.3)$$

where parameters a , b , c and d depend on the specific carbon pool:

Carbon pool(s)	a	b	c	d
Surface/belowground metabolic	4.8	0.5	1.14	0.7
Surface/belowground structural	4.0	0.5	1.1	0.7
Surface active	4.0	0.5	1.1	0.7
Belowground active	4.8	0.5	1.14	0.7
Surface/belowground slow	4.0	0.5	1.1	0.7
Belowground passive	3.0	0.5	1.1	0.7

18.2.2 Effect of temperature

The effect of temperature on decomposition, $f(T)$, follows the same general structure as the effect of pH, but with fixed coefficients (i.e. $a = 15.7$, $b = 0.56$, $c = 1.46$ and $d = 0.031$) (Del Grosso et al., 2005):

$$f(T) = 0.56 + \frac{1.46}{\pi} \cdot \tan^{-1}[0.0309 \cdot \pi \cdot (T - 15.7)] \quad (18.4)$$

For most carbon pools, the temperature value is that corresponding to the first (top) soil layer. For decomposition of snags, the temperature within the canopy is used.

18.2.3 Effect of moisture

The effect of moisture is modelled using (Kelly et al., 2000):

$$f(\theta_{\text{rel}}) = \left(\frac{\theta_{\text{rel}} - b}{a - b} \right)^{\frac{d \cdot (b - a)}{a - c}} \cdot \left(\frac{\theta_{\text{rel}} - c}{a - c} \right)^d \quad (18.5)$$




Figure 18.1: Effect of pH on decomposition rates, for different carbon pools.




Figure 18.2: Effect of temperature on decomposition rates.

where θ_{rel} is moisture relative to saturation, and here parameters a , b , c and d depend on soil texture (Kelly et al., 2000):

Soil texture	a	b	c	d
Fine texture	0.6	1.27	0.0012	2.84
Coarse texture	0.55	1.7	-0.007	3.22




Figure 18.3: Effect of moisture relative to saturation (θ_{rel}) on decomposition rates, for fine textured and coarse textured soil.

For most carbon pools, relative moisture is estimated from the moisture of the first (top) soil layer. In the case of snag carbon pools, relative moisture is estimated from a ratio of moisture content in equilibrium with air relative humidity and moisture content at saturation.

18.2.4 Effect of anaerobiosis

Decomposition of belowground pools includes an effect for anaerobic soils, represented by a rate factor $f(O_2)$ that varies between zero and one and that decreases decomposition rates as soil becomes increasingly waterlogged.

18.3 Decomposition process of leaf and fine root litter, coarse woody debris and standing dead wood

18.3.1 Leaf and fine root litter

When leaf or fine root litter is produced, the corresponding carbon is divided into structural and metabolic pools (fig. 15.5). Metabolic litter is labile material that decomposes quickly, whereas structural material contains cellulose and lignin and decomposes more slowly. The fraction of leaf and fine root litter that is added to the corresponding surface or belowground metabolic pool is determined using:

$$f_{\text{metabolic}} = 0.85 - 0.0013(\text{lignin}/N) \quad (18.6)$$

A higher lignin/N ratio thus results in more structural litter. In turn, structural litter pools have a slower decaying rate (by default $k_{\text{base,leaf-struct}} = 2\text{y}^{-1}$ and $k_{\text{base,fineroot-struct}} = 4.9\text{y}^{-1}$ for leaves and fine roots, respectively) compared to their corresponding metabolic pools (by default $k_{\text{base,surface-metabolic}} = 8\text{y}^{-1}$ and $k_{\text{base,belowground-metabolic}} = 18.5\text{y}^{-1}$ for surface and belowground pools, respectively).

There are as many leaf and fine root structural pools as plant species, because decomposition rates depend on lignin content. In contrast, there is only one surface metabolic pool and one belowground metabolic pool.

18.3.1.1 Leaf structural pools

Leaf structural pools are species-specific and receive carbon from partitioning of leaf litter. The rate of decomposition of a leaf structural pool i with leaf lignin content $f_{\text{leaf-lignin},i}$ is:

$$k_{\text{leaf-struct},i} = k_{\text{base, leaf-struct}} \cdot f(T) \cdot f(\theta_{\text{rel}}) \cdot f(pH) \cdot \exp(-3 \cdot f_{\text{leaf-lignin},i}) \quad (18.7)$$

The loss of carbon from a leaf structural pool ($\Delta C_{\text{leaf-struct},i}$) is apportioned between heterotrophic respiration ($HR_{\text{leaf-struct},i}$) and the carbon flux to the surface active ($F_{\text{leaf-struct,surface-active},i}$) and surface slow ($F_{\text{leaf-struct,surface-slow},i}$) pools:

$$HR_{\text{leaf-struct},i} = \Delta C_{\text{leaf-struct},i} \cdot (f_{\text{leaf-lignin},i} \cdot 0.30 + (1.0 - f_{\text{leaf-lignin},i}) \cdot 0.45) \quad (18.8)$$

$$F_{\text{leaf-struct,surface-active},i} = \Delta C_{\text{leaf-struct},i} \cdot (1.0 - f_{\text{leaf-lignin},i}) \cdot (1 - 0.45) \quad (18.9)$$

$$F_{\text{leaf-struct,surface-slow},i} = \Delta C_{\text{leaf-struct},i} \cdot f_{\text{leaf-lignin},i} \cdot (1 - 0.30) \quad (18.10)$$

18.3.1.2 Surface metabolic pool

The surface metabolic pool receives carbon from partitioning of leaf litter. The rate of decomposition of the surface metabolic pool is:

$$k_{\text{surface-metabolic}} = k_{\text{base,surface-metabolic}} \cdot f(T) \cdot f(\theta_{\text{rel}}) \cdot f(pH) \quad (18.11)$$

The loss of carbon from the surface metabolic pool ($\Delta C_{\text{surface-metabolic}}$) is apportioned between heterotrophic respiration ($HR_{\text{surface-metabolic}}$) and the carbon flux to the surface active ($F_{\text{surface-metabolic,surface-active}}$) pool:

$$HR_{\text{surface-metabolic}} = \Delta C_{\text{surface-metabolic}} \cdot 0.55 \quad (18.12)$$

$$F_{\text{surface-metabolic,surface-active}} = \Delta C_{\text{surface-metabolic}} \cdot (1 - 0.55) \quad (18.13)$$

18.3.1.3 Fine root structural pools

Fine root structural pools are species-specific and receive carbon from fine root litter. The rate of decomposition of a fine root structural pool i with fine root lignin content $f_{\text{fineroot-lignin},i}$ is:

$$k_{\text{fineroot-struct},i} = k_{\text{base, fineroot-struct}} \cdot f(T) \cdot f(\theta_{\text{rel}}) \cdot f(pH) \cdot \exp(-3 \cdot f_{\text{fineroot-lignin},i}) \cdot f(O_2) \quad (18.14)$$

The loss of carbon from a fine root structural pool ($\Delta C_{\text{fineroot-struct},i}$) is apportioned between heterotrophic respiration ($HR_{\text{fineroot-struct},i}$) and the carbon flux to the belowground active ($F_{\text{fineroot-struct,belowground-active},i}$) and belowground slow ($F_{\text{fineroot-struct,belowground-slow},i}$) pools:

$$HR_{\text{fineroot-struct},i} = \Delta C_{\text{fineroot-struct},i} \cdot (f_{\text{fineroot-lignin},i} \cdot 0.30 + (1.0 - f_{\text{leaf-lignin},i}) \cdot 0.55) \quad (18.15)$$

$$F_{\text{fineroot-struct,belowground-active},i} = \Delta C_{\text{fineroot-struct},i} \cdot (1.0 - f_{\text{fineroot-lignin},i}) \cdot (1 - 0.55) \quad (18.16)$$

$$F_{\text{fineroot-struct,belowground-slow},i} = \Delta C_{\text{fineroot-struct},i} \cdot f_{\text{fineroot-lignin},i} \cdot (1 - 0.30) \quad (18.17)$$

18.3.1.4 Belowground metabolic pool

The belowground metabolic pool receives carbon from partitioning of fine root litter as well as from root exudation (RE). The rate of decomposition of the belowground metabolic pool is:

$$k_{\text{belowground-metabolic}} = k_{\text{base,belowground-metabolic}} \cdot f(T) \cdot f(\theta_{rel}) \cdot f(pH) \cdot f(O_2) \quad (18.18)$$

The loss of carbon from the belowground metabolic pool ($\Delta C_{\text{belowground-metabolic}}$) is apportioned between heterotrophic respiration ($HR_{\text{belowground-metabolic}}$) and the carbon flux to the belowground active ($F_{\text{belowground-metabolic,belowground-active}}$) pool:

$$HR_{\text{belowground-metabolic}} = \Delta C_{\text{belowground-metabolic}} \cdot 0.55 \quad (18.19)$$

$$F_{\text{belowground-metabolic,belowground-active}} = \Delta C_{\text{belowground-metabolic}} \cdot (1 - 0.55) \quad (18.20)$$

18.3.2 Coarse roots

Coarse root carbon pools (in $gC \cdot m^{-2}$) include belowground woody material of all sizes, created from mortality of belowground plant structures. The rate of decomposition of a coarse root pool i with lignin content $f_{\text{woody-lignin},i}$ is:

$$k_{\text{coarseroot},i} = k_{\text{base, coarseroot}} \cdot f(T) \cdot f(\theta_{rel}) \cdot \exp(-3 \cdot f_{\text{woody-lignin},i}) \cdot f(O_2) \quad (18.21)$$

And the loss of carbon ($\Delta C_{\text{coarseroot},i}$) is apportioned between heterotrophic respiration ($HR_{\text{coarseroot},i}$) and the carbon flux to the belowground active ($F_{\text{coarseroot, belowground-active},i}$) and belowground slow ($F_{\text{coarseroot, belowground-slow},i}$) pools:

$$HR_{\text{coarseroot},i} = \Delta C_{\text{coarseroot},i} \cdot (f_{\text{woody-lignin},i} \cdot 0.30 + (1.0 - f_{\text{woody-lignin},i}) \cdot 0.45) \quad (18.22)$$

$$F_{\text{coarseroot, belowground-active},i} = \Delta C_{\text{coarseroot},i} \cdot (1.0 - f_{\text{woody-lignin},i}) \cdot (1 - 0.45) \quad (18.23)$$

$$F_{\text{coarseroot, belowground-slow},i} = \Delta C_{\text{coarseroot},i} \cdot f_{\text{woody-lignin},i} \cdot (1 - 0.30) \quad (18.24)$$

18.3.3 Downed woody debris

Downed woody debris is created via snag fall or twig litterfall. Three types of carbon pools are considered for downed woody debris (fig. 15.5): **twigs** (diameters lower than 0.635 cm; 1h fuels), **small branches** (diameters between 0.635 cm and 7.5 cm; 10h and 100h fuels; in $gC \cdot m^{-2}$) and **large wood** (diameters larger than 7.5 cm; 1000h and 10000h fuels; in $gC \cdot m^{-2}$). In contrast to litter, there is no separation between structural and metabolic components, since all carbon is considered to be structural.

The rate of decomposition of a downed dead pool i with lignin content $f_{\text{woody-lignin},i}$ is:

$$k_i = k_{\text{base,downed-woody}} \cdot f(T) \cdot f(\theta_{rel}) \cdot \exp(-3 \cdot f_{\text{woody-lignin},i}) \quad (18.25)$$

where $k_{\text{base,downed-woody}}$ is the base daily decomposition rate for downed woody elements (which is higher for twigs, than small branches, in turn larger than large wood); temperature and moisture factors are the same as for leaf litter.

The loss of carbon from downed coarse wood ($\Delta C_{\text{downed-woody},i}$) is apportioned between heterotrophic respiration ($HR_{\text{downed-woody},i}$) and the carbon flux to the surface active ($F_{\text{downed-woody,surface-active},i}$) and surface slow ($F_{\text{downed-woody,surface-slow},i}$) pools:

$$HR_{\text{downed-woody},i} = \Delta C_{\text{downed-woody},i} \cdot (f_{\text{woody-lignin},i} \cdot 0.30 + (1.0 - f_{\text{woody-lignin},i}) \cdot 0.45) \quad (18.26)$$

$$F_{\text{downed-woody,surface-active},i} = \Delta C_{\text{downed-woody},i} \cdot (1.0 - f_{\text{woody-lignin},i}) \cdot (1 - 0.45) \quad (18.27)$$

$$F_{\text{downed-woody,surface-slow},i} = \Delta C_{\text{downed-woody},i} \cdot f_{\text{woody-lignin},i} \cdot (1 - 0.30) \quad (18.28)$$

18.3.4 Standing dead wood

Standing dead wood is created via mortality of plant aboveground structures or sapwood senescence (in the case of shrubs). Two types carbon pools are considered for standing dead wood (fig. 15.5): **small branches** (diameters between 0.635 cm and 7.5 cm; 10h and 100h fuels; in $gC \cdot m^{-2}$) and **large wood** (diameters larger than 7.5 cm; 1000h and 10000h fuels; in $gC \cdot m^{-2}$).

The rate of decomposition of a standing dead pool i with lignin content $f_{\text{woody-lignin},i}$ is:

$$k_{\text{standing-woody},i} = k_{\text{standing-woody,base}} \cdot f(T) \cdot f(\theta_{rel}) \cdot \exp(-3 \cdot f_{\text{woody-lignin},i}) \quad (18.29)$$

where $k_{\text{standing-woody,base}}$ is the base daily decomposition rate for standing woody elements (which is higher for small branches than large wood); and atmosphere (or canopy in the case of advanced simulations) air temperature is used for temperature effects (i.e. $T = T_{air}$ or $T = T_{can}$). Moisture relative to saturation (θ_{rel}) is estimated as the ratio of actual fuel moisture to its maximum value (both with units of percent of water mass to dry mass):

$$\theta_{rel} = \theta_{simard}(T_{air}, RH_{air}) / \theta_{max} \quad (18.30)$$

where θ_{simard} is the equilibrium moisture content (Simard 1968) of wood, according to an empirical relationship with air temperature and air relative humidity (Viney, 1991) and θ_{max} is the maximum fuel moisture content, which depends on wood density (assuming the density of water as $1 g \cdot cm^{-3}$):

$$\theta_{max} = 100 \cdot \left(\frac{1}{\rho_{wood}} - \frac{1}{\rho_{ws}} \right) \quad (18.31)$$

where ρ_{wood} is wood density ($g \cdot cm^{-3}$) and $\rho_{ws} = 1.53 g \cdot cm^{-3}$ is the density of the wood substance (a constant). Since θ_{simard} ranges between 0% and 30%, whereas θ_{max} ranges between 45% ($\rho_{wood} = 0.9$) and 135% ($\rho_{wood} = 0.5$), the θ_{rel} ratio is frequently lower than the same ratio assessed from soil moisture, resulting in slower decomposition rates.

Similarly to the downed dead wood, the loss of carbon from standing dead pools ($\Delta C_{\text{standing-woody},i}$) is apportioned between heterotrophic respiration ($HR_{\text{standing-woody},i}$) and the carbon flux to the surface active ($F_{\text{standing-woody,surface-active},i}$) and surface slow ($F_{\text{standing-woody,surface-slow},i}$) pools:

$$HR_{\text{standing-woody},i} = \Delta C_{\text{standing-woody},i} \cdot (f_{\text{woody-lignin},i} \cdot 0.30 + (1.0 - f_{\text{woody-lignin},i}) \cdot 0.45) \quad (18.32)$$

$$F_{\text{standing-woody,surface-active},i} = \Delta C_{\text{standing-woody},i} \cdot (1.0 - f_{\text{woody-lignin},i}) \cdot (1 - 0.45) \quad (18.33)$$

$$F_{\text{standing-woody,surface-slow},i} = \Delta C_{\text{standing-woody},i} \cdot f_{\text{woody-lignin},i} \cdot (1 - 0.30) \quad (18.34)$$

18.4 Decomposition in soil organic carbon pools

Soil organic matter is represented by pools of active, slow and passive carbon that represent material with increasingly longer turnover time. These pools have a surface and belowground component, except for the passive pool, which occurs belowground only (fig. 15.5). The active pools represent live soil microbes and microbial products and decomposes rapidly. The slow pool includes decay-resistant plant material from

the lignin portion of structural litter and soil-stabilized material from the active pool. Carbon fluxes exist between the active and slow pools of the surface and belowground layers, as well as between the active and passive belowground pools. In addition, a carbon flux connection between surface and belowground is included via physical mixing between slow pools. The following details the carbon losses, heterotrophic respiration and fluxes for and between pools.

18.4.1 Active surface pool

The active surface pool receives carbon from surface structural pools, the metabolic surface pool and from the slow surface pool. The rate of decomposition of the active surface pool is:

$$k_{\text{surface-active}} = k_{\text{base,surface-active}} \cdot f(T) \cdot f(\theta_{rel}) \cdot f(pH) \quad (18.35)$$

The loss of carbon from the surface active pool ($\Delta C_{\text{surface-active}}$) is apportioned between heterotrophic respiration ($HR_{\text{surface-active}}$) and the carbon flux to the surface slow pool ($F_{\text{surface-active,surface-slow}}$):

$$HR_{\text{surface-active}} = \Delta C_{\text{surface-active}} \cdot 0.60 \quad (18.36)$$

$$F_{\text{surface-active,surface-slow}} = \Delta C_{\text{surface-active}} \cdot (1 - 0.60) \quad (18.37)$$

18.4.2 Slow surface pool

The slow surface pool receives carbon from decomposition of surface structural litter and from the active surface pool. The rate of decomposition of the slow surface pool is:

$$k_{\text{surface-slow}} = (k_{\text{base,surface-slow}} \cdot f(pH) + k_{\text{mixing}}) \cdot f(T) \cdot f(\theta_{rel}) \quad (18.38)$$

where k_{mixing} is the rate of carbon flux from slow surface to slow belowground due to mixing. The loss of carbon from the surface slow pool ($\Delta C_{\text{surface-slow}}$) is apportioned between heterotrophic respiration ($HR_{\text{surface-slow}}$), the carbon flux to back to the surface active pool ($F_{\text{surface-slow,surface-active}}$) and the carbon flux to the belowground slow pool ($F_{\text{surface-slow,belowground-slow}}$):

$$HR_{\text{surface-slow}} = \Delta C_{\text{surface-slow}} \cdot (1 - k_{\text{mix,frac}}) \cdot 0.55 \quad (18.39)$$

$$F_{\text{surface-slow,surface-active}} = \Delta C_{\text{surface-slow}} \cdot (1 - k_{\text{mix,frac}}) \cdot (1 - 0.55) \quad (18.40)$$

$$F_{\text{surface-slow,belowground-slow}} = \Delta C_{\text{surface-slow}} \cdot k_{\text{mix,frac}} \quad (18.41)$$

where $k_{\text{mix,frac}} = k_{\text{mixing}} / (k_{\text{base,surface-slow}} \cdot f(pH) + k_{\text{mixing}})$.

18.4.3 Active belowground pool

The active belowground pool receives carbon from belowground structural and metabolic pools as well as from the slow and passive belowground pools. The rate of decomposition of the active belowground pool is:

$$k_{\text{belowground-active}} = k_{\text{base,belowground-active}} \cdot f(T) \cdot f(\theta_{rel}) \cdot f(pH) \cdot f_{\text{texture}} \cdot f(O_2) \quad (18.42)$$

where $f_{\text{texture}} = 0.25 + 0.75 \cdot P_{\text{sand}}$ and P_{sand} is the proportion of sand.

The loss of carbon from the belowground active pool ($\Delta C_{\text{belowground-active}}$) is apportioned between heterotrophic respiration ($HR_{\text{belowground-active}}$), the carbon flux to the belowground slow pool ($F_{\text{belowground-active,belowground-slow}}$) and the carbon flux to the belowground passive pool ($F_{\text{belowground-active,belowground-passive}}$), as a function of P_{clay} :

$$HR_{\text{belowground-active}} = \Delta C_{\text{belowground-active}} \cdot (0.17 + 0.68 \cdot P_{\text{clay}}) \quad (18.43)$$

$$F_{\text{belowground-active,belowground-passive}} = \Delta C_{\text{belowground-active}} \cdot (0.003 + 0.032 \cdot P_{\text{clay}}) \cdot f_{\text{anaer}} \quad (18.44)$$

$$F_{\text{belowground-active,belowground-slow}} = \Delta C_{\text{belowground-active}} - HR_{\text{belowground-active}} - F_{\text{belowground-active,belowground-passive}} \quad (18.45)$$

where $f_{\text{anaer}} = (1 + 5 \cdot (1 - f(O_2)))$. Thus, the decomposition rate of the belowground active pool increases and the respiration fraction of carbon losses also increases along with the sand fraction.

18.4.4 Slow belowground pool

The slow belowground pool receives carbon from belowground structural pools, from the slow surface pool as well as from the active belowground pool. The rate of decomposition of the slow belowground pool is:

$$k_{\text{belowground-slow}} = k_{\text{base,belowground-slow}} \cdot f(T) \cdot f(\theta_{\text{rel}}) \cdot f(pH) \cdot f(O_2) \quad (18.46)$$

The loss of carbon from the belowground slow pool ($\Delta C_{\text{belowground-slow}}$) is apportioned between heterotrophic respiration ($HR_{\text{belowground-slow}}$), the carbon flux to back to the belowground active pool ($F_{\text{belowground-slow,belowground-active}}$) and the carbon flux to the belowground passive pool ($F_{\text{belowground-slow,belowground-passive}}$):

$$HR_{\text{belowground-slow}} = \Delta C_{\text{belowground-slow}} \cdot 0.55 \quad (18.47)$$

$$F_{\text{belowground-slow,belowground-active}} = \Delta C_{\text{belowground-slow}} \cdot (1 - (0.003 + 0.009 \cdot P_{\text{clay}}) \cdot f_{\text{anaer}} - 0.55) \quad (18.48)$$

$$F_{\text{belowground-slow,belowground-passive}} = \Delta C_{\text{belowground-slow}} \cdot (0.003 + 0.009 \cdot P_{\text{clay}}) \cdot f_{\text{anaer}} \quad (18.49)$$

where $f_{\text{anaer}} = (1 + 5 \cdot (1 - f(O_2)))$.

18.4.5 Passive belowground pool

The passive belowground pool receives carbon from the slow belowground pool. The rate of decomposition of the passive belowground pool is:

$$k_{\text{belowground-passive}} = k_{\text{base,belowground-passive}} \cdot f(T) \cdot f(\theta_{\text{rel}}) \cdot f(pH) \cdot f(O_2) \quad (18.50)$$

The loss of carbon from the passive slow pool ($\Delta C_{\text{belowground-passive}}$) is apportioned between heterotrophic respiration ($HR_{\text{belowground-passive}}$), the carbon flux to back to the belowground active pool ($F_{\text{belowground-passive,belowground-active}}$):

$$HR_{\text{belowground-passive}} = \Delta C_{\text{belowground-passive}} \cdot 0.55 \cdot f(O_2) \quad (18.51)$$

$$F_{\text{belowground-passive,belowground-active}} = \Delta C_{\text{belowground-passive}} \cdot (1 - 0.55 \cdot f(O_2)) \quad (18.52)$$

Part V

Forest dynamics modelling

Chapter 19

Forest dynamics model

This chapter provides an overview of a forest dynamics model, which builds on the previous models and allows simulating the recruitment, growth, mortality and management of woody plant cohorts in a forest stand. The model was described in De Cáceres et al. (2023) and is run using function `fordyn()` for a set of years.

19.1 Design principles

The design of the forest dynamics model is, to a large degree, inherited from the energy/water balance and growth models it builds on. Readers should refer to former sections to learn the design of the basic water balance (see 3.1), advanced water/energy balance (see 7.2) or growth/mortality processes (see 15.1). Regeneration, from either seed germination and establishment or via resprouting, forest management and snag fall are explicitly simulated at the level of `fordyn()` and hence their design will be the focus of this chapter, with implementation details described in chapter 20. More generally, the processes implemented at the level of `fordyn()` are those that are represented at annual resolution, as opposed to the daily (or subdaily) process resolution of the previous models.

19.1.1 Recruitment from seeds

Recruitment of saplings involves a number of processes (flowering and pollination, fruit/seed production, dispersal, storage, seed predation, germination, seedling establishment and survival until the sapling stage). All these processes have their own biotic and abiotic drivers (Price et al., 2001), so that modelling becomes a challenging task. Processes leading to sapling recruitment are frequently extremely simplified or their mechanisms ignored in many forest models (Price et al., 2001). The design of the forest dynamics model with respect to recruitment follows that of many gap models. Local seed production is considered in a binary way, where plants are considered fertile and able to produce viable seeds if they reach a given height (different for shrubs and trees). Alternatively, the user can specify a set of species whose seeds arrive to the target stand via simulation control parameters or dispersal processes (see chapter 22). A seed bank exists in the forest stand, where seed levels are relative (up to 100%) and result from the interplay between seed rain and seed mortality. Germination is scheduled once a year (assumed to be in winter), and results in age cohorts entering a seedling bank. A constant probability of germination determines is used as a statistical surrogate of processes affecting germination success. Once successfully germinated, the germinant enters the most vulnerable life-stage of a tree, characterized by the highest mortality of the whole plant life-cycle. During the years between germination and recruitment the amount of seedlings decreases due to environmental filtering, depending on a set of regeneration *thresholds*. Typically, regeneration thresholds concern environmental conditions, although some models also consider ungulate browsing (Wehrli et al., 2007). In our case we focus on three environmental drivers limiting the transition from seedlings to saplings:

1. Tolerance to low temperatures, indicated by the mean temperature of the coldest month.

2. Drought tolerance, indicated by the annual moisture index or the daily water potential experienced by seedlings/saplings (Marañón et al., 2004; Lloret et al., 2009).
3. Shade tolerance, indicated by the percentage of photosynthetic active radiation reaching the ground (Marañón et al., 2004).

While aging, seedling cohorts progressively increase their rooting depth, which influences their chances of survival in front of drought (Padilla and Pugnaire, 2007). Aboveground growth is not explicitly modelled for cohorts in the seedling bank. When they reach a pre-specified age, seedling/saplings leave the seedling bank and recruitment as functional cohorts occurs. This entails ingrowth into a given diameter for trees, typically 1 *cm*, or into a given height for shrubs. Maximum recruitment densities and plant size of recruited individuals are specified via species parameters. Importantly, trees recruited are subject to self-thinning processes before attaining the diameter of *adult* ingrowth (typically 7.5 *cm* diameter; see 17.4.1).

19.1.2 Resprouting

Resprouting is a common feature of Mediterranean species. Most, if not all, Mediterranean broadleaved species are able to resprout from buds protected in branches, subterranean structures (e.g. burl, taproot, lateral roots) or in the root collar after a disturbance destroys the aerial part. Yet, some differences in the resprouting ability (i.e. survival) exist regarding the type of disturbance and the species affected.

In medfate, resprouting is assumed happen from buds at the stump/stool level after a given disturbance has led to aboveground plant die-back. Resprouting can occur as a response to different disturbances. For example, resprouting occurs after clipping (e.g. if management is applied), after a fire, or after cavitation has led to dessication of above-ground plant organs.

Regarding disturbance type, differences in survivorship through resprouting mostly depends on the intensity of the disturbance experienced, with an increasing occurrence of mortality after disturbances in the following order: *browsing*, *cutting*, *drought* and *fire* (Espelta et al., 1999). Fire is the disturbance that causes greatest mortality, probably because it may physically destroy part of the bud bank due to lethal temperatures or directly charring it (Pausas et al., 2016). Moreover, resprouting vigor also depends on the size of the plant (Moreira et al., 2012). According to different literature sources, the number of resprouts initially produced is approximately 2 per 1 *cm*² of stump area (Retana et al., 1992; García-Jiménez et al., 2017). Hence, in medfate the final density of resprouts depends on the disturbance that caused resprouting (via survivorship) and on the diameter of the *parent* plant, but in all cases the resprouts inherit the root system of their parent.

Like saplings originated from seeds, resprouts are subject to self-thinning processes before attaining the diameter of ingrowth (see 17.4.1).

19.1.3 Forest management

Forest management is an optional process in `fordyn()` simulations. Furthermore, management actions need to be defined in an external function to be supplied by the user. This design was chosen because there are multiple potential management strategies, and different traditions are followed in different countries. To work properly within `fordyn()` simulations, the supplied management function needs to accept a `forest` object as input, as well as a list of management parameters, and it has to return the reduction in tree density, the reduction in shrub cover and the management parameters to be applied in subsequent calls.

Although the former design allows users to tailor management functions to their simulation needs, the package includes `defaultManagementFunction()` to facilitate the simulation of these process in many situations, and which is briefly described here. The function implements two different management models:

- *Irregular management*: An uneven-aged stand is managed using thinning operations each time a threshold value of a chosen stand-level metric (such as basal area or Hart-Becking index) is trespassed. Thinning operations can focus on trees of specific diameter classes.
- *Regular management*: An even-aged (monospecific) stand is managed in cycles where thinning (preparatory) cuts are followed by final (regeneration) cuts. Thinning operations are similar as those of irregular management. One or several final cuts can be scheduled, the first starting whenever mean

diameter surpasses a chosen threshold, and the following after a chosen number of years. Optionally, tree planting (of a chosen species) can be scheduled to occur after the last final cut.

The former models apply cuts on tree cohorts irrespective of their species. The default management function applies shrub clearing each time there is an operation on trees, resulting in removal of any shrub cover above a given maximum value.

19.1.4 Snag fall

Snag fall based on a simplification of the model by Vanderwel et al. (2006), where 5-yr. probability of snag fall depends on DBH, decay class and species identity. Following the results of Oberle et al. (2018), we defined species identity effects as a function of wood durability. Following the distribution of years of death determined by Aakala et al. (2008) for snags corresponding to different age classes, we defined snag decay class as a function of age.

19.2 State variables

The main state variables of the forest dynamic model are those conforming the structure and composition of the forest stand, i.e. the set of woody cohorts (either trees or shrubs) and their attributes (height, density, DBH, cover, etc.). Additionally, the seed species identity and relative abundance in the seed bank are also state variables of the model. Since the model performs calls to the `growth()` model, many other state variables are defined for intra-annual simulations (see 15.4). When including forest management action, additional state variables may be defined as management parameters.

19.3 Process scheduling

The `fordyn()` model divides the period to be simulated in years, which is the top-level time step of simulations. Given an input `forest` object, the function first initializes the input for function `growth()`. For each year to be simulated the model performs the following steps:

1. Calls function `growth()` to simulate daily water/carbon balance, growth and mortality processes (sub-daily processes may also be involved in `transpirationMode = "Sperry"` or `transpirationMode = "Sureau"`). See section 15.4 for details of growth scheduling.
2. If a management function is supplied, calls this function and apply the resulting reductions in tree density and shrub cover.
3. If required, simulates seed production, seed bank mortality, seed rain, recruitment from seeds and/or resprouting (see chapter 20). In a spatial context, seed rain will include seeds dispersed from other forest stands 22.
4. Removes tree (or shrub) cohorts whose remaining density (resp. cover) is lower than a specified threshold.
5. Merges surviving cohorts with recruitment in the `forest` object and prepares the input of function `growth()` for the next annual time step.
6. Store current status of the `forest` object and update output tables/summaries.

19.4 Inputs and outputs

An important difference between `fordyn()` and the previous simulation functions is that it does not require a specific input object, as in `spwb()` or `growth()` functions. In other words, soil, vegetation, meteorology and control inputs are directly introduced as parameters to the function call to `fordyn()`.

19.4.1 Soil, vegetation and meteorology

Soil

Soil input requirements are the same as for the former models and were fully described in section 2.3.

Vegetation

Unlike the former models, vegetation input for **fordyn** are objects of the class **forest**, which were described in section 2.4.2.

Meteorological input

The minimum weather variables required to run the model are min/max temperatures (T_{min} and T_{max}), min/max relative humidity (RH_{min} and RH_{max}), precipitation (P) and solar radiation (Rad). Wind speed (u) is also needed, but the user may use missing values if not available (a default value will be used in this case). Wind speed is assumed to have been measured at a specific height above the canopy (by default at 2 m). Atmospheric CO_2 concentration (C_{atm}) may also be specified, but if missing a default constant value is assumed, which is taken from the control parameters. Definitions and units of these variables were given in section 2.5.

19.4.2 Vegetation functional parameters

The forest dynamics model requires many functional parameters to be specified for plant cohorts. Some of them depend on whether the basic or advanced water balance is adopted, whereas others are inherited from the growth model. Here we report functional parameters needed in addition to those necessary for the growth model (see 15.5.2).

All of them concern the simulation of recruitment and are specified in the species parameter table (i.e. **SpParams**).

Symbol	Units	R	Description
H_{seed}	cm	SeedProductionHeight	Height for seed production
SM	mg	SeedMass	Seed dry mass
SL	yr	SeedLongevity	Bank average longevity
$Disp_{dist}$	m	DispersalDistance	Dispersal parameter for dispersal kernel
$Disp_{shape}$		DispersalShape	Dispersal parameter for dispersal kernel
P_{recr}	[0-1]	ProbRecr	Probability of recruitment from seeds within the bioclimatic limits imposed by temperature, moisture and light thresholds
TCM_{recr}	°C	MinTempRecr	Minimum average temperature (Celsius) of the coldest month for successful recruitment from seeds
MI_{recr}		MinMoistureRecr	Minimum value of the moisture index (annual precipitation over annual PET) for successful recruitment from seeds
$FPAR_{recr}$	%	MinFPARRecr	Minimum percentage of PAR at the ground level for successful recruitment from seeds
$N_{tree,recr}$	$ind \cdot ha^{-1}$	RecrTreeDensity	Density of tree recruits from seeds.
$N_{tree,ingrowth}$	$ind \cdot ha^{-1}$	IngrowthTreeDensity	Density of trees reaching ingrowth DBH.
$DBH_{tree,recr}$	cm	RecrTreeDBH	DBH for tree recruits from seeds or resprouting (e.g. 1 cm).
$DBH_{tree,ingrowth}$	cm	IngrowthTreeDBH	DBH for trees (e.g. 7.5 cm).
$H_{tree,recr}$	cm	RecrTreeHeight	Height for tree recruits from seeds or resprouting
$Cover_{shrub,recr}$	%	RecrShrubCover	Recruitment cover for shrubs
$H_{shrub,recr}$	cm	RecrShrubHeight	Recruitment height for shrubs
$Resp_{fire}$		RespFire	Number of resprouts per stem after fire disturbance
$Resp_{dist}$		RespDist	Number of resprouts per stem after undefined disturbance (typically desiccation)
$Resp_{clip}$		RespClip	Number of resprouts per stem after clipping

19.4.3 Control parameters

Control parameters modulate the overall behavior of **fordyn** simulations, which extend the parameters used for growth simulations (see section 15.5.3). First, there are parameters that regulate the application of seed production, seed bank dynamics, recruitment from seeds, resprouting and the removal of cohorts with few individuals:

- **applySeedBankDynamics** [= TRUE]: Boolean flag to indicate that seed bank dynamics (seed production, seed bank mortality and seed rain) need to be simulated.
- **applyRecruitment** [= TRUE]: Boolean flag to indicate that recruitment from seeds is allowed.
- **applyResprouting** [= TRUE]: Boolean flag to indicate that resprouting is allowed.
- **recruitmentMode** [= "annual/stochastic"]: String describing how recruitment from seeds is applied. Current accepted values are "annual/deterministic", "annual/stochastic", "daily/deterministic" or "daily/stochastic".
- **removeEmptyCohorts** [= FALSE]: Boolean flag to indicate the removal of cohorts whose density is too low.
- **minimumTreeCohortDensity** [= 1]: Threshold of tree density resulting in cohort removal.
- **minimumShrubCohortCover** [= 0.01]: Threshold of shrub cover resulting in cohort removal.
- **dynamicallyMergeCohorts** [= TRUE]: Boolean flag to indicate that cohorts should be merged when possible. This option speeds up calculations but results in a loss of cohort identity and reinitialization of many state variables.
- **keepCohortsWithID** [= TRUE]: Boolean flag to indicate that cohorts having a non-missing value in a column **ObsID** (if present) should not be merged or removed.

Next, a few parameters control the production/arrival of seeds:

- **seedRain** [= NULL]: Vector of species names whose seed rain is to be added to seed bank, regardless of local seed production.
- **seedProductionTreeHeight** [= 300]: Default minimum tree height for producing seeds (when species parameter **SeedProductionHeight** is missing).
- **seedProductionShrubHeight** [= 30]: Default minimum shrub height for producing seeds (when species parameter **SeedProductionHeight** is missing).

Then we have default parameters determining whether recruitment occurs:

- **probRecr** [= 0.05]: Default annual probability of germination (when species parameter **ProbRecr** is missing).
- **minTempRecr** [= 0]: Default threshold of minimum average temperature of the coldest month necessary for recruiting (when species parameter **MinTempRecr** is missing).
- **minMoistureRecr** [= 0.3]: Default threshold of minimum moisture index (annual precipitation over annual ETP) necessary for recruiting (when species parameter **MinMoistureRecr** is missing).
- **minFPARRecr** [= 10]: Default threshold of minimum fraction of PAR (in %) reaching the ground necessary for recruiting (when species parameter **MinFPARRecr** is missing).

Finally, there are a set of parameters specifying default values for recruited cohort attributes:

- **recrTreeDBH** [= 1]: Default DBH (cm) for recruited trees (when species parameter **RecrTreeDBH** is missing).
- **recrTreeDensity** [= 100]: Default maximum density (ind · ha⁻¹) for recruited trees (when species parameter **RecrTreeDensity** is missing).
- **recrTreeHeight** [= 100]: Default height (cm) for recruited trees (when species parameter **RecrTreeHeight** is missing).
- **recrShrubCover** [= 1]: Default maximum cover (%) for recruited shrubs (when species parameter **RecrShrubCover** is missing).
- **recrShrubHeight** [= 100]: Default height (cm) for recruited shrubs (when species parameter **RecrShrubHeight** is missing).

19.4.4 Model output

Element	Description
StandSummary	A data frame with stand-level summaries (leaf area index, tree basal area, tree density, shrub cover, etc.) at the beginning of the simulation and after each simulated year.
SpeciesSummary	A data frame with species-level summaries (leaf area index, tree basal area, tree density, shrub cover, etc.) at the beginning of the simulation and after each simulated year.
CohortSummary	A data frame with cohort-level summaries (leaf area index, tree basal area, tree density, shrub cover, etc.) at the beginning of the simulation and after each simulated year.
TreeTable	A data frame with tree-cohort data (species, density, diameter, height, etc.) at the beginning of the simulation (if any) and after each simulated year.
DeadTreeTable	A data frame with dead tree-cohort data (species, density, diameter, height, etc.) at the beginning of the simulation and after each simulated year.
CutTreeTable	A data frame with cut tree-cohort data (species, density, diameter, height, etc.) per each simulated year.
ShrubTable	A data frame with shrub-cohort data (species, density, cover, height, etc.) at the beginning of the simulation and after each simulated year.
DeadShrubTable	A data frame with dead shrub-cohort data (species, density, cover, height, etc.) at the beginning of the simulation (if any) and after each simulated year.
CutShrubTable	A data frame with cut shrub-cohort data (species, density, cover, height, etc.) per each simulated year.
ForestStructure	A list with the <code>forest</code> object of the stand at the beginning of the simulation and after each simulated year.
GrowthResults	A list with the results of calling function <code>growth</code> (i.e., see 15.5.4) for each simulated year.
ManagementArgs	If management is considered, a list of management arguments to be used in another call to <code>fordyn()</code> .
NextInputObject	An object of class <code>growthInput</code> to be used in a subsequent simulation.
NextForestObject	An object of class <code>forest</code> to be used in a subsequent simulation.

19.5 Applications

Chapter 20

Regeneration

20.1 Seed production and seed bank dynamics

The model considers **mortality of seeds** in the seed bank before adding new seeds. Annual seed bank mortality is simulated for each species using an exponential decay function, driven by the species-specific seed longevity (SL):

$$Seeds_{t+1} = Seeds_t \cdot \exp(-1/SL) \quad (20.1)$$

To determine seed production, the model determines the **seed rain** in the stand by determining which cohorts have total plant heights above maturity thresholds for trees and shrubs. This is done, for each cohort, by comparing its height with the species-specific parameter seed production height (H_{seed}). Missing values of H_{seed} are given values from control variables **seedProductionTreeHeight** and **seedProductionShrubHeight** for trees and shrubs, respectively. In addition to locally produced seeds, the user can use control parameter **seedRain** to specify a list of species names whose seeds arrive to the stand. Despite the origin, in non-spatial simulations the seed bank relative amount of species with seed rain is set to 100%, assuming that there is enough seed production to allow normal recruitment. However, when simulations are conducted in a spatial context, the species identity and relative amount of seeds is determined via dispersal sub-model (see chapter 22).

20.2 Recruitment from seeds

Actual **sapling recruitment** (i.e. recruitment of small trees, typically 1 cm diameter) depends on environmental conditions in the stand. Specifically, the model calculates, for the year that ended, the mean temperature of the coldest month (TCM), the moisture index (MI) and the fraction of photosynthetic active radiation reaching the ground, given the current structure ($FPAR$). These values are compared to species specific parameter thresholds TCM_{recr} , MI_{recr} and $FPAR_{recr}$, respectively. More specifically, a given species j can recruit only if $TCM > TCM_{recr,j}$, $MI > MI_{recr,j}$ and $FPAR > FPAR_{recr,j}$. These filters do not ensure recruitment, as it is assumed that multiple processes can further determine the death of recruits. A probability of recruitment P_{recr} is used to represent these additional processes. When simulation of recruitment is *stochastic*, a given species will recruit if (in addition to the bioclimatic limits) a Bernoulli draw falls below P_{recr} and, if so, the initial density or cover will be fixed. If recruitment is deterministic, then the probability of recruitment is used to multiply the recruitment density or cover.

Tree recruitment density, diameter and height are determined by parameters $N_{tree,recr}$, $DBH_{tree,recr}$ and $H_{tree,recr}$, respectively; whereas cover and height of shrub recruitment is determined by parameters $Cover_{recr}$ and $H_{shrub,recr}$, respectively. Density/cover of recruits is reduced depending on the relative amount of seeds in the seed bank (i.e. maximum density or cover will occur if seed bank levels are 100%). If stochastic

simulation of recruitment is requested, then density or cover values are considered mean values of a Poisson distribution. Note that density of tree recruits will decrease during the years after recruitment due to a **self-thinning** process, until DBH reaches $DBH_{tree,ingrowth}$ where N will be $N_{tree,ingrowth}$, as explained in 17.4.1.

20.3 Resprouting

Currently, resprouting only occurs in the model if plant cohorts have been **cut**, **burned** or they died of **desiccation** (future model versions will include resprouting after fire impacts). In other words, resprouting does not occur for baseline mortality, self-thinning of recruits or due to starvation.

Resprouting survivorship

The model first determines **resprouting survivorship**, which depends on the disturbance type and, in the case of fire, on species identity. Unfortunately, we lack information on the mortality caused in resprouting by cutting or drought for the vast majority of species except for holm oak (*Quercus ilex*). Thus, we suggest applying tentatively to all species the mortality values obtained in different studies for this species: 2.5% after browsing (Espelta et al., 2006), 4% after cutting (Retana et al., 1992), 5% after drought (Lloret et al., 2004). Larger differences among species have been reported for fire (Espelta et al., 2012). This pattern may be linked to inter-specific differences in the size of the bud bank, the degree of bud protection or the amount of stored resources in belowground organs (burl, taproot). Accordingly, default values for survivorship are $Resp_{clip} = 0.96$ (for cutting) and $Resp_{dist} = 0.95$ (for dessication), whereas no default value is given for $Resp_{fire}$. Naturally, the number of surviving trees (stumps) is given by the number of trees affected by the disturbance multiplied by the survivorship rate:

$$N_{surv} = N_{mort,dist} \cdot Resp_{dist} \quad (20.2)$$

for a generic disturbance, but the same equation would apply for cutting, dessication or fire.

Resprouting vigor

For those individuals that survive the disturbance event and are able to resprout, the number of resprouts present in a given moment is a matter of the stump area and age. According to different literature sources, the number of resprouts initially produced is approximately 1.82 per 1 cm² of stump area, but owing to the intra- and inter-individual competition (self-thinning), the number of resprouts will decrease with time following an exponential equation (Retana et al. 1999. Espelta et al. 1999, 2013):

$$N_{resp} = N_{surv} \cdot \pi \cdot (DBH/2)^2 \cdot 1.82 \cdot 10^{-0.053 \cdot x} \quad (20.3)$$

where DBH is diameter at breast height of the *parent* tree cohort, in cm, $x = 5$ is time in years, N_{resp} is the number of resprouts, N_{surv} is the number of parent trees that survived.

Initial characteristics and self-thinning

Tree resprouts have all a starting diameter $DBH_{tree,recr}$ and a height $H_{tree,recr}$, but the root system is the same of the *parent* tree cohort. Analogously to recruitment from seeds, the initial density of tree resprout will decrease during the years after the initial resprouting due to a **self-thinning** process, until DBH reaches $DBH_{tree,ingrowth}$ where N will be $N_{tree,ingrowth}$, as explained in 17.4.1.

Resprouting in shrubs

In the case of shrub species, the same operations are done, but on cover values instead of density values. Shrub height of resprouts is assumed to be $H_{shrub,recr}$.

Part VI

Landscape/regional modelling

Chapter 21

Watershed hydrology

This chapter describes the distributed watershed water balance sub-model implemented in functions `spwb_land()`, `growth_land()` and `fordyn_land()` of package **medfateland** (Fig. 1.3).

21.1 Design principles

Watersheds are described in raster (i.e. gridded) mode, each cell representing a patch of vegetation (or another land cover) in a catchment. Five land cover types are allowed:

- a. *wildland*: forests, shrublands or grasslands.
- b. *agriculture*: agricultural lands.
- c. *rock*: rock outcrops.
- d. *artificial*: urban areas.
- e. *water*: water bodies.

Watershed water balance simulations extend the simulation of soil water balance to the landscape scale. Hence, the design of most vertical hydrological processes is the same as those of forest water balance. However, cells classified as rock, artificial or water lack any soil and must be treated differently. Additional water compartments and (spatial transport) processes are represented in watershed water balance simulations. Each cell in the watershed may include the following: a snow-pack compartment, one or several soil layers (including a rocky layer down to several meters depth), and a groundwater compartment normally beyond the reach of plant roots for most landscape cells.

Similar to other models such as TETIS (Francés et al., 2007), three lateral flows are considered between adjacent cells:

- (i) Overland surface flows from upper elevation cells to lower elevation ones.
- (ii) Lateral saturated soil flows (i.e. interflow) between adjacent cells.
- (iii) Lateral groundwater flow (i.e. baseflow) between adjacent cells.

Routing of overland surface flows is modeled following T-HYDRO Ostendorf and Reynolds (1993). In large watersheds not all water entering the river channel will end up reaching the watershed outlet the same day, so it is necessary to perform channel routing. If the user specifies a river network, river channel routing is performed by calculating an average flow velocity of the whole channel network, according to Manning's equation. Subsurface flows, including both soil saturated flow and groundwater flow, are modeled following the kinematic wave approximation of DSHVM Wigmosta et al. (1994).

Split-parameter parametrization, following Francés et al. (2007), allows reducing the parameters to be calibrated for practical applications.

21.2 State variables

Distributed simulation of rainfall-runoff processes implies that state variables are defined for each cell of the watershed. For this kind of simulations, state variables describing the water content of different compartments are the most important. In the case of *wildland* and *agriculture* cells these are:

- W_s , the proportion of soil moisture in relation to field capacity for each soil layer s in the cell.
- S_{snow} the snow water equivalent (mm) of the snow pack storage over the cell surface.
- $S_{aquifer}$ the water content (mm) of the cell's aquifer lying over the undisturbed bedrock.

Additional cell state variables in *wildland* cells concern the water status (or other state variables) of plant cohorts in the cell, and they were described in chapters 3 and 7. In the case of cells of other land cover types (i.e. *rock*, *artificial* or *water*) soils are not defined, and neither are soil state variables. In this case, only S_{snow} and $S_{aquifer}$ are defined.

21.3 Water balance

In distributed watershed simulations water balance can be defined both at the cell level and at the watershed level.

21.3.1 Cell level water balance

At the cell level, daily variations in water content can occur in the *snowpack*, *soil* or *groundwater* compartments:

1. Variations in snowpack water equivalent content follows the balance between snowfall (Ps) and snow melt (Sm) eq. (3.2):

$$\Delta S_{snow} = Ps - Sm \quad (21.1)$$

2. Variations in soil water content (ΔV_{soil}) follow (3.6):

$$\Delta V_{soil} = (If + Lw + Cr) - (Dd + Se + Es + Tr_{herb} + Tr_{woody}) \quad (21.2)$$

where the terms were defined in 3.3. Here, the lateral flows (i.e. Lw) are specially important, compared to spatially-uncoupled simulations. They are defined as $Lw = \Delta S_{lat}$, the variation in water content derived from the balance between soil lateral inputs and outputs:

$$Lw = \Delta S_{lat} = S_{lat,in} - S_{lat,out} \quad (21.3)$$

3. Variations in the aquifer water content of a cell are summarized by:

$$\Delta S_{aquifer} = Dd - Cr - Exf - Dal - \Delta S_{base} \quad (21.4)$$

where Dd is deep drainage from the above-lying soil, Cr is capillary rise into the above-lying soil, Exf is the aquifer exfiltration (occurring when the water table reaches the soil surface), Dal is drainage loss to a deeper aquifer (not connected to watershed outlets) and ΔS_{base} is the variation in water content of the aquifer derived from the balance between groundwater lateral inputs and outputs.

$$\Delta S_{base} = S_{base,in} - S_{base,out} \quad (21.5)$$

21.3.2 Watershed water balance

At the watershed level, separate water balances can again be defined for the average water content of snow pack, soil or groundwater compartments. These result from averaging water balances across cells. Additionally, a water balance is defined regarding the overall water content in the watershed.

1. Changes in the average snow pack water equivalent over cells is the result of balancing precipitation as snow (P_s) and snow melt (S_m), both flows averaged over cells:

$$\Delta \hat{S}_{snow} = \hat{P}_s - \hat{S}_m \quad (21.6)$$

where $\hat{P}_s$ and $\hat{S}_m$ are the average snow fall and snow melt over cells.

2. Changes in the average soil moisture ($\Delta \hat{V}_{soil}$) are the result of pooling soil inputs and outputs over cells, which yields:

$$\Delta \hat{V}_{soil} = (\hat{I}f + \hat{C}r) - (\hat{D}d + \hat{S}e + \hat{E}s + \hat{T}r_{woody} + \hat{T}r_{herb}) \quad (21.7)$$

where $\hat{I}f$, $\hat{C}r$, $\hat{D}d$, $\hat{S}e$, $\hat{E}s$, $\hat{T}r_{woody}$ and $\hat{T}r_{herb}$ are the average (over cells) of infiltration, capillary rise, soil evaporation, plant transpiration and herb transpiration, respectively. Lateral saturated soil flows are not included as they cancel out at the watershed level.

3. Changes in the average of cell aquifer water content ($\Delta \hat{S}_{aquifer}$) are the result of balancing averages for deep drainage ($\hat{D}d$) from soils, capillarity rise to soils ($\hat{C}r$), aquifer exfiltration ($\hat{E}xf$) and loss to a deeper aquifer ($\hat{D}al$):

$$\Delta \hat{S}_{aquifer} = \hat{D}d - \hat{C}r - \hat{E}xf - \hat{D}al \quad (21.8)$$

4. If we integrate the three water compartments, water balance at the watershed level is given by:

$$\Delta S_{watershed} = \hat{P}r - (\hat{I}n + \hat{E}s + \hat{T}r_{woody} + \hat{T}r_{herb}) - \hat{E}xp - \hat{D}al \quad (21.9)$$

where $\Delta S_{watershed} = \Delta \hat{S}_{snow} + \Delta \hat{V}_{soil} + \Delta \hat{S}_{aquifer}$ is the change in water content in the watershed and $\hat{E}xp$ is the water exported as runoff to catchment outlet cells (note that the watershed balance equation assumes that all water reaching river channel has reached network outlets).

21.4 Process scheduling

The model can perform water balance simulations on the whole watershed or its subwatersheds, the latter allowing parallelization. If subwatersheds are considered, the results need to be aggregated at the end of water balance simulations.

For every day to be simulated in a watershed or subwatershed, the model performs the following steps:

1. Calculation of soil lateral flows between adjacent cells, according to sub-steps if required by the user (see 21.6.1) and determine the balance of soil lateral flows (i.e. determine ΔS_{lat}) for each cell.
2. Process cell vertical processes, including overland runoff-runon dynamics:
 - a. In cells that lack soil (i.e. *rock*, *artificial* and *water*). For *rock* and *artificial* cells, subsurface flows are not considered, runon and snow pack dynamics is still processed in these cells. In *artificial* cells all runon, rainfall input or snow melt becomes runoff to be passed downhill. In *rock* cells, water can infiltrate directly to the aquifer, where a maximum rate is confronted with the runon/rainfall/snowmelt input. In the case of *water* cells, snow pack dynamics are also processed, but all liquid water (from rain or snow melt) is poured (as deep drainage) down to the aquifer below.
 - b. In cells having soil (i.e. *wildland* or *agriculture*), cells are processed accounting for subsurface flows, runon from upslope cells, and capillarity from the groundwater level, which complete eqs. (21.1) and (21.2). All processes mentioned in 3.4 are taken into account. In cells that are watershed outlet or belong to channel capillarity from the aquifer is excluded to avoid the confusion (and instability) between local soil saturation excess and aquifer exfiltration.
 - c. In any kind of cell, after vertical flows are considered any surface runoff generated is distributed as overland flow to neighbor downhill cells (see 21.6.3). If a cell is part of the channel network, its runoff is instead stored for latter processing of channel routing (see 21.6.4). If a cell does not have downhill neighbors (i.e outlet cell) and is not in a channel, its runoff becomes water exported from the watershed (see 21.6.4).

3. Calculation of groundwater lateral flows between adjacent cells (see 21.6.2) and apply changes in cell aquifer water content due to the balance of groundwater lateral flows (ΔS_{base} in eq. (21.4)) and vertical flows between soils and the aquifer below, i.e. deep drainage and capillarity rise.
4. Apply drainage losses from the aquifer towards a deeper aquifer not connected to watershed outlets (21.6.5), completing eq. (21.4).

Channel routing is optional, depending on whether a channel network has been supplied. Channel routing is performed after water balance simulations of the watershed (or its subwatersheds) are finished. For each day to be simulated and each channel cell, the model adds the water that reached the channel during water balance simulations to the volume of water in the network of the corresponding watershed outlet. After that the model applies channel routing, which can result in a channel backlog of water for the next day (see details 21.6.4).

21.5 Inputs and outputs

21.5.1 Gridded inputs

Hydrological distributed models demand large amounts of data, information and parameters in order to accurately represent the spatial variability of the main hydrological processes and weather inputs. As mentioned in design principles, watersheds are described in raster (i.e. gridded) mode, each cell representing a patch of vegetation (or another land cover) in a catchment. Rasters do not need to be (should not be) complete, since the model is intended to be used on individual catchments with defined boundaries.

Given a set of grid cells representing the watershed at a given spatial resolution, input data for `spwb_land()` and `growth_land()` should be specified using `sf` objects. These were described in section 2.1.2. Each row in the `sf` object has to correspond to a single raster cell, but there can be raster cells that are not described in the `sf` object and, hence, correspond to empty cells.

21.5.2 Watershed meteorology input

Simulation of watershed hydrology requires specifying weather variables for all cells and the simulation period considered. Functions `spwb_land()` and `growth_land()` allow weather to be specified in three ways:

- If a single data frame of daily weather is provided, the model assumes that all watershed cells experience the same weather.
- A different data frame of daily weather may be provided for each cell, using column `meteo` of the `sf` object (see 2.1.2).
- If an interpolator object of class `stars` is provided, the model will call function `interpolate_data()` for every day to be simulated (see `meteoland` package documentation).

21.5.3 Watershed hydraulic control parameters

Distributed hydrological models suffer from the need to specify many parameters. Frances et al. (2007) demonstrated that by using a proper parameter structure, it is possible to obtain an excellent automatic calibration of a distributed conceptual model while maintaining the spatial variability of parameters. In particular, they showed how to define effective model parameters for those quantities normally estimated at the point scale, so that model parameters and estimates usually suffer from being *uncommensurable*. Effective model parameters can in this situation be defined as a *correction function* of the corresponding hydrological characteristic. If the correction function takes into account the model and input errors, the temporal and spatial scale effects and also the hydrological characteristics estimation error, it is reasonable to assume the correction function for each parameter will be common to all cells within the watershed. Frances et al. (2007) suggest using a correction factor for each hydraulic parameter, in a strategy called *split-parameter*. From the calibration point of view, a very important consequence of the split-parameter approach is that the number of variables to be adjusted is reduced dramatically.

Our model follows Frances et al. (2007) in that correction factors are defined for three hydraulic parameters with spatial variation across the watershed:

- $K_{sat,vert}$ saturated soil conductivity for vertical fluxes into, within and out of the soil.
- $K_{sat,lat}$ saturated soil conductivity for soil lateral (horizontal) fluxes.
- $K_{sat,base}$ bedrock conductivity for groundwater lateral flows.

The effective model parameter for each of these is defined for the spatial (depending on the grid resolution) and temporal (daily) scales of the model. Parameters $K_{sat,vert}$ and $K_{sat,lat}$ are split into the input soil saturated conductivity value K_{sat} (in the case of $K_{sat,lat}$, that corresponding to the topsoil), assumed to have been estimated at a given spatial and temporal scale, and the corresponding correction factor, i.e. R_{vert} or R_{lat} . Analogously, $K_{sat,base}$ is split between an input bedrock conductivity value ($K_{bedrock}$, supplied as **bedrock_conductivity** layer) and the correction factor R_{base} . The values for these correction factors should be the target of calibration procedures, assuming that the spatial heterogeneity of effective parameters is adequately covered by the soil and bedrock spatial inputs.

The following table describes the full set of watershed hydraulic control parameters:

Symbol	R param	Description
R_{vert}	R_localflow	Multiplier of soil vertical conductivity.
	interflow	Logical flag to include saturated soil flows in simulations.
R_{lat}	R_interflow	Multiplier of soil lateral conductivity (eq. (21.12)).
n_{lat}	n_interflow	Power exponent regulating the effect of soil saturation level on soil lateral transmissivity (eq. (21.12)).
	baseflow	Logical flag to include groundwater lateral flows in simulations.
R_{base}	R_baseflow	Multiplier of groundwater lateral conductivity (eq. (21.14)).
n_{base}	n_baseflow	Power exponent regulating the effect of aquifer height on groundwater lateral transmissivity (eq. (21.14)).
	n_manning	Manning's roughness coefficient for channel routing calculations.
	free_drainage_outlets	A boolean flag to prevent water table effects into local soil water balance of outlet/channel cells (included for numerical stability).
	num_daily_substeps	Number of daily sub-steps for interflow calculations.
	subwatersheds	A boolean flag to define watershed subunits (useful for parallelization).
	max_overlap	Maximum proportion of overlapping cells for subwatersheds to be considered independent. Lower values will normally produce larger subunits.
	rock_max_infiltration	Maximum infiltration rate for rocky cells ($\text{mm} \cdot \text{day}^{-1}$).
	deep_aquifer_loss	Maximum daily loss rate from watershed aquifer towards a deeper aquifer not connected to outlets ($\text{mm} \cdot \text{day}^{-1}$).

21.5.4 Model outputs

Distributed hydrological models can produce very detailed spatial and temporal outputs, hence consuming lots of memory resources. While the model runs at a daily scale, functions **spwb_land()** and **growth_land()** can be asked to retain spatial outputs and summaries at coarser temporal scales (i.e. weekly, monthly, etc.). The output of functions **spwb_land()** and **growth_land()** is an S3 list with the following data elements:

Element	Description
<code>watershed_control</code>	A list with the watershed control parameters used in the simulation.
<code>sf</code>	An object of class <code>sf</code> containing the geometries, final state and temporal summary for each of the landscape cells (values in mm/day, mm/week, mm/month or mm/year depending on summary frequency). Additionally, it can contain full water balance simulation results for specific cells targetted by the user.
<code>watershed_balance</code>	A data frame with spatially-averaged (over all watershed cells) components of the daily water balance (in mm/day).
<code>watershed_soil_balance</code>	A data frame with spatially-averaged components of the daily water balance over cells that have defined soils (i.e. <i>wildland</i> and <i>agriculture</i> cells) (in mm/day).
<code>channel_export_m3s</code>	A matrix with daily values of runoff (in m ³ /s) reaching each of the channel cells of the landscape (useful for channel processing with an external model).
<code>outlet_export_m3s</code>	A matrix with daily runoff (in m ³ /s) at each of the outlet cells of the landscape. Each outlet drains its own subset of cells (sometimes including channel routing), so the overall daily watershed outflow corresponds to the sum of values in the corresponding row.

Note that while simulations can be performed for sub-watersheds separately, at present the model does not produce results at the sub-watershed level. However, the partitioning of sub-watersheds is included in the output, so that users can cross it with `outlet_export_m3s` to get the discharge generated in each sub-watershed.

21.6 Process details

21.6.1 Soil saturated lateral flow

Soil saturated lateral flows follow the saturated subsurface flow model of Wigmosta et al. (1994), which is based on a kinematic wave approximation. Each grid cell can exchange water with its eight adjacent neighbors.

Local hydraulic gradients are approximated by local ground surface slopes, so that each cell will generally receive water from its upslope neighbors and discharge to its downslope neighbors. The rate of saturated flow from cell i to its downgradient neighbors is equal to (Wigmosta and Lettenmaier, 1999):

$$q_{ij} = -T_i \cdot \tan(\beta_{i,j}) \cdot w_j \quad \beta_{i,j} < 0 \quad (21.10)$$

$$q_{ij} = 0 \quad \beta_{i,j} \geq 0 \quad (21.11)$$

where $q_{i,k}$ is the flux rate (in *m³/day*) from cell i in the direction towards neighbor j , T_i is the transmissivity (in *m²/day*) at cell i , $\beta_{i,j}$ is the water table slope in the direction of j and w_j is the width (in *m*) of cell j . The transmissivity function for soil saturated lateral flux in cell i ($T_{lat,i}$) is specified using:

$$T_{sat,i} = \frac{R_{lat} \cdot K_{sat,i} \cdot Z_{soil,i}}{n_{lat}} \cdot (1 - Z_{sat,i}/Z_{soil,i})^{n_{lat}} \quad (21.12)$$

where $K_{sat,i}$ is the saturated conductivity of the first (top) soil layer in cell i , $Z_{soil,i}$ is the soil depth at cell i , $Z_{sat,i}$ is the depth of saturated water at cell i (see 2.3.4) and n_{lat} is a power exponent regulating the responsiveness of transmissivity to the level of saturation. R_{lat} is the correction factor (a watershed parameter) for the fact that $K_{sat,i}$ is assessed from soil samples, whereas the parameter in the model is defined at the scale of a grid cell (Francés et al., 2007), as explained above.

Soil saturated lateral flux inputs and outputs for a given cell i are balanced to determine increase or decrease in soil moisture:

$$\Delta S_{lat,i} = S_{lat,in} - S_{lat,out} = \sum_j q_{lat,ji} - \sum_j q_{lat,ij} \quad (21.13)$$

Hence, soil layers are not considered separately in the estimation of soil saturated flows and balance. For their source/sink effect on local soil water balance, however, the resulting lateral flow is divided between soil layers according to the water volume present (i.e. layers with larger amounts of water are assumed to exchange water more easily than layers that are drier, regardless of the soil volume they occupy).

21.6.2 Groundwater lateral flow

Like the preceeding, groundwater lateral flows follow the saturated subsurface flow model of Wigmosta et al. (1994), which is based on a kinematic wave approximation. Each grid cell can exchange water with its eight adjacent neighbors.

The transmissivity function for groundwater lateral flow in cell i ($T_{base,i}$) is analogous to the soil saturated lateral flows:

$$T_{base,i} = \frac{R_{base} \cdot K_{bedrock,i} \cdot Z_{bedrock,i}}{n_{base}} \cdot (1 - Z_{aquifer,i}/Z_{bedrock,i})^{n_{base}} \quad (21.14)$$

where $K_{bedrock,i}$ is the bedrock conductivity at cell i , R_{base} is the correction factor for base flows, $Z_{bedrock,i}$ is the depth of the unaltered bedrock, $Z_{aquifer,i}$ is the depth of the aquifer water table and n_{base} is again a power exponent regulating the responsiveness of transmissivity to the aquifer level.

Groundwater lateral flux inputs and outputs for a given cell i are balanced to determine increase or decrease of the aquifer balance:

$$\Delta S_{base,i} = S_{base,in} - S_{base,out} = \sum_j q_{base,ji} - \sum_j q_{base,ij} \quad (21.15)$$

Aquifer update due to lateral flows is coupled to the vertical balance due to deep drainage or capillarity rise from/to the overlying soil. If the depth to aquifer becomes negative (i.e. the aquifer reaches the soil surface) for an outlet or channel cell, then exfiltration (Exf) occurs.

21.6.3 Overland runoff

To simulate surface runoff routing from one cell to the other, the approach of Ostendorf and Reynolds (1993) is used, as in SIERRA (Mouillot et al., 2001). Overland water lateral transport for a given day occurs instantaneously (i.e. no velocities are calculated) and depends on topography only. The model determines cell neighbors following the queen rule (up to eight neighbors per cell). The proportion of surface water runoff of cell i that will be added to the input (runon) of a neighboring cell j is Ostendorf and Reynolds (1993):

$$q_{ij} = \frac{\Delta z_{ij}/L_{ij}}{\sum_j \Delta z_{ij}/L_{ij}} \quad (21.16)$$

if $\Delta z_{ij} = z_i - z_j > 0$, that is, if the difference in elevation between the two cells is positive (i.e. if $z_j < z_i$). Otherwise there is no overland transport from i to j , i.e. $q_{ij} = 0$. L_{ij} indicates the distance between cell i and j (which depends on cell size and on whether the neighboring cell j is diagonal to cell i). The summation of the denominator is done only for neighbors at lower elevation, so that $\sum_i q_{ij} = 1$. The table of q_{ij} values is calculated when initializing distributed watershed objects.

Every day, cells are processed in order from higher to lower elevation. For a given cell i , runoff is (Ru_i) includes infiltration excess and saturation excess (aquifer exfiltration is processed separately when dealing with groundwater flows). This runoff is divided among the neighboring cells at lower elevation. The runon of a neighbor j , Ro_j is updated as:

$$Ro_j = Ro_j + Ru_i \cdot q_{ij} \quad (21.17)$$

Note that a given cell j can receive overland flux from more than one neighbor.

21.6.4 Channel routing and water exported from the watershed

A special situation arises in overland runoff processing when processing cells that do not have downhill neighbors defined (i.e. where $q_{ij} = 0$ for all j), either because: (a) they are in a hollow surface (all neighbors have higher elevations); and/or (b) because they belong to a specified river channel network. In the first case (a), cells are called *outlet cells* in the model, and the runoff they generate contributes directly to watershed runoff exported (*Exp*), without any channel routing process. In the second case (b), cells are called *channel cells*, and channel routing is performed as explained in the next paragraph.

At the time of initialization, the model determines for each channel cell which is the corresponding outlet cell (this is also a *channel*, but located in the boundary of the catchment) and how far it is in terms of the number of cells. Additionally, a numeric vector is created to store the backlog of every channel outlet (i.e. to store the volume of water, in cubic meters, that is in the channel network). Then, during overland runoff processing, the runoff of a channel cell is stored for subsequent processing of channel routing.

Channel routing is performed after water balance simulations of the watershed (or its subwatersheds) are finished. For each day to be simulated and each channel cell, the model first adds the water that reached the channel during water balance simulations to the current backlog of the corresponding watershed outlet. Then, the model processes channel routing for each channel outlet as follows:

1. Find the water stream elevation corresponding to the total volume in the channel network.
2. Determine the wet area, perimeter and average slope (across wet cells) in the channel network.
3. Calculate average stream velocity (m/s) in the channel network, using Manning's equation and `n_manning` control parameter.
4. For every channel network cell whose elevation is lower than the water stream elevation, calculate the corresponding water volume and determine the time to outlet. Time to outlet depends on average stream velocity, the distance in number of channel cells and cell width. If the time to outlet less than one day, add the water volume to the outlet discharge of the current day. If time to outlet is more than one day, this volume will be kept as outlet backlog for the next day to be processed.

21.6.5 Losses towards a deeper aquifer

Daily losses from the aquifer towards a deeper aquifer not connected to watershed outlets are simulated by means of a daily maximum loss rate (which can be specified either via the watershed control parameter called `deep_aquifer_loss` or using an input `sf` column with the same name) that is compared with the current aquifer content. The minimum of those values is subtracted from the current aquifer content.

Chapter 22

Seed dispersal

This chapter describes the seed dispersal sub-model implemented in functions `fordyn_scenario()` and `fordyn_land()` of package **medfateland** (Fig. 1.3).

22.1 Design principles

Given their sessile lifestyle, dispersal is the only process through which plants can spread in a landscape. Different parts of plants can be dispersed (seeds, fruits, branches or even the entire plant), but here it is assumed that seeds are the dispersed elements. Mathematically, dispersal can be addressed via process-based models (e.g. Nathan et al. (2001)). However, it is more commonly addressed via empirical dispersal kernels, which describe the statistical distribution of dispersal distances in a population (Nathan et al., 2012). More specifically, the dispersal kernel is a probability density function describing the distribution of post-dispersal locations relative to a source point. While many different equations can be used and compared to describe dispersal (Nathan et al., 2012), here we focus on the exponential power kernel proposed by Clark et al. (1998), which has shown a good performance compared to other kernels (Bullock et al., 2017) and has been used for other dynamic vegetation models (Snell, 2014). The kernel is isotropic, meaning that is equal in all directions. Therefore, one can consider dispersal either from source or sink locations.

When implementing a dispersal kernel in a dynamic forest simulation model, an important issue to overcome is the fact that forest stands in the target landscape are often separated by distances that are larger (e.g. 500 m, 1 km or even more) than the average area that they represent (normally forest plots have a radius between 10 and 25 m). This entails that there is a uncertainty in the model regarding the presence of seed sources from forest stands at distances between the two scales. In other words, the model does not know whether there are forest patches that can act as seed sources, between a given target forest stand and its nearest neighbors (Fig. 22.1). Ignoring this fact can lead to underestimating colonization. We address this issue by considering that seed sources from unavailable forest stands can be estimated from the set of stands that are represented in the data set, using weights inverse to their distance to the location of the unavailable stand. This is illustrated in Fig. 22.1 below with the area with unavailable stands in grey and distances represented using red and black arrows:

22.2 Process scheduling

Seed dispersal is considered once a year. Process scheduling in the dispersal sub-model is rather straightforward:

1. Mortality of seeds is first performed for all stands (see 20.1).
2. Seed production is simulated in all stands with woody vegetation (see 20.1).
3. For each target forest stand, seed dispersal (including local seed rain) is simulated, leading to a replenishment of its seed bank.




Figure 22.1: Example of sparse forest stand distribution and how to deal with dispersal in this situation. The target stand where seed rain is to be determined is represented using a red dot. Blue dots are the remaining forest stands. The upper figure shows the spatial distribution of stands, whereas the figure below represents the number of stands per distance classes. In both cases, grey is used to identify an area with unavailable data.

22.3 Process details

The exponential power kernel depends on species-specific parameters $Disp_{dist}$ and $Disp_{shape}$ and is calculated for a given distance r using (Clark et al., 1998):

$$f(r) = \frac{1}{N} \cdot \exp \left[- \left(\frac{r}{Disp_{dist}} \right)^{Disp_{shape}} \right] \quad (22.1)$$

where N is the normalizing constant:

$$N = \frac{2 \cdot \pi \cdot Disp_{dist}^2 \cdot \Gamma(2/Disp_{shape})}{Disp_{shape}} \quad (22.2)$$

and $\Gamma()$ is the Gamma function.

Let i be the stand corresponding to seed sink location (e.g. the red dot in Fig. 22.1). We begin by computing the geographic distances between i and all the other stands using geographic coordinates x and y (assuming these are in meters).

$$d(i, j) = \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2} \quad (22.3)$$

Let us now consider a given distance value $r > 0$ from i (e.g. the grey band in Fig. 22.1). We define the probability of choosing a given stand j for a location at this distance from i as follows:

$$p_i(r, j) = \frac{1}{M} \cdot \frac{1}{|r - d(i, j)|} \quad (22.4)$$

where M is an appropriate normalizing factor so that:

$$\sum_{j=1}^n p_i(r, j) = 1 \quad (22.5)$$

Note that this last equation assumes that the density of spatial points is even across space. If not, the sum should be lower than one.

Given the above definitions and assuming that seed bank of a given species should be refilled to 100% if and only if all plots are seed sources for that species, the percentage of seeds of species s reaching the target location i , i.e. $P_{s,i}$, is estimated using:

$$P_{s,i} = 100 \cdot 2\pi \cdot \int_{r>0}^{r_{\max}} r \cdot f(r) \cdot \left(\sum_{j=1}^n I(s, j) \cdot p_i(r, j) \right) \cdot \delta r \quad (22.6)$$

where $r_{\max}$ is the maximum allowed dispersal distance and $I(s, j) \in 0, 1$ is an indicator variable of the availability of seed sources of species s in stand j .

Part VII

Static modules

Chapter 23

Allometric models

The purpose of this chapter is to describe how certain plant structural attributes are estimated in **medfate** using allometric relationships. These static models are used to initialize vegetation input parameters from forest plot data before simulations (in functions **spwbInput()** and **growthInput()**), to update vegetation structure during growth simulations or to calculate foliar biomass for fuel assessments.

23.1 Input data

23.1.1 Forest plot data

As explained in section 2.4, **medfate** has been specially designed to work with forest inventory plots with data in form of plant cohorts. The tree/shrub cohort attributes required to apply allometric models are the following:

Symbol	Units	R	Description	trees	shrubs
SP_i		Species	Species identity	Y	Y
H_i	cm	Height	Average tree or shrub height	Y	Y
N_i	ind · ha ⁻¹	N	Density of tree individuals	Y	N
DBH_i	cm	DBH	Tree diameter at breast height	Y	N
$Cover_i$	%	Cover	Shrub percent cover	N	Y

Most allometric functions require data in form of a **forest** object (see section 2.4.2).

23.1.2 Species parameters

The following table shows the allometric coefficients needed for the calculations detailed in this chapter.

Symbol	Units	R param	Description
$a_{fbt}, b_{fbt}, c_{fbt}$		a_fbt, b_fbt, c_fbt	Coefficients to calculate foliar biomass of an individual tree
a_{ash}, b_{ash}		a_ash, b_ash	Coefficients relating the square of shrub height with shrub area
a_{bsh}, b_{bsh}		a_bsh, b_bsh	Coefficients relating crown volume with dry weight of shrub individuals
$a_{cr}, b_{1cr}, b_{2cr}, b_{3cr}, c_{1cr}, c_{2cr}$		a_cr, b_1cr, b_2cr, b_3cr, c_1cr, c_2cr	Coefficients to calculate crown ratio of trees

Symbol	Units	R param	Description
cr	[0-1]	cr	Ratio between crown length and total height for shrubs
$r_{6.35}$		r635	Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm
SLA	$m^2 \cdot kg^{-1}$	SLA	Specific leaf area

23.2 Allometric relationships

23.2.1 Leaf biomass

Leaf biomass calculations are done differently for trees and shrub cohorts. They are calculated from **forest** objects using function `plant_foliarBiomass()`.

Tree cohorts

Foliar biomass for a single tree of cohort i ($FB_{tree,i}$; in kg) is calculated using:

$$FB_{tree,i} = a_{fbt} \cdot DBH_i^{b_{fbt}} \cdot e^{c_{fbt} \cdot BAL_i} \cdot e^{-0.0001 \cdot N_i} \quad (23.1)$$

where DBH_i is the diameter of the tree (in cm), BAL_i is the cumulative basal area ($m^2 \cdot ha^{-1}$) of trees having an equal or larger diameter (including the target tree), N_i is the density of the cohort, and a_{fbt} , b_{fbt} and c_{fbt} are species-specific regression coefficients. The first two determine the relationship between tree diameter and its foliar biomass, whereas the third one leads to reductions of foliar biomass because of over-shading and self-shading. Factor $e^{-0.001 \cdot N_i}$ reduces foliar biomass for cohorts with very high density (such as very dense regeneration after a fire). The foliar biomass of the whole tree cohort (FB_i ; in $kg \cdot m^{-2}$) is obtained multiplying tree foliar biomass by tree density (N_i ; in $ind. \cdot ha^{-1}$):

$$FB_i = FB_{tree,i} \cdot (N_i/10000) \quad (23.2)$$

Shrub cohorts

To calculate the leaf biomass of a shrub cohort, we first determine $A_{sh,i}$, the area (in cm^2) occupied by one average individual of height H_i (in cm), using the relationship:

$$A_{sh,i} = a_{ash} \cdot H_i^{b_{ash}} \quad (23.3)$$

where a_{ash} and b_{ash} are species-specific parameters. The model then estimates the dry weight of leaves and branches up to 6.35mm in diameter ($B_{sh,i}$, in kg) of this average individual (i.e. fine fuel biomass), using an allometric relationship with shrub crown phytovolume assuming a cylinder (in cm^3):

$$B_{sh,i} = a_{bsh} \cdot (A_{sh,i} \cdot H_i)^{b_{bsh}} \cdot e^{-0.235 \cdot LAI_{trees}^{live}} \quad (23.4)$$

where a_{bsh} and b_{bsh} are species-specific parameters. Here LAI_{trees}^{live} ($m^2 \cdot m^{-2}$) is the sum of leaf area index values of tree cohorts, which induces a reduction of $B_{sh,i}$ similar to the one described for tree foliar biomass by BAL_i . This reduction is important for biomass and leaf area estimation of shrubs in forests, since it has been observed that light availability affects the growth, architecture and bulk density of the understory (Pimont et al., 2018).

Shrub density (N_i ; in $ind. \cdot m^{-2}$) can be grossly estimated from percent cover (C_i , in percent) and $A_{sh,i}$ (in cm^2):

$$N_i = \frac{C_i/100}{A_{sh,i}/10000} \quad (23.5)$$

The fine fuel biomass of a shrub cohort (W_i , in $kg \cdot m^{-2}$) is simply the product of $B_{sh,i}$ (kg of dry weight) and N_i :

$$W_i = B_{sh,i} \cdot N_i \quad (23.6)$$

Foliar biomass (in $kg \cdot m^{-2}$) can be obtained using the species-specific ratio $r_{6.35,i}$:

$$FB_i = W_i / r_{6.35,i} \quad (23.7)$$

If not known, $r_{6.35,i}$ can be set to a default value of 2 (equivalent to 50% of weight corresponding to leaves).

Herbaceous layer

Foliar biomass of the herbaceous layer (which is equal to its fine fuel loading) is estimated from herbaceous cover ($Cover_{herb}$; in %) and height (H_{herb} ; in cm) assuming a constant bulk density coefficient:

$$FB_{herb} = 0.014 \cdot Cover_{herb} \cdot (H_{herb}/100.0) \cdot e^{-0.235 \cdot LAI_{woody}^{live}} \quad (23.8)$$

where the factor $e^{-0.235 \cdot LAI_{woody}^{live}}$ is analogous to that of shrubs (eq. (23.4)), but using the overall leaf area index of woody cohorts as driving variable.

23.2.2 Leaf area and LAI

Leaf area index (LAI in $m^2 \cdot m^{-2}$) of a given cohort i can be calculated from its foliar biomass (in $kg \cdot m^{-2}$) by using a *specific leaf area* coefficient (SLA , in $m^2 \cdot kg^{-1}$) that in **medfate** is species-specific:

$$LAI_i = FB_i \cdot SLA \quad (23.9)$$

The leaf area (LA in m^2) of an individual of the cohort is found by dividing it by plant density (N_i , in $ind. \cdot ha^{-1}$) and multiplying by 10000 $m^2 \cdot ha^{-1}$:

$$LA_i = 10000 \cdot LAI_i / N_i \quad (23.10)$$

LAI values of woody cohorts are calculated from **forest** objects using function **plant_LAI()**.

Leaf area index of the herbaceous layer (LAI_{herb}) is estimated from FB_{herb} by applying a $SLA = 9$ value, is limited to $LAI_{herb} = 2 m^2 \cdot m^{-2}$, and can be calculated from **forest** objects using function **herb_LAI()**.

23.2.3 Crown vertical dimensions

Crown base height, i.e. the height where the first living branch of trees or shrubs occurs, is an important parameter to know the vertical distribution of leaves (see section 2.4.3.3). Crown base height of a plant cohort i (i.e., $H_{crown,i}$, in cm) is related to the total height of the cohort (H_i , in cm) and its crown ratio (CR_i), which is the ratio between crown length and plant height:

$$H_{crown,i} = (1 - CR_i) \cdot H_i \quad (23.11)$$

In the case of shrubs the crown ratio CR_i is normally an input (species-specific) parameter (cr). In the case of trees, the crown ratio is modelled as a function of tree size and stand competition, following a modification of the logistic equation of Hasenauer and Monserud (1996) :

$$CR_i = \frac{1}{1 + e^{-(a_{cr} + b_{1cr} \cdot HD_i + b_{2cr} \cdot (H_i/100) + b_{3cr} \cdot DBH_i^2 + c_{1cr} \cdot BAL_i + c_{2cr} \cdot \ln(CCF_i))}} \quad (23.12)$$

where $HD_i = H_i / (100 \cdot DBH_i)$ is the height to diameter ratio (in $m \cdot cm^{-1}$), H_i is the tree height, DBH_i is the diameter, CCF_i is the crown competition factor and a_{cr} , b_{1cr} , b_{2cr} , b_{3cr} , c_{1cr} and c_{2cr} are species-specific parameters. The crown competition factor is in turn calculated using (Krajicek et al., 1961):

$$CCF_i = \sum_i N_i \cdot \pi \cdot (CW_i/2)^2 / 100 \quad (23.13)$$

where N_i is the tree density and CW_i is the crown width (in m) assuming an open-grown tree, estimated from an allometric relationship with tree diameter:

$$CW_i = a_{cw} \cdot DBH_i^{b_{cw}} \quad (23.14)$$

where again a_{cw} and b_{cw} are species-specific parameters.

Crown ratios can be calculated from forest objects using function `plant_crownRatio()`. Similar calculations are performed to estimate crown length and crown base height, with functions `plant_crownLength()` and `plant_crownBaseHeight()`, respectively.

Chapter 24

Post-simulation procedures

24.1 Water use efficiency

Chapter 25

Wind extinction

In this chapter we describe a few wind extinction models that are implemented in **medfate**. Wind extinction is relevant: (a) to determine convective heat exchanges between the soil and the canopy; (b) to estimate wind speed and energy balance at the leaf level for different plant cohorts; and (c) for fire behaviour calculations.

25.1 Wind speed at the top of the canopy

Input wind speed (u) is assumed to represent the speed of wind at 6 m (20 feet) above the canopy. Following Albini and Baughman (1979) the wind speed (in $m \cdot s^{-1}$) at the top of the canopy is:

$$u_{top} = \frac{(1.01857 \cdot u) \cdot 0.4265092 \cdot H_{canopy}}{\log(20 + 1.181102 \cdot H_{top})} \quad (25.1)$$

where H_{top} is the canopy top height in m .

25.2 Wind extinction profile

The wind extinction profile, i.e the wind speed at any height z in m , can be calculated following Massman (1987):

$$\beta_{stand} = \frac{4.0 \cdot 0.2 \cdot LAI_{stand}^{phi}}{0.16 \cdot 1.5^2} \quad (25.2)$$

$$u(z) = u_{top} \cdot \sqrt{\frac{\cosh(\beta_{stand} \cdot z / H_{top})}{\cosh(\beta_{stand})}} \quad (25.3)$$

Chapter 26

Fuel characteristics and fire behaviour

26.1 Overview

Functions `fuel_FCCS()` and `fire_FCCS()` allow calculating potential fire behaviour for forest inventory plots. Formulation of fuel characteristics and fire behaviour is an adaptation of the Fuel Characteristics Classification System [FCCS; Prichard et al. (2013)]. In FCCS, fuelbed is divided into six strata, including canopy, shrub, herbaceous vegetation, dead woody materials, leaf litter and ground fuels. All except ground fuels are considered here. The intensity of burning depends on several factors, including topography, wind conditions, fuel structure and its moisture content, which is determined from antecedent and current meteorological conditions. A modification of the Rothermel's (1972) model is used to calculate the intensity of surface fire reaction (in kW/m^2) and the rate of fire spread (in m/min) of surface fires assuming a steady-state fire. Both quantities are dependent on fuel characteristics, windspeed and direction, and topographic slope and aspect.

Fuel and fire behavior calculations provide the following results:

- 1) Fuel characteristics by stratum.
- 2) Surface fire behavior (i.e. reaction intensity, rate of spread, fireline intensity and flame length);
- 3) Crown fire behavior.
- 4) Fire potential ratings of surface fire behavior and crown fire behavior.

The following figure provide an overview of the steps to calculate surface/crown fire behavior and fire potentials from fuel characteristics.

26.2 Input data

26.2.1 Forest plot data

As explained in section 2.4, **medfate** has been specially designed to work with forest inventory plots. The tree/shrub cohort attributes required to apply allometric models are same as for chapter 23:

Symbol	Units	R	Description	trees	shrubs
SP_i		Species	Species identity	Y	Y
H_i	<i>cm</i>	Height	Average tree or shrub height	Y	Y
N_i	<i>ind · ha⁻¹</i>	N	Density of tree individuals	Y	N
DBH_i	<i>cm</i>	DBH	Tree diameter at breast height	Y	N
$Cover_i$	%	Cover	Shrub percent cover	N	Y

Cohorts are not distinguished for the herbaceous stratum, and the variables needed are:




Figure 26.1: Overview of the steps involved in fire behavior and fire potential calculations starting from fuel properties

Symbol	Units	R	Description
C_{he}	%	<code>herbCover</code>	Herbaceous percent cover
H_{he}	cm	<code>herbHeight</code>	Mean herb height

Finally, the model also requires the percent cover of trees in the canopy (C_{ca}). This is easily available from forest inventory data, but could also be derived from the description of tree cohorts.

Fire behaviour functions require data in form of a `forest` object (see section 2.4.2).

26.2.2 Species parameters

The following functional parameters are required for each species:

Symbol	Units	R	Description
L_{shape}	Categorical	<code>LeafShape</code>	Leaf shape: “Broad”, “Needle”, “Linear”, “Scale”, “Spines” or “Succulent”
L_{size}	Categorical	<code>LeafSize</code>	Leaf size: “Small” (< 225 mm), “Medium” (> 225 mm & < 2025 mm) or “Large” (> 2025 mm)
$r_{6.35}$		<code>r635</code>	Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm
ρ_{wood}	$g \cdot cm^{-3}$	<code>WoodDensity</code>	Density of wood tissue
ρ_{leaf}	$g \cdot cm^{-3}$	<code>LeafDensity</code>	Density of leaf tissue
σ_i	$m^2 \cdot m^{-3}$	<code>SAV</code>	Surface-area-to-volume ratio of the small fuel (1h) fraction (leaves and branches < 6.35mm)
h	$kJ \cdot kg^{-1}$	<code>HeatContent</code>	High fuel heat content.
LD	years	<code>LeafDuration</code>	Leaf duration (in years)
LI	%	<code>PercentLignin</code>	Percentage of lignin in leaves

Leaf shape and size categories are used to determine leaf litter fuel types.

26.2.3 Other inputs

Other inputs may be given by expert opinion or they may be calculated from another model. Specifically, for each plant cohort (and for any day of application) the fire behaviour model requires:

- P_{dead} : Proportion of the plant that is dead.
- PMC : Fuel moisture content (in percent of dry weight).

Analogously, the same variables are needed for the herbaceous stratum.

- $P_{dead,he}$: Proportion of herb fuels that respond to humidity changes as 1-h dead fuels.
- $PMC_{live,he}$: Fuel moisture content of live herb fuels (in percent of dry weight).

The model also needs the following input parameters:

- PMC_{dead} : the moisture content of 1-h dead fuels (in percent of dry weight).
- U : Midflame windspeed (in $m \cdot s^{-1}$).
- S : Slope (in percent).

26.3 Fuel characteristics

26.3.1 Fuel strata

The Fuel Characteristics Classification System (FCCS) on which this document is based, defines six fuel strata (Prichard et al., 2013):

- *Canopy*: Trees, snags and ladder fuels.
- *Shrubs*: Primary and secondary layers.
- *Non-woody vegetation (herbs)*: grasses, sedges, rushes and forbs.
- *Woody fuels*: All downed and dead wood, sound wood, rotten wood and stumps.
- *Litter-lichen-moss*: Lichen, litter and moss layers.
- *Ground fuels*: Duff, basal accumulation and squirrel middens.

Shrubs, herbs and woody fuels are constitute the **upper surface fuels**, whereas herbs and woody fuels alone constitute the **lower surface fuels**. FCCS summarizes and calculates characteristics for each fuelbed stratum and layer. Our model estimates fuel loading and characteristics for canopy, shrub, non-woody vegetation, as well as fine (1h) woody fuels and litter fuels. Larger woody fuels (10h or 100h) could be considered if information about forest management actions is available. Ground fuels are not included here.

26.3.2 Cohort fuel loading

Here we consider as burnable fuels foliage and branches up to 6.35 mm = 0.25 in in diameter. The same consideration applies to both trees and shrubs. They are calculated from **forest** objects using function `plant_fuelLoading()`.

Tree cohorts

Fine fuel loading for a tree cohort (W_i ; in $kg \cdot m^{-2}$), including its leaves and branches with diameter up to 6.35 mm = 0.25 in, is calculated from foliar biomass (FB_i , see eq. (23.2)) using:

$$W_i = r_{6.35,i} \cdot FB_i \quad (26.1)$$

where $r_{6.35,i}$ is the ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm in diameter for the species of cohort i . The biomass corresponding to branches of less than < 6.35 mm (SBB_i , also in $kg \cdot m^{-2}$) is obtained by subtraction:

$$SBB_i = (r_{6.35,i} - 1) \cdot FB_i \quad (26.2)$$

Whereas W_i is the cohort loading variable influencing fire behavior, FB_i and SBB_i are cohort variables used to estimate fine dead woody and leaf litter loadings.

Shrub cohorts

Our procedure to estimate shrub fuel loading differs from Prichard et al. (2013) because they calculate first total biomass of the shrub species and then consider the percentage of total weight that corresponds to leaves

and small branches. In our case, we estimate fine fuel loading (W_i , in $kg \cdot m^{-2}$) and foliar biomass (FB_i , in $kg \cdot m^{-2}$) of shrubs from eqs. (23.6) and (23.7). Biomass of small branches (in $kg \cdot m^{-2}$) can be obtained from :

$$SBB_i = W_i - FB_i \quad (26.3)$$

26.3.3 Vertical distribution of cohort fuels

Vertical distribution of fine fuels are distributed between the crown base height ($H_{crown,i}$; in cm) and the total height (H_i , in cm) following a **truncated Gaussian distribution**, as done for the distribution of leaves (see section 2.4.3.3). Crown base height of trees is calculated as explained in 23.2.3. The loading of a cohort that occurs within a given height interval of limits H_1 and H_2 is calculated as:

$$W_i(H_1, H_2) = W_i \cdot p_i(H_1, H_2) \quad (26.4)$$

where $p_i(H_1, H_2)$ is the proportion of the crown of cohort i that corresponds to the height interval (H_1, H_2).

26.3.4 Fuel bulk density profile

Knowing at which height fuels are placed, the **fuel bulk density profile** (Reinhardt et al., 2006) is defined for any given interval (H_1, H_2) as the bulk density (kg/m^3) of fine fuels corresponding to that interval:

$$BDP(H_1, H_2) = \frac{\sum_i W_i(H_1, H_2)}{H_2 - H_1} \quad (26.5)$$

Canopy bulk density normally ranges between 0 and $0.4 kg/m^3$ (Scott and Reinhardt, 2002). Sando and Wick (1972) arbitrarily defined canopy base height as the lower vertical 0.3-m section with a weight greater than $0.01124 kg/m^3$. A user-defined threshold t_{BDP} (in kg/m^3) in 0.1-m sections is used to differentiate the surface fuelbed from canopy fuels. Using this threshold the model calculates the following three heights (Reinhardt et al., 2006):

- *Shrub stratum base height*, H_{sb} (in cm): the minimum height between 0 and 2 m where fuel bulk density is larger than t_{BDP} .
- *Shrub stratum top height*, H_{st} (in cm): the maximum height between 0 and 2 m where fuel bulk density larger than t_{BDP} . With this definition h_s cannot be higher than 2 m (corresponding to fuel model 4 in Anderson 1982).
- *Canopy base height*, H_{cb} (in cm): In terms of its consequences to crown fire initiation, canopy base height can be defined as the lowest height above the ground at which there is sufficient canopy fuel to propagate fire vertically through the canopy. It is calculated as the minimum height over H_{st} when the bulk density starts again to be larger than t_{BDP} .
- *Canopy top height*, H_{ct} (in cm): the maximum height where bulk density is larger than t_{BDP} .
- *Canopy gap*, H_{gap} (in cm): the difference between H_{cb} and H_{st} . The canopy gap is used to calculate crown initiation potential.

Figure 26.2 illustrates the definition and analysis of the fuel bulk density profile for a given forest stand (see function `vprofile_fuelBulkDensity()`). Following Mitsopoulos and Dimitrakopoulos (2007), a threshold $t_{BDP} = 0.04$ is used to determine shrub and canopy heights.

26.3.5 Fuel loading (w) and fuel depth (δ)

26.3.5.1 Canopy stratum

Canopy loading (in $kg \cdot m^{-2}$) is the sum of (tree and shrub) cohort loadings above 2 m (i.e. 200 cm):

$$w_{ca} = \sum_i w_{i,ca} = \sum_i W_i(200, \infty) \quad (26.6)$$




Figure 26.2: Bulk density profile of an example forest

where $w_{i,ca}$ is the canopy stratum loading of cohort i . Canopy depth (in m) is defined as the average of tree (or shrub) crown lengths above 2 m, weighted by the loadings of cohorts in the canopy:

$$\delta_{ca} = \frac{1}{100} \cdot \frac{\sum_i w_{i,ca} \cdot (H_i - H_{b,i}) \cdot p_{i,ca}}{\sum_i w_{i,ca}} \quad (26.7)$$

where the proportion of a tree (or shrub) cohort in the canopy stratum is $p_{i,ca} = p_i(200, \infty)$.

26.3.5.2 Shrub stratum

Shrub loading (in $kg \cdot m^{-2}$) is the sum of (tree and shrub) cohort loadings between the ground and 2 m (i.e. 200 cm):

$$w_{sh} = \sum_i w_{i,sh} = \sum_i W_i(0, 200) \quad (26.8)$$

where $w_{i,sh}$ is the shrub stratum loading of cohort i . The depth of the shrub stratum (in m) is defined as the average of tree (or shrub) crown lengths below 2 m, weighted by the loadings of cohorts in the shrub stratum:

$$\delta_{sh} = \frac{1}{100} \cdot \frac{\sum_i w_{i,sh} \cdot (H_i - H_{b,i}) \cdot p_{i,sh}}{\sum_i w_{i,sh}} \quad (26.9)$$

where the proportion of a shrub (or tree) cohort in the shrub stratum is $p_{i,sh} = p_i(0, 200)$.

26.3.5.3 Non-woody stratum

Herb percent cover and average herb height are transformed into herbaceous loading ($kg \cdot m^{-2}$) using (ref piropinos):

$$w_{he} = 0.014 \cdot C_{he} \cdot (H_{he}/100) \quad (26.10)$$

The depth of the herbaceous stratum (in m) is simply the mean height of herbs:

$$\delta_{he} = H_{he}/100 \quad (26.11)$$

26.3.5.4 Woody and litter strata

In FCCS (Prichard et al., 2013), woody surface loading includes several fuel sizes. However, when calculating surface fire behavior w_{wo} includes 100% of 1h fuels, 25% of 10h fuels and 12.5% of 100h fuels, which represents the material available for flaming combustion. Obtaining loading estimates for 10h- and 100h-fuels is very difficult without field fuel sampling. However, we might estimate 1h woody fuels and leaf litter from standing biomass of small branches ($< 6.35\text{mm}$) and leaves for trees and shrubs. Hence, our treatment of surface woody fuels includes only fine (1h) fuels.

Assuming a continuous input of litter, the variation in accumulated litter is described by a simple differential equation (Birk and Simpson, 1980):

$$\frac{dX}{dt} = L - k \cdot X \quad (26.12)$$

where k is the decay constant, L is the rate of litterfall and X is the litter mass accumulated in the forest floor. Assuming that litter mass has reached a steady state, X can be estimated as the ratio between L and k . If litterfall is estimated as the total foliar biomass divided by leaf duration, the amount of steady state leaf litter corresponding to each tree and shrub cohort can be estimated using:

$$w_{li,i} = \frac{FB_i}{LD(SP_i) \cdot k_i} \quad (26.13)$$

where FB_i is the foliar biomass of cohort i , $LD(SP_i)$ is the species-specific average leaf duration (in years) and k_i is the rate of decay of leaves of cohort i , which is given by the regression model of Meentemeyer (1978):

$$k_i = (-0.5365 + 0.00241 \cdot AET) - (-0.01586 + 0.000056 \cdot AET) \cdot LI(SP_i) \quad (26.14)$$

where $LI(SP_i)$ is the species-specific percentage of lignin content in leaves and AET is actual evapotranspiration (default $AET = 1000\text{mm}$). Litter loadings are summed for four litter types (short pine needles, long pine needles, other conifers, broadleaves). In the case of fine dead woody materials (small fallen branches), loading of small branches is taken as woody litter and it is assumed that small branch litterfall occurs at the same time as leaf litterfall (i.e. according to leaf duration):

$$w_{wo} = \sum_i w_{wo,i} = \sum_i \frac{SBB_i}{LD(SP_i) \cdot k_{wo}} \quad (26.15)$$

where $k_{wo} = 0.95\text{y}^{-1}$ is a constant rate of decomposition for small branches.

In FCCS, the depth of woody and LLM strata are inputs. In our case the depth of the woody and litter strata are estimated from the corresponding fuel loadings:

$$\delta_{wo} = w_{wo} / \rho_{b,wo} \quad (26.16)$$

$$\delta_{li} = w_{li} / \rho_{b,li} \quad (26.17)$$

where $\rho_{b,wo}$, $\rho_{b,li}$ are the woody and litter bulk density (in $\text{kg} \cdot \text{m}^{-3}$), respectively. Litter bulk density $\rho_{b,li}$ is calculated as a weighted average of litter types:

$$\rho_{b,li} = \frac{\sum_k \rho_{b,k} \cdot w_{li,k}}{\sum_k w_{li,k}} \quad (26.18)$$

where k indicates litter type. The bulk density for litter types are [Prichard et al. (2013); Table 1]:

$$\rho_{b,shortneedlepine} = \rho_{b,longneedlepine} = \rho_{b,otherconifer} = 1.65\text{lb} \cdot \text{ft}^{-3} = 26.43\text{kg} \cdot \text{m}^{-3} \quad (26.19)$$

$$\rho_{b,hardwood} = 0.83\text{lb} \cdot \text{ft}^{-3} = 13.30\text{kg} \cdot \text{m}^{-3} \quad (26.20)$$

26.3.6 Other fuel characteristics

All the following characteristics are calculated in metric units (although British units are indicated to qualify specific values for compatibility).

26.3.6.1 Particle density (ρ_p)

Particle density is the ratio of dry weight over volume for fuel particles (in $kg \cdot m^{-3}$). When species have different values, particle density averages for shrub and canopy strata can be obtained as:

$$\rho_{p,sh} = \frac{\sum_i w_{i,sh} \cdot \rho_p(SP_i)}{\sum_i w_{i,sh}} \quad (26.21)$$

$$\rho_{p,ca} = \frac{\sum_i w_{i,ca} \cdot \rho_p(SP_i)}{\sum_i w_{i,ca}} \quad (26.22)$$

where $\rho_p(SP_i)$ is the species-specific particle density (in $kg \cdot m^{-3}$), which can be obtained from wood tissue (ρ_{wood}) and leaf tissue (ρ_{leaf}) densities, using $r_{6.35}$ to weight them:

$$\rho_p(SP_i) = 1000 \cdot (\rho_{leaf}(SP_i) \cdot f_{leaves,vol} + \rho_{wood}(SP_i) \cdot (1 - f_{leaves,vol})) \quad (26.23)$$

where $f_{leaves,vol}$ is the volumetric fraction of leaves with respect to branchlets:

$$f_{leaves,vol} = \left[1.0 + (r_{6.35}(SP_i) - 1.0) \cdot \frac{\rho_{wood}(SP_i)}{\rho_{leaf}(SP_i)} \right]^{-1} \quad (26.24)$$

Dead woody and litter particle densities, i.e. $\rho_{p,wo}$ and $\rho_{p,li}$, are obtained averaging wood tissue and leaf tissue densities across cohorts:

$$\rho_{p,wo} = \frac{\sum_i w_i \cdot \rho_{wood}(SP_i)}{\sum_i w_i} \quad (26.25)$$

$$\rho_{p,li} = \frac{\sum_i w_i \cdot \rho_{leaf}(SP_i)}{\sum_i w_i} \quad (26.26)$$

$$(26.27)$$

Finally, herb particle density, $\rho_{p,he}$, is set to a default value $\rho_{p,he} = 400 kg \cdot m^{-3} = 25 lb \cdot ft^{-3}$ (Prichard et al., 2013).

26.3.6.2 Particle volume (PV)

Particle volume is defined as the volume of particles per surface area (in $m^3 \cdot m^{-2}$). Is calculated as dry weight loading divided by particle density. If species have different particle density values, the particle volume for canopy (PV_{ca}) and shrub (PV_{sh}) strata can be calculated using:

$$PV_{ca} = \sum_i PV_{i,ca} = \sum_i w_{i,ca} / \rho_p(SP_i) \quad (26.28)$$

$$PV_{sh} = \sum_i PV_{i,sh} = \sum_i w_{i,sh} / \rho_p(SP_i) \quad (26.29)$$

where $PV_{i,ca}$ and $PV_{i,sh}$ are the particle volume of cohort i in the canopy and shrub strata, respectively. The particle volume for woody and herb strata are simply:

$$PV_{wo} = w_{wo} / \rho_{p,wo} \quad (26.30)$$

$$PV_{he} = w_{he} / \rho_{p,he} \quad (26.31)$$

The particle volume for the litter stratum is the sum of particle volume of litter components:

$$PV_{li} = \sum_i PV_{li,k} = \sum_i w_{li,k} / \rho_{p,li} \quad (26.32)$$

26.3.6.3 Packing ratio (β)

The proportion of fuelbed stratum volume occupied by fuel particles is an important factor to predict fire behavior. At low packing ratios (low particle density) fire intensity is limited by excessive heat loss. At high

packing ratios (high particle density), lack of oxygen limits combustion. The packing ratios for the canopy and shrub stratum (β_{ca} and β_{sh} ; dimensionless) are given by:

$$\beta_{ca} = \frac{PV_{ca}}{\delta_{ca}} \quad (26.33)$$

$$\beta_{sh} = \frac{PV_{sh}}{\delta_{sh}} \quad (26.34)$$

where $w_{i,ca}$ and $w_{i,sh}$ are the contribution of cohort i to canopy and shrub strata loading (in $kg \cdot m^{-2}$), respectively, and $\rho_p(SP_i)$ is the particle density (in $kg \cdot m^{-3}$) of fuels in cohort i . The packing ratio for the herbaceous, woody and litter strata are:

$$\beta_{he} = \frac{PV_{he}}{\delta_{he}} \quad (26.35)$$

$$\beta_{wo} = \frac{PV_{wo}}{\delta_{wo}} = \frac{\rho_{b,wo}}{\rho_{p,wo}} \quad (26.36)$$

$$\beta_{li} = \frac{PV_{li}}{\delta_{li}} = \frac{\rho_{b,li}}{\rho_{p,li}} \quad (26.37)$$

Note that the packing ratio expressions for woody and litter strata as a ratio of bulk and particle density arises as a consequence of how fuel depth and particle volume are estimated.

26.3.6.4 Surface-area-to-volume ratio (σ)

The surface-area-to-volume ratio (in $m^2 \cdot m^{-3}$) for the canopy or shrub strata are calculated using weighted averages:

$$\sigma_{ca} = \frac{\sum_i w_{i,ca} \cdot \sigma(SP_i)}{\sum_i w_{i,ca}} \quad (26.38)$$

$$\sigma_{sh} = \frac{\sum_i w_{i,sh} \cdot \sigma(SP_i)}{\sum_i w_{i,sh}} \quad (26.39)$$

where $w_{i,ca}$ and $w_{i,sh}$ are the contribution of cohort i to canopy and shrub strata loading (in $kg \cdot m^{-2}$), respectively, and $\sigma(SP_i)$ is the species-specific surface-area-to-volume ratio. The surface-area-to-volume ratio of herbs is assumed constant $\sigma_{he} = 11483m^2 \cdot m^{-3} = 3500ft^2 \cdot ft^{-3}$ and that of small (1-h) woody fuels is $\sigma_{wo} = 1601.05m^2 \cdot m^{-3} = 488ft^2 \cdot ft^{-3}$. The surface-area-to-volume ratio for the litter stratum is:

$$\sigma_{li} = \frac{\sum_k w_{li,k} \cdot \sigma_k}{\sum_k w_{li,k}} \quad (26.40)$$

and the surface-area-to-volume ratio for litter types are:

$$\sigma_{shortneedlepine} = 6562m^2 \cdot m^{-3} = 2000ft^2 \cdot ft^{-3} \quad (26.41)$$

$$\sigma_{longneedlepine} = 4921m^2 \cdot m^{-3} = 1500ft^2 \cdot ft^{-3} \quad (26.42)$$

$$\sigma_{otherconifer} = 8202m^2 \cdot m^{-3} = 2500ft^2 \cdot ft^{-3} \quad (26.43)$$

$$\sigma_{hardwood} = 8202m^2 \cdot m^{-3} = 2500ft^2 \cdot ft^{-3} \quad (26.44)$$

26.3.6.5 Fuel area index (FAI)

The fuel area index (FAI) is the total fuel surface area per unit of ground area (unitless). It is analogous to leave area index (LAI), and it is used to calculate FCCS fire potentials (Schaaf et al., 2007). For shrub and canopy strata, FAI is calculated as:

$$FAI_{ca} = \sum_i FAI_{i,ca} = \sum_i PV_{i,ca} \cdot \sigma(SP_i) \quad (26.45)$$

$$FAI_{sh} = \sum_i FAI_{i,sh} = \sum_i PV_{i,sh} \cdot \sigma(SP_i) \quad (26.46)$$

where $FAI_{i,ca}$ and $FAI_{i,sh}$ are the FAI of cohort i in the canopy and shrub strata, respectively. The FAI of herbs and woody strata are given by:

$$FAI_{he} = PV_{he} \cdot \sigma_{he} \quad (26.47)$$

$$FAI_{wo} = PV_{wo} \cdot \sigma_{wo} \quad (26.48)$$

For the litter layer, FAI is calculated as a sum of FAI for litter components:

$$FAI_{li} = \sum_k FAI_{li,k} = \sum_k PV_{li,k} \cdot \sigma_k \quad (26.49)$$

26.3.6.6 Moisture content (FMC)

Fuel moisture content (FMC in percent of dry weight) is averaged across cohorts composing the shrub or canopy strata, to obtain $FMC_{live,sh}$ and $FMC_{live,ca}$:

$$FMC_{live,sh} = \frac{\sum_i w_{i,sh} \cdot FMC_i}{\sum_i w_{i,sh}} \quad (26.50)$$

$$FMC_{live,ca} = \frac{\sum_i w_{i,ca} \cdot FMC_i}{\sum_i w_{i,ca}} \quad (26.51)$$

Live fuel moisture of herb stratum ($FMC_{live,he}$) is an input. The moisture of dead plant in the canopy and shrub layers ($FMC_{dead,ca}$ and $FMC_{dead,sh}$), the moisture of dead herbs ($FMC_{dead,he}$), as well as that of litter (FMC_{li}) and woody (FMC_{wo}) strata are all assumed equal to the moisture of 1-h dead fuels, which is an input of the model.

26.3.6.7 Proportion of dead fuel (P_{dead})

Woody and litter strata are dead fuels, but for canopy, shrub and herb strata the proportion of fuels that are dead are variable. The proportion of dead fuels in the herbaceous stratum ($P_{dead,he}$) is an input of the model, but for the shrub and canopy strata these are calculated from the proportion of dead fuels in each cohort:

$$P_{dead,sh} = \frac{\sum_i w_{i,sh} \cdot P_{dead,i}}{\sum_i w_{i,sh}} \quad (26.52)$$

$$P_{dead,ca} = \frac{\sum_i w_{i,ca} \cdot P_{dead,i}}{\sum_i w_{i,ca}} \quad (26.53)$$

26.3.6.8 Low heat content (h)

The low fuel heat content of each surface fuel stratum (in $kJ \cdot kg^{-1}$) is used for the calculation of reaction intensity. Heat content values are adjusted for live foliar moisture content in canopy, shrub and herb strata; and are left to the default value for woody and litter strata:

$$h_{ca} = h_{ca,def} - (M_{live,ca}/100) \cdot V \quad (26.54)$$

$$h_{sh} = h_{sh,def} - (M_{live,sh}/100) \cdot V \quad (26.55)$$

$$h_{he} = h_{def} - (M_{live,he}/100) \cdot V \quad (26.56)$$

$$h_{wo} = h_{li} = h_{def} \quad (26.57)$$

where $h_{def} = 18608 kJ \cdot kg^{-1} = 8000 Btu \cdot lb^{-1}$ is the default low heat content value for herbs, woody and litter strata, and $V = 2596 kJ \cdot kg^{-1} = 1116 Btu \cdot lb$ is the latent heat of vaporisation of water. The default low heat of contents for the canopy and shrub strata ($h_{ca,def}$ and $h_{sh,def}$) are calculated as a weighted average across cohorts:

$$h_{ca,def} = \frac{\sum_i w_{i,ca} \cdot h(SP_i)}{\sum_i w_{i,ca}} \quad (26.58)$$

$$h_{sh,def} = \frac{\sum_i w_{i,sh} \cdot h(SP_i)}{\sum_i w_{i,sh}} \quad (26.59)$$

where $h(SP_i)$ is a species-specific low heat content value.

26.3.6.9 Reactive volume (RV)

The volume per surface unit ($m^3 \cdot m^{-2}$) that would be involved in flaming combustion.

$$RV_{sh} = w_{shrub}/\rho_{p,sh} \quad (26.60)$$

$$RV_{he} = w_{he}/\rho_{p,he} \quad (26.61)$$

$$RV_{wo} = w_{wo}/\rho_{p,wo} \quad (26.62)$$

$$RV_{li} = \min(w_{li}, w_{\max,li})/\rho_{p,li} \quad (26.63)$$

In the case of litter, the flame loading is limited by $w_{\max,li}$, the maximum loading that would be consumed in the flaming stage of combustion, calculated as a weighted average of litter types:

$$w_{\max,li} = \frac{\sum_k w_{\max,k} \cdot w_{li,k}}{\sum_k w_{li,k}} \quad (26.64)$$

where k indicates litter type. The maximum combustion loadings for litter types are [Prichard et al. (2013); Table 2]:

$$w_{\max,shortneedlepine} = w_{\max,otherconifer} = 0.3248kg \cdot m^{-2} = 2900lb \cdot ac^{-1} \quad (26.65)$$

$$w_{\max,longneedlepine} = 0.6496kg \cdot m^{-2} = 5800lb \cdot ac^{-1} \quad (26.66)$$

$$w_{\max,hardwood} = 0.3472kg \cdot m^{-2} = 3100lb \cdot ac^{-1} \quad (26.67)$$

26.3.7 Unit conversion of fuel characteristics

FCCS calculations employ empirical equations that were derived in British units system. Hence, all the fuel characteristics and model inputs that are in metric units have to be translated into British units prior to fire behaviour calculations:

- Loading: $1kg \cdot m^{-2} = 0.204918lb \cdot ft^{-2}$
- Depths: $1m = 3.2808399ft$
- Particle density and bulk density: $1kg \cdot m^{-3} = 0.06242796lb \cdot ft^{-3}$
- Particle volume and reactive volume: $1m^3 \cdot m^{-2} = 3.2808399ft^3 \cdot ft^{-2}$
- Surface-to-area-volume ratio: $1m^2 \cdot m^{-3} = 0.3048ft^2 \cdot ft^{-3}$
- Heat content: $1kJ \cdot kg^{-1} = 0.429922614Btu \cdot lb^{-1}$
- Wind speed: $1m \cdot s^{-1} = 2.23693629mph$

26.4 Surface fire behavior

26.4.1 Surface rate of spread (R)

In the Rothermel (1972) model, surface rate of spread is defined as the ratio of heat source (i.e. the surface fire energy propagated to unburned fuels) to surface fuel heat sink (i.e. the energy required to preheat fuels). Owing to the difference in packing ratio between the litter stratum and the other surface fuels, litter-dominated fuelbeds may have substantially different spread rates than other fuelbeds. For this reason, in FCCS the rate of spread (in $ft \cdot min^{-1}$) is calculated separately for litter stratum and the final rate of spread is the maximum of the rate of spread of all surface fuels and that of the litter stratum. Rate of spread is also limited to a maximum based in windspeed and slope.

$$R = \min(WindSlopeCap, \max(R_{surf}, R_{litter})) \quad (26.68)$$

The surface fuel and litter fuel rates of spread are given by the application of Rothermel's (1972) equation to each case:

$$R_{surf} = \frac{I_{R,surf} \cdot \xi_{surf} \cdot (1 + \phi_W + \phi_S)}{q_{surf}} \quad (26.69)$$

$$R_{litter} = \frac{I_{R,litter} \cdot \xi_{litter} \cdot (1 + \phi_W + \phi_S)}{q_{litter}} \quad (26.70)$$

where $I_{R,surf}$ and $I_{R,litter}$ are the reaction intensities (in $Btu \cdot ft^{-2} \cdot min^{-1}$), ξ_{surf} and ξ_{litter} are the propagating flux ratios, q_{surf} and q_{litter} are the heat sinks. Finally, ϕ_W and ϕ_S are the slope and wind modifiers. All of them are explained in the following sections. The maximum rate of spread calculated from windspeed and slope is:

$$WindSlopeCap = 88 \cdot U \cdot (1 + \phi_S) \quad (26.71)$$

where U is windspeed (in mph) and 88 is a conversion factor (from mph to ft/min).

26.4.1.1 Reaction intensity (I_R)

Reaction intensity of surface fuels (in $Btu \cdot ft^{-2} \cdot min^{-1}$) is calculated as the sum of component reaction intensities of the four different surface fuel strata, whereas the reaction intensity in the litter uses this strata alone:

$$I_{R,surf} = I_{R,sh} + I_{R,he} + I_{R,wo} + I_{R,li} \quad (26.72)$$

$$I_{R,litter} = I_{R,li} \quad (26.73)$$

Each component reaction intensity is calculated using:

$$I_{R,sh} = (\eta_{\beta'_{allsurf}})^{A_{sh}} \cdot \Gamma'_{max,sh} \cdot w_{sh} \cdot h_{sh} \cdot \eta_{FMC,sh} \cdot \eta_{K,sh} \quad (26.74)$$

$$I_{R,he} = (\eta_{\beta'_{lowsurf}})^{A_{he}} \cdot \Gamma'_{max,he} \cdot w_{he} \cdot h_{he} \cdot \eta_{FMC,he} \cdot \eta_{K,he} \quad (26.75)$$

$$I_{R,wo} = (\eta_{\beta'_{lowsurf}})^{A_{wo}} \cdot \Gamma'_{max,wo} \cdot w_{wo} \cdot h_{wo} \cdot \eta_{FMC,wo} \cdot \eta_{K,wo} \quad (26.76)$$

$$I_{R,li} = (\eta_{\beta'_{litter}})^{A_{li}} \cdot \Gamma'_{max,li} \cdot w_{li} \cdot h_{li} \cdot \eta_{FMC,li} \cdot \eta_{K,li} \quad (26.77)$$

In the above equations, w_{sh} , w_{he} , w_{wo} and w_{li} are the loadings of the corresponding shrub, herb, woody and litter strata, respectively. These quantities were defined in previous sections, as were the corresponding low heat fuel contents (h_{sh} , h_{he} , h_{wo} and h_{li}). Mineral damping coefficient (η_K ; dimensionless) is set to the same value (corresponding to the conventional value for silica-free ash content of 1%) for all strata:

$$\eta_{K,sh} = \eta_{K,he} = \eta_{K,wo} = \eta_{K,li} = 0.42 \quad (26.78)$$

In the following subsections, we describe the calculation of the remaining variables for each stratum: reaction efficiency ($\eta_{\beta'}$), Rothermel's A parameter, maximum reaction velocity (Γ'_{max}) and moisture damping coefficient (η_{FMC}).

26.4.1.2 Reaction efficiency ($\eta_{\beta'}$)

Reaction efficiency (between 0 and 1) represents the damping effect of inefficiently packed fuels in the reaction intensity. Because shrubs rarely burn without lower surface fuels, the reaction efficiency of the surface layer ($\eta_{\beta'_{allsurf}}$) includes shrubs, herbs and woody fuels. Low surface fuels may carry flames without involving shrubs, so are assumed to burn with a single reaction efficiency ($\eta_{\beta'_{lowsurf}}$) determined by the combined characteristics of herb and woody fuel strata. Both are calculated similarly:

$$\eta_{\beta'_{allsurf}} = \beta'_{allsurf} \cdot e^{1-\beta'_{allsurf}} \quad (26.79)$$

$$\eta_{\beta'_{lowsurf}} = \beta'_{lowsurf} \cdot e^{1-\beta'_{lowsurf}} \quad (26.80)$$

where $\beta'_{allsurf}$ and $\beta'_{lowsurf}$ are the relative packing ratios corresponding to all surface fuels and low surface fuels, respectively. Relative packing ratios (β' ; dimensionless) are defined as the ratio of optimum depth (δ_{opt}) to effective depth (δ_{eff}):

$$\beta'_{allsurf} = \delta_{opt,allsurf} / \delta_{eff,allsurf} \quad (26.81)$$

$$\beta'_{lowsurf} = \delta_{opt,lowsurf} / \delta_{eff,lowsurf} \quad (26.82)$$

Optimum depth is the depth (in ft) at which fuels are optimally packed for maximum reaction intensity:

$$\delta_{opt,allsurf} = PV_{allsurf} + OptAirVol_{allsurf} \quad (26.83)$$

$$\delta_{opt,lowsurf} = PV_{lowsurf} + OptAirVol_{lowsurf} \quad (26.84)$$

where $PV_{allsurf}$ and $PV_{lowsurf}$ are the volume of particles (in $ft^3 \cdot ft^{-2}$) for all surface fuels and low surface fuels, respectively, given by:

$$PV_{allsurf} = PV_{sh} + PV_{he} + PV_{wo} \quad (26.85)$$

$$PV_{lowsurf} = PV_{he} + PV_{wo} \quad (26.86)$$

$OptAirVol_{allsurf}$ and $OptAirVol_{lowsurf}$ are the volume of air space (in $ft^3 \cdot ft^{-2}$) between fuel particles that would result in maximum reaction intensity:

$$OptAirVol_{allsurf} = 45 \cdot (RV_{sh} + RV_{he} + RV_{wo}) \quad (26.87)$$

$$OptAirVol_{lowsurf} = 45 \cdot (RV_{he} + RV_{wo}) \quad (26.88)$$

On the other hand, effective depths of all surface fuels and low surface fuels (in ft) are calculated as their depth, weighted by the reactive volume (and percentage cover in FCCS):

$$\delta_{eff,allsurf} = \frac{(RV_{sh} \cdot \delta_{sh}) + (RV_{he} \cdot \delta_{he}) + (RV_{wo} \cdot \delta_{wo})}{RV_{sh} + RV_{he} + RV_{wo}} \quad (26.89)$$

$$\delta_{eff,lowsurf} = \frac{(RV_{he} \cdot \delta_{he}) + (RV_{wo} \cdot \delta_{wo})}{RV_{he} + RV_{wo}} \quad (26.90)$$

Reaction efficiency of the litter stratum is determined separately from the other strata. It is defined as the average of reaction efficiency across litter types, calculated using loadings as weights:

$$\eta_{\beta'_{litter}} = \frac{\sum_k \eta_{\beta'_k} \cdot w_{li,k}}{\sum_k w_{li,k}} \quad (26.91)$$

where k indicates litter type. The reaction efficiencies of litter types are [Prichard et al. (2013); Table 2]:

$$\eta_{\beta'_{shortneedlepine}} = \eta_{\beta'_{otherconifer}} = 0.18 \quad (26.92)$$

$$\eta_{\beta'_{longneedlepine}} = 0.27 \quad (26.93)$$

$$\eta_{\beta'_{hardwood}} = 0.11 \quad (26.94)$$

26.4.1.3 Rothermel's A

A dimensionless coefficient that modifies reaction's efficiency (eq. (26.80)) to account for lower sensitivity of reaction efficiency to relative packing ratio in flash fuels:

$$A_{wo} = A_{li} = 1.0 \quad (26.95)$$

$$A_{sh} = 133 \cdot \sigma_{sh}^{-0.7913} \quad (26.96)$$

$$A_{he} = 133 \cdot \sigma_{he}^{-0.7913} \quad (26.97)$$

where σ_{sh} and σ_{he} have to be expressed in $ft^2 \cdot ft^{-3}$; Values 133 and -0.7913 are empirical constants (Rothermel, 1972).

26.4.1.4 Maximum reaction velocity ($\Gamma'_{\max}$)

The reaction velocity (in min^{-1}) that would exist at optimum fuelbed depth with no fuel moisture or mineral content.

$$\Gamma'_{\max,sh} = 9.495 \cdot \frac{\sigma_{sh}}{\sigma_{wo}} \quad (26.98)$$

$$\Gamma'_{\max,he} = 9.495 \cdot \frac{\sigma_{he}}{\sigma_{wo}} \quad (26.99)$$

$$\Gamma'_{\max,wo} = 9.495 \quad (26.100)$$

$$\Gamma'_{\max,li} = 15 \quad (26.101)$$

where $\sigma_{wo} = 488 \text{ ft}^2 \cdot \text{ft}^{-3} = 1601.05 \text{ m}^2 \cdot \text{m}^{-3}$ is the surface-to-area-volume ratio typical of small woody fuels. In Prichard et al. (2013) σ_{sh} is defined as the average of shrub foliar surface-to-area-volume ratio and σ_{wo} , but in our case $\sigma(SP_i)$ for each species includes both leaves and small branches. Eq. (26.101) represent a significant departure from Rothermel (1972) maximum reaction velocity, and are also different from Sandberg et al. (2007).

26.4.1.5 Moisture damping coefficient (η_{FMC})

Moisture damping reduces reaction velocity and hence reaction intensity (eq. (26.73)). It is calculated for each stratum using the following regression equations:

$$\eta_{FMC, \text{live}, sh} = \left[1 - 2.59 \cdot \left(\frac{FMC_{\text{live}, sh}}{X_{\text{live}, sh}} \right) \right] + \left[5.11 \cdot \left(\frac{FMC_{\text{live}, sh}}{X_{\text{live}, sh}} \right)^2 \right] - \left[3.52 \cdot \left(\frac{FMC_{\text{live}, sh}}{X_{\text{live}, sh}} \right)^3 \right] \quad (26.102)$$

$$\eta_{FMC, \text{dead}, sh} = \left[1 - 2.59 \cdot \left(\frac{FMC_{\text{dead}, sh}}{X_{\text{dead}, sh}} \right) \right] + \left[5.11 \cdot \left(\frac{FMC_{\text{dead}, sh}}{X_{\text{dead}, sh}} \right)^2 \right] - \left[3.52 \cdot \left(\frac{FMC_{\text{dead}, sh}}{X_{\text{dead}, sh}} \right)^3 \right] \quad (26.103)$$

$$\eta_{FMC, \text{live}, he} = \left[1 - 2.59 \cdot \left(\frac{FMC_{\text{live}, he}}{X_{\text{live}, he}} \right) \right] + \left[5.11 \cdot \left(\frac{FMC_{\text{live}, he}}{X_{\text{live}, he}} \right)^2 \right] - \left[3.52 \cdot \left(\frac{FMC_{\text{live}, he}}{X_{\text{live}, he}} \right)^3 \right] \quad (26.104)$$

$$\eta_{FMC, \text{dead}, he} = \left[1 - 2.59 \cdot \left(\frac{FMC_{\text{dead}, he}}{X_{\text{dead}, he}} \right) \right] + \left[5.11 \cdot \left(\frac{FMC_{\text{dead}, he}}{X_{\text{dead}, he}} \right)^2 \right] - \left[3.52 \cdot \left(\frac{FMC_{\text{dead}, he}}{X_{\text{dead}, he}} \right)^3 \right] \quad (26.105)$$

$$\eta_{FMC, wo} = \left[1 - 2.59 \cdot \left(\frac{FMC_{wo}}{X_{wo}} \right) \right] + \left[5.11 \cdot \left(\frac{FMC_{wo}}{X_{wo}} \right)^2 \right] - \left[3.52 \cdot \left(\frac{FMC_{wo}}{X_{wo}} \right)^3 \right] \quad (26.106)$$

$$\eta_{FMC, li} = \left[1 - 2.59 \cdot \left(\frac{FMC_{li}}{X_{li}} \right) \right] + \left[5.11 \cdot \left(\frac{FMC_{li}}{X_{li}} \right)^2 \right] - \left[3.52 \cdot \left(\frac{FMC_{li}}{X_{li}} \right)^3 \right] \quad (26.107)$$

where moisture contents of extinctions were arbitrarily set to $X_{\text{dead}, sh} = X_{\text{dead}, he} X_{wo} = X_{li} = 25$, $X_{\text{live}, sh} = 180$ and $X_{\text{live}, he} = 120$ in Sandberg et al. (2007). As it can be seen in the equations above, in the case of shrub and herb strata, moisture damping of live and dead fuels are differentiated. Average values are found after accounting for the proportion of live and dead material:

$$\eta_{FMC, sh} = \eta_{FMC, \text{live}, sh} \cdot (1 - P_{\text{dead}, sh}) + \eta_{FMC, \text{dead}, sh} \cdot P_{\text{dead}, sh} \quad (26.108)$$

$$\eta_{FMC, he} = \eta_{FMC, \text{live}, he} \cdot (1 - P_{\text{dead}, he}) + \eta_{FMC, \text{dead}, he} \cdot P_{\text{dead}, he} \quad (26.109)$$

26.4.1.6 Propagating flux ratio (ξ)

The propagating flux ratio (dimensionless) is the proportion of the reaction intensity (eq. (26.73)) that contributes to the forward rate of spread, estimated using an empirical regression:

$$\xi_{\text{surf}} = 0.03 + 2.5 \cdot \min \left[0.06, \frac{RV_{sh} + RV_{he} + RV_{wo} + RV_{li}}{\delta_{\text{surf heatsink}}} \right] \quad (26.110)$$

$$\xi_{\text{litter}} = 0.03 + 2.5 \cdot \min \left[0.06, \frac{RV_{li}}{\delta_{li}} \right] \quad (26.111)$$

where $\delta_{surfheatsink}$ is the depth of surface heat sink (in *ft*), which in Prichard et al. (2013) is calculated as the sum of strata depths weighted by their relative cover. In our case we weighted stratum depths as in the calculation of effective depth ($\delta_{eff,allsurf}$), but considering all four strata:

$$\delta_{surfheatsink} = \frac{(RV_{sh} \cdot \delta_{sh}) + (RV_{he} \cdot \delta_{he}) + (RV_{wo} \cdot \delta_{wo}) + (RV_{li} \cdot \delta_{li})}{RV_{sh} + RV_{he} + RV_{wo} + RV_{li}} \quad (26.112)$$

26.4.2 Heat sink (q)

Like reaction intensity, the heat sink term (in $Btu \cdot ft^{-3}$) of the rate of spread equation is calculated in FCCS for each fuel stratum and then summed:

$$q_{surf} = q_{sh} + q_{he} + q_{wo} + q_{li} \quad (26.113)$$

$$q_{litter} = q_{li} \quad (26.114)$$

where the heat sink for each stratum is:

$$q_{sh} = \eta_{\beta'_{surf}} \cdot \frac{RV_{sh} \cdot \rho_{p,sh} \cdot Qig_{sh}}{\min(\delta_{sh}, 1ft)} \quad (26.115)$$

$$q_{he} = \eta_{\beta'_{lowsurf}} \cdot \frac{RV_{he} \cdot \rho_{p,he} \cdot Qig_{he}}{\min(\delta_{he}, 1ft)} \quad (26.116)$$

$$q_{wo} = \eta_{\beta'_{lowsurf}} \cdot \frac{RV_{wo} \cdot \rho_{p,wo} \cdot Qig_{wo}}{\min(\delta_{wo}, 1ft)} \quad (26.117)$$

$$q_{li} = \eta_{\beta'_{li}} \cdot \frac{RV_{li} \cdot \rho_{p,li} \cdot Qig_{li}}{\min(\delta_{li}, 1ft)} \quad (26.118)$$

Where $\rho_{p,sh}$, $\rho_{p,he}$, $\rho_{p,wo}$ and $\rho_{p,li}$ are the particle densities (in $lb \cdot ft^{-3}$) of each fuel stratum; and RV_{sh} , RV_{he} , RV_{wo} , and RV_{li} are the reactive volumes of each fuel stratum. Unlike in Sandberg et al. (2007), the calculated heat sink is corrected by the reaction-efficiency term ($\eta_{\beta'_{surf}}$, $\eta_{\beta'_{lowsurf}}$ or $\eta_{\beta'_{li}}$), and the effective depth of each stratum included is limited to 1ft, based on the assumption that it is not necessary to preheat more than one 1ft of depth within a stratum to achieve ignition.

Heat of pre-ignition (Qig ; in $Btu \cdot lb^{-1}$) is the amount of heat required to ignite 1lb of fuel. It is calculated by stratum as a weighted average of live and dead fuels in shrubs and herbs.

$$Qig_{sh} = Qig_{live,sh} \cdot (1 - P_{dead,sh}) + Qig_{dead,sh} \cdot P_{dead,sh} \quad (26.119)$$

$$Qig_{he} = Qig_{live,he} \cdot (1 - P_{dead,he}) + Qig_{dead,he} \cdot P_{dead,he} \quad (26.120)$$

Whereas $Qig_{live,sh}$ and $Qig_{live,he}$ are corrected by fuel moisture, $Qig_{dead,sh}$, $Qig_{dead,he}$ and the other strata (Qig_{wo} and Qig_{li}) are assumed a constant value:

$$Qig_{live,sh} = 250 + (V \cdot (M_{live,sh}/100)) \quad (26.121)$$

$$Qig_{live,he} = 250 + (V \cdot (M_{live,he}/100)) \quad (26.122)$$

$$Qig_{dead,sh} = Qig_{dead,he} = Qig_{wo} = Qig_{li} = 250 \quad (26.123)$$

where $250Btu/lb$ is the heat of preignition of dry cellulose and $V = 1116Btu/lb$ is the latent heat of vaporization.

26.4.3 Wind and slope coefficients (ϕ_W and ϕ_S)

Wind and slope coefficients modify the heat source term of the rate of spread equation. Owing to differences in fuel characteristics and boundary conditions between the litter stratum and other surface fuel strata, in FCCS wind and slope coefficients are calculated separately for the litter stratum. The wind and slope

coefficients terms in the rate of spread equation are a weighted average of litter and surface wind and slope coefficients using the relative contribution to reaction intensity as weights:

$$\phi_W = (1 - I_{R,litter}/I_{R,surf}) \cdot \phi_{W,surf} + (I_{R,litter}/I_{R,surf}) \cdot \phi_{W,litter} \quad (26.124)$$

$$\phi_S = (1 - I_{R,litter}/I_{R,surf}) \cdot \phi_{S,surf} + (I_{R,litter}/I_{R,surf}) \cdot \phi_{S,litter} \quad (26.125)$$

Wind coefficients are calculated using:

$$\phi_{W,surf} = 8.8 \cdot \beta_{surf}'^{-E} \cdot (U/BMU)^B \quad (26.126)$$

$$\phi_{W,litter} = 8.8 \cdot \beta_{litter}'^{-E} \cdot (U/BMU)^B \quad (26.127)$$

where U is the input midflame windspeed (in $ft \cdot min^{-1}$), $BMU = 352 ft \cdot min^{-1}$ is the benchmark midflame windspeed, β_{surf}' is the relative packing ratio (eq. 26.82), B is the exponential response of wind coefficient to windspeed ($B = 1.2$ in Sandberg et al. (2007)), and E is the exponential term representing the mild effect of large fuels in reducing the accelerating effect of wind on fire spread by attenuating wind flow, given by:

$$E = 0.55 - 0.2 \cdot \frac{FAI_{sh} + FAI_{he}}{FAI_{sh} + FAI_{he} + FAI_{wo}} \quad (26.128)$$

E is assumed to be the same for both all surface fuels and litter fuels.

Slope coefficients are calculated using the empirical equation of Rothermel (1972), applied to all surface fuels and litter fuels:

$$\phi_{S,surf} = 5.275 \cdot (S/100)^2 \cdot (\beta_{sh} + \beta_{he} + \beta_{wo})^{-0.3} \quad (26.129)$$

$$\phi_{S,litter} = 5.275 \cdot (S/100)^2 \cdot \beta_{li}^{-0.3} \quad (26.130)$$

where S is the slope (in percent) and β is the packing ratio (not relative!) of fuels.

26.4.4 Fireline intensity (I_B) and flame length (FL)

Byram's fireline intensity (I_B) is the rate of heat release per unit of fire edge (in $Btu \cdot ft^{-1} \cdot min^{-1}$), and in FCCS is calculated as (Albini, 1976):

$$I_B = I_{R,surf} \cdot (R \cdot t_R) \quad (26.131)$$

where $I_{R,surf}$ is the surface reaction intensity, R is the rate of spread and t_R is the flame residence time, which is defined as the time (in min) fuels contribute to propagating flux and is estimated as Albini (1976):

$$t_R = 192 \cdot \frac{(I_{R,sh} \cdot RT_{sh}) + (I_{R,he} \cdot RT_{he}) + (I_{R,wo} \cdot RT_{wo}) + (I_{R,li} \cdot RT_{li})}{I_{R,surf}} \quad (26.132)$$

where RT is the reaction thickness, the approximate thickness (in ft) of a fuel element shell that contributes to reaction intensity. In FCCS, reaction thickness is estimated as $RT = 0.0028 ft$ for thermally thick fuel elements (Sandberg et al., 2007). When the diameter of a fuel element is less than twice the reaction thickness, the entire fuel element contributes to reaction intensity. Reaction thickness values for each stratum are given by:

$$RT_{sh} = \min(0.0028, 2/\sigma_{sh}) \quad (26.133)$$

$$RT_{he} = \min(0.0028, 2/\sigma_{he}) \quad (26.134)$$

$$RT_{wo} = \min(0.0028, 2/\sigma_{wo}) \quad (26.135)$$

$$RT_{li} = \min(0.0028, 2/\sigma_{li}) \quad (26.136)$$

Flame length is defined as the distance (in ft) between the flame tip and the midpoint of the flame depth at the base of the flame, and is calculated (Byram, 1959):

$$FL = 0.45 \cdot (I_B/60)^{0.46} \quad (26.137)$$

where 60.0 is a factor to convert from $Btu \cdot ft^{-1} \cdot min^{-1}$ to $Btu \cdot ft^{-1} \cdot s^{-1}$.

26.5 Crown fire behavior

Crown fire behavior is difficult to model and actual rates of spread are not possible to predict. Here we mainly follow the approach given in FCCS (Prichard et al., 2013), although in our case the canopy is not subdivided into layers (overstory, midstory and understory).

26.5.1 Crown fire rate of spread (R_{crown})

The rate of spread of crown fires is estimated by using a modification of Rothermel's equation:

$$R_{crown} = \frac{I_{R,crown} \cdot \xi_{crown} \cdot WAF}{q_{crown}} = \frac{(I_{R,surf} + I_{R,ca}) \cdot \xi_{crown} \cdot WAF}{q_{surf} + q_{ca}} \quad (26.138)$$

where $I_{R,surf}$ is the surface reaction intensity, $I_{R,ca}$ is the canopy reaction intensity, ξ_{crown} is the propagating flux ratio in the canopy, q_{surf} is the surface heat sink and q_{ca} is the canopy heat sink. Note that reaction intensities and heat sinks of canopy and surface fuels are added for the application of Rothermel's equation. Other modifications include the exclusion of slope effects and the consideration of wind effects through a wind adjustment factor (WAF).

Crown propagating flux ratio (ξ_{crown} ; in $Btu \cdot ft^{-3}$) represents the proportion of the crown reaction intensity that contributes to crown fire's forward rate of spread:

$$\xi_{crown} = 1 - e^{\left(-\frac{FAI_{ca}}{4\delta_{ca}}\right)} \quad (26.139)$$

where FAI_{ca} is the fuel area index of the canopy, and δ_{ca} is the canopy depth (in ft). Wind adjustment factor (WAF) is defined as:

$$WAF = \frac{U/\sqrt{U^2 + VS^2}}{BMU/\sqrt{BMU^2 + VS^2}} \quad (26.140)$$

where U is the input (midflame) windspeed (in $ft \cdot min^{-1}$), $BMU = 352 ft \cdot min^{-1}$ is the benchmark windspeed and $VS = 900 ft \cdot min^{-1}$ is the vertical stack velocity.

The following two subsections detail the calculation of canopy reaction intensity ($I_{R,ca}$) and canopy heat sink (q_{ca}).

26.5.2 Canopy reaction intensity ($I_{R,ca}$)

Reaction intensity of canopy fuels (in $Btu \cdot ft^{-2} \cdot min^{-1}$) is estimated as:

$$I_{R,ca} = (\eta_{\beta'_{ca}})^{A_{ca}} \cdot \Gamma'_{max,ca} \cdot w_{ca} \cdot h_{ca} \cdot \eta_{FMC,ca} \cdot \eta_{K,ca} \quad (26.141)$$

where $A_{ca} = 133 \cdot \sigma_{ca}^{-0.7913}$ is Rothermel's A coefficient, $\Gamma'_{max,ca} = 15 min^{-1}$ is the maximum reaction velocity of the canopy, w_{ca} is the loading of canopy fuels (in $lb \cdot ft^{-2}$), h_{ca} is the heat content of the canopy fuels (in $Btu \cdot lb^{-1}$) and $\eta_{K,ca} = 0.42$ is the mineral damping coefficient of the canopy stratum. Moisture damping coefficient for the canopy ($\eta_{FMC,ca}$) is estimated as done for shrub and herb strata:

$$\eta_{FMC, live, ca} = \left[1 - 2.59 \cdot \left(\frac{FMC_{live, ca}}{X_{live, ca}}\right)\right] + \left[5.11 \cdot \left(\frac{FMC_{live, ca}}{X_{live, ca}}\right)^2\right] - \left[3.52 \cdot \left(\frac{FMC_{live, ca}}{X_{live, ca}}\right)^3\right] \quad (26.142)$$

$$\eta_{FMC, dead, ca} = \left[1 - 2.59 \cdot \left(\frac{FMC_{dead, ca}}{X_{dead, ca}}\right)\right] + \left[5.11 \cdot \left(\frac{FMC_{dead, ca}}{X_{dead, ca}}\right)^2\right] - \left[3.52 \cdot \left(\frac{FMC_{dead, ca}}{X_{dead, ca}}\right)^3\right] \quad (26.143)$$

$$\eta_{FMC, ca} = \eta_{FMC, live, ca} \cdot (1 - P_{dead, ca}) + \eta_{FMC, dead, ca} \cdot P_{dead, ca} \quad (26.144)$$

where moisture contents of extinctions were arbitrarily set to $X_{dead, ca} = 25$ and $X_{live, ca} = 180$.

The reaction efficiency in the canopy ($\eta_{\beta'_{canopy}}$) represents the damping effect of inefficiently packed fuels in the canopy:

$$\eta_{\beta'_{canopy}} = \beta'_{canopy} \cdot e^{1-\beta'_{canopy}} \quad (26.145)$$

where β'_{canopy} is the relative packing ratio in the canopy:

$$\beta'_{canopy} = \delta_{opt,canopy} / \delta_{eff,canopy} \quad (26.146)$$

where the effective depth is $\delta_{eff,canopy} = \delta_{ca}$ (in *ft*) and the optimum canopy depth is calculated using:

$$\delta_{opt,canopy} = 0.4 \cdot FAI_{ca} + \beta_{ca} \cdot (\delta_{ca} \cdot C_{ca} / 100) \quad (26.147)$$

where C_{ca} is the percent cover of the canopy, FAI_{ca} is the fuel area index of the canopy, β_{ca} is the packing ratio of canopy fuels and δ_{ca} is the canopy depth (in *ft*).

26.5.3 Canopy heat sink (q_{ca})

Canopy heat sink (in $Btu \cdot ft^{-3}$) is estimated using:

$$q_{ca} = \frac{0.5 \cdot FAI_{ca} \cdot RT_{ca} \cdot \rho_{p,ca} \cdot Qig_{ca}}{(C_{ca} / 100) \cdot \delta_{ca}} \quad (26.148)$$

where C_{ca} is the percent cover of the canopy, FAI_{ca} is the fuel area index of the canopy, $RT_{ca} = \min(0.0028, 2/\sigma_{ca})$ is the reaction thickness of the canopy stratum (in *ft*), $\rho_{p,ca}$ is the particle density of the canopy (in $lb \cdot ft^{-3}$), δ_{ca} is the canopy depth (in *ft*) and Qig_{ca} is the heat of pre-ignition of the canopy stratum (in $Btu \cdot lb^{-1}$), which is calculated as a weighted average of live and dead fuels:

$$Qig_{live,ca} = 250 + (V \cdot (M_{live,ca} / 100)) \quad (26.149)$$

$$Qig_{dead,ca} = 250 \quad (26.150)$$

$$Qig_{ca} = Qig_{live,ca} \cdot (1 - P_{dead,ca}) + Qig_{dead,ca} \cdot P_{dead,ca} \quad (26.151)$$

where $250Btu/lb$ is the heat of preignition of dry cellulose and $V = 1116Btu/lb$ is the latent heat of vaporization of water.

26.5.4 Fireline intensity ($I_{B,crown}$) and flame length (FL_{crown})

Byram's fireline intensity for crown fires is estimated using:

$$I_{B,crown} = I_{R,crown} \cdot (R_{crown} \cdot t_{R,crown}) \quad (26.152)$$

where $I_{R,crown}$ is the crown reaction intensity (i.e. the sum of canopy and surface reaction intensities), R_{crown} is the rate of crown fire spread and $t_{R,crown}$ is the flame residence time, estimated as:

$$t_R = 192 \cdot RT_{ca} \quad (26.153)$$

where $RT_{ca} = \min(0.0028, 2/\sigma_{ca})$ is the reaction thickness of the canopy. As for surface fires, flame length is calculated using:

$$FL_{crown} = 0.45 \cdot (I_{B,crown} / 60)^{0.46} \quad (26.154)$$

where 60.0 is a factor to convert from $Btu \cdot ft^{-1} \cdot min^{-1}$ to $Btu \cdot ft^{-1} \cdot s^{-1}$.

26.6 Fire potentials

26.6.1 Surface fire behavior potentials

The **surface fire behavior potential** (*SFP*; between 0 and 9) is an index defined as the maximum of spread potential (*SP*) and flame length (*FL*) potential indices (both between 0 and 9):

$$SFP = \max(SP, FP) \quad (26.155)$$

Spread potential is derived from R (in $ft \cdot min^{-1}$), and *flame length potential* is derived from FL (in *ft*), both quantities being calculated at benchmark environmental conditions:

$$SP = \min[9, R^{1/2}] \quad (26.156)$$

$$FP = \min[9, FL^{1/2}] \quad (26.157)$$

26.6.2 Crown fire behavior potentials

The *crown fire summary potential* (CPF) combines three subpotentials into a single index value between 0 and 9. It places more emphasis on crown fire initiation (IC) and rate of spread (RC) than to crown-to-crown transmissivity (TC):

$$CPF = 0.4286 \cdot (IC + (TC/3) + RC) \quad (26.158)$$

where 0.4286 is used to limit CPF between 0 and 9.

The *crown fire initiation potential* (IC) represents the likelihood of a surface fire torching into single or multiple trees. IC is based on the work by Van Wagner. If $FAI_{ca} = 0$ then $IC = 0$. Otherwise it is calculated as:

$$IC = \min \left[9, 4 \cdot \left(\frac{I_B/60}{I_c} \right)^{0.2} \right] \quad (26.159)$$

where I_B is the surface fireline intensity (60 is used to convert it to $Btu \cdot ft^{-1} \cdot s^{-1}$) and I_c is Van Wagner's critical fireline intensity (Scott and Reinhardt, 2002):

$$I_c = 0.288894658 \cdot [0.01 \cdot (H_{gap}/100) \cdot (460 + 25.9 \cdot M_{live,ca})]^{1.5} \quad (26.160)$$

where $M_{live,ca}$ is the moisture content of the canopy (in percent of dry weight), 0.288894658 is used to convert from $kJ \cdot m^{-1} \cdot s^{-1}$ to $Btu \cdot ft^{-1} \cdot s^{-1}$ and H_{gap} is the canopy gap (in cm) determined from the analysis of the bulk density profile.

The *crown-to-crown transmissivity potential* (TC) is set to zero if $FAI_{ca} < TFAI/(3 \cdot \pi)$, where $TFAI$ is a threshold for FAI calculated as:

$$TFAI = A_q \cdot e^{-0.0019 \cdot U} \quad (26.161)$$

where $A_q = 3.2868$ if $\sigma_{ca} > 2000 ft^2 \cdot ft^{-3}$ and $A_q = 2.6296$ otherwise. If $FAI_{ca} > TFAI/(3 \cdot \pi)$, then TC is calculated as:

$$TC = \min [9, 10 \cdot TC_q] \quad (26.162)$$

where TC_q is the efficiency of crown-to-crown heat transfer, as a proportion of maximum efficiency at 100 % canopy cover:

$$TC_q = \frac{(\max(0.0, C_{ca} \cdot WAF - 40))^{0.3}}{(100 \cdot WAF - 40)^{0.3}} \quad (26.163)$$

In this last equation 40 represents the threshold of canopy cover necessary to initiate dependent crown spread, and 0.3 is a coefficient describing the assumed effect of crown cover on transmissivity at benchmark windspeed. The canopy adjustment ratio WAF is added to modulate transmissivity depending on windspeed.

Finally, the *crown fire rate of spread potential* (RC) is defined as:

$$RC = \min [9, 2.5 \cdot R_{crown}^{1/e}] \quad (26.164)$$

where R_{crown} is the rate of spread (in $ft \cdot min^{-1}$) of the crown fire.

26.7 Unit conversion of outputs

The following factors are used to express fire behavior outputs to metric units:

- *Fire spread rates:* $1 ft \cdot min^{-1} = 0.3048 m \cdot min^{-1}$
- *Flame length:* $1 ft = 0.3048 m$
- *Reaction intensity:* $1 Btu \cdot ft^{-2} \cdot min^{-1} = 11.3484 kJ \cdot m^{-2} \cdot min^{-1}$
- *Heat sink:* $1 Btu \cdot ft^{-3} = 37.2589458 kJ \cdot m^{-3}$
- *Fireline intensity:* $1 Btu \cdot ft^{-1} \cdot min^{-1} = 0.0576911555 kW \cdot m^{-1}$

26.8 Applications

Function `fuel_FCCS()` has been used to estimate understory and overstory fuel properties for forests over large areas by applying it on national forest inventory plots (Sánchez-Pinillos et al., 2019, 2021). Function `fire_FCCS()` has been employed to estimate relationship between forest management and potential fire behaviour in several publications (Vilà-Vilardell et al., 2023; Miezi et al., 2022) and to project wildfire risk under climate scenarios (Ameztegui et al., 2024).

Appendix A

Inbuilt parameter estimation

A.1 Introduction

Package **medfate** has been designed to allow simulations requiring a minimum set of vegetation functional parameters. This entails that several other parameters have to be estimated automatically (via inbuilt procedures) before starting simulations. Inbuilt parameter estimation is done in functions `spwbInput()` and `growthInput()`, with the user controlling the process through the species parameter table input (e.g., `SpParamsMED`) and the object `control` (see default control values in `defaultControl()`).

A.2 Strict, scaled and imputable parameters

Different kinds of vegetation functional parameters can be distinguished according to whether inbuilt parameter estimation is possible and how it is conducted:

- **Strictly-required parameters** are those for which there are no inbuilt estimation procedures implemented in the initialization functions. Hence, either values in the species parameter table input are non-missing or suitable values need to be specified before running simulation models. Since **medfate ver. 2.3**, only plant/leaf classification parameters and plant size parameters are strict. The remaining ones can be estimated from other parameters. This facilitates having a functional species parameter table, because only a set of parameters have to be strictly filled, from either soft trait databases or forest inventory data.
- **Scaled parameters** are functional parameters that cannot be defined at the species level, because they need to be estimated taking into account the size and structure of the plant cohort. These are not normally defined at the level of species parameter table. Specific `control` parameters are used to determine how scaling is performed.
- **Imputable parameters** parameters are those for which the initialization routines can provide default values or estimations derived from relationships with other parameters. Parameter imputation is conducted if control parameter `fillMissingSpParams = TRUE`. Sometimes, default parameter values are also specified in the `control` object.

The following tables describe how the different functional parameters are dealt with, grouped by function. Links are given to the chapter subsections where scaling and/or imputation procedures are described.

Plant/leaf classification

Symbol	R	Description	Strict	Scaled	Imputable
GF	GrowthForm	Growth form, defined depending on the treatment in forest inventory plots (Tree, Shrub or Tree/Shrub)	Yes	No	No
LF	LifeForm	Raunkiaer life form	Yes	No	No
L_{shape}	LeafShape	Leaf type (Linear, Needle, Broad, Scale, Spines or Succulent)	Yes	No	No
L_{size}	LeafSize	Leaf size (Small, Medium, Large)	Yes	No	No
L_{pheno}	PhenologyType	Phenology type	Yes	No	No

Plant size

Symbol	R	Description	Strict	Scaled	Imputable
H_{max}	Hmed	Maximum plant height	Yes	No	No
H_{med}	Hmed	Median plant height	Yes	No	No
Z_{50}	Z50	Depth above which 50% of the fine root mass is located	No	No	A.3.2
Z_{95}	Z95	Depth above which 95% of the fine root mass is located	Yes	No	No
Z_{100}	Z100	Depth above which 100% of the fine root mass is located	No	No	A.3.2

Allometric coefficients

Symbol	R	Description	Strict	Scaled	Imputable
a_{ash}, b_{ash}	a_ash, b_ash	Coefficients relating the square of shrub height with shrub area	No	No	A.3.3
a_{bsh}, b_{bsh}	a_bsh, b_bsh	Coefficients relating crown volume with dry weight of shrub individuals	No	No	A.3.3
cr	cr	Ratio between crown length and total height for shrubs	No	No	A.3.3
$a_{fbt}, b_{fbt}, c_{fbt}$	a_fbt, b_fbt, c_fbt	Coefficients to calculate foliar biomass of an individual tree	No	No	A.3.4
$a_{cr}, b_{1cr}, b_{2cr}, b_{3cr}, c_{1cr}, c_{2cr}$	a_cr, b_1cr, b_2cr, b_3cr, c_1cr, c_2cr	Coefficients to calculate crown ratio of trees	No	No	A.3.4
a_{cw}, b_{cw}	a_cw, b_cw	Regression coefficients used to calculate the crown width of trees	No	No	A.3.4
$f_{HD,min}$	fHDmin	Minimum height-to-diameter ratio	No	No	A.3.4
$f_{HD,max}$	fHDmax	Maximum height-to-diameter ratio	No	No	A.3.4

Leaf phenology

Symbol	R	Description	Strict	Scaled	Imputable
LD	LeafDuration	Average duration of leaves	No	No	A.3.10
$t_{0,eco}$	t0gdd	Degree days corresponding to leaf budburst	No	No	A.3.10
S_{eco}^*	Sgdd	Degree days corresponding to leaf budburst	No	No	A.3.10
T_{eco}	Tbgdd	Base temperature for the calculation of degree days to leaf budburst	No	No	A.3.10
S_{sen}^*	Ssen	Degree days corresponding to leaf senescence	No	No	A.3.10
Ph_{sen}	Phsen	Photoperiod corresponding to start counting senescence degree-days	No	No	A.3.10
T_{sen}	Tbsen	Base temperature for the calculation of degree days to leaf senescence	No	No	A.3.10
x_{sen}	xsen	Discrete values, to allow for any absent/proportional/more than proportional effects of temperature on senescence	No	No	A.3.10
y_{sen}	ysen	Discrete values, to allow for any absent/proportional/more than proportional effects of photoperiod on senescence	No	No	A.3.10

Plant anatomy

Symbol	R	Description	Strict	Scaled	Imputable
$1/H_v$	A12As	Ratio of leaf area to sapwood area	No	No	A.3.8
RLR	Ar2A1	Fine root area to leaf area ratio	No	No	A.3.9
LW	LeafWidth	Leaf width	No	No	A.3.5
SLA	SLA	Specific leaf area	No	No	A.3.5
ρ_{leaf}	LeafDensity	Leaf tissue density	No	No	A.3.6
ρ_{wood}	WoodDensity	Wood tissue density	No	No	A.3.6
$\rho_{fineroot}$	FineRootDensity	Fine root tissue density	No	No	A.3.6
$f_{conduits}$	conduit2Sapwood	Proportion of sapwood corresponding to xylem conduits	No	No	A.3.8
SRL	SRL	Specific fine root length	No	No	A.3.7
RLD	RLD	Fine root length density	No	No	A.3.7
$r_{6.35}$	r635	Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm	No	No	A.3.5

Radiation balance and water interception

Symbol	R	Description	Strict	Scaled	Imputable
k_b	kDIR	Direct light extinction coefficient	No	No	A.3.12
k_{PAR}	kPAR	PAR extinction coefficient	No	No	A.3.12
α_{SWR}	alphaSWR	Short-wave radiation leaf absorbance coefficient	No	No	A.3.12
γ_{SWR}	gammaSWR	Short-wave radiation leaf reflectance (albedo)	No	No	A.3.12
s_{water}	g	Crown water storage capacity	No	No	A.3.12

Hydraulics, transpiration, photosynthesis

Symbol	R	Description	Strict	Scaled	Imputable
$T_{max,LAI}$	Tmax_LAI	Empirical coefficient relating LAI with the ratio of maximum transpiration over potential evapotranspiration	No	No	A.3.11
$T_{max,sqLAI}$	Tmax_LAI ^{sq}	Empirical coefficient relating squared LAI with the ratio of maximum transpiration over potential evapotranspiration	No	No	A.3.11
WUE_{max}	WUE	Water use efficiency at VPD = 1kPa and without light or CO2 limitations	No	No	A.3.11
WUE_{PAR}	WUE_par	Coefficient describing the progressive decay of WUE with lower light levels	No	No	A.3.11
WUE_{CO2}	WUE_co2	Coefficient for WUE dependency on atmospheric CO2 concentration	No	No	A.3.11
WUE_{VPD}	WUE_vpd	Coefficient for WUE dependency on vapor pressure deficit	No	No	A.3.11
$\Psi_{extract}$	Psi_Extract	The water potential at which plant transpiration is 50% of its maximum	No	No	A.3.11
Ψ_{critic}	Psi_Critic	The water potential corresponding to 50% of stem xylem cavitation	No	No	A.3.17
g_{swmin}	Gwmin	Minimum stomatal conductance to water vapour	No	No	A.3.13
g_{swmax}	Gwmax	Maximum stomatal conductance to water vapour	No	No	A.3.13
$J_{max,298}$	Jmax298	Maximum rate of electron transport at 298K	No	No	A.3.18
$V_{max,298}$	Vmax298	Rubisco's maximum carboxylation rate at 298K	No	No	A.3.18
$K_{stem,max,ref}$	Kmax_stem_xylem	Maximum stem sapwood reference conductivity per leaf area unit	No	No	A.3.15
$K_{root,max,ref}$	Kmax_root_xylem	Maximum root sapwood reference conductivity per leaf area unit	No	No	A.3.15
$k_{leaf,max}$	VCleaf_kmax	Maximum leaf hydraulic conductance	No	A.4.2	A.3.16
$k_{stem,max}$	VCstem_kmax	Maximum stem hydraulic conductance	No	A.4.1	No
$k_{root,max,s}$	VCroot_kmax	Maximum root hydraulic conductance for each soil layer	No	A.4.3	No
$k_{rhizo,max,s}$	VGrhizo_kmax	Maximum hydraulic conductance of the rhizosphere for each soil layer	No	A.4.4	No
c_{leaf}, d_{leaf}	VCleaf_c, VCleaf_d	Parameters of the vulnerability curve for leaves	No	No	A.3.17
c_{stem}, d_{stem}	VCstem_c, VCstem_d	Parameters of the vulnerability curve for stem xylem	No	No	A.3.17
c_{root}, d_{root}	VCroot_c, VCroot_d	Parameters of the vulnerability curve for root xylem	No	No	A.3.17

Plant water storage

Symbol	R	Description	Strict	Scaled	Imputable
ϵ_{leaf}	LeafEPS	Modulus of elasticity of leaves	No	No	A.3.14
ϵ_{stem}	StemEPS	Modulus of elasticity of symplastic xylem tissue	No	No	A.3.14

Symbol	R	Description	Strict	Scaled	Imputable
$\pi_{0,leaf}$	LeafPI0	Osmotic potential at full turgor of leaves	No	No	A.3.14
$\pi_{0,stem}$	StemPI0	Osmotic potential at full turgor of symplastic xylem tissue	No	No	A.3.14
$f_{apo,leaf}$	LeafAF	Apoplastic fraction in leaf tissues	No	No	A.3.14
$f_{apo,stem}$	StemAF	Apoplastic fraction in stem tissues	No	No	A.3.14
V_{leaf}	Vleaf	Leaf water capacity per leaf area unit	No	A.4.5	No
$V_{sapwood}$	Vsapwood	Sapwood water capacity per leaf area unit	No	A.4.5	No

Growth and mortality

Symbol	R	Description	Strict	Scaled	Imputable
N_{leaf}	Nleaf	Leaf nitrogen concentration per dry mass	No	No	A.3.19
$N_{sapwood}$	Nsapwood	Sapwood nitrogen concentration per dry mass	No	No	A.3.19
$N_{fineroot}$	Nfineroot	Fine root nitrogen concentration per dry mass	No	No	A.3.19
MR_{leaf}	RERleaf	Leaf respiration rate at 20 °C	No	No	A.3.19
$MR_{sapwood}$	RERSapwood	Living sapwood (parenchymatic tissue) respiration rate at 20 °C	No	No	A.3.19
$MR_{fineroot}$	RERfineroot	Fine root respiration rate at 20 °C	No	No	A.3.19
$RGR_{leaf,max}$	RGRleaf,max	Maximum leaf area daily growth rate, relative to sapwood area	No	No	A.3.20
$RGR_{cambium,max}$	RGRsapwood,max	Maximum tree daily sapwood growth rate relative to cambium perimeter length	No	No	A.3.20
$RGR_{sapwood,max}$	RGRsapwood,max	Maximum shrub daily sapwood growth rate relative to sapwood area	No	No	A.3.20
$RGR_{fineroot,max}$	RGRfineroot,max	Maximum daily fine root relative growth rate	No	No	A.3.20
$SR_{sapwood}$	SRsapwood	Daily sapwood senescence rate	No	No	A.3.21
$SR_{fineroot}$	SRfineroot	Daily fine root senescence rate	No	No	A.3.21
$RSSG$	RSSG	Minimum relative starch for sapwood growth	No	No	A.3.22
C_{wood}	WoodC	Wood carbon content per dry weight	No	No	A.3.23
$P_{mort,base}$	MortalityBaseline	Baseline deterministic proportion or probability specifying the baseline reduction of cohort's density occurring in a year	No	No	A.3.24

Recruitment

Symbol	R	Description	Strict	Scaled	Imputable
H_{seed}	SeedProductionHeight	Minimum height for seed production	No	No	A.3.25
TCM_{recr}	MinTempRecr	Minimum average temperature (Celsius) of the coldest month for successful recruitment	No	No	A.3.25
MI_{recr}	MinMoistureRecr	Minimum value of the moisture index for successful recruitment	No	No	A.3.25
$FPAR_{recr}$	MinFPARRecr	Minimum percentage of PAR at the ground level for successful recruitment	No	No	A.3.25

Symbol	R	Description	Strict	Scaled	Imputable
DBH_{recr}	RecrTreeDBH	Recruitment DBH for trees	No	No	A.3.25
$H_{tree,recr}$	RecrTreeHeight	Recruitment height for trees	No	No	A.3.25
$N_{tree,recr}$	RecrTreeDensity	Recruitment density for trees	No	No	A.3.25
$Cover_{shrub,recr}$	RecrShrubCover	Recruitment cover for shrubs	No	No	A.3.25
$H_{shrub,recr}$	RecrShrubHeight	Recruitment height for shrubs	No	No	A.3.25
$Z50_{recr}$	RecrZ50	Soil depth corresponding to 50% of fine roots for recruitment	No	No	A.3.25
$Z95_{recr}$	RecrZ95	Soil depth corresponding to 95% of fine roots for recruitment	No	No	A.3.25

Flammability

Symbol	R	Description	Strict	Scaled	Imputable
ρ_p	PD	Density of fuel particles	No	No	A.3.26
σ	SAV	Surface-area-to-volume ratio of the small fuel (1h) fraction (leaves and branches < 6.35mm)	No	No	A.3.26
h	HeatContent	High fuel heat content.	No	No	A.3.26
LI	PercentLignin	Percentage of lignin in leaves	No	No	A.3.26

A.3 Imputation of missing values

A.3.1 Genus-level imputation

Given that plant traits have some degree of phylogenetic conservatism, parameter imputation of species parameter values is performed primarily by looking for the corresponding genus-average, which should be available in **SpParams**. This genus-based imputation can be controlled using **fillMissingWithGenusParams**, which by default is **TRUE** (setting it to **FALSE** is not normally recommended). Should the parameter be also missing at the genus level, **medfate** will try an imputation based on trait coordination or family-level average values, depending on the case. In the following subsections we report the percentage of species-level variance that is accounted for by genus-level or family-level averages, and detail trait-coordination relationships when used for imputation.

The following figure summarizes the percentage of missing values in **SpParamsMED** for different model parameters and the other model parameters used for the imputation of missing values:

A.3.2 Rooting depth

Parameter Z_{95} is a strict parameter, but Z_{50} and Z_{100} can be imputed when missing, using the following formulae:


$$Z_{50} = \exp(\log(Z_{95})/1.4) \quad (\text{A.1})$$

$$Z_{100} = \exp(\log(Z_{95})/0.95) \quad (\text{A.2})$$

Note that Z_{100} will be imputed only if **truncateRootDistribution** = **TRUE** in the control parameters.

A.3.3 Shrub allometric coefficients

Missing shrub allometric coefficients are filled using information from Raunkiaer's life form and maximum plant height (H_{max}).



Life form	H_{max}	a_{ash}	b_{ash}	a_{bsh}	b_{bsh}	cr
Chamaephyte	[any]	24.5888	1.1662	0.7963	0.3762	0.8076
Phanerophyte	< 300 cm	1.0083	1.8700	0.7900	0.6942	0.6630
Phanerophyte	> 300 cm	5.8458	1.4944	0.3596	0.7138	0.7190
(Hemi)cryptophyte	[any]	24.5888	1.1662	0.7963	0.3762	0.9500

Allometric coefficients were taken from De Cáceres et al. (2019).

A.3.4 Tree allometric coefficients

Missing tree allometric coefficients are replaced with values depending on whether the plant species is a gymnosperm or an angiosperm:

Parameter	Gymnosperm	Angiosperm
a_{fbt}	0.1300	0.0527
b_{fbt}	1.2285	1.5782
c_{fbt}	-0.0147	-0.0066
a_{cw}	0.747	0.839
b_{cw}	0.672	0.735
a_{cr}	1.995	1.506
b_{1cr}	-0.649	-0.706
b_{2cr}	-0.020	-0.078
b_{3cr}	-0.00012	0.00018
c_{1cr}	-0.004	-0.007
c_{2cr}	-0.159	0.000
fHD_{min}	80	40
fHD_{max}	120	140

A.3.5 Leaf width, specific leaf area and fine foliar ratio

Leaf width (LW), specific leaf area (SLA) and the ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm ($r_{6.35}$) are key anatomical parameters. When missing from species parameter table, default estimates for these parameters are obtained from combinations of leaf shape and leaf size:

Leaf shape	Leaf size	SLA	LW	$r_{6.35}$
Broad	Large	16.039	6.898	2.278
Broad	Medium	11.499	3.054	2.359
Broad	Small	9.540	0.644	3.026
Linear	Large	5.522	0.639	3.261
Linear	Medium	4.144	0.639	3.261
Linear	Small	13.189	0.639	3.261
Needle	[any]	9.024	0.379	1.716
Spines	[any]	9.024	0.379	1.716
Scale	[any]	4.544	0.101	1.483

These estimates have been obtained by averaging species-level values across combinations of the categorical variables.

Table A.1: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of leaf density, wood density and fine root density.

Trait	spp.	Family (%)	Genus (%)	Total (%)
LeafDensity	4258	14.89	28.98	43.9
WoodDensity	14682	23.24	50.82	74.1
FineRootDensity	1490	8.76	32.86	41.6

Table A.2: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of specific fine root length.

Trait	spp.	Family (%)	Genus (%)	Total (%)
SRL	2093	8.48	25.87	34.3

A.3.6 Tissue density

Imputation of the dry weight density of leaves, wood and fine roots (in $g \cdot cm^{-3}$) is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set `medfate:::trait_family_means`). The percentage of species-level explained variance of these imputations is reported in the following table:

If the family is not any of those in the table, default values are $\rho_{leaf} = 0.7$ and $\rho_{wood} = 0.652$ and $\rho_{fineroot} = 0.165$.

A.3.7 Specific root length and root length density

Imputation of specific fine root length and fine root length density is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation of specific fine root length is performed using taxonomic family (internal data set `medfate:::trait_family_means`). The percentage of species-level explained variance of these imputations is reported in the following table:

Default value for specific fine root length is $3870 \text{ cm} \cdot g^{-1}$. Fine root length density is imputed a value $10 \text{ cm} \cdot \text{cm}^{-3}$ when missing at the genus level.

A.3.8 Huber value and ratio of conduits to sapwood

Imputation of the **A12As**, the inverse of the Huber value ($1/Hv$) and the fraction of sapwood area corresponding to xylem conduits (`conduit2sapwood`; $f_{conduits}$) is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set `medfate:::trait_family_means`). The percentage of species-level explained variance of these imputations is reported in the following table:

If there is no information derived from taxonomic family for **A12As**, a default value is given depending on leaf shape and leaf size:

Table A.3: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of leaf area to sapwood area and fraction of sapwood corresponding to xylem elements.

Trait	spp.	Family (%)	Genus (%)	Total (%)
conduit2sapwood	651	28.14	42.64	70.8
A12As	1330	6.66	54.69	61.4

Leaf shape	Leaf size	A12As
Broad	Large	4768.7
Broad	Medium	2446.1
Broad	Small	2284.9
Linear	Large	2156.0
Linear	Medium	2156.0
Linear	Small	2156.0
Needle	[any]	2751.7
Scale	[any]	1696.6

Missing values for $f_{conduits}$, the fraction of sapwood corresponding to conduits are derived from taxonomic family (see table above). If information from taxonomic family is missing, default values are $f_{conduits} = 0.7$ (i.e. 30% of parenchyma) for angiosperms, and $f_{conduits} = 0.925$ (i.e. 7.5% of parenchyma) for gymnosperms (Plavcová and Jansen, 2015).

A.3.9 Fine root to leaf area ratio

When missing, the fine root area to leaf area ratio is given a default value of $RLR = 1 \text{ m}^2 \cdot \text{m}^{-2}$.

A.3.10 Leaf phenology

When missing, leaf duration (**LeafDuration**) is assigned a value of 1 year for winter-deciduous species and 2.41 years for the remaining leaf phenology types. Missing values for the number of days required to complete bud formation (**BudFormationDays**) are assigned 20 days.

Current default values for leaf budburst parameters are the same regardless of the leaf phenology type:

Phenology type	t0gdd	Sgdd	Ugdd	Tbgdd
Evergreen	50	200	300	0
Winter deciduous	50	200	300	0
Winter semi-deciduous	50	200	300	0

Current default values for leaf senescence parameters are the same regardless of the leaf phenology type:

Phenology type	Ssen	Phsen	Tbsen	xsen	ysen
Evergreen	8268	12.5	28.5	2	2
Winter deciduous	8268	12.5	28.5	2	2
Winter semi-deciduous	8268	12.5	28.5	2	2

Leaf senescence values were derived for deciduous broad-leaved forests by Delpierre et al. (2009).

A.3.11 Basic transpiration and water-use efficiency

When the basic soil water balance model is used, $T_{max,LAI}$ and $T_{max,sqLAI}$ are species-specific parameters that regulate the maximum transpiration of plant cohorts (see 6.1.1). When these parameters are missing from **SpParams** table, they are given default values $T_{max,LAI} = 0.134$ and $T_{max,sqLAI} = -0.006$, according to Granier et al. (1999).

When maximum water use efficiency (WUE_{max}) is missing, it is given a value of $WUE_{max} = 7.55$. By default, the coefficient describing the decay of water use efficiency with lower light levels is given a default

Table A.4: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of leaf minimum conductance to water vapour.

Trait	spp.	Family (%)	Genus (%)	Total (%)
Gswmin	465	-6.34	82.84	76.5

value of $WUE_{PAR} = 0.2812$, and the coefficient regulating the relationship between gross photosynthesis and CO₂ concentration is given a default $WUE_{CO_2} = 0.0028$.

When missing, the water potential corresponding to 50% of transpiration ($\Psi_{extract}$) is estimated by calculating the water potential corresponding to the loss leaf turgor (Ψ_{tlp}), using equation (10.10) from Bartlett et al. (2012). The parameters of the leaf pressure-volume curve needed for applying equation (10.10) may be themselves estimated (see A.3.14). Note that Ψ_{tlp} has been found to be highly correlated to Ψ_{gs50} , the water potential corresponding to 50% of stomatal conductance (Bartlett et al., 2016).

A.3.12 Radiation balance and water interception

Default value for direct light extinction is $k_b = 0.8$. Default values for diffuse radiation extinction, absorbance, reflectance and water interception parameters depend on the leaf shape:

Leaf shape	k_{PAR}	α_{SWR}	γ_{SWR}	s_{water}
Broad	0.55	0.70	0.18	0.5
Linear	0.45	0.70	0.15	0.8
Needle/Scale	0.50	0.70	0.14	1.0

where k_{PAR} is the diffuse PAR extinction coefficient, α_{SWR} is the short-wave radiation leaf absorbance coefficient, γ_{SWR} is the short-wave radiation leaf reflectance (albedo) and s_{water} is the crown water storage capacity per LAI unit.

A.3.13 Stomatal conductance

Minimum stomatal conductance

Imputation of the minimum and maximum conductance to water vapour (g_{swmin} ; in $mol\ H_2O \cdot s^{-1} \cdot m^{-2}$) is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set **medfate::trait_family_means**). The percentage of species-level explained variance of these imputations is reported in the following table.

If there is no information derived from taxonomic family, $g_{swmin} = 0.0049$.

Maximum stomatal conductance

Imputation of g_{swmax} is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set **medfate::trait_family_means**). The percentage of species-level explained variance of these imputations is reported in the following table:

If there is no information derived from taxonomic family, $g_{swmax} = 0.200$.

Decades of theory have focused on the basis of anatomical maximum conductance to water vapour (g_{swmax} ; in $mol\ H_2O \cdot s^{-1} \cdot m^{-2}$), with stomatal density and size being the main predictors (Sack and Buckley, 2016; Ochoa et al., 2024), usually incorporating assumptions about allometries among guard cell dimensions. This is frequently used to infer maximum stomatal conductance of the fossil record (Franks and Beerling,

Table A.5: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of maximum conductance to water vapour.

Trait	spp.	Family (%)	Genus (%)	Total (%)
Gsw_q99	2718	0.98	60.57	61.6

Table A.6: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of leaf pressure-volume parameters.

Trait	spp.	Family (%)	Genus (%)	Total (%)
LeafPI0	550	7.03	20.75	27.8
LeafEPS	367	16.45	37.95	54.4
LeafAF	118	18.22	26.19	44.4

2009). However, anatomical (theoretical) estimates of g_{swmax} are often much higher the relationship than operational (observed physiological) g_{swmax} (McElwain et al., 2016), for several reasons (Sack and Buckley, 2016). The anatomical g_{swmax} is therefore a theoretical value estimating the maximum stomatal diffusion capacity. Alternatively, operational g_{swmax} has been shown to be coordinated with $k_{l,max}$ (Franks, 2006) and the density of veins in leaves (McElwain et al., 2016).

A.3.14 Pressure-volume curves

Parameters of the pressure-volume curve (i.e. $\pi_{0,stem}$ and ϵ_{stem}) for leaf and stem symplastic tissue are required for each species.

When parameters for stem tissue are missing, **medfate** estimates them from wood density following Christoffersen et al. (2016):

$$\pi_{0,stem} = 0.52 - 4.16 \cdot \rho_{wood} \quad (\text{A.3})$$

$$\epsilon_{stem} = \sqrt{1.02 \cdot e^{8.5 \cdot \rho_{wood}} - 2.89} \quad (\text{A.4})$$

while the apoplastic fraction of stem is assumed $f_{apo,stem} = f_{conduits}$ (see A.3.8).

Imputation of leaf pressure-volume parameters, i.e. $\pi_{0,leaf}$ (**LeafPI0**), ϵ_{leaf} (**LeafEPS**) and $f_{apo,leaf}$ (**LeafAF**), is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set **medfate::trait_family_means**). The percentage of species-level explained variance of these imputations is reported in the following table.

If family-level values are missing, following Bartlett et al. (2012) average values for Mediterranean climate leaves are taken as defaults, i.e. $\pi_{0,leaf} = -2$ MPa, $\epsilon_{leaf} = 17$, whereas a 29% leaf apoplastic fraction is assumed (i.e. $f_{apo,leaf} = 0.29$).

A.3.15 Stem and root maximum hydraulic conductivity

Tissue-level maximum conductivity parameters (i.e. $K_{stem,max,ref}$ and $K_{root,max,ref}$) are not direct parameters to simulation functions. Instead, they are scaled to estimate stem- and root-level hydraulic conductances (i.e. $k_{stem,max}$ and $k_{root,max}$) using plant size (see A.4.1 and A.4.3 for details). $K_{stem,max,ref}$ and $K_{root,max,ref}$ are supplied via species parameter table and missing values can therefore occur.

Imputation of $K_{stem,max,ref}$ (**Ks** or **Kmax_stemxylem**) is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using

Table A.7: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of maximum stem hydraulic conductivity.

Trait	spp.	Family (%)	Genus (%)	Total (%)
Ks	1965	3.28	66.19	69.5

Table A.8: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of leaf maximum hydraulic conductance.

Trait	spp.	Family (%)	Genus (%)	Total (%)
kleaf	308	5.41	44.27	49.7

taxonomic family (internal data set `medfate::trait_family_means`). The percentage of species-level explained variance of these imputations is reported in the following table.

If family-level values are missing, suitable $K_{stem,max,ref}$ values are decided according to combinations of taxon group (either Angiosperm or Gymnosperm), growth form (either tree or shrub) and leaf phenology (Maherali et al., 2004):

Group	Growth form	Leaf phenology	$K_{stem,max,ref}$
Angiosperm	Tree	Winter-(semi)deciduous	1.58
Angiosperm	Shrub	Winter-(semi)deciduous	1.55
Angiosperm	Tree/Shrub	Evergreen	2.43
Gymnosperm	Tree	any	0.48
Gymnosperm	Shrub	any	0.24

Following Oliveras et al. (2003), missing values for $K_{root,max,ref}$ are assumed to be four-times the values given or estimated for $K_{stem,max,ref}$.

A.3.16 Leaf maximum hydraulic conductance

Leaf maximum hydraulic conductance ($k_{l,max}$, in $mmol \cdot m^{-2} \cdot s^{-1} \cdot MPa^{-1}$) is not scaled, unlike stem hydraulic conductance, but an input parameter that should be provided for each species. Imputation of $k_{l,max}$ is first performed at the genus level using the corresponding row of `SpParams`. The percentage of species-level explained variance of these imputations is reported in the following table.

When genus-level averages are also missing, leaf maximum hydraulic conductance is estimated from a relationship with maximum stomatal conductance (g_{swmax}), following Franks (2006) (original coefficients were modified for better fit):

$$k_{l,max} = (g_{swmax}/0.015)^{1/1.3} \quad (A.5)$$

Note that values for g_{swmax} may also be imputed (see A.3.13), and that this imputation is uncertain, so that coordination with g_{swmax} should be avoided. Alternatively, $k_{l,max}$ has also been related to maximum photosynthetic capacity and leaf nitrogen content (Scoffoni et al., 2016), and to leaf xylem vulnerability (Yan et al., 2020). Focusing on its relationships with anatomical features, leaf maximum hydraulic conductance is known to be coordinated with vein density (also known as vein length per unit area), of which 80% of its length comes from minor veins (Scoffoni et al., 2016).

Note that leaf hydraulic conductance is influenced by petiole and vein traits that affect the hydraulic conductance of the leaf xylem (leaf apoplastic conductance) and of the pathways outside of the xylem, that is, through the bundle sheath and the mesophyll (extra-xylem or symplastic conductance) (Sack and Holbrook, 2006; Sack and Scoffoni, 2013). In `medfate`, leaf extra-xylary conductance is assumed to contribute, by default, to 50% of total maximum leaf hydraulic conductance (see control parameter `fractionLeafSymplasm`). Vein architecture contributes to both xylary and extra-xylary components: the former because of wider and more numerous xylem conduits increase water conductance; the latter because minor vein density decreases the distance for water to move from the xylem to the transpiration site (Sack and Scoffoni, 2013).

A.3.17 Xylem vulnerability

Table A.9: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of stem P50.

Trait	spp.	Family (%)	Genus (%)	Total (%)
VCstem_P50	1942	18.79	39.82	58.6

Table A.10: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of root P50.

Trait	spp.	Family (%)	Genus (%)	Total (%)
VCroot_P50	137	22.39	44.92	67.3

Imputation of $\Psi_{50,stem}$ is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set **medfate::trait_family_means**). The percentage of species-level explained variance of these imputations is reported in the following table.

If family-level values is missing, a suitable estimate of $\Psi_{50,stem}$ the water potential corresponding to 50% of conductance loss, can be obtained from Maherali et al. (2004) according to combinations of taxon group (either Angiosperm or Gymnosperm), growth form (either tree or shrub) and leaf phenology:

Group	Growth form	Leaf phenology	$\Psi_{50,stem}$
Angiosperm	Tree/Shrub	Winter-(semi)deciduous	-2.34
Angiosperm	Tree	Evergreen	-1.51
Angiosperm	Shrub	Evergreen	-5.09
Gymnosperm	Tree	any	-4.17
Gymnosperm	Shrub	any	-8.95

$\Psi_{50,stem}$ estimates are taken for parameter Ψ_{critic} , in the case of the basic water balance model.

Water potentials corresponding to 12% and 88% PLC ($\Psi_{12,stem}$ and $\Psi_{88,stem}$), when missing are not estimated using genus-level averages, in order to avoid inconsistencies in the stem vulnerability curve (i.e., with respect to $\Psi_{50,stem}$). When $\Psi_{88,stem}$ is missing, a regression equation is used to estimate it from $\Psi_{50,stem}$ ($R^2_{adj} = 0.633$; data compiled using **forestables**):

$$\Psi_{88,stem} = 1.43637 \cdot \Psi_{50,stem} \quad (A.6)$$

Similarly, when $\Psi_{12,stem}$ is missing, a similar regression equation is used to estimate it from $\Psi_{50,stem}$ ($R^2_{adj} = 0.724$; data compiled using **forestables**):

$$\Psi_{12,stem} = 0.626191 \cdot \Psi_{50,stem} \quad (A.7)$$

Finally, estimates of the Weibull function (c_{stem} and d_{stem}) are obtained from $\Psi_{50,stem}$ and $\Psi_{88,stem}$ via function **hydraulics_psi2Weibull()**.

Root vulnerability

Imputation of $\Psi_{50,stem}$ is first performed at the genus level using the corresponding row of **SpParams**. The percentage of species-level explained variance of imputations using genus averages is reported in the following table.

Note that vulnerability curves for root xylem are less common than for stem xylem. If genus-level averages are missing, a linear relationship is used to estimate $\Psi_{50,root}$ from $\Psi_{50,stem}$ ($R^2_{adj} = 0.499$; data compiled using **forestables**):

$$\Psi_{50,root} = 0.627 \cdot \Psi_{50,stem} \quad (A.8)$$

Finally, $\Psi_{88,stem}$ and the Weibull vulnerability parameters are obtained as explained for stems.

Water potentials corresponding to 12% and 88% root PLC ($\Psi_{12,root}$ and $\Psi_{88,root}$), when missing are not estimated using genus-level averages, in order to avoid inconsistencies in the root vulnerability curve (i.e.,

Table A.11: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of leaf P50.

Trait	spp.	Family (%)	Genus (%)	Total (%)
VCleaf_P50	634	10.37	39.65	50

with respect to $\Psi_{50,root}$). Similarly to the stem vulnerability curve, when $\Psi_{88,root}$ is missing, a regression equation is used to estimate it from $\Psi_{50,root}$ ($R_{adj}^2 = 0.742$; data compiled using **forestables**):

$$\Psi_{88,root} = 1.57274 \cdot \Psi_{50,root} \quad (A.9)$$

Similarly, when $\Psi_{12,root}$ is missing, a similar regression equation is used to estimate it from $\Psi_{50,root}$ ($R_{adj}^2 = 0.804$):

$$\Psi_{12,root} = 0.52103 \cdot \Psi_{50,root} \quad (A.10)$$

Finally, estimates of the Weibull function (c_{root} and d_{root}) are obtained from $\Psi_{50,root}$ and $\Psi_{88,root}$ via function `hydraulics_psi2Weibull()`.

Leaf vulnerability

Imputation of $\Psi_{50,leaf}$ is first performed at the genus level using the corresponding row of **SpParams**. The percentage of species-level explained variance of imputations using genus averages is reported in the following table.

Vulnerability curves for leaf xylem are also less common than for stem xylem. If genus-level averages values are missing, a relationship calibrated with data from Bartlett et al. (2016) is used to estimate $\Psi_{50,leaf}$ from the water potential at turgor loss point (Ψ_{tlp}):

$$\Psi_{50,leaf} = 0.2486 + 0.9944 \cdot \Psi_{tlp} \quad (A.11)$$

Water potentials corresponding to 12% and 88% leaf PLC ($\Psi_{12,leaf}$ and $\Psi_{88,leaf}$), when missing are not estimated using genus-level averages, in order to avoid inconsistencies in the leaf vulnerability curve (i.e., with respect to $\Psi_{50,leaf}$). Similarly to the stem vulnerability curve, when $\Psi_{88,leaf}$ is missing, a regression equation is used to estimate it from $\Psi_{50,leaf}$ ($R_{adj}^2 = 0.765$; data compiled using **forestables**):

$$\Psi_{88,leaf} = 1.28837 \cdot \Psi_{50,leaf} \quad (A.12)$$

Similarly, when $\Psi_{12,leaf}$ is missing, a similar regression equation is used to estimate it from $\Psi_{50,leaf}$ ($R_{adj}^2 = 0.766$):

$$\Psi_{12,leaf} = 0.76389 \cdot \Psi_{50,leaf} \quad (A.13)$$

Finally, estimates of the Weibull function (c_{leaf} and d_{leaf}) are obtained from $\Psi_{50,leaf}$ and $\Psi_{88,leaf}$ via function `hydraulics_psi2Weibull()`.

A.3.18 Photosynthesis rates

Rubisco's maximum carboxylation rate at 25°C ($V_{max,298}$, in $\mu\text{molCO}_2 \cdot \text{s}^{-1} \cdot \text{m}^{-2}$) is a required input parameter for each species (**Vmax298**). When missing, the work by Walker et al. (2014) suggests that suitable estimates can be derived from SLA and N_{area} , the latter being the nitrogen concentration per leaf area:

$$V_{max,298} = e^{1.993 + 2.555 \cdot \log(N_{area}) - 0.372 \cdot \log(SLA) + 0.422 \cdot \log(N_{area}) \cdot \log(SLA)} \quad (A.14)$$

In turn, imputation for SLA is explained in A.3.5, whereas values for N_{area} are determined from N_{leaf} and SLA , being N_{leaf} estimated as indicated in A.3.19. Would N_{leaf} and SLA values be both unavailable, a default value of $100 \mu\text{molCO}_2 \cdot \text{s}^{-1} \cdot \text{m}^{-2}$ is used for $V_{max,298}$ imputation.

Table A.12: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of tissue nitrogen concentration.

Trait	spp.	Family (%)	Genus (%)	Total (%)
Nleaf	11920	17.43	38.78	56.2
Nsapwood	65	34.11	16.79	50.9
Nfineroot	1793	12.57	39.50	52.1

When the maximum rate of electron transport at the same temperature ($J_{max,298}$) is not provided by the user, it can be estimated from $V_{max,298}$ using (Walker et al., 2014):

$$J_{max,298} = e^{1.197+0.847 \cdot \log(V_{max,298})} \quad (\text{A.15})$$

A.3.19 Maintenance respiration rates

Imputation of maintenance respiration rates for leaves, sapwood and fine roots (RER_{leaf} , $RER_{sapwood}$ and $RER_{fineroot}$; all in $g\text{ gluc} \cdot g\text{ dry}^{-1} \cdot \text{day}^{-1}$) is first performed at the genus level using the corresponding row of **SpParams**. When missing at the genus, they are estimated from corresponding tissue nitrogen concentrations (N_{leaf} , $N_{sapwood}$ and $N_{fineroot}$; all in $mg\text{ N} \cdot g\text{ dry}^{-1}$) following the equations given by Reich et al. (2008) (after appropriate unit conversion):

$$RER_{leaf} = e^{0.691+1.639 \cdot \log(N_{leaf})} \quad (\text{A.16})$$

$$RER_{sapwood} = e^{1.024+1.344 \cdot \log(N_{sapwood})} \quad (\text{A.17})$$

$$RER_{fineroot} = e^{0.980+1.352 \cdot \log(N_{fineroot})} \quad (\text{A.18})$$

where in the previous equations nitrogen concentrations are in $mmol\text{ N} \cdot g\text{ dry}^{-1}$ and respiration rates in $nmol\text{ CO}_2 \cdot g\text{ dry}^{-1} \cdot s^{-1}$.

In turn, imputation of tissue nitrogen concentration is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set **medfate::trait_family_means**). The percentage of species-level explained variance of these imputations is reported in the following table.

When family values are also missing, default tissue-averaged nitrogen concentrations are given: $N_{leaf} = 20.088$, $N_{sapwood} = 3.9791$ and $N_{fineroot} = 12.207$.

Default **control** values ($MR_{leaf} = 0.00260274$), sapwood ($MR_{sapwood} = 6.849315e - 05$) and fine roots ($MR_{fineroot} = 0.002054795$) were used in previous model versions, derived from Ogle and Pacala (2009), but these are no longer used because of easier parameterization using tissue nitrogen concentration.

A.3.20 Relative growth rates

When missing at the species parameter table, maximum relative growth rates for leaves, sapwood and fine roots are taken from **control** parameters. Default values are provided in 15.5.3.

A.3.21 Senescence rates

When missing at the species parameter table, senescence rates for sapwood and fine roots are taken from **control** parameters. The default daily senescence rate for fine roots is $0.001897231\text{ day}^{-1}$ which corresponds to a 50% annual turnover rate (Gill and Jackson, 2000).

A.3.22 Relative starch for sapwood growth

When missing at the species parameter table, the minimum relative starch for sapwood growth is taken from **control** parameters. Default value is provided in 15.5.3.

Table A.13: Number of species with available data and percentage of variance explained individually by family and genus, and total explained variance for taxonomically-based imputations of wood carbon content.

Trait	spp.	Family (%)	Genus (%)	Total (%)
WoodC	614	17.91	27.76	45.7

A.3.23 Wood carbon

Imputation of C_{wood} is first performed at the genus level using the corresponding row of **SpParams**. When genus-level averages are also missing, imputation is performed using taxonomic family (internal data set **medfate::trait_family_means**). The percentage of species-level explained variance of these imputations is reported in the following table.

If family-level values are also missing, default value of $C_{wood} = 0.5 \text{ g C} \cdot \text{g dry}^{-1}$ is used.

A.3.24 Mortality baseline rate

When missing at the species parameter table, the mortality baseline rate is taken from **control** parameters. Default value is provided in 15.5.3.

A.3.25 Recruitment

Imputation of missing values for recruitment is specified via **control** parameters. Default values are provided in 19.4.3.

A.3.26 Flammability

Default values for the surface-area-to-volume ratio (σ), fuel heat content (h) and lignin percent (LI) are defined from leaf size and leaf shape as follows:

Leaf shape	Leaf size	σ	h	LI
Broad	Large	5740	19740	15.50
Broad	Medium	4039	19825	20.21
Broad	Small	4386	20062	22.32
Linear/Needle	Large	3620	18250	24.52
Linear/Needle	Medium	4758	21182	24.52
Linear/Needle	Small	6697	21888	18.55
Spines	[any]	6750	20433	14.55
Scale	[any]	1120	20504	14.55

Default value for the density of fuel particles (ρ_p) is $400 \text{ kg} \cdot \text{m}^{-3}$.

A.4 Scaling size-related parameters

A.4.1 Stem xylem maximum conductance

Estimation of maximum stem conductance ($k_{s,max}$, in $\text{mmol} \cdot \text{m}^{-2} \cdot \text{s}^{-1} \cdot \text{MPa}^{-1}$) is done by function **hydraulics_maximumStemHydraulicConductance()** and follows the work by Savage et al. (2010), Olson et al. (2014) and Christoffersen et al. (2016). Calculations are based on tree height and two species-specific parameters: maximum sapwood reference conductivity ($K_{s,max,ref}$) and the ratio of leaf area to sapwood area (A_l/A_s ; **A12As** in **SpParamsMED**), i.e. the inverse of the Huber value H_v .


The reference value for maximum sapwood conductivity $K_{s,max,ref}$ is assumed to have been measured on a *terminal branch* of a plant of known height H_{ref} . If our target plant is very different in height, the conduits of terminal branches will have different radius and hence conductivity. We correct the reference conductivity to the target plant height using the following empirical relationship, developed by Olson et al. (2014) between tree height and diameter of conduits for angiosperms and the equation described by Christoffersen et al. (2016):

$$2 \cdot r_{int,H} = 10^{1.257+(0.24 \cdot \log_{10}(H))} \quad (\text{A.19})$$

$$2 \cdot r_{int,ref} = 10^{1.257+(0.24 \cdot \log_{10}(H_{ref}))} \quad (\text{A.20})$$

$$K_{s,max,cor} = K_{s,max,ref} \cdot (r_{int,H}/r_{int,ref})^2 \quad (\text{A.21})$$

Where $r_{int,H}$ is the radius of conduits for a terminal branch of a tree of height H and $r_{int,ref}$ is the corresponding radius for a tree of height H_{ref} (H and H_{ref} are measured in m). The form of the empirical relationship by Olson et al. (2014) is:



Let's consider an example for a *Quercus ilex* target tree of 4m height and where species-specific conductivity $K_{s,max,ref} = 0.77$ is the apical value for trees of $H_{ref} = 6.6$ m (in `medfate`, values of H_{ref} are taken from median height values; see parameter `Hmed` in `SpParamsMED`). The corrected conductivity for a tree of height 4 m will be a bit lower than that of the reference height:

```
xylem_kmax = 0.77
H = 400 # in cm
Href = 660 # in cm
f = hydraulics_referenceConductivityHeightFactor(Href, H);
f
```

```
## [1] 0.7863352
```

```
xylem_kmax_cor = xylem_kmax * f
xylem_kmax_cor
```

```
## [1] 0.6054781
```


Once the reference conductivity is corrected, the maximum stem conductance without accounting for conduit taper is:

$$k_{s,max,notaper} = \frac{1000}{0.018} \frac{K_{s,max,cor} \cdot A_s}{H \cdot A_l} \quad (\text{A.22})$$

where H is the tree height (here in m), A_s is the sapwood area, A_l is the leaf area and $1000/0.018$ is a factor used to go from kg to mmol. The ratio $A_l/A_s = 1/H_v$ is a fixed species parameter in soil water balance calculations (see parameter `A12As`), but becomes variable when simulating plant growth. Let's assume that *Quercus ilex* the leaf to sapwood area ratio is $A_l/A_s = 2512$. The maximum (leaf-specific) stem conductance without taper ($k_{s,max,notaper}$) for the tree of 4 m height is then:

```
A12As = 2512
```

```
kstemmax = hydraulics_maximumStemHydraulicConductance(xylem_kmax,
                                                         Href, A12As, H, taper = FALSE)
kstemmax
```

```
## [1] 3.347698
```

In order to consider taper of xylem conduits we calculate the whole-tree conductance per unit leaf area ($k_{s,max,taper}$) as described in Christoffersen et al. (2016):

$$k_{s,max,taper} = \frac{1000}{0.018} \cdot \frac{K_{s,max,pet} \cdot A_s}{H \cdot A_l} \cdot \chi_{tap:notap,ag}(H) \quad (\text{A.23})$$

where $K_{s,max,pet}$ is the conductivity at the petiole level and $\chi_{tap:notap,ag}(H)$ is the taper factor accounting for the decrease in the xylem conduits diameter with the height, from the petiole to base of the trunk, which mitigates the negative effects of height on the hydraulic safety. The conductivity at the petiole level is obtained from $K_{s,max,ref}$ using again:


$$K_{s,max,pet} = K_{s,max,ref} \cdot (r_{int,pet}/r_{int,ref})^2 \quad (\text{A.24})$$

where $r_{int,pet}$ is the radius of the petiole in the model of Savage et al. (2010). Christoffersen et al. (2016) use $r_{int,pet} = 10 \mu m$ but we define it as the radius of apical conduits in a tree of 1 m height:

```
hydraulics_terminalConduitRadius(100.0)
```

```
## [1] 9.035871
```

$\chi_{tap:notap,ag}(H)$ is calculated as described in the Appendix 1 section of Christoffersen et al. (2016) (see also Savage et al. (2010)). The following figure shows the value of $\chi_{tap:notap,ag}$ for different heights:



Note that, since $\chi_{tap:notap,ag}(1) = 3.82$ (indicated using grey dashed lines in the last figure), the equation of maximum conductance with taper would give a higher conductance than the equation without taper for a tree of 1 m height, which is supposed to have a conductance equal to conductivity. To solve this issue we define the taper factor as $\chi_{tap:notap,ag}(H)/\chi_{tap:notap,ag}(1)$:


$$k_{s,max,taper} = \frac{1000}{0.018} \cdot \frac{K_{s,max,pet} \cdot A_s}{H \cdot A_l} \cdot \frac{\chi_{tap:notap,ag}(H)}{\chi_{tap:notap,ag}(1)} \quad (A.25)$$

The maximum stem conductance with taper ($k_{s,max,taper}$) of a *Q. ilex* tree of 4 m height, calculated with this second equation, is:

```
kstemmax_tap = hydraulics_maximumStemHydraulicConductance(xylem_kmax,
    Href, A12As, H, taper = TRUE)
kstemmax_tap
```

```
## [1] 4.764396
```

The next two plots show the variation of $k_{s,max}$ for *Q. ilex* depending on the tree height and with/without considering taper of conduits. The plot on the right (both axes in log) show the slope of the dependency of conductance with height in both cases:



A.4.2 Leaf maximum hydraulic conductance

A.4.3 Root xylem maximum hydraulic conductance

To obtain maximum root xylem conductance ($k_{r,max}$, in $\text{mmol} \cdot \text{m}^{-2} \cdot \text{s}^{-1} \cdot \text{MPa}^{-1}$), one option taken by Christoffersen et al. (2016) is to assume that minimum stem resistance (inverse of maximum conductance) represents a fixed proportion of the minimum total tree (stem+root) resistance. A value 0.625 (i.e. 62.5%) suggested by these authors leads to maximum total tree conductance for our *Q. ilex* tree being:

```
ktot = kstemmax*0.625
ktot
```

```
## [1] 2.092311
```

and the maximum root xylem conductance would be therefore:

```
krootmax = 1/((1/ktot)-(1/kstemmax))
krootmax
```

```
## [1] 5.579497
```

Now, we need to divide total maximum conductance of the root system xylem among soil layers we need weights inversely proportional to the length of transport distances (Sperry et al., 2017). Vertical transport lengths can be calculated from soil depths and radial spread can be calculated assuming cylinders with volume proportional to the proportions of fine root biomass. Let's assume a soil with three layers:

```
s <- soil(defaultSoilParams(3))
d <- s$widths
d
```

```
## [1] 300 700 1000
```

The proportion of fine roots in each layer, assuming a linear dose response model, will be:

```
Z50 <- 200
Z95 <- 1500
v1 <- root_ldrDistribution(Z50,Z95, NA, d)
v1
```

```
##           [,1]      [,2]      [,3]
## [1,] 0.6661036 0.2784153 0.05548106
```

Having this information, the calculation of root length (i.e. the sum of vertical and radial lengths) to each layer (L_j) is done using function `root_coarseRootLengths()`:

```
rfc <- c(20,50,70)
Vol <- root_coarseRootSoilVolumeFromConductance(1.0, 2500,krootmax,
                                                v1, d, rfc)
l <- root_coarseRootLengthsFromVolume(Vol, v1, d, rfc)
l
```

```
## [1] 4503.445 3250.363 3085.055
```

where lengths are in mm. The proportion of total root xylem conductance corresponding to each layer (w_j) is given by `root_xylemConductanceProportions()`:

```
w1 <- hydraulics_rootxylemConductanceProportions(v1, l)
w1
```

```
## [1] 0.09131268 0.15767645 0.75101087
```

Xylem conductance proportions can be quite different than the fine root biomass proportions. This is because radial lengths are largest for the first top layers and vertical lengths are largest for the bottom layers. The maximum root xylem conductances of each layer will be the product of maximum total conductance of root xylem and weights:

```
w1*krootmax
```

```
## [1] 0.5094788 0.8797553 4.1902629
```

The maximum root xylem conductances of each layer would be:

```
krootmaxvec = w1*krootmax
krootmaxvec
```


```
## [1] 0.5094788 0.8797553 4.1902629
```

and the fraction of total xylem resistance due to stem would be:

```
(1/kstemmax)/((1/kstemmax)+(1/krootmax))
```

```
## [1] 0.625
```


In contrast with the approach of Christoffersen et al. (2016), in this approach the root maximum conductance depends root length and distribution, and is not a fixed fraction of stem maximum conductance. Assuming constant root length, then the proportion of total resistance due to the stem will increase with tree height (Magnani et al., 2000):



where the horizontal gray line indicates the value of 62.5%. Of course rooting depth also increases with tree age, but young trees have higher root-to-shoot ratios than older ones. Hence, a root maximum conductance that is not fixed but increases with age seems a priori more realistic. Moreover, Christoffersen et al. (2016) justify the value of 62.5% from a study which quantified total aboveground and belowground resistance in tropical trees (Fisher et al., 2006) under near-saturated (wet season) conditions, but values of belowground resistance reported in this study for wet conditions and trees of 30 m height are around 13%, which equals to 87% fraction of aboveground resistance. On the other hand, while rooting depths are limited by soil depth, lateral root length increases with age and, hence, the model could be made more realistic if this is taken into account and the curve above would probably saturate at lower percentages.

A.4.4 Rhizosphere maximum hydraulic conductance

Maximum rhizosphere conductance ($k_{rh,max}$, in $mmol \cdot m^{-2} \cdot s^{-1} \cdot MPa^{-1}$) is difficult to measure directly, as it depends on the rhizosphere (i.e. fine root) surface in each soil layer, and will probably always be a parameter to be calibrated. Instead of trying to estimate rhizosphere surface from root architecture (Sperry et al., 1998), we follow Sperry et al. (2016) and determine the maximum rhizosphere conductance in each layer from an inputted ‘average percentage rhizosphere resistance’. The percentage of continuum resistance corresponding to the rhizosphere is calculated from the vulnerability curves of stem, root and rhizosphere at the same water potential. The average resistance is found by evaluating the percentage for water potential values between 0 and Ψ_{crit} . The following figure illustrates how the supply function, for different soil water potentials, is affected by increasing values of the average percentage of rhizosphere resistance:



Sperry et al. (2016) found average percentages of rhizosphere resistance around 67%, but these exceptionally-high values were probably a consequence of using an unsegmented supply function (i.e. single vulnerability curve for roots, stem and leaves). If we specify a 15% of average resistance in the rhizosphere (see parameter `averageFracRhizosphereResistance` in function `defaultControl()`), the maximum rhizosphere conductance values for the three layers are found calling:

```
krmax = rep(0,3)
krmax[1] = hydraulics_findRhizosphereMaximumConductance(15,
  s$VG_n[1], s$VG_alpha[1],
  krootmax, rootc, rootd,
  kstemmax, stemc, stemd,
  kleafmax, leafc, leafd, -2, 0)
krmax[2] = hydraulics_findRhizosphereMaximumConductance(15,
  s$VG_n[2], s$VG_alpha[2],
  krootmax, rootc, rootd,
  kstemmax, stemc, stemd,
  kleafmax, leafc, leafd, -2, 0)
krmax[3] = hydraulics_findRhizosphereMaximumConductance(15,
  s$VG_n[3], s$VG_alpha[3],
  krootmax, rootc, rootd,
  kstemmax, stemc, stemd,
  kleafmax, leafc, leafd, -2, 0)
krmax
```

```
## [1] 48037860 48037860 48037860
```

The values are the same because the texture of the three layers is the same in this case. If we take into account root distribution, actual maximum rhizosphere conductance values are:

```
krmax*v1
```

```
##           [,1]      [,2]      [,3]
## [1,] 31998192 13374476 2665192
```

A.4.5 Plant water storage capacity

The water storage capacity of sapwood tissue per leaf area unit ($V_{sapwood}$; in $l \cdot m^{-2}$) can be estimated as the product of stem height (H in m) and Huber value (H_v ; ratio of sapwood area to leaf area in $m^2 \cdot m^{-2}$) times a factor to account for the non-cylindrical shape (<http://www.fao.org/forestry/17109/en/>):

$$V_{sapwood} = 10^3 \cdot 0.48 \cdot H \cdot H_v \cdot \Theta_{sapwood} \quad (A.26)$$

$\Theta_{sapwood}$ is sapwood porosity (cm^3 of water per cm^3 of sapwood tissue), which can be estimated from wood density (ρ_{wood} ; in $g \cdot cm^{-3}$):

$$\Theta_{sapwood} = 1 - (\rho_{wood}/1.54) \quad (A.27)$$

where the density of wood substance can be assumed to be fixed and equal to $1.54 g \cdot cm^{-3}$ (Dunlap, 1914). For example, wood densities ranging from 0.443 to $1.000 g \cdot cm^{-3}$ result in sapwood porosity values between 0.35 and 0.71.

Water storage capacity of leaf tissue per leaf area unit (V_{leaf} ; in $l \cdot m^{-2}$) can be estimated as the product of specific leaf area (SLA; in $m^2 \cdot kg^{-1}$) and leaf density (ρ_{leaf} ; in $g \cdot cm^{-3}$):

$$V_{leaf} = \frac{1}{SLA \cdot \rho_{leaf}} \cdot \Theta_{leaf} \quad (A.28)$$

Θ_{leaf} is leaf porosity (cm^3 of water per cm^3 of leaf tissue), which can be estimated from leaf density:

$$\Theta_{leaf} = 1 - (\rho_{leaf}/1.54) \quad (A.29)$$

where the density of wood substance can be assumed to be fixed and equal to $1.54 g \cdot cm^{-3}$ (Dunlap 1914).

For example, let's calculate the stem and leaf water capacity for a Q. ilex tree of 15 m height:

```
ld = 0.7
SLA = 5.870
moisture_leafWaterCapacity(SLA, ld)
```

```
## [1] 0.1327463
```

A.4.6 Horizontal root overlap

A.4.6.1 Basic water balance

A given plant cohort i will have its roots in layer s partitioned among different water pools. We thus need to define $fr_{i,s,j}$, the (horizontal) proportion of fine roots of cohort i in layer s of the water pool j , with the restriction that:

$$\sum_j fr_{i,s,j} = 1 \quad \forall i, s \quad (A.30)$$

It is important to realize that proper estimation of $fr_{i,s,j}$ is challenging when we do not have explicit plant coordinates, root lateral widths, etc. For this reason, an intuitive approach is followed here based on the following two premises:

- The amount of overlap between roots of different plants should monotonically increase along with the LAI of the stand (i.e. LAI_{stand}^{live}).
- The amount of overlap between roots of different plants at a given soil depth should increase/decrease with the vertical proportion of roots at that depth.

The specific formulation we chose for $fr_{i,s,j}$ is:

$$fr_{i,s,j} = f_{pool,j} \cdot (1 - \exp(-f_{overlap} \cdot LAI_{stand}^{live})) \cdot \left(\frac{\sqrt{FRP_{i,s} \cdot FRP_{j,s}}}{\max_s(FRP_{i,s})} \right) \quad (A.31)$$

for all $j \neq i$, where $f_{overlap}$ is an overlap factor (a control parameter called `poolOverlapFactor`). For $j = i$ then we simply have:

$$fr_{i,s,i} = 1 - \sum_j fr_{i,s,j} \quad (\text{A.32})$$

Note that if $f_{overlap} = 0$ then $fr_{i,s,j} = 1$ if $j = i$ and zero otherwise (i.e. plants exploit their corresponding water pools only). For very large values of LAI_{stand}^{live} and/or $f_{overlap}$ we have that $(1 - \exp(-f_{overlap} \cdot LAI_{stand}^{live})) = 0$ and $fr_{i,s,j} = f_{pool,j}$ (neglecting vertical differences), so that plants exploit the each water pool in the same proportion as the fraction of stand occupied by the pool (i.e. overlap is maximum).

Appendix B

Symbols

The following tables list all symbols used in this document, along with their units and definition. When symbols are input for **medfate** model functions, the R name of those parameters in the package (either in data frame **SpParamsMED**, soil input data frame, or the output of functions such as **spwbInput()**) is also indicated.

B.1 Dimensions

Symbol	Units	R	Description
Δt_{step}	s	<code>86400/ndailysteps</code>	Sub-daily temporal step
$\Delta t_{substep}$	s	<code>86400/(ndailysteps*ndailyubsteps)</code>	Sub-daily substep
c	(count)		Number of plant cohorts
l	(count)		Number of vertical layers
Δz	m	<code>verticalLayerSize/100</code>	Width of vertical layers
S	(count)		Number of soil layers
d_s	mm	<code>widths</code>	Width of soil layer s
Z_s	mm	<code>widths</code>	Depth of soil layer s
Z_{soil}	mm		Overall soil depth

B.2 Soils

Symbol	Units	R	Description
$P_{clay,s}$	%	<code>clay</code>	Percent of clay in soil layer s
$P_{sand,s}$	%	<code>sand</code>	Percent of sand in soil layer s
OM_s	%	<code>om</code>	Percentage of organic mater per dry weight in soil layer s
BD_s	$g \cdot cm^{-3}$	<code>bd</code>	Bulk density in soil layer s
$P_{rocks,s}$	%	<code>rfc</code>	Percentage of rock fragment content in soil layer s
θ_s	$m^3 \cdot m^{-3}$		Volumetric moisture in soil layer s
Ψ_s	MPa		Water potential in soil layer s
Ψ_{fc}	MPa		Water potential at field capacity
A_s, B_s			Parameters of the Saxton pedotransfer functions for soil layer s
$\theta_{sat,s}$	$m^3 \cdot m^{-3}$	<code>VG_theta_sat</code>	Volumetric moisture at saturation for soil layer s
$\theta_{fc,s}$	$m^3 \cdot m^{-3}$		Volumetric moisture at field capacity (-0.033 MPa) for soil layer s

Symbol	Units	R	Description
$\theta_{wp,s}$	$m^3 \cdot m^{-3}$		Volumetric moisture at wilting point (-1.5 MPa) for soil layer s
$\theta_{res,s}$	$m^3 \cdot m^{-3}$	VG_theta_res	Residual volumetric moisture for soil layer s
n_s		VG_n	Parameter n of the Van Genuchten (1980) model for soil layer s
α_s	MPa^{-1}	VG_alpha	Parameter α of the Van Genuchten (1980) model for soil layer s
$K_{sat,s}$	$mmol \cdot s^{-1} \cdot m^{-1} \cdot MPa^{-1}$	Ksat	Saturated soil conductivity for soil layer s .
$K_{unsat,s}$	$mmol \cdot s^{-1} \cdot m^{-1} \cdot MPa^{-1}$	Kunsat	Unsaturated soil conductivity for soil layer s .
$P_{macro,s}$	%	macro	Percentage of macroporosity corresponding to soil layer s .
γ_{soil}	$mm \cdot day^{-1}$	Gsoil	The maximum daily evaporation from soil
Z_{sat}	mm		Depth to saturation
V_s	mm		Water volume in soil layer s
V_{soil}	mm		Overall water volume in the soil
$V_{fc,soil}$	mm		Water retention capacity of the whole soil
W_s	[0-1]	W	Proportion of moisture in relation to field capacity in soil layer s
$W_{i,s}$	[0-1]	W	Proportion of moisture in relation to field capacity in soil layer s within fraction of stand area covered by cohort i
$W_{rhizo,i,s}$	[0-1]		Proportion of moisture in relation to field capacity in soil layer s within the rhizosphere of cohort i
S_{snow}	mm	SWE	Snow water equivalent of the snow pack storage over the soil surface

B.3 Plant/leaf classification

Symbol	Units	R	Description
GF	Categorical	GrowthForm	“Tree”, “Shrub” or “Tree/Shrub”
LF	Categorical	LifeForm	Raunkiaer life form
L_{shape}	Categorical	LeafShape	Leaf shape: “Broad”, “Needle”, “Linear”, “Scale”, “Spines” or “Succulent”
L_{size}	Categorical	LeafSize	Leaf size: “Small” (< 225 mm), “Medium” (> 225 mm & < 2025 mm) or “Large” (> 2025 mm)
L_{pheno}	Categorical	PhenologyType	Leaf phenology type

B.4 Vegetation

Symbol	Units	R	Description
SP_i	(count)	Species	Species identity of cohort i
$A_{sh,i}$	cm^2		Area occupied by an average shrub individual of cohort i
$B_{sh,i}$	kg		Fine-fuel biomass of an average shrub individual of cohort i
H_i	cm	Height	Average tree or shrub height of cohort i

Symbol	Units	R	Description
$H_{crown,i}$	cm		Crown base height (i.e. the height corresponding to the first living branch) of cohort i
CCF_i			Crown competition factor of cohort i
CW_i	m		Crown width that a tree of cohort i would have in open-ground conditions
CR_i	[0-1]	CR	Crown ratio (i.e. ratio between crown length and plant height) of cohort i
N_i	ind · ha ⁻¹	N	Density of tree individuals of cohort i
DBH_i	cm	DBH	Diameter at breast height of trees in cohort i
$Cover_i$	%	Cover	Percent cover of shrubs in cohort i
H_{herb}	%	herbHeight	Average height (in cm) of herbaceous vegetation
$Cover_{herb}$	%	herbCover	Percent cover of herbaceous vegetation
FB_i	kg · m ⁻²		Foliar biomass of cohort i
$FB_{tree,i}$	kg		Foliar biomass of a single tree of cohort i
ϕ_i	[0-1]		Fraction of maximum leaf area currently expanded for cohort i
LAI_i^{live}	m ² · m ⁻²	LAI_live	(Maximum) leaf area index (one-side leaf area per surface area of the stand) of woody cohort i
LAI_i^{dead}	m ² · m ⁻²	LAI_dead	Dead leaf area index (one-side dead leaf area per surface area of the stand) of woody cohort i
LAI_i^ϕ	m ² · m ⁻²	LAI_expanded	Current expanded leaf area index (one-side expanded leaf area per surface area of the stand) of woody cohort i
LAI_i^{all}	m ² · m ⁻²		Total leaf area index (live or dead) of woody cohort i
LAI_{woody}^{live}	m ² · m ⁻²		Maximum leaf area index of living leaves of woody vegetation in the stand
LAI_{woody}^{dead}	m ² · m ⁻²		Leaf area index of dead leaves of woody vegetation in the stand
LAI_{woody}^ϕ	m ² · m ⁻²		Current expanded leaf area index of live leaves of woody vegetation in the stand
LAI_{woody}	m ² · m ⁻²		Total leaf area index (live or dead) of woody vegetation in the stand
LAI_{herb}	m ² · m ⁻²		Leaf area index of herbaceous vegetation in the stand
LAI_{stand}	m ² · m ⁻²		Total leaf area index of the stand
LA_i	m ²		Leaf area of an individual of cohort i
$FRP_{i,s}$	[0-1]	V[i,s]	The proportion of fine roots of cohort i in soil layer s
$FRA_{i,s}$	m ²		Fine root area of cohort i in soil layer s
$L_{radial,i,s}$	m		Radial length of coarse roots of cohort i in soil layer s
W_i	kg · m ⁻²		Fine fuel biomass of cohort i
$Z_{50,i}$	mm	Z50	Depth above which 50% of the fine root mass of cohort i is located
$Z_{95,i}$	mm	Z95	Depth above which 95% of the fine root mass of cohort i is located

B.5 Meteorology

Symbol	Units	R	Description
DOY		DOY	Day of the year
T_{mean}	°C	MeanTemperature	Mean daily temperature
T_{min}	°C	MinTemperature	Minimum daily temperature

Symbol	Units	R	Description
T_{max}	$^{\circ}\text{C}$	MaxTemperature	Maximum daily temperature
T_{base}	$^{\circ}\text{C}$		Base temperature for calculation of GDD
RH_{min}	%	MinRelativeHumidity	Minimum daily relative humidity
RH_{max}	%	MaxRelativeHumidity	Maximum daily relative humidity
P	$L \cdot m^{-2} = mm$	Precipitation	Precipitation (including rainfall and snow)
Pr	$L \cdot m^{-2} = mm$	Rainfall	Liquid water precipitation (rainfall)
Ps	$L \cdot m^{-2} = mm$	Snow	Snow precipitation
PET	$L \cdot m^{-2} = mm$	PET	Potential evapotranspiration, preferably calculated using Penman's equation
Rad	$MJ \cdot m^{-2}$	Radiation	Solar radiation after accounting for clouds
u	$m \cdot s^{-1}$	WindSpeed	Wind speed
P_{atm}	kPa		Atmospheric pressure
ρ_{air}	$kg \cdot m^{-3}$		Air density
T_{atm}	$^{\circ}\text{C}$	Tatm	Atmospheric (above-canopy) air temperature
T_{can}	$^{\circ}\text{C}$	Tcan	Canopy air temperature
$T_{air,j}$	$^{\circ}\text{C}$	Tair	Air temperature in canopy layer j
$T_{soil,s}$	$^{\circ}\text{C}$	Tsoil.s	Temperature of soil layer s
e_{atm}	kPa		Atmospheric (above-canopy) water vapor pressure
$e_{air,j}$	kPa	VPair	Water vapor pressure in canopy layer j
C_{atm}	kPa	Catm	Atmospheric (above-canopy) CO_2 concentration
$C_{air,j}$	kPa	VPair	CO_2 concentration in canopy layer j
u_j	$m \cdot s^{-1}$		Wind speed at canopy layer j

B.6 Anatomy

Symbol	Units	R	Description
H_v	$m^2 \cdot m^{-2}$	1/A12As	Huber value (ratio of sapwood area to leaf area)
LW	cm	LeafWidth	Leaf width
SLA	$m^2 \cdot kg^{-1}$	SLA	Specific leaf area
ρ_{leaf}	$g \cdot cm^{-3}$	LeafDensity	Leaf tissue density
ρ_{wood}	$g \cdot cm^{-3}$	WoodDensity	Wood tissue density
$\rho_{fineroot}$	$g \cdot cm^{-3}$	FineRootDensity	Fine root tissue density
$\Theta_{sapwood}$	$m^3 \cdot m^{-3}$		Sapwood porosity (volume of empty spaces over total volume)
Θ_{leaf}	$m^3 \cdot m^{-3}$		Leaf porosity (volume of empty spaces over total volume)
SRL	$cm \cdot g^{-1}$	SRL	Specific root length
RLD	$cm \cdot cm^{-3}$	RLD	Fine root length density (i.e. density of root length per soil volume)
$r_{6.35}$		r635	Ratio between the weight of leaves plus branches and the weight of leaves alone for branches of 6.35 mm

B.7 Radiation

Symbol	Units	R	Description
$k_{PAR,i}$	(unitless)	kPAR	PAR extinction coefficient for cohort i
$k_{SWR,i}$	(unitless)		SWR extinction coefficient for cohort i
k_{LWR}	(unitless)		LWR extinction coefficient
k_b	(unitless)		Extinction coefficient for direct light for cohort i
$k_{d,i}$	(unitless)		Extinction coefficient for diffuse light for cohort i (equal to $k_{PAR,i}$ or $k_{SWR,i}$)
$\alpha_{SWR,i}$	[0-1]	alphaSWR	Short-wave radiation absorbance coefficient for cohort i
$\alpha_{PAR,i}$	[0-1]		PAR absorbance coefficient for cohort i
$\gamma_{SWR,i}$	[0-1]	gammaSWR	Short-wave radiation leaf reflectance (albedo) for cohort i
$\gamma_{SWR,soil}$	[0-1]		Short-wave radiation soil reflectance
$\gamma_{PAR,i}^{PAR}$	[0-1]		PAR leaf reflectance for cohort i
L_{ground}^{PAR}	[0-1]	LgroundPAR	Fraction of PAR reaching the ground
L_{ground}^{SWR}	[0-1]		Fraction of SWR reaching the ground
I_{beam}	$W \cdot m^{-2}$		Instantaneous direct shortwave irradiance from the atmosphere
I_{dif}	$W \cdot m^{-2}$		Instantaneous diffuse shortwave irradiance from the atmosphere
L_{atm}	$W \cdot m^{-2}$		Instantaneous longwave radiation from the atmosphere
$I_{beam,j}$	$W \cdot m^{-2}$		Instantaneous direct shortwave irradiance at the top of canopy layer j
$I_{dif,j}$	$W \cdot m^{-2}$		Instantaneous diffuse shortwave irradiance at the top of canopy layer j
$I_{beam,soil}$	$W \cdot m^{-2}$		Instantaneous direct shortwave irradiance reaching the soil
$I_{dif,soil}$	$W \cdot m^{-2}$		Instantaneous diffuse shortwave irradiance reaching the soil
$LAI_{i,j}^{sunlit}$	$m^2 \cdot m^{-2}$		Leaf area index of sunlit leaves of cohort i in layer j .
$LAI_{i,j}^{shade}$	$m^2 \cdot m^{-2}$		Leaf area index of shade leaves of cohort i in layer j .
$\Phi_{i,j}^{sunlit}$	$W \cdot m^{-2}$		Short-wave radiation absorbed by sunlit leaves of cohort i in layer j , per leaf area unit.
$\Phi_{i,j}^{shade}$	$W \cdot m^{-2}$		Short-wave radiation absorbed by sunlit leaves of cohort i in layer j , per leaf area unit.
$K_{abs,i,j}^{sunlit}$	$W \cdot m^{-2}$		Short-wave radiation absorbed by sunlit foliage of cohort i in layer j , per ground area unit.
$K_{abs,i,j}^{shade}$	$W \cdot m^{-2}$		Short-wave radiation absorbed by sunlit foliage of cohort i in layer j , per ground area unit.

B.8 Water balance

Symbol	Units	R	Description
Ps	mm	Snow	Precipitation as snow
Pr	mm	Rain	Precipitation as rainfall
Sm	mm	Snowmelt	Snowmelt
Pr_{net}	mm	NetRain	Net rainfall
In	mm	Interception	Interception loss
Ru	mm	Runoff	Water exported from the stand as runoff
Ro	mm	Runon	Water imported to the stand as runon
Dd	mm	DeepDrainage	Water exported from the stand as deep drainage
Es	mm	SoilEvaporation	Evaporation from the soil surface
Tr_{woody}	mm	Transpiration	Woody plant transpiration
Tr_{herb}	mm	HerbTranspiration	Herbaceous plant transpiration

Symbol	Units	R	Description
Tr_i	mm		Transpiration of woody plant cohort i

B.9 Energy balance

Symbol	Units	R	Description
$K_{abs,can}$	$W \cdot m^{-2}$	SWRcan	Atmosphere shortwave radiation absorbed by the canopy
$K_{abs,j}$	$W \cdot m^{-2}$		Atmosphere shortwave radiation absorbed by canopy layer j
$K_{abs,soil}$	$W \cdot m^{-2}$	SWRsoil	Atmosphere shortwave radiation absorbed by the soil
$L_{net,can}$	$W \cdot m^{-2}$	LWRcan	Canopy net longwave radiation
$L_{net,j}$	$W \cdot m^{-2}$		Net long-wave radiation of canopy layer j
$L_{net,soil}$	$W \cdot m^{-2}$	LWRsoil	Soil net longwave radiation
$H_{can,atm}$	$W \cdot m^{-2}$	Hcan	Turbulent heat exchange between the canopy and the atmosphere
$H_{can,soil}$	$W \cdot m^{-2}$	Hcansoil	Turbulent heat exchange between the canopy and the soil
H_j	$W \cdot m^{-2}$		Sensible heat flux between canopy layer j and the leaves it contains
LE_{can}	$W \cdot m^{-2}$	LEcan	Energy released as latent heat from the canopy towards the atmosphere
LE_j	$W \cdot m^{-2}$		Energy released as latent heat by canopy layer j
LE_{soil}	$W \cdot m^{-2}$	LEsoil	Energy released as latent heat from the soil
TC_{LAI}	$J \cdot m^{-2} \cdot K^{-1}$	thermalcapacityCAn	Canopy thermal capacitance per LAI unit
TC_{can}	$J \cdot m^{-2} \cdot K^{-1}$		Canopy thermal capacitance
TC_j	$J \cdot m^{-2} \cdot K^{-1}$		Thermal capacitance of canopy layer j
$TC_{soil,s}$	$J \cdot m^{-2} \cdot K^{-1}$		Thermal capacitance of soil layer s
$VHC_{soil,s}$	$J \cdot m^{-3} \cdot K^{-1}$		Volumetric heat capacity of soil in layer s

B.10 Forest hydrology

Symbol	Units	R	Description
P_G	mm		Amount of rainfall needed to saturate the canopy for a given event
S_{canopy}	mm	Cm	Canopy water storage capacity
ER_{ratio}	(unitless)		Ratio between evaporation rate and rainfall rate
C_{canopy}	[0-1]		Canopy cover
PE_{soil}	mm		Evaporation demand from the soil
SE_{soil}	mm		Evaporation supply from the soil
t_{soil}	day		Time needed to evaporate the current water deficit in the first soil layer
PET_{herb}	mm		Potential evapotranspiration in the herb layer

B.11 Plant hydraulics

Symbol	Units	R	Description
$K_{stem,max,ref}$	$kg \cdot s^{-1} \cdot m^{-1} \cdot MPa^{-1}$	Kmax_stemxylem	Maximum stem sapwood reference conductivity per leaf area unit
$K_{root,max,ref}$	$kg \cdot s^{-1} \cdot m^{-1} \cdot MPa^{-1}$	Kmax_rootxylem	Maximum root sapwood reference conductivity per leaf area unit
$k_{stem,max}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	VCstem_kmax	Maximum whole-stem conductance (per leaf area unit)
$k_{root,max}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	VCroot_kmax	Maximum root conductance (per leaf area unit)
$k_{rhizo,max}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$		Maximum rhizosphere conductance (per leaf area unit)
$k_{leaf,max}$	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$	VCleaf_kmax	Maximum leaf conductance (per leaf area unit)
c_{leaf}, d_{leaf}	(unitless), MPa	VCleaf_c, VCleaf_d	Parameters of the vulnerability curve for leaves
c_{root}, d_{root}	(unitless), MPa	VCroot_c, VCroot_d	Parameters of the vulnerability curve for root xylem
c_{stem}, d_{stem}	(unitless), MPa	VCstem_c, VCstem_d	Parameters of the vulnerability curve for stem xylem
Ψ	MPa		Water potential in a given water compartment/segment
Ψ_P	MPa		Turgor water potential in a given water compartment/segment
Ψ_S	MPa		Osmotic (solute) water potential in a given water compartment
Ψ_{cav}	MPa		Minimum water potential experienced by xylem in previous steps (cavitation)
Ψ_{canopy}	MPa		Canopy water potential
Ψ_{leaf}	MPa		Leaf water potential
$\Psi_{rootcrown}$	MPa		Water potential at the root crown
Ψ_{stem}	MPa		Water potential at the end (highest part) of the stem
PLC	[0-1]		Proportion of conductance loss
PLC_{stem}	[0-1]		Proportion of conductance loss in stem xylem tissue
p_{root}	[0-1]	pRootDisc	Relative root conductance leading to hydraulic disconnection from a soil layer
E_i	$mmol \cdot s^{-1} \cdot m^{-2}$		Steady-state water flow through a hydraulic segment i
k_i	$mmol \cdot s^{-1} \cdot m^{-2} \cdot MPa^{-1}$		Hydraulic conductance function for segment i
Ψ_{up}	MPa		Upstream water potential
Ψ_{down}	MPa		Downstream water potential

B.12 Photosynthesis

Symbol	Units	R	Description
$WUE_{\max}$	$gC \cdot mm^{-1}$	WUE	Water use efficiency at VPD = 1kPa and without light or CO2 limitations
WUE_{PAR}		WUE_par	Coefficient describing the progressive decay of WUE with lower light levels
WUE_{CO2}		WUE_co2	Coefficient for WUE dependency on atmospheric CO2 concentration
WUE_{VPD}		WUE_vpd	Coefficient for WUE dependency on vapor pressure deficit
$J_{\max}$	$\mu mol e \cdot m^{-2} \cdot s^{-1}$	Jmax	Maximum rate of electron transport at current leaf temperature
$V_{\max,298}$	$\mu mol CO_2 \cdot s^{-1} \cdot m^{-2}$	Vmax	Rubisco's maximum carboxylation rate at current leaf temperature
$J_{\max,298}$	$\mu mol e \cdot m^{-2} \cdot s^{-1}$	Jmax298	Maximum rate of electron transport at 298K
$V_{\max,298}$	$\mu mol CO_2 \cdot s^{-1} \cdot m^{-2}$	Vmax298	Rubisco's maximum carboxylation rate at 298K
T_{leaf}	$^{\circ}C$		Leaf temperature.
u_{leaf}	$m \cdot s^{-1}$		Leaf-level wind speed.
VPD_{leaf}	kPa		Leaf vapour pressure deficit.
g_w	$mol H_2O \cdot s^{-1} \cdot m^{-2}$		Leaf diffusive conductance to water vapor
g_c	$mol CO_2 \cdot s^{-1} \cdot m^{-2}$		Leaf diffusive conductance to carbon dioxide
g_{sw}	$mol H_2O \cdot s^{-1} \cdot m^{-2}$		Leaf stomatal conductance to water vapour
$g_{sw,min}$	$mol H_2O \cdot s^{-1} \cdot m^{-2}$	Gswmin	Minimum stomatal conductance to water vapour
$g_{sw,max}$	$mol H_2O \cdot s^{-1} \cdot m^{-2}$	Gswmax	Maximum stomatal conductance to water vapour
Φ_{PAR}^{leaf}	$W \cdot m^{-2}$		Photosynthetically active radiation absorbed per leaf area
Q_{PAR}^{leaf}	$\mu mol photon \cdot m^{-2} \cdot s^{-1}$		PAR photon flux density per leaf area
A_c	$\mu mol CO_2 \cdot s^{-1} \cdot m^{-2}$		Rubisco-limited photosynthesis rate
A_e	$\mu mol CO_2 \cdot s^{-1} \cdot m^{-2}$		Electron transport-limited photosynthesis rate
A	$\mu mol CO_2 \cdot s^{-1} \cdot m^{-2}$		Leaf gross photosynthesis rate
A_n	$\mu mol CO_2 \cdot s^{-1} \cdot m^{-2}$		Leaf net photosynthesis rate

B.13 Plant water content

Symbol	Units	R	Description
$LFMC_{\max}$	%	maxFMC	Maximum live fuel moisture content, corresponding to fine fuels (< 6.35 mm twigs and leaves).
ϵ_{leaf}	MPa	LeafEPS	Modulus of elasticity of leaves
ϵ_{stem}	MPa	StemEPS	Modulus of elasticity of symplastic xylem tissue
$\pi_{0,leaf}$	MPa	LeafPIO	Osmotic potential at full turgor of leaves
$\pi_{0,stem}$	MPa	StemPIO	Osmotic potential at full turgor of symplastic xylem tissue
RWC	[0-1]		Relative water content

Symbol	Units	R	Description
RWC_{sym}	[0-1]		Relative water content in the symplasm fraction of a tissue
RWC_{apo}	[0-1]		Relative water content in the apoplasm fraction of a tissue
$V_{segment}$	$l \cdot m^{-2}$		Water capacity of a segment (leaf or stem)
V_{leaf}	$l \cdot m^{-2}$	Vleaf	Leaf water capacity per leaf area unit
$V_{sapwood}$	$l \cdot m^{-2}$	Vsapwood	Sapwood water capacity per leaf area unit

B.14 Stomatal regulation

Symbol	Units	R	Description
$\theta_1(\Psi_{leaf}),$ $\theta_2(\Psi_{leaf})$	[0-1]		Cost functions 1 and 2
$\beta(\Psi_{leaf})$	[0-1]		Gain function
$Profit_1(\Psi_{leaf}),$ $Profit_2(\Psi_{leaf})$	[0-1]		Profit functions 1 and 2
E_{sunlit}	$mmol \cdot s^{-1} \cdot m^{-2}$		Instantaneous transpiration flow rate for sunlit leaves
E_{shade}	$mmol \cdot s^{-1} \cdot m^{-2}$		Instantaneous transpiration flow rate for shade leaves

B.15 Carbon pools

Symbol	Units	R	Description
$V_{storage,leaf,i}$	$L \cdot ind^{-1}$		Leaf (carbon) storage volume for an individual of cohort i
$V_{storage,sapwood,i}$	$L \cdot ind^{-1}$		Sapwood (carbon) storage volume for an individual of cohort i
$SS_{leaf,i}$	$mol\ gluc \cdot L^{-1}$	SugarLeaf	Sugar concentration in leaves of cohort i
$SS_{sapwood,i}$	$mol\ gluc \cdot L^{-1}$	SugarSapwood	Sugar concentration in sapwood of cohort i
$ST_{leaf,i}$	$mol\ gluc \cdot L^{-1}$	StarchLeaf	Starch concentration in leaves of cohort i
$ST_{sapwood,i}$	$mol\ gluc \cdot L^{-1}$	StarchSapwood	Starch concentration in sapwood of cohort i
S_{plant}	$mol\ gluc \cdot ind^{-1}$		Total labile carbon storage in a plant individual
B_{leaf}	$g\ dry \cdot ind^{-1}$		Leaf dry biomass in a plant individual (for respiration costs)
$B_{sapwood}$	$g\ dry \cdot ind^{-1}$		Sapwood dry biomass in a plant individual
$B_{living,sapwood}$	$g\ dry \cdot ind^{-1}$		Sapwood dry biomass corresponding to (living) parenchymatic tissues in a plant individual (for respiration costs)
$B_{fineroot}$	$g\ dry \cdot ind^{-1}$		Fine root dry biomass in a plant individual (for respiration costs)

B.16 Carbon balance

Symbol	Units	R	Description
M_{leaf}	$mol\ gluc \cdot day^{-1}$		Leaf daily maintenance respiration in a plant individual
$M_{sapwood}$	$mol\ gluc \cdot day^{-1}$		Sapwood daily maintenance respiration in a plant individual
$M_{fineroot}$	$mol\ gluc \cdot day^{-1}$		Fine root daily maintenance respiration in a plant individual
MR_{leaf}	$g\ gluc \cdot g\ dry^{-1} \cdot day^{-1}$	RERleaf	Leaf respiration rate at 20 °C
$MR_{sapwood}$	$g\ gluc \cdot g\ dry^{-1} \cdot day^{-1}$	RERsapwood	Living sapwood (parenchymatic tissue) respiration rate at 20 °C
$MR_{fineroot}$	$g\ gluc \cdot g\ dry^{-1} \cdot day^{-1}$	RERfineroot	Fine root respiration rate at 20 °C
G_{leaf}	$mol\ gluc \cdot day^{-1}$		Leaf daily growth respiration in a plant individual
$G_{sapwood}$	$mol\ gluc \cdot day^{-1}$		Sapwood daily growth respiration in a plant individual
$G_{fineroot}$	$mol\ gluc \cdot day^{-1}$		Fine root daily growth respiration in a plant individual
F_{phloem}	$mol\ gluc \cdot day^{-1}$		Daily phloem transport of sugars from leaves to sapwood
SC_{leaf}	$mol\ gluc \cdot day^{-1}$		Daily conversion from leaf sugar to leaf starch
$SC_{sapwood}$	$mol\ gluc \cdot day^{-1}$		Daily conversion from sapwood sugar to sapwood starch
TS_{leaf}	$mol\ gluc \cdot day^{-1}$		Daily translocation of leaf sugars (prior to senescence)
$TS_{sapwood}$	$mol\ gluc \cdot day^{-1}$		Daily translocation of sapwood sugars (prior to senescence)
TT_{leaf}	$mol\ gluc \cdot day^{-1}$		Daily translocation of leaf starch (prior to senescence)
RE_{leaf}	$mol\ gluc \cdot day^{-1}$		Daily root exudation of leaf carbon
$RE_{sapwood}$	$mol\ gluc \cdot day^{-1}$		Daily root exudation of sapwood carbon

B.17 Growth, senescence and mortality

Symbol	Units	R	Description
$RGR_{leaf,max}$	$m^2 \cdot cm^{-2} \cdot day^{-1}$	RGRleafmax	Maximum leaf area daily growth rate, relative to sapwood area
$RGR_{sapwood,max}$	$cm^2 \cdot cm^{-2} \cdot day^{-1}$	RGRsapwoodmax	Maximum daily sapwood relative growth rate (in sapwood area basis)
$RGR_{fineroot,max}$	$g\ dry \cdot g\ dry^{-1} \cdot day^{-1}$	RGRfinerootmax	Maximum daily fine root relative growth rate
LA^{target}	m^2		Leaf area allocation target

Symbol	Units	R	Description
ΔLA_{alloc}	m^2		Leaf area increase dictated by the difference between leaf area allocation target and current leaf area
ΔLA_{source}	m^2		Leaf area increase according to leaf metabolic carbon availability
ΔLA_{sink}	m^2		Leaf area increase dictated by sink limitations (leaf relative growth rate and relative cell expansion rate)
ΔLA	m^2		Actual leaf area increase
ΔSA_{source}	cm^2		Sapwood area increase according to sapwood storage carbon availability
ΔSA_{sink}	cm^2		Sapwood area increase dictated by sink limitations
ΔSA	cm^2		Actual sapwood area increase
$B_{fineroot,target}$	$g\ dry$		Fine root biomass allocation target
$\Delta B_{fineroot,alloc}$	$g\ dry$		Fine root biomass increase dictated by the difference between fine root biomass allocation target and current fine root biomass
$\Delta B_{fineroot,source}$	$g\ dry$		Fine root biomass increase according to sapwood metabolic carbon availability
$\Delta B_{fineroot,sink}$	$g\ dry$		Fine root biomass increase dictated by sink limitations (fine root relative growth rate and relative cell expansion rate)
$\Delta B_{fineroot}$	$g\ dry$		Actual fine root biomass increase

Bibliography

- Aakala, T., Kuuluvainen, T., Gauthier, S., and De Grandpré, L. (2008). Standing dead trees and their decay-class dynamics in the northeastern boreal old-growth forests of Quebec. *Forest Ecology and Management*.
- Adams, H. D., Park Williams, A., Xu, C., Rauscher, S. A., Jiang, X., and McDowell, N. G. (2013). Empirical and process-based approaches to climate-induced forest mortality models. *Frontiers in Plant Science*, 4(NOV):1–5.
- Adams, H. D., Zeppel, M. J., Anderegg, W. R., Hartmann, H., Landhäusser, S. M., Tissue, D. T., Huxman, T. E., Hudson, P. J., Franz, T. E., Allen, C. D., Anderegg, L. D., Barron-Gafford, G. A., Beerling, D. J., Breshears, D. D., Brodribb, T. J., Bugmann, H., Cobb, R. C., Collins, A. D., Dickman, L. T., Duan, H., Ewers, B. E., Galiano, L., Galvez, D. A., Garcia-Forner, N., Gaylord, M. L., Germino, M. J., Gessler, A., Hacke, U. G., Hakamada, R., Hector, A., Jenkins, M. W., Kane, J. M., Kolb, T. E., Law, D. J., Lewis, J. D., Limousin, J. M., Love, D. M., Macalady, A. K., Martínez-Vilalta, J., Mencuccini, M., Mitchell, P. J., Muss, J. D., O'Brien, M. J., O'Grady, A. P., Pangle, R. E., Pinkard, E. A., Piper, F. I., Plaut, J. A., Pockman, W. T., Quirk, J., Reinhardt, K., Ripullone, F., Ryan, M. G., Sala, A., Sevanto, S., Sperry, J. S., Vargas, R., Vennetier, M., Way, D. A., Xu, C., Yopez, E. A., and McDowell, N. G. (2017). A multi-species synthesis of physiological mechanisms in drought-induced tree mortality. *Nature Ecology and Evolution*, 1(9):1285–1291. Publisher: Springer US.
- Albini, F. and Baughman, R. (1979). Estimating windspeeds for predicting wildland fire behavior. *USDA Forest Service, Intermountain Forest and Range Experiment Station, Research Paper*, pages INT–RP–221. ISBN: INT-221.
- Albini, F. A. (1976). Estimating wildfire behavior and effects. *For. Serv. Gen. Tech. Rep. INT-30*, 4235(789273912):44 ST – Guidelines and sample protocol for sampli. ISBN: GTR-INT-30 Publisher: USDA Forest Service.
- Ameztegui, A., Cabon, A., De Caceres, M., and Coll, L. (2017). Managing stand density to enhance the adaptability of Scots pine stands to climate change: A modelling approach. *Ecological Modelling*, 356:141–150. Publisher: Elsevier B.V.
- Ameztegui, A., Coll, L., Cáceres, M. D., and Morán-Ordóñez, A. (2024). Disturbance impacts on Mediterranean forests across climate and management scenarios. *Journal of Environmental Management*, 371:123193.
- Anderegg, W. R., Wolf, A., Arango-Velez, A., Choat, B., Chmura, D. J., Jansen, S., Kolb, T., Li, S., Meinzer, F. C., Pita, P., Resco de Dios, V., Sperry, J. S., Wolfe, B. T., and Pacala, S. (2018). Woody plants optimise stomatal behaviour relative to hydraulic risk. *Ecology Letters*, 21(7):968–977.
- Anten, N. P. R. and Bastiaans, L. (2016). The use of canopy models to analyze light competition among plants. In Hikosaka, K., Niinemets, ., and Anten, N. P. R., editors, *Canopy photosynthesis: from basics to application*, pages 379–398. Springer.
- Bajguz, A. and Piotrowska-Niczyporuk, A. (2023). Biosynthetic Pathways of Hormones in Plants. *Metabolites*, 13(8):884.

- Balaguer-Romano, R., De Cáceres, M., and Espelta, J. M. (2025). Second-Growth Forests Exhibit Higher Sensitivity to Dry and Wet Years than Long-Existing Ones. *Ecosystems*, 28(1):6.
- Balaguer-Romano, R., Díaz-Sierra, R., De Cáceres, M., Cunill-Camprubí, ., Nolan, R. H., Boer, M. M., Voltas, J., and Resco de Dios, V. (2022). A semi-mechanistic model for predicting daily variations in species-level live fuel moisture content. *Agricultural and Forest Meteorology*, 323(December 2021):109022.
- Balaguer-Romano, R., Díaz-Sierra, R., De Cáceres, M., Voltas, J., Boer, M. M., and Resco de Dios, V. (2023). Modeling fuel moisture dynamics under climate change in Spain's forests. *Fire Ecology*, 19(1):65.
- Baldocchi, D. (1994). An analytical solution for coupled leaf photosynthesis and stomatal conductance models. *Tree Physiology*, 14:1069–1079.
- Bartlett, M. K., Klein, T., Jansen, S., Choat, B., and Sack, L. (2016). The correlations and sequence of plant stomatal, hydraulic, and wilting responses to drought. *Proceedings of the National Academy of Sciences of the United States of America*, 113(46):13098–13103.
- Bartlett, M. K., Scoffoni, C., and Sack, L. (2012). The determinants of leaf turgor loss point and prediction of drought tolerance of species and biomes: a global meta-analysis. *Ecology letters*, 15(5):393–405.
- Baumgarten, F., Aitken, S., Vitasse, Y., Guy, R. D., and Wolkovich, E. (2026). (In)determinacy in Woody Plants: Limits and Opportunities for Timing Growth in a Changing Climate. *Ecology Letters*, 29(6):e70435.
- Bernacchi, C. J., Singaas, E. L., Pimentel, C., Portis, A. R., and Long, S. P. (2001). Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, Cell and Environment*, 24(2):253–259. ISBN: 0140-7791.
- Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J. (2011). The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes. *Geoscientific Model Development*, 4(3):677–699. arXiv: 1011.1669v3 ISBN: 1991-959X.
- Birk, E. M. and Simpson, R. W. (1980). Steady state and the continuous input model of litter accumulation and decomposition in Australian eucalypt forests. *Ecology*, 61(3):481–485. ISBN: 0012-9658.
- Blackman, C. J., Pfautsch, S., Choat, B., Delzon, S., Gleason, S. M., and Duursma, R. A. (2016). Toward an index of desiccation time to tree mortality under drought. *Plant Cell and Environment*, 39(10):2342–2345.
- Bonan, G. (2019). *Climate change and terrestrial ecosystem modeling*. Cambridge University Press, Cambridge, UK.
- Bonan, G. B., Williams, M., Fisher, R. A., and Oleson, K. W. (2014). Modeling stomatal conductance in the earth system: Linking leaf water-use efficiency and water transport along the soil-plant-atmosphere continuum. *Geoscientific Model Development*, 7(5):2193–2222.
- Boughton, W. (1989). A review of the USDA SCS curve number method. *Australian Journal of Soil Research*, 27:511–523.
- Brotons, L., Aquilué, N., de Cáceres, M., Fortin, M.-J., and Fall, A. (2013). How fire history, fire suppression practices and climate change affect wildfire regimes in Mediterranean landscapes. *PloS one*, 8(5):e62392.
- Bugmann, H., Seidl, R., Hartig, F., Bohn, F., Brûna, J., Cailleret, M., François, L., Heinke, J., Henrot, A. J., Hickler, T., Hülsmann, L., Huth, A., Jacquemin, I., Kollas, C., Lasch-Born, P., Lexer, M. J., Merganič, J., Merganičová, K., Mette, T., Miranda, B. R., Nadal-Sala, D., Rammer, W., Rammig, A., Reineking, B., Roedig, E., Sabaté, S., Steinkamp, J., Suckow, F., Vacchiano, G., Wild, J., Xu, C., and Reyer, C. P. (2019). Tree mortality submodels drive simulated long-term forest dynamics: assessing 15 models from the stand to global scale. *Ecosphere*, 10(2):e02616.
- Bullock, J. M., Mallada González, L., Tamme, R., Götzenberger, L., White, S. M., Pärtel, M., and Hooftman, D. A. P. (2017). A synthesis of empirical plant dispersal kernels. *Journal of Ecology*, 105(1):6–19.

- Butler, B. W. and Dickinson, M. B. (2010). Tree Injury and Mortality in Fires: Developing Process-Based Models. *Fire Ecology*, 6(1):55–79.
- Byram, G. (1959). Combustion of forest fuels. In Davis, K., editor, *Forest fire: control and use*, pages 61–89. McGraw-Hill, New York.
- Bär, A., Michaletz, S. T., and Mayr, S. (2019). Fire effects on tree physiology. *New Phytologist*, 223(4):1728–1741.
- Cabon, A., Ameztegui, A., Anderegg, W. R. L., Martínez-Vilalta, J., and De Cáceres, M. (2024). Probing the interplay of biophysical constraints and photosynthesis to model tree growth. *Agricultural and Forest Meteorology*, 345:109852.
- Cabon, A., Fernández-de-Uña, L., Gea-Izquierdo, G., Meinzer, F. C., Woodruff, D. R., Martínez-Vilalta, J., and De Cáceres, M. (2020a). Water potential control of turgor-driven tracheid enlargement in Scots pine at its xeric distribution edge. *New Phytologist*, 225(1):209–221.
- Cabon, A., Peters, R. L., Fonti, P., Martínez-Vilalta, J., and De Cáceres, M. (2020b). Temperature and water potential co-limit stem cambial activity along a steep elevational gradient. *New Phytologist*, 226(5):1325–1340.
- Campbell, G. S. and Norman, J. M. (1998). *An introduction to environmental biophysics*. Springer-Verlag, New York.
- Cannell, M. G. and Thornley, J. H. (2000). Modelling the Components of Plant Respiration: Some Guiding Principles. *Annals of Botany*, 85(1):45–54.
- Carsel, R. F. and Parrish, R. S. (1988). Developing joint probability distributions of soil water retention characteristics. *Water Resources Research*, 24(5):755–769.
- Cho, N., Kang, S., Agossou, C., Kim, E., and Lim, J.-H. (2022). Modeling temporal variations of non-structural carbohydrate (NSC) storages across biomes. *Forest Ecology and Management*, 508:120033.
- Christoffersen, B. O., Gloor, M., Fauset, S., Fyllas, N. M., Galbraith, D. R., Baker, T. R., Rowland, L., Fisher, R. A., Binks, O. J., Sevanto, S. A., Xu, C., Jansen, S., Choat, B., Mencuccini, M., McDowell, N. G., and Meir, P. (2016). Linking hydraulic traits to tropical forest function in a size-structured and trait-driven model (TFS v.1-Hydro). *Geoscientific Model Development Discussions*, 9(June):4227–4255.
- Chuine, I., Garcia de Cortazar-Atauri, I., Kramer, K., and Hänninen, H. (2013). Plant Development Models. In Schwartz, M., editor, *Phenology: An Integrative Environmental Science*, pages 275–293. Springer Science, Dordrecht.
- Clark, J. S., Macklin, E., and Wood, L. (1998). Stages and spatial scales of recruitment limitation in Southern Appalachian forests. *Ecological Monographs*, 68(2):213–235.
- Cochard, H., Pimont, F., Ruffault, J., and Martin-StPaul, N. (2021). SurEau: a mechanistic model of plant water relations under extreme drought. *Annals of Forest Science*, 78(2). Publisher: Annals of Forest Science ISBN: 1359502101067.
- Collatz, G. J., Ball, J. T., Grivet, C., and Berry, J. A. (1991). Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer. *Agricultural and Forest Meteorology*, 54(2-4):107–136. ISBN: 0168-1923.
- Collins, D. B. G. and Bras, R. L. (2007). Plant rooting strategies in water-limited ecosystems. *Water Resources Research*, 43(6):W06407.
- Cox, P. M., Betts, R. A., Bunton, C. B., Essery, R. L. H., Rowntree, P. R., and Smith, J. (1999). The impact of new land surface physics on the GCM simulation of climate and climate sensitivity. *Climate Dynamics*, 15(3):183–203. ISBN: 0930-7575.

- De Cáceres, M., Casals, P., Gabriel, E., and Castro, X. (2019). Scaling-up individual-level allometric equations to predict stand-level fuel loading in Mediterranean shrublands. *Annals of Forest Science*, 76(3):87. Publisher: Annals of Forest Science ISBN: 1359501908.
- De Cáceres, M., Martin-StPaul, N., Turco, M., Cabon, A., and Granda, V. (2018). Estimating daily meteorological data and downscaling climate models over landscapes. *Environmental Modelling & Software*, 108(August):186–196. Publisher: Elsevier.
- De Cáceres, M., Martínez-Vilalta, J., Coll, L., Llorens, P., Casals, P., Poyatos, R., Pausas, J. G., and Brotons, L. (2015). Coupling a water balance model with forest inventory data to predict drought stress: the role of forest structural changes vs. climate changes. *Agricultural and Forest Meteorology*, 213:77–90. Publisher: Elsevier B.V.
- De Cáceres, M., Mencuccini, M., Martin-StPaul, N., Limousin, J.-M., Coll, L., Poyatos, R., Cabon, A., Granda, V., Forner, A., Valladares, F., and Martínez-Vilalta, J. (2021). Unravelling the effect of species mixing on water use and drought stress in Mediterranean forests: A modelling approach. *Agricultural and Forest Meteorology*, 296:108233.
- De Cáceres, M., Molowny-Horas, R., Cabon, A., Martínez-Vilalta, J., Mencuccini, M., García-Valdés, R., Nadal-Sala, D., Sabate, S., Morin, X., D’Adamo, F., Batllori, E., and Ameztegui, A. (2023). MEDFATE 2.9.3: A trait-enabled model to simulate Mediterranean forest function and dynamics at regional scales. *Geoscientific Model Development*, 16:3165–3201.
- De Pury, D. G. G. and Farquhar, G. D. (1997). Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models. *Plant, Cell and Environment*, 20(5):537–557. ISBN: 0140-7791.
- Deguchi, A., Hattori, S., and Park, H.-T. (2006). The influence of seasonal changes in canopy structure on interception loss: Application of the revised Gash model. *Journal of Hydrology*, 318(1-4):80–102.
- Del Grosso, S. J., Parton, W. J., Mosier, A. R., Holland, E. A., and Et., A. (2005). Modeling soil CO₂ emissions from ecosystems. *Biogeochemistry*.
- Delpierre, N., Dufrêne, E., Soudani, K., Ulrich, E., Cecchini, S., Boé, J., and François, C. (2009). Modelling interannual and spatial variability of leaf senescence for three deciduous tree species in France. *Agricultural and Forest Meteorology*, 149(6-7):938–948. ISBN: 0168-1923.
- Dharssi, I., Vidale, P. L., Verhoef, A., MacPherson, B., Jones, C., and Best, M. (2009). New soil physical properties implemented in the Unified Model at PS18. (528):9–12.
- Dietze, M. C., Sala, A., Carbone, M. S., Czimczik, C. I., Mantooth, J. A., Richardson, A. D., and Vargas, R. (2014). Nonstructural Carbon in Woody Plants. *Annual Review of Plant Biology*, 65(1):667–687. ISBN: 1543-5008.
- Dunlap, F. (1914). Density of wood substance and porosity of wood. *Journal of Agricultural Research*, II(6):423–428.
- Duursma, R. A., Blackman, C. J., López, R., Martin-StPaul, N. K., Cochard, H., and Medlyn, B. E. (2018). On the minimum leaf conductance: its role in models of plant water use, and ecological and environmental controls. *New Phytologist*, 221(September):693–705.
- Espelta, J. M., Barbati, A., Quevedo, L., Tárrega, R., Navascués, P., Bonfil, C., Peguero, G., Fernández-Martínez, M., and Rodrigo, A. (2012). Post-Fire Management of Mediterranean Broadleaved Forests. In Moreira, F., Arianoutsou, M., Corona, P., and De las Heras, J., editors, *Post-Fire Management and Restoration of Southern European Forests*, Managing Forest Ecosystems, pages 171–194. Springer Netherlands, Dordrecht.
- Espelta, J. M., Habrouk, A., and Retana, J. (2006). Response to natural and simulated browsing of two Mediterranean oaks with contrasting leaf habit after a wildfire. *Annals of Forest Science*, 63(4):441–447. Publisher: EDP Sciences.

- Espelta, J. M., Sabaté, S., and Retana, J. (1999). Resprouting Dynamics. In Rodà, F., Retana, J., Gracia, C. A., and Bellot, J., editors, *Ecology of Mediterranean Evergreen Oak Forests*, Ecological Studies, pages 61–73. Springer, Berlin, Heidelberg.
- Fatichi, S., Leuzinger, S., and Körner, C. (2014). Moving beyond photosynthesis: From carbon source to sink-driven vegetation modeling. *New Phytologist*, 201(4):1086–1095. ISBN: 1469-8137.
- Federer, C. (1982). Transpirational supply and demand: plant, soil, and atmospheric effects evaluated by simulation. *Water Resources Research*, 18(2):355–362.
- Fisher, R. A., Williams, M., Do Vale, L. R., Da Costa, A. L., and Meir, P. (2006). Evidence from Amazonian forests is consistent with a model of isohydric control of leaf water potential. *Plant, Cell and Environment*, 29(2):151–165.
- Flerchinger, G. N., Xiao, W., Sauer, T. J., and Yu, Q. (2009). Simulation of within-canopy radiation exchange. *NJAS - Wageningen Journal of Life Sciences*, 57(1):5–15.
- Francés, F., Vélez, J. I., and Vélez, J. J. (2007). Split-parameter structure for the automatic calibration of distributed hydrological models. *Journal of Hydrology*, 332(1-2):226–240.
- Franks, P. J. (2006). Higher rates of leaf gas exchange are associated with higher leaf hydrodynamic pressure gradients. *Plant, Cell and Environment*, 29(4):584–592.
- Franks, P. J. and Beerling, D. J. (2009). Maximum leaf conductance driven by CO₂ effects on stomatal size and density over geologic time. *Proceedings of the National Academy of Sciences*, 106(25):10343–10347.
- Friend, A. D., Stevens, A. K., Knox, R. G., and Cannell, M. G. R. (1997). A process-based, terrestrial biosphere model of ecosystem dynamics (Hybrid v3.0). *Ecological Modelling*, 95:249–287. ISBN: 0304-3800.
- Fyllas, N. M. and Troumbis, A. Y. (2009). Simulating vegetation shifts in north-eastern Mediterranean mountain forests under climatic change scenarios. *Global Ecology and Biogeography*, 18(1):64–77.
- Först, P., Werner, F., and Delgado, A. (2002). On the pressure dependence of the viscosity of aqueous sugar solutions. *Rheologica Acta*, 41(4):369–374.
- García-Jiménez, R., Palmero-Iniesta, M., and Espelta, J. (2017). Contrasting Effects of Fire Severity on the Regeneration of *Pinus halepensis* Mill. and Resprouter Species in Recently Thinned Thickets. *Forests*, 8(3):55.
- Gash, J., Lloyd, C., and Lachaud, G. (1995). Estimating sparse forest rainfall interception with an analytical model. *Journal of Hydrology*, 170(79-86).
- Genuchten, M. V. (1980). A closed-form equation for predicting the hydraulic conductivity of unsaturated soils. *Soil science society of America journal*, 44:892–898.
- Gifford, R. M. (2003). Plant respiration in productivity models: Conceptualisation, representation and issues for global terrestrial carbon-cycle research. *Functional Plant Biology*, 30(2):171–186.
- Gill, R. A. and Jackson, R. B. (2000). Global patterns of root turnover for terrestrial ecosystems. *New Phytologist*, 147(1):13–31. __eprint: <https://nph.onlinelibrary.wiley.com/doi/pdf/10.1046/j.1469-8137.2000.00681.x>.
- Granier, A., Bréda, N., Biron, P., and Villette, S. (1999). A lumped water balance model to evaluate duration and intensity of drought constraints in forest stands. *Ecological Modelling*, 116:269–283. ISBN: 3338339402.
- Granier, A., Reichstein, M., Bréda, N., Janssens, I., Falge, E., Ciais, P., Grünwald, T., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Facini, O., Grassi, G., Heinesch, B., Ilvesniemi, H., Keronen, P., Knohl, A., Köstner, B., Lagergren, F., Lindroth, A., Longdoz, B., Loustau, D., Mateus, J., Montagnani, L., Nys, C., Moors, E., Papale, D., Peiffer, M., Pilegaard, K., Pita, G., Pumpanen, J., Rambal, S., Rebmann, C., Rodrigues, A., Seufert, G., Tenhunen, J., Vesala, T., and Wang, Q. (2007). Evidence for soil water control

- on carbon and water dynamics in European forests during the extremely dry year: 2003. *Agricultural and Forest Meteorology*, 143(1-2):123–145.
- Green, H. W. and Ampt, G. A. (1911). Studies on Soil Physics. *The Journal of Agricultural Science*, 4(1):1–24.
- Guillemot, J., Martin-Stpaul, N. K., Dufrêne, E., François, C., Soudani, K., Ourcival, J. M., and Delpierre, N. (2015). The dynamic of the annual carbon allocation to wood in European tree species is consistent with a combined source-sink limitation of growth: Implications for modelling. *Biogeosciences*, 12(9):2773–2790. ISBN: 1227732015.
- Hammond, W. M., Yu, K., Wilson, L. A., Will, R. E., Anderegg, W. R., and Adams, H. D. (2019). Dead or dying? Quantifying the point of no return from hydraulic failure in drought-induced tree mortality. *New Phytologist*.
- Hasenauer, H. and Monserud, R. A. (1996). A crown ratio model for Austrian Forests. *Forest Ecology and Management*, 84(1-3):49–60. ISBN: 0378-1127.
- Haverkamp, R., Vauclin, M., Touma, J., Wierenga, P. J., and Vachaud, G. (1977). A Comparison of Numerical Simulation Models For One-Dimensional Infiltration. *Soil Science Society of America Journal*, 41(2):285–294.
- Hawkes, C. (2000). Woody plant mortality algorithms: Description, problems and progress. *Ecological Modelling*, 126(2-3):225–248. ISBN: 0304-3800.
- Hikosaka, K., Kumagai, T., and Ito, A. (2016). Modeling Canopy Photosynthesis. pages 239–268.
- Hood, S. M., Varner, J. M., Van Mantgem, P., and Cansler, C. A. (2018). Fire and tree death: Understanding and improving modeling of fire-induced tree mortality. *Environmental Research Letters*, 13(11). Publisher: IOP Publishing.
- Hölttä, T., Cochard, H., Nikinmaa, E., and Mencuccini, M. (2009). Capacitive effect of cavitation in xylem conduits: Results from a dynamic model. *Plant, Cell and Environment*, 32(1):10–21. ISBN: 1365-3040.
- Hölttä, T., Lintunen, A., Chan, T., Mäkelä, A., and Nikinmaa, E. (2017). A steady-state stomatal model of balanced leaf gas exchange, hydraulics and maximal source-sink flux. *Tree physiology*, 37(7):851–868. arXiv: 1611.06654 ISBN: 2076792171.
- Jarvis, N. J., Jansson, P., Dik, P. E., and Messing, I. (1991). Modelling water and solute transport in macroporous soil. I. Model description and sensitivity analysis. *Journal of Soil Science*, 42(1):59–70.
- Jarvis, P. G. (1976). The Interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 273(927):593–610. ISBN: 0080-4622.
- Javoy, T., Perret, S., Couteau, C., and Perot, T. (2025). Impact of defoliation on tree growth for sessile oak and Scots pine following heat-drought events. *Forest Ecology and Management*, 598:123243.
- Karavani, A., De Cáceres, M., Martínez de Aragón, J., Bonet, J., and de Miguel, S. (2018). Effect of climatic and soil moisture conditions on mushroom productivity and related ecosystem services in Mediterranean pine stands facing climate change. *Agricultural and Forest Meteorology*, 248:432–440.
- Katul, G. G., Mahrt, L., Poggi, D., and Sanz, C. (2004). One- and two-equation models for canopy turbulence. *Boundary-Layer Meteorology*, 113(1):81–109.
- Keane, R., Austin, M., Field, C., and Huth, A. (2001). Tree mortality in gap models: Application to climate change. *Climatic Change*, 51:509–540.
- Kelly, R. H., Parton, W. J., Hartman, M. D., Stretch, L. K., Ojima, D. S., and Schimel, D. S. (2000). Intra-annual and interannual variability of ecosystem processes in shortgrass steppe. *Journal of Geophysical Research: Atmospheres*, 105(D15):20093–20100. _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2000JD900259>.

- Kergoat, L. (1998). A model for hydrological equilibrium of leaf area index on a global scale. *Journal of Hydrology*, 212-213:268–286.
- Krajicek, J., Brinkman, K., and Gingrich, S. (1961). Crown competition-a measure of density. *Forest Science*, 7(1):35–42.
- Kursar, T. A., Engelbrecht, B. M., Burke, A., Tyree, M. T., El Omari, B., and Giraldo, J. P. (2009). Tolerance to low leaf water status of tropical tree seedlings is related to drought performance and distribution. *Functional Ecology*, 23(1):93–102.
- Körner, C. (2015). Paradigm shift in plant growth control. *Current Opinion in Plant Biology*, 25:107–114.
- Körner, C. (2019). No need for pipes when the well is dry-a comment on hydraulic failure in trees. *Tree physiology*, 39(5):695–700.
- Lai, C.-t., Katul, G., Oren, R., Ellsworth, D., and Schäfer, K. (2000). Modeling CO₂ and water vapor turbulent flux distributions within a forest canopy. *Journal of Geophysical Research: Atmospheres*, 105(D21):26333–26351.
- Landsberg, J. and Waring, R. (1997). A generalised model of forest productivity using simplified concepts of radiation-use efficiency, carbon balance and partitioning. *Forest Ecology and Management*, 95(3):209–228.
- Larsbo, M., Roulier, S., Stenemo, F., Kasteel, R., and Jarvis, N. (2005). An Improved Dual-Permeability Model of Water Flow and Solute Transport in the Vadose Zone. *Vadose Zone Journal*, 4(2):398–406.
- Leij, F. J., Alves, W. J., Genuchten, M. V., and Williams, J. R. (1996). *The UNSODA Unsaturated Soil Hydraulic Database User's Manual Version 1.0*. Issue: August.
- Leuning, R. (2002). Temperature dependence of two parameters in a photosynthesis model. *Plant, Cell and Environment*, 25(9):1205–1210. ISBN: 0140-7791.
- Lindner, M., Sievänen, R., and Pretzsch, H. (1997). Improving the simulation of stand structure in a forest gap model. *Forest Ecology and Management*, 95(2):183–195.
- Liu, S. (2001). Evaluation of the Liu model for predicting rainfall interception in forests world-wide. *Hydrological Processes*, 15(12):2341–2360.
- Lloret, F., Peñuelas, J., Prieto, P., Llorens, L., and Estiarte, M. (2009). Plant community changes induced by experimental climate change: Seedling and adult species composition. *Perspectives in Plant Ecology, Evolution and Systematics*, 11(1):53–63.
- Lloret, F., Siscart, D., and Dalmases, C. (2004). Canopy recovery after drought dieback in holm-oak Mediterranean forests of Catalonia (NE Spain). *Global Change Biology*, 10(12):2092–2099.
- Ma, Y. and Liu, H. (2019). An Advanced Multiple-Layer Canopy Model in the WRF Model With Large-Eddy Simulations to Simulate Canopy Flows and Scalar Transport Under Different Stability Conditions. *Journal of Advances in Modeling Earth Systems*, 11(7):2330–2351.
- Magnani, F., Mencuccini, M., and Grace, J. (2000). Age-related decline in stand productivity: The role of structural acclimation under hydraulic constraints. *Plant, Cell and Environment*, 23(3):251–263. ISBN: 0140-7791.
- Maherali, H., Pockman, W., and Jackson, R. (2004). Adaptive variation in the vulnerability of woody plants to xylem cavitation. *Ecology*, 85(8):2184–2199.
- Manoli, G., Huang, C.-w., Bonetti, S., Domec, J.-c., Marani, M., and Katul, G. (2017). Competition for light and water in a coupled soil-plant system. *Advances in Water Resources*, 108:216–230. Publisher: Elsevier Ltd.
- Marañón, T., Zamora, R., Villar, R., Zavala, M. A., Quero, J., Pérez-Ramos, I., Mendoza, I., and Castro, J. (2004). Regeneration of tree species and restoration under constrained Mediterranean habitats: Field and glasshouse experiments. *International Journal of Ecology and Environmental Sciences*, 30(3):187–196.

- Martin-StPaul, N., Delzon, S., and Cochard, H. (2017). Plant resistance to drought depends on timely stomatal closure. *Ecology Letters*, 20(11):1437–1447.
- Martinez-Vilalta, J., Anderegg, W. R. L., Sapes, G., and Sala, A. (2019). Greater focus on water pools may improve our ability to understand and anticipate drought-induced mortality in plants. *New Phytologist*, 223(1):22–32.
- Martínez-Vilalta, J., Sala, A., Asensio, D., Galiano, L., Hoch, G., Palacio, S., Piper, F. I., and Lloret, F. (2016). Dynamics of non-structural carbohydrates in terrestrial plants: a global synthesis. *Ecological Monographs*, 86(4):495–516.
- Massman, W. (1987). A comparative study of some mathematical models of the mean wind structure and aerodynamic drag of plant canopies. *Boundary-Layer Meteorology*, 40(1-2):179–197.
- McDowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., Plaut, J., Sperry, J. S., West, A., Williams, D. G., and Yezpe, E. A. (2008). Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? *The New Phytologist*, 178(4):719–39.
- McDowell, N. G. (2011). Mechanisms linking drought, hydraulics, carbon metabolism, and vegetation mortality. *Plant Physiology*, 155(3):1051–1059. ISBN: 0032-0889.
- McDowell, N. G., Sapes, G., Pivovarov, A., Adams, H. D., Allen, C. D., Anderegg, W. R. L., Arend, M., Breshears, D. D., Brodrigg, T., Choat, B., Cochard, H., De Cáceres, M., De Kauwe, M. G., Grossiord, C., Hammond, W. M., Hartmann, H., Hoch, G., Kahmen, A., Klein, T., Mackay, D. S., Mantova, M., Martínez-Vilalta, J., Medlyn, B. E., Mencuccini, M., Nardini, A., Oliveira, R. S., Sala, A., Tissue, D. T., Torres-Ruiz, J. M., Trowbridge, A. M., Trugman, A. T., Wiley, E., and Xu, C. (2022). Mechanisms of woody-plant mortality under rising drought, CO₂ and vapour pressure deficit. *Nature Reviews Earth & Environment*, 3:294–308. ISBN: 0123456789.
- McElwain, J. C., Yiotis, C., and Lawson, T. (2016). Using modern plant trait relationships between observed and theoretical maximum stomatal conductance and vein density to examine patterns of plant macroevolution. *New Phytologist*, 209(1):94–103. ISBN: 1469-8137.
- McMurtrie, R. E., Rook, D. A., and Kelliher, F. M. (1990). Modelling the yield of *Pinus radiata* on a site limited by water and nitrogen. *Forest Ecology and Management*, 30(1-4):381–413. ISBN: 0378-1127.
- Medlyn, B. E., Loustau, D., and Delzon, S. (2002). Temperature response of parameters of a biochemically based model of photosynthesis. I. Seasonal changes in mature maritime pine (*Pinus pinaster* Ait.). *Plant, Cell and Environment*, 25(9):1155–1165. ISBN: 1365-3040.
- Mencuccini, M. and Grace, J. (1995). Climate influences the leaf area/sapwood area ratio in Scots pine. *Tree Physiology*, 15(1):1–10.
- Michaletz, S. T. and Johnson, E. A. (2006). A heat transfer model of crown scorch in forest fires. *Canadian Journal of Forest Research*, 36(11):2839–2851.
- Michaletz, S. T. and Johnson, E. A. (2008). A biophysical process model of tree mortality in surface fires. *Canadian Journal of Forest Research*, 38(7):2013–2029.
- Miezite, L. E., Ameztegui, A., De Cáceres, M., Coll, L., Morán-Ordóñez, A., Vega-García, C., and Rodrigues, M. (2022). Trajectories of Wildfire Behavior Under Climate Change. Can Forest Management Mitigate the Increasing Hazard? *Journal of Environmental Management*, 322:116134.
- Miralles, D. G., Gash, J. H., Holmes, T. R. H., de Jeu, R. A. M., and Dolman, A. J. (2010). Global canopy interception from satellite observations. *Journal of Geophysical Research*, 115(D16):D16122.
- Mitsopoulos, I. D. and Dimitrakopoulos, A. P. (2007). Allometric equations for crown fuel biomass of Aleppo pine (*Pinus halepensis* Mill.) in Greece. *International Journal of Wildland Fire*, 16(5):642–647. ISBN: 1448-5516

- Moore, A. D. and Noble, I. R. (1990). An individualistic model of vegetation stand dynamics. *Journal of Environmental Management*, 31(1):61–81.
- Moreira, B., Tormo, J., and Pausas, J. G. (2012). To resprout or not to resprout: factors driving intraspecific variability in resprouting. *Oikos*, 121(10):1577–1584.
- Morán-Ordóñez, A., Ameztegui, A., De Cáceres, M., De-Miguel, S., Lefèvre, F., Brotons, L., and Coll, L. (2020). Future trade-offs and synergies among ecosystem services in Mediterranean forests under global change scenarios. *Ecosystem Services*, 45(January):101174. Publisher: Elsevier B.V.
- Mouillot, F., Rambal, S., and Joffre, R. (2002). Simulating climate change impacts on fire frequency and vegetation dynamics in a Mediterranean-type ecosystem. *Global Change Biology*, 8(5):423–437. Publisher: Wiley Online Library.
- Mouillot, F., Rambal, S., and Lavorel, S. (2001). A generic process-based Simulator for mediterranean landscApes (SIERRA): design and validation exercises. *Forest Ecology and Management*, 147(1):75–97.
- Nathan, R., Klein, E., Robledo-Arnuncio, J. J., and Revilla, E. (2012). Dispersal kernels: review. In Clobert, J., Baguette, M., Benton, T. G., and Bullock, J. M., editors, *Dispersal Ecology and Evolution*, pages 186–210. Oxford University Press.
- Nathan, R., Safriel, U. N., and Noy-Meir, I. (2001). Field validation and sensitivity analysis of a mechanistic model for tree seed dispersal by wind. *Ecology*, 82(2):374–388.
- Oberle, B., Ogle, K., Zanne, A. E., and Woodall, C. W. (2018). When a tree falls: Controls on wood decay predict standing dead tree fall and new risks in changing forests. *PLOS ONE*, 13(5):e0196712. Publisher: Public Library of Science.
- Ochoa, M. E., Henry, C., John, G. P., Medeiros, C. D., Pan, R., Scoffoni, C., Buckley, T. N., and Sack, L. (2024). Pinpointing the causal influences of stomatal anatomy and behavior on minimum, operational, and maximum leaf surface conductance. *Plant Physiology*, 196(1):51–66.
- Ogle, K. and Pacala, S. W. (2009). A modeling framework for inferring tree growth and allocation from physiological, morphological and allometric traits. *Tree Physiology*, 29(4):587–605.
- Oliveras, I., Martínez-Vilalta, J., Jiménez-Ortiz, T., Lledó, M. J., Escarré, A., and Piñol, J. (2003). Hydraulic architecture of *Pinus halepensis*, *P. pinea* and *Tetraclinis articulata* in a dune ecosystem of Eastern Spain. *Plant Ecology*, pages 131–141.
- Olson, M. E., Anfodillo, T., Rosell, J. A., Petit, G., Crivellaro, A., Isnard, S., León-Gómez, C., Alvarado-Cárdenas, L. O., and Castorena, M. (2014). Universal hydraulics of the flowering plants: Vessel diameter scales with stem length across angiosperm lineages, habits and climates. *Ecology Letters*, 17(8):988–997. ISBN: 1461-0248.
- Ostendorf, B. and Reynolds, J. F. (1993). Relationships between a terrain-based hydrologic model and patch-scale vegetation patterns in an arctic tundra landscape. *Landscape Ecology*, 8(4):229–237. ISBN: 0921-2973.
- Padilla, F. M. and Pugnaire, F. I. (2007). Rooting depth and soil moisture control Mediterranean woody seedling survival during drought. *Functional Ecology*, 21(3):489–495.
- Parton, W. J., Hartman, M., Ojima, D., and Schimel, D. (1998). DAYCENT and its land surface submodel: description and testing. *Global and Planetary Change*, 19(1-4):35–48.
- Parton, W. J., Scurlock, J. M., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S., Kirchner, T., Menaut, J. ., Seastedt, T., Garcia Moya, E., Kamnalrut, A., and Kinyamario, J. I. (1993). Observations and modeling of biomass and soil organic matter dynamics for the grassland biome worldwide. *Global Biogeochemical Cycles*, 7(4):785–809.
- Parton, W. J., Stewart, J. W. B., and Cole, C. V. (1988). Dynamics of C, N, P and S in grassland soils: a model. *Biogeochemistry*, 5(1):109–131.

- Pausas, J. G., Pratt, R. B., Keeley, J. E., Jacobsen, A. L., Ramirez, A. R., Vilagrosa, A., Paula, S., Kaneakua-Pia, I. N., and Davis, S. D. (2016). Towards understanding resprouting at the global scale. *New Phytologist*, 209(3):945–954.
- Peris-Llopis, M., Mola-Yudego, B., Berninger, F., Garcia-Gonzalo, J., and González-Olabarria, J. R. (2024). Impact of species composition on fire-induced stand damage in Spanish forests. *Scientific Reports*, 14(1):8594. Publisher: Nature Publishing Group.
- Pimont, F., Dupuy, J. L., and Rigolot, E. (2018). A simple model for shrub-strata-fuel dynamics in *Quercus coccifera* L. communities. *Annals of Forest Science*, 75(2):1–9. Publisher: Annals of Forest Science.
- Plavcová, L. and Jansen, S. (2015). The Role of Xylem Parenchyma in the Storage and Utilization of Nonstructural Carbohydrates. In *Functional and Ecological Xylem Anatomy*, pages 209–234. Springer International Publishing, Cham.
- Prentice, I. C., Sykes, M. T., and Cramer, W. (1993). A simulation model for the transient effects of climate change on forest landscapes. *Ecological Modelling*, 65(1-2):51–70.
- Prescott, C. E., Grayston, S. J., Helmisaari, H.-S., Kaštovská, E., Körner, C., Lambers, H., Meier, I. C., Millard, P., and Ostonen, I. (2020). Surplus Carbon Drives Allocation and Plant–Soil Interactions. *Trends in Ecology & Evolution*, 35(12):1110–1118.
- Price, D., Zimmermann, N., and Meer, P. V. D. (2001). Regeneration in gap models: Priority issues for studying forest responses to climate change. *Climatic Change*, 51:475–508.
- Prichard, S. J., Sandberg, D. V., Ottmar, R. D., Eberhardt, E., Andreu, A., Eagle, P., and Swedin, K. (2013). Classification System Version 3.0: Technical Documentation. Technical report.
- Rahimi, A. A., Sepaskhah, A. R., and Ahmadi, S. H. (2011). Evaluation of different methods for the prediction of saturated hydraulic conductivity in tilled and untilled soils. *Archives of Agronomy and Soil Science*, 57(8):899–914.
- Rasche, L., Fahse, L., Zingg, A., and Bugmann, H. (2012). Enhancing gap model accuracy by modeling dynamic height growth and dynamic maximum tree height. *Ecological Modelling*, 232:133–143. Publisher: Elsevier B.V.
- Reich, P. B., Tjoelker, M. G., Pregitzer, K. S., Wright, I. J., Oleksyn, J., and Machado, J. L. (2008). Scaling of respiration to nitrogen in leaves, stems and roots of higher land plants. *Ecology Letters*, 11(8):793–801.
- Reinhardt, E., Lutes, D., and Scott, J. (2006). FuelCalc: A method for estimating fuel characteristics. Technical report. Publication Title: USDA Forest Service Proceedings RMRS-P-41.
- Retana, J., Riba, M., Castell, C., and Espelta, J. M. (1992). Regeneration by sprouting of holm-oak (*Quercus ilex*) stands exploited by selection thinning. *Vegetatio*, 99(1):355–364.
- Richardson, A. D., Carbone, M. S., Keenan, T. F., Czimczik, C. I., Hollinger, D. Y., Murakami, P., Schaberg, P. G., and Xu, X. (2013). Seasonal dynamics and age of stemwood nonstructural carbohydrates in temperate forest trees. *New Phytologist*, 197(3):850–861.
- Ritchie, J. (1972). Model for predicting evaporation from a row crop with incomplete cover. *Water resources research*, 8:1204–1213.
- Roces-Díaz, J. V., Vayreda, J., De Cáceres, M., García-Valdés, R., Banqué-Casanovas, M., Morán-Ordóñez, A., Brotons, L., De-Miguel, S., and Martínez-Vilalta, J. (2021). Temporal changes in Mediterranean forest ecosystem services are driven by stand development, rather than by climate-related disturbances. *Forest Ecology and Management*, 480(July 2020):118623. Publisher: Elsevier.
- Rolo, V. and Moreno, G. (2019). Shrub encroachment and climate change increase the exposure to drought of Mediterranean wood-pastures. *Science of The Total Environment*, 660:550–558. Publisher: Elsevier B.V.

- Rothermel, R. (1972). A mathematical model for predicting fire spread in wildland fuels. *USDA Forest Service Research Paper INT USA*. Publisher: USDA Forest Service, Intermountain Forest and Range Experiment Station.
- Ruffault, J., Martin-StPaul, N. K., Duffet, C., Goge, F., and Mouillot, F. (2014). Projecting future drought in Mediterranean forests: bias correction of climate models matters! *Theoretical and Applied Climatology*, 117(1-2):113–122.
- Ruffault, J., Martin-StPaul, N. K., Rambal, S., and Mouillot, F. (2013). Differential regional responses in drought length, intensity and timing to recent climate changes in a Mediterranean forested ecosystem. *Climatic Change*, 117(1-2):103–117.
- Ruffault, J., Pimont, F., Cochard, H., Dupuy, J.-L., and Martin-StPaul, N. (2022). SurEau-Ecos v2.0: a trait-based plant hydraulics model for simulations of plant water status and drought-induced mortality at the ecosystem level. *Geoscientific Model Development*, 15(14):5593–5626.
- Rötzer, T., Häberle, K. H., Kallenbach, C., Matyssek, R., Schütze, G., and Pretzsch, H. (2017). Tree species and size drive water consumption of beech/spruce forests - a simulation study highlighting growth under water limitation. *Plant and Soil*, 418(1-2):337–356. Publisher: Plant and Soil.
- Sack, L. and Buckley, T. N. (2016). The Developmental Basis of Stomatal Density and Flux. *Plant Physiology*, 171(4):2358–2363.
- Sack, L. and Holbrook, N. M. (2006). Leaf Hydraulics. *Annual Review of Plant Biology*, 57(1):361–381. arXiv: 1011.1669 ISBN: 1040-2519.
- Sack, L. and Scoffoni, C. (2013). Leaf venation: structure, function, development, evolution, ecology and applications in the past, present and future. *New Phytologist*, 198(4):983–1000.
- Sancho-Knapik, D., Ferrio, J. P., Gil-Pelegrín, E., López-Ballesteros, A., and Peguero-Pina, J. J. (2026). *Viscum album* shares hydraulic traits but causes a water uncoupling despite the adjustments of its host *Pinus sylvestris*. *Tree Physiology*, 46(3):tpag020.
- Sandberg, D. V., Riccardi, C. L., and Schaaf, M. D. (2007). Reformulation of Rothermel’s wildland fire behaviour model for heterogeneous fuelbeds. *Canadian Journal of Forest Research*, 37:2438–2455.
- Sando, R. W. and Wick, C. H. (1972). A method of evaluating crown fuels in forest stands. Research Paper NC-84. *USDA Forest Service Research Paper NC-84*, (NC-84):10 pp.
- Saponaro, V., De Cáceres, M., Dalmonech, D., D’Andrea, E., Vangi, E., and Collalti, A. (2024). Assessing the combined effects of forest management and climate change on carbon and water fluxes in European beech forests. *Forest Ecosystems*, page 100290.
- Savage, V. M., Bentley, L. P., Enquist, B. J., Sperry, J. S., Smith, D. D., Reich, P. B., and von Allmen, E. I. (2010). Hydraulic trade-offs and space filling enable better predictions of vascular structure and function in plants. *Proceedings of the National Academy of Sciences of the United States of America*, 107(52):22722–7. ISBN: 1012194108.
- Saxton, K., Rawls, W. J., Romberger, J., and Papendick, R. (1986). Estimating generalized soil-water characteristics from texture. *Soil Science Society of America Journal*, 50:1031–1036.
- Saxton, K. E. and Rawls, W. J. (2006). Soil water characteristic estimates by texture and organic matter for hydrologic solutions. *Soil Science Society of America Journal*, 70(5):1569.
- Schaaf, M. D., Sandberg, D. V., Schreuder, M. D., and Riccardi, C. L. (2007). A conceptual framework for ranking crown fire potential in wildland fuelbeds. *Canadian Journal of Forest Research*, 37:2464–2478.
- Schenk, H. J. and Jackson, R. B. (2002). Rooting depths, lateral root spreads and below-ground/above-ground allometries of plants in water-limited ecosystems. *Journal of Ecology*, 90(3):480–494.
- Schume, H., Jost, G., and Hager, H. (2004). Soil water depletion and recharge patterns in mixed and pure forest stands of European beech and Norway spruce. *Journal of Hydrology*, 289(1-4):258–274.

- Scoffoni, C., Chatelet, D. S., Pasquet-kok, J., Rawls, M., Donoghue, M. J., Edwards, E. J., and Sack, L. (2016). Hydraulic basis for the evolution of photosynthetic productivity. *Nature Plants*, 2(6):16072. Publisher: Nature Publishing Group.
- Scott, J. H. and Reinhardt, E. D. (2002). Estimating canopy fuels in conifer forests. *Fire Management Today*, 62(4):45–50. ISBN: 0194-214X.
- Sellers, P. J. (1985). Canopy reflectance, photosynthesis and transpiration. *International Journal of Remote Sensing*, 6(8):1335–1372.
- Shinozaki, K., Yoda, K., Hozumi, K., and Kira, T. (1964). A quantitative analysis of plant form - the pipe model theory. I. Basic analysis. *Japanese Journal of Ecology*, 14:97–105.
- Sitch, S., Smith, B., Prentice, I. C., Arneth, a., Bondeau, a., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S. (2003). Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology*, 9(2):161–185.
- Snell, R. S. (2014). Simulating long-distance seed dispersal in a dynamic vegetation model. *Global Ecology and Biogeography*, 23(1):89–98.
- Sperry, J. S., Adler, F. R., Campbell, G. S., and Comstock, J. P. (1998). Limitation of plant water use by rhizosphere and xylem conductance: results from a model. *Plant, Cell & Environment*, 21:347–359.
- Sperry, J. S. and Love, D. M. (2015). What plant hydraulics can tell us about responses to climate-change droughts. *New Phytologist*, 207(1):14–27. ISBN: 0028-646X.
- Sperry, J. S., Venturas, M. D., Anderegg, W. R. L., Mencuccini, M., Mackay, D. S., Wang, Y., and Love, D. M. (2017). Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost. *Plant, Cell & Environment*, 40(6):816–830.
- Sperry, J. S., Wang, Y., Wolfe, B. T., Mackay, D. S., Anderegg, W. R. L., McDowell, N. G., and Pockman, W. T. (2016). Pragmatic hydraulic theory predicts stomatal responses to climatic water deficits. *New Phytologist*, 212:577–589.
- Spicer, R. (2005). Senescence in Secondary Xylem: Heartwood Formation as an Active Developmental Program. *Vascular Transport in Plants*, pages 457–475. ISBN: 9780120884575.
- Spitters, C. J. T., Toussaint, H., and Goudriaan, J. (1986). Separating the diffuse and direct components of global radiation and its implications for modeling canopy photosynthesis. I. Components of incoming radiation. *Agricultural and Forest Meteorology*, 38:231–242. ISBN: 0168-1923.
- Stephan, K., Miller, M., and Dickinson, M. B. (2010). First-Order Fire Effects on Herbs and Shrubs: Present Knowledge and Process Modeling Needs. *Fire Ecology*, 6(1):95–114.
- Stolf, R., Thurler, ., Oliveira, O., Bacchi, S., and Reichardt, K. (2011). Method to estimate soil macroporosity and microporosity based on sand content and bulk density. *Revista Brasileira de Ciências do Solo*, 35(1):447–459.
- Sánchez-Dávila, J., De Cáceres, M., Vayreda, J., and Retana, J. (2024). Regional patterns and drivers of modelled water flows along environmental, functional, and stand structure gradients in Spanish forests. *Hydrol. Earth Syst. Sci.*, 28:3037–3050.
- Sánchez-Dávila, J., De Cáceres, M., Vayreda, J., and Retana, J. (2026). Recent water cycle changes in Spanish forests are driven by stand structure more than climatic changes. *Journal of Environmental Management*, 402:129070.
- Sánchez-Pinillos, M., De Cáceres, M., Ameztegui, A., and Coll, L. (2019). Temporal dimension of forest vulnerability to fire along successional trajectories. *Journal of Environmental Management*, 248(March):109301. Publisher: Elsevier.

- Sánchez-Pinillos, M., De Cáceres, M., Casals, P., Alvarez, A., Beltrán, M., Pausas, J. G., Vayreda, J., and Coll, L. (2021). Spatial and temporal variations of overstory and understory fuels in Mediterranean landscapes. *Forest Ecology and Management*, 490(February):119094.
- Tjoelker, M. G., Oleksyn, J., and Reich, P. B. (2001). Modelling respiration of vegetation: Evidence for a general temperature-dependent Q10. *Global Change Biology*, 7(2):223–230.
- Tyree, M. T. and Yang, S. (1990). Water-storage capacity of Thuja, Tsuga and Acer stems measured by dehydration isotherms - The contribution of capillary water and cavitation. *Planta*, 182(3):420–426. ISBN: 0032-0935.
- Tóth, B., Weynants, M., Nemes, A., Makó, A., Bilas, G., and Tóth, G. (2015). New generation of hydraulic pedotransfer functions for Europe. *European Journal of Soil Science*, 66(1):226–238.
- Urli, M., Porte, A. J., Cochard, H., Guengant, Y., Burlett, R., and Delzon, S. (2013). Xylem embolism threshold for catastrophic hydraulic failure in angiosperm trees. *Tree Physiology*, 33(7):672–683.
- Van Lier, Q. D. J., Neto, D. D., and Metselaar, K. (2009). Modeling of transpiration reduction in van genuchten-mualem type soils. *Water Resources Research*, 45(2):1–9. ISBN: 00431397 (ISSN).
- Vanderwel, M. C., Caspersen, J. P., and Woods, M. E. (2006). Snag dynamics in partially harvested and unmanaged northern hardwood forests. *Canadian Journal of Forest Research*, 36(11):2769–2779.
- Vilà-Vilardell, L., De Cáceres, M., Piqué, M., and Casals, P. (2023). Prescribed fire after thinning increased resistance of sub-Mediterranean pine forests to drought events and wildfires. *Forest Ecology and Management*, 527:120602.
- Viney, N. (1991). A review of fine fuel moisture modelling. *International Journal of Wildland Fire*, 1(4):215. ISBN: 1448-5516.
- Walker, A. P., Beckerman, A. P., Gu, L., Kattge, J., Cernusak, L. A., Domingues, T. F., Scales, J. C., Wohlfahrt, G., Wullschlegel, S. D., and Woodward, F. I. (2014). The relationship of leaf photosynthetic traits - Vmax and Jmax - to leaf nitrogen, leaf phosphorus, and specific leaf area: A meta-analysis and modeling study. *Ecology and Evolution*, 4(16):3218–3235. ISBN: 0018655769.
- Watanabe, T. and Mizutani, K. (1996). Model study on micrometeorological aspects of rainfall interception over an evergreen broad-leaved. *Agricultural and Forest Meteorology*, 80:195–214.
- Wehrli, A., Weisberg, P. J., Schönenberger, W., Brang, P., and Bugmann, H. (2007). Improving the establishment submodel of a forest patch model to assess the long-term protective effect of mountain forests. *European Journal of Forest Research*, 126(1):131–145.
- Wigmosta, M., Vail, L., and Lettenmaier, D. (1994). A distributed hydrology-vegetation model for complex terrain. *Water Resources Research*, 30:1665–1679.
- Wigmosta, M. S. and Lettenmaier, D. P. (1999). A comparison of simplified methods for routing topographically driven subsurface flow. *Water Resources Research*, 35(1):255–264.
- Wolf, A., Anderegg, W. R. L., and Pacala, S. W. (2016). Optimal stomatal behavior with competition for water and risk of hydraulic impairment. *Proceedings of the National Academy of Sciences*, 113(46):E7222–E7230.
- Yan, C.-L., Ni, M.-Y., Cao, K.-F., and Zhu, S.-D. (2020). Leaf hydraulic safety margin and safety–efficiency trade-off across angiosperm woody species. *Biology Letters*, 16(11):20200456.
- Zweifel, R., Bangerter, S., Rigling, A., and Sterck, F. J. (2012). Pine and mistletoes: how to live with a leak in the water flow and storage system? *Journal of Experimental Botany*, 63(7):2565–2578.