

## 1. Science Justification

### 1.1. Cluster Mass Mapping

The mass distribution in galaxy clusters is a crucial test of our cosmological paradigm. Total halo masses probe the amplitude of matter fluctuations and growth of structure in the Universe, and the abundance of substructure within halos is sensitive to the history of hierarchical assembly and potentially the interaction cross section of dark matter (e.g., Natarajan et al. 2002a,b; Voit 2005; Clowe et al. 2006). Gravitational lensing has played an important role in mapping dark matter within halos, primarily by exploiting the distortion of images of background galaxies. On small scales inside lensing clusters, strong distortions produce extended arcs and multiple images enabling the production of detailed mass maps, while on larger scales the distortions are weaker and require averaging the shapes of many background galaxies over a broad area to extract a shear signal.

We propose to map the projected mass distribution in the massive cluster Abell 2261 with a new lensing technique that uses kinematic measurements of background sources to drastically improve the signal-to-noise (S/N). The typical weak lensing shear signal averaged within the virial radius of a massive cluster ( $\gamma_t \sim 0.05$ ) is small compared to the distribution of intrinsic shapes in imaging (which has scatter  $\sigma_\gamma \sim 0.25$ ). Kinematic information can be used to infer the intrinsic shape and orientation of the background galaxies, potentially reducing the shape noise per galaxy by a factor of ten. This enables  $S/N \gtrsim 1$  measurements of shear with individual galaxies. With such precision we will improve the spatial resolution of mass maps and extend the precision of strong lensing out to the weak regime near the virial radius of the cluster. We discuss this technique and its potential for improving both the precision and accuracy of future large-scale lensing experiments in the next section. Additionally, the repeat slit spectra taken for hundreds of sources behind A2261 will provide two-dimensional kinematics of emission line disk galaxies, greatly enhancing the sample size at that redshift for studying evolution of galaxy kinematics and the Tully-Fisher relation (TFR).

### 1.2. Kinematic Weak Lensing

The utility of kinematic maps for weak lensing was originally described by Blain (2002) and Morales (2006), who forecasted constraints from high S/N, high spatial resolution observations with future radio arrays. The basic idea for the kinematic shear observables is described in Figure ???. In an image, an inclined rotating circular disk has elliptical isophotes. When the image of this galaxy is sheared, the isophotes remain elliptical (in the weak shear limit,  $|\gamma_t| \ll 1$ ) with a new axis ratio and position angle. New photometric axes are inferred from this ellipse in the sheared image, and information about the original axes is lost. The case is different however, with kinematic measurements. The unsheared circular disk has kinematic axes that are perpendicular to one

another and are aligned with the unsheared photometric axes. This cross shape becomes skewed when the velocity map is sheared; the kinematic axes are no longer perpendicular and they are misaligned with the photometric axes inferred from the sheared isophotal ellipse in the imaging data.

Full two-dimensional kinematic maps are not required for this measurement, and our approach adds to the previous discussion by incorporating information from the TFR, an empirical scaling relation for disk galaxies between their luminosity and circular velocity  $v_{\text{circ}}$ . With slit spectroscopy, the measured amplitude of a rotation curve is related to the true circular velocity by  $v_{\text{obs}} = v_{\text{circ}} \sin i$ , where  $i$  is the inclination of the disk toward the observer. Thus the offset between  $v_{\text{obs}}$  and  $v_{\text{circ}}$  predicted from the TFR gives a kinematic estimate of  $\sin i$ . Typically observers estimate  $\sin i$  from the ellipticity of the image. The difference between the photometric and kinematic estimates of  $\sin i$  gives an estimate of the shear.

For ideal circular disks, the shear can be measured to arbitrary precision, limited only by the spatial and spectral resolution of the data. In practice, the intrinsic scatter of the TFR limits the precision of the shear measurement. After inclination correction, estimates for the intrinsic fractional scatter in  $v_{\text{circ}}$  at fixed luminosity or stellar mass are typically  $0.05 - 0.06$  dex both locally and to  $z \sim 1.3$  (Reyes et al. 2011; Miller et al. 2011). For a properly weighted estimator of the galaxy shape based on the TFR offset, this scatter maps to an uncertainty in the intrinsic galaxy ellipticity of only  $\sigma_{\epsilon, \text{TFR}} = 0.015$ . This should be compared to the intrinsic shape noise for pure disks  $\sigma_{\epsilon, \text{disks}} = 0.59$  and for real imaging surveys  $\sigma_{\epsilon, \text{im}} \approx 0.4$ , and is related to the shear noise through  $\sigma_{\epsilon}/\sigma_{\gamma} \approx 2(1 - \sigma_{\epsilon}^2)$  (Bernstein & Jarvis 2002; Hirata et al. 2004). This quantity is sensitive to measurement errors in  $v_{\text{circ}}$ , as shown in Figure 1; in general, rotation curve errors of  $< 10$  km/s reduce the disk galaxy shape noise by an order of magnitude.

### 1.3. A Pilot Study for Future Dark Energy Experiments

Weak gravitational lensing has been widely touted as a powerful probe of cosmology. It is a major science driver for several large imaging surveys, such as the Dark Energy Survey (DES) the Large Synoptic Survey Telescope (LSST), and space missions such as Euclid and the Wide-Field Infrared Survey Telescope (WFIRST). Weak lensing is weak, however, and all current measurements are limited by the scatter in galaxy shapes, which is an order of magnitude or more greater than the typical weak lensing signal. This means that weak lensing analyses must average over every available galaxy image, pushing the analysis to include faint and poorly-resolved galaxies for which unbiased measurements of galaxy shapes are difficult. Precision estimates of galaxy redshifts from photometry is crucial for these analyses, as the current generation of ongoing surveys will entail sample sizes of a few  $\times 10^8$  galaxies, which is two orders of magnitude beyond the size of the largest existing spectroscopic samples. Percent-level biases in the photometric redshifts will have a large impact on the error budgets for this next generation of surveys.

Here we propose a pilot study for a new weak lensing measurement technique that uses the Tully-Fisher relation (TFR) to obviate the photo-z and shear calibration problems while offering a very large reduction in the effective lensing noise.

#### 1.4. 2-d Disk Kinematics at $z \sim 0.5$

#### REFERENCES

- Bernstein, G. M., & Jarvis, M. 2002, *AJ*, 123, 583
- Blain, A. W. 2002, *ApJ*, 570, L51
- Clowe, D., Bradač, M., Gonzalez, A. H., Markevitch, M., Randall, S. W., Jones, C., & Zaritsky, D. 2006, *ApJ*, 648, L109
- Hirata, C. M., et al. 2004, *MNRAS*, 353, 529
- Miller, S. H., Bundy, K., Sullivan, M., Ellis, R. S., & Treu, T. 2011, *ApJ*, 741, 115
- Morales, M. F. 2006, *ApJ*, 650, L21
- Natarajan, P., Kneib, J.-P., & Smail, I. 2002a, *ApJ*, 580, L11
- Natarajan, P., Loeb, A., Kneib, J.-P., & Smail, I. 2002b, *ApJ*, 580, L17
- Reyes, R., Mandelbaum, R., Gunn, J. E., Pizagno, J., & Lackner, C. N. 2011, *MNRAS*, 417, 2347
- Voit, G. M. 2005, *Reviews of Modern Physics*, 77, 207

## **2. Technical Remarks**

### **2.1. Targets and Exposures**

### **2.2. Backup Program**

### **2.3. Supplementary Observations**

### **2.4. Status of Previously Approved Keck Programs**

### **2.5. Path to Science from Observations**

### **2.6. Technical Concerns**

### **2.7. Experience and Publications**

### **2.8. Resources and Publication Timescale**




Fig. 1.— Effective shape noise  $\sigma_{TF}$  as a function of the measurement error on the disk circular velocity.