

Generalized Additive Models & Functional Gradient Boosting with Geometrically Designed (GeD) Splines: Application to Insurance Data

Dimitrina S. Dimitrova¹, Vladimir K. Kaishev¹ and Emilio Sáenz Guillén (presenter)¹

Insurance Data Science Conference, 17 - 18 June 2024.

¹ Faculty of Actuarial Science and Insurance, Bayes Business School.

Email: emilio.saenz-guillen@bayes.city.ac.uk

Insurance

Data

Science

Motivation

GeDS estimation method

Generalized Additive Models with GeDS

Functional Gradient Boosting with GeDS

Insurance Data Application

Motivation

- ✱ **Geometrically Designed Splines (GeDS)** (Kaishev et al., [2016](#), Dimitrova et al., [2023](#)), — accurate and efficient tool for regression problems with one or two covariates and large datasets.

Motivation

- ✱ **Geometrically Designed Splines (GeDS)** (Kaishev et al., [2016](#), Dimitrova et al., [2023](#)), — accurate and efficient tool for regression problems with one or two covariates and large datasets.
- ✱ GeD spline methodology is extended further by:
 1. **GAM-GeDS**: encompassing **Generalized Additive Models (GAM)**, thereby making GeDS highly multivariate.
 2. **FGB-GeDS**: incorporating **Functional Gradient Boosting (FGB)**, improving the construction of the underlying spline regression model.

Motivation

- ✱ **Geometrically Designed Splines (GeDS)** (Kaishev et al., 2016, Dimitrova et al., 2023), — accurate and efficient tool for regression problems with one or two covariates and large datasets.
- ✱ GeD spline methodology is extended further by:
 1. **GAM-GeDS**: encompassing **Generalized Additive Models (GAM)**, thereby making GeDS highly multivariate.
 2. **FGB-GeDS**: incorporating **Functional Gradient Boosting (FGB)**, improving the construction of the underlying spline regression model.

Implemented in the R package **GeDS**, available from CRAN:
<https://cran.r-project.org/package=GeDS>

Motivation

GeDS estimation method

Generalized Additive Models with GeDS

Functional Gradient Boosting with GeDS

Insurance Data Application

GeDS estimation method

Free-knot spline regression technique based on a ***residual-driven (locally-adaptive) knot insertion scheme*** that produces a piecewise linear spline fit, over which ***smoother higher order spline fits*** are subsequently built.

GeDS estimation method

Free-knot spline regression technique based on a ***residual-driven (locally-adaptive) knot insertion scheme*** that produces a piecewise linear spline fit, over which ***smoother higher order spline fits*** are subsequently built.

GeDS method unfolds into two phases:

- **STAGE A** constructs a least squares linear spline fit to the data.
 - ▶ Starting with a straight-line, LS fit, which is then sequentially “broken” by iteratively introducing knots at those points ‘where the fit deviates most from the underlying functional shape determined by the data’.

GeDS estimation method

Free-knot spline regression technique based on a ***residual-driven (locally-adaptive) knot insertion scheme*** that produces a piecewise linear spline fit, over which ***smoother higher order spline fits*** are subsequently built.

GeDS method unfolds into two phases:

- **STAGE A** constructs a least squares linear spline fit to the data.
 - ▶ Starting with a straight-line, LS fit, which is then sequentially “broken” by iteratively introducing knots at those points ‘where the fit deviates most from the underlying functional shape determined by the data’.
- **STAGE B**
 - ▶ Builds smoother higher order spline fits using Schoenberg’s variation diminishing spline (VDS) approximation to the linear fit from Stage A.
 - ▶ For each higher spline order (quadratic, cubic...), compute the *averaging knot location* and re-estimate the spline coefficients by LS.

Motivation

GeDS estimation method

Generalized Additive Models with GeDS

Functional Gradient Boosting with GeDS

Insurance Data Application

GAM-GeDS

The **Generalized Additive Model (GAM)** assumes the response variable, $Y \sim E.F.$, and relates its conditional expectation, $\mu = E[Y|X]$, to the predictor variables, X_1, \dots, X_P , via a link function $g(\cdot)$:

$$g(\mu) = \alpha + \sum_{j=1}^P f_j(X_j), \text{ with } \mathbb{E}[f_j(X_j)] = 0, \quad j = 1, \dots, P \quad (1)$$

Hastie and Tibshirani, [1990](#) — *local-scoring* and *backfitting* algorithms in conjunction with scatterplot smoothers, to fit GAMs.

GAM-GeDS

The **Generalized Additive Model (GAM)** assumes the response variable, $Y \sim E.F.$, and relates its conditional expectation, $\mu = E[Y|X]$, to the predictor variables, X_1, \dots, X_P , via a link function $g(\cdot)$:

$$g(\mu) = \alpha + \sum_{j=1}^P f_j(X_j), \text{ with } \mathbb{E}[f_j(X_j)] = 0, \quad j = 1, \dots, P \quad (1)$$

Hastie and Tibshirani, 1990 — *local-scoring* and *backfitting* algorithms in conjunction with scatterplot smoothers, to fit GAMs.

GAM with GeD Splines: Local-scoring algorithm using GeD splines as the function smoothers, f_j , within the backfitting algorithm.

Motivation

GeDS estimation method

Generalized Additive Models with GeDS

Functional Gradient Boosting with GeDS

Insurance Data Application

FGB-GeDS

- **Functional Gradient Boosting** (Friedman, 2001): ensemble machine learning technique that iteratively combines multiple simple models ('weak-learners'), each striving to enhance the performance of the previous accumulative model.
 - *Component-wise Gradient Boosting* (Bühlmann and Yu, 2003; Schmid and Hothorn, 2008): boosting algorithm for fitting additive models, inherently performing variable selection; implemented in **mboost** package (boosting with P-splines).

FGB-GeDS

- **Functional Gradient Boosting** (Friedman, 2001): ensemble machine learning technique that iteratively combines multiple simple models ('weak-learners'), each striving to enhance the performance of the previous accumulative model.
 - **Component-wise Gradient Boosting** (Bühlmann and Yu, 2003; Schmid and Hothorn, 2008): boosting algorithm for fitting additive models, inherently performing variable selection; implemented in **mboost** package (boosting with P-splines).


✱ FGB with GEDS base-learners

- ➡ Flexible control of the strength of the base-learners:
 1. Weak GeDS initial learner + few boosting iterations with strong GeDS learners.
 2. Boosting iterations with weak GeDS learners based on single knot addition with memory.
- ➡ Optimal number of boosting iterations determined by a **stopping rule** based on a ratio of consecutive deviances.
- ➡ Final boosted fit expressed as a **single spline model.**

Simulated Data Application


Consider the function:

$$f_1(x) = 40 \frac{x}{1 + 100x^2} + 4, \quad x \in (-2, 2)$$




➔ Generate 1,000 random samples, $\{X_i, Y_i\}_{i=1}^N$ with $Y_i \sim N(\mu_i, \sigma)$ with $\sigma = 0.2$, $\mu_i = \eta_i = f_1(X_i)$ and $X_i \sim U[-2, 2]$, $i = 1, \dots, N$, where $N = 500$.

{ **GeDS** : 10 int. knots
FGB-GeDS : initial learner with 2 int. knots + 1 boosting iter. with 8 int. knots
mboost : 10,000 boosting iter. with 36 knots p/iter.




And setting **mboost** to have 10 int. knots p/boosting iter. instead:




Real Data Application

- High pressure neutron barium-iron arsenide (BaFe_2As_2) powder diffraction data (Kimber et al., 2009), with number of observations $N = 1151$.


NGeDS
282 knots
MSE: 59,385.34



NGeDSboost
1 knot p/boosting iter.,
284 boosting iter.
MSE: 59,175.96



mboost
285 knots p/boosting iter.,
10,000 boosting iter.
MSE: 2,760,848.65



Motivation

GeDS estimation method

Generalized Additive Models with GeDS

Functional Gradient Boosting with GeDS

Insurance Data Application


Insurance Data Application

Motorcycle insurance data `swmotorcycle` available through the R package `CASdatasets` (Dutang and Charpentier, 2020).

—→ We follow Delong et al., 2021 and model **gamma claim sizes**:

- ① Gamma GLM regression + Gamma Neural Network regression.
 - ② `mboost`: FGB with P-splines.
 - ③ GAM-GeDS.
 - ④ FGB-GeDS.
- *Response*: `ClaimAmount/ClaimNb`, i.e., the average claim size.
 - *Covariates*: `OwnerAge`; `Gender`; `Area`, `RiskClass`; `VehAge`.
 - *Train/Test split*: **80%/20%**.
- Simulate 100 different splits of data.

GLM/GAM Models



Boosting Models













Table 1: GLM/GAM Models

| | Gamma Deviance | | Time (sec.) | Internal knots (OwnerAge+VehAge) |
|--------------------|----------------|-----------|-------------|-------------------------------------|
| | Train Data | Test Data | | |
| GLM | 1.585727 | 1.694797 | 0.008708 | - |
| GLM NN | 1.719903 | 1.859394 | 167.224576 | - |
| GAM-GeDS quadratic | 1.557612 | 1.686492 | 0.671260 | 5 |

Table 2: Boosting Models

| | Gamma Deviance | | Time (sec.) | Internal knots p/boosting iter. (OwnerAge+VehAge) | Boosting iterations |
|---|----------------|-----------|-------------|---|------------------------|
| | Train Data | Test Data | | | |
| mboost | 1.610290 | 1.676810 | 0.156095 | 4 | 100 |
| FGB-GeDS linear (2 starting knots) | 1.575972 | 1.648345 | 0.130963 | 2 | 1 |
| FGB-GeDS w/mem. linear (1 starting knot) | 1.575536 | 1.667158 | 0.129040 | 1 | 3 |

-  Bühlmann, P., & Yu, B. (2003). Boosting with the l2 loss. *Journal of the American Statistical Association*, 98(462), 324–339. <https://doi.org/10.1198/016214503000125>
-  Delong, E., Lindholm, M., & Wüthrich, M. V. (2021). Making tweedie's compound poisson model more accessible. *European Actuarial Journal*, 11(1), 185–226. <https://doi.org/10.1007/s13385-021-00264-3>
-  Dimitrova, D. S., Kaishev, V. K., Lattuada, A., & Verrall, R. J. (2023). Geometrically designed variable knot splines in generalized (non-)linear models. *Applied Mathematics and Computation*, 436, 127493. <https://doi.org/https://doi.org/10.1016/j.amc.2022.127493>
-  Dutang, C., & Charpentier, A. (2020). *Casdatasets: Insurance datasets* [R package version 1.0-11].
-  Friedman, J. H. (2001). Greedy function approximation: A gradient boosting machine.. *The Annals of Statistics*, 29(5), 1189–1232. <https://doi.org/10.1214/aos/1013203451>
-  Hastie, T., & Tibshirani, R. (1990). Generalized additive models. *Monographs on statistics and applied probability. Chapman & Hall*, 43, 335.
-  Kaishev, V. K., Dimitrova, D. S., Haberman, S., & Verrall, R. J. (2016). Geometrically designed, variable knot regression splines. *Computational Statistics*, 31(3), 1079–1105. <https://doi.org/10.1007/s00180-015-0621-7>
-  Kimber, S. A. J., Kreyssig, A., Zhang, Y.-Z., Jeschke, H. O., Valentí, R., Yokaichiya, F., Colombier, E., Yan, J., Hansen, T. C., Chatterji, T., McQueeney, R. J., Canfield, P. C., Goldman, A. I., & Argyriou, D. N. (2009). Similarities between structural distortions under pressure and chemical doping in superconducting BaFe2As2. *Nature Materials*, 8(6), 471–475.
-  Schmid, M., & Hothorn, T. (2008). Boosting additive models using component-wise p-splines. *Computational Statistics & Data Analysis*, 53(2), 298–311. <https://doi.org/https://doi.org/10.1016/j.csda.2008.09.009>