

First Model Problem

- Poisson Equation in 2D: $\nabla^2 \varphi = f$ or $\frac{\partial^2 \varphi}{\partial x^2} + \frac{\partial^2 \varphi}{\partial y^2} = f(x, y)$

Step 1: Define Solution Domain

Step 2: Define Mesh

Step 3: Approximate Spatial Derivatives

Step 4: Substitute into PDE

Step 5: Incorporate Boundary Conditions

Step 5: Incorporate Boundary Conditions

- Example #2:

- Neumann at x_0 : $\left. \frac{\partial \varphi}{\partial x} \right|_{x_0} = g(y)$

- approximate derivative by one-sided finite difference formula

$$\left. \frac{\partial \varphi}{\partial x} \right|_{0,j} = \frac{-3\varphi_{0,j} + 4\varphi_{1,j} - \varphi_{2,j}}{2\Delta}$$

- solve for $\varphi_{0,j}$


$$\varphi_{0,j} = \frac{1}{3} (-2\Delta g_j + 4\varphi_{1,j} - \varphi_{2,j})$$

$$\varphi_{2,j} - 4\varphi_{1,j} + \varphi_{0,j} + \varphi_{1,j+1} + \varphi_{1,j-1} = \Delta^2 f_{1,j}$$

- substitute into finite difference formula at $i=1$


$$\frac{2}{3}\varphi_{2,j} - \frac{8}{3}\varphi_{1,j} + \varphi_{1,j+1} + \varphi_{1,j-1} = \Delta^2 f_{i,j} + \frac{2}{3}\Delta g_j$$

➡ Neumann boundary conditions modify both sides!



Step 6: Matrix Form (only for illustration, never code!)


- Example:
 - use Dirichlet boundary conditions
 - $M=N=4 \Rightarrow 3 \times 3$ interior points
- ➡ the solution vector ϕ covers 2D space
⇒ need to define a sorting



Step 6: Matrix Form (only for illustration, never code!)

- Example:
 - use Dirichlet boundary conditions
 - $M=N=4 \Rightarrow 3 \times 3$ interior points
- ➡ the solution vector φ covers 2D space
⇒ need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$




Step 6: Matrix Form (only for illustration, never code!)

- Example:
 - use Dirichlet boundary conditions
 - $M=N=4 \Rightarrow 3 \times 3$ interior points
- ➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$

$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$



Step 6: Matrix Form (only for illustration, never code!)

- Example:


- use Dirichlet boundary conditions
- $M=N=4 \Rightarrow 3 \times 3$ interior points

➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$

$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$

$$(3,1): \quad -4\varphi_{3,1} + \varphi_{2,1} + \varphi_{3,2} = \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0}$$



Step 6: Matrix Form (only for illustration, never code!)

- Example:

- use Dirichlet boundary conditions
- $M=N=4 \Rightarrow 3 \times 3$ interior points


➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$

$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$

$$(3,1): \quad -4\varphi_{3,1} + \varphi_{2,1} + \varphi_{3,2} = \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0}$$

$$(1,2): \quad \varphi_{2,2} - 4\varphi_{1,2} + \varphi_{1,3} + \varphi_{1,1} = \Delta^2 f_{1,2} - \varphi_{0,2}$$



Step 6: Matrix Form (only for illustration, never code!)

- Example:

- use Dirichlet boundary conditions
- $M=N=4 \Rightarrow 3 \times 3$ interior points

➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting


$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$

$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$

$$(3,1): \quad -4\varphi_{3,1} + \varphi_{2,1} + \varphi_{3,2} = \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0}$$

$$(1,2): \quad \varphi_{2,2} - 4\varphi_{1,2} + \varphi_{1,3} + \varphi_{1,1} = \Delta^2 f_{1,2} - \varphi_{0,2}$$

$$(2,2): \quad \varphi_{3,2} - 4\varphi_{2,2} + \varphi_{1,2} + \varphi_{2,3} + \varphi_{2,1} = \Delta^2 f_{2,2}$$



Step 6: Matrix Form (only for illustration, never code!)

- Example:

- use Dirichlet boundary conditions
- $M=N=4 \Rightarrow 3 \times 3$ interior points

➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$


$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$

$$(3,1): \quad -4\varphi_{3,1} + \varphi_{2,1} + \varphi_{3,2} = \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0}$$

$$(1,2): \quad \varphi_{2,2} - 4\varphi_{1,2} + \varphi_{1,3} + \varphi_{1,1} = \Delta^2 f_{1,2} - \varphi_{0,2}$$

$$(2,2): \quad \varphi_{3,2} - 4\varphi_{2,2} + \varphi_{1,2} + \varphi_{2,3} + \varphi_{2,1} = \Delta^2 f_{2,2}$$

$$(3,2): \quad -4\varphi_{3,2} + \varphi_{2,2} + \varphi_{3,3} + \varphi_{3,1} = \Delta^2 f_{3,2} - \varphi_{4,2}$$



Step 6: Matrix Form (only for illustration, never code!)

- Example:

- use Dirichlet boundary conditions
- $M=N=4 \Rightarrow 3 \times 3$ interior points

➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$

$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$


$$(3,1): \quad -4\varphi_{3,1} + \varphi_{2,1} + \varphi_{3,2} = \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0}$$

$$(1,2): \quad \varphi_{2,2} - 4\varphi_{1,2} + \varphi_{1,3} + \varphi_{1,1} = \Delta^2 f_{1,2} - \varphi_{0,2}$$

$$(2,2): \quad \varphi_{3,2} - 4\varphi_{2,2} + \varphi_{1,2} + \varphi_{2,3} + \varphi_{2,1} = \Delta^2 f_{2,2}$$

$$(3,2): \quad -4\varphi_{3,2} + \varphi_{2,2} + \varphi_{3,3} + \varphi_{3,1} = \Delta^2 f_{3,2} - \varphi_{4,2}$$

$$(1,3): \quad \varphi_{2,3} - 4\varphi_{1,3} + \varphi_{1,2} = \Delta^2 f_{1,3} - \varphi_{0,3} - \varphi_{1,4}$$



Step 6: Matrix Form (only for illustration, never code!)

- Example:

- use Dirichlet boundary conditions
- $M=N=4 \Rightarrow 3 \times 3$ interior points

➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$

$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$

$$(3,1): \quad -4\varphi_{3,1} + \varphi_{2,1} + \varphi_{3,2} = \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0}$$


$$(1,2): \quad \varphi_{2,2} - 4\varphi_{1,2} + \varphi_{1,3} + \varphi_{1,1} = \Delta^2 f_{1,2} - \varphi_{0,2}$$

$$(2,2): \quad \varphi_{3,2} - 4\varphi_{2,2} + \varphi_{1,2} + \varphi_{2,3} + \varphi_{2,1} = \Delta^2 f_{2,2}$$

$$(3,2): \quad -4\varphi_{3,2} + \varphi_{2,2} + \varphi_{3,3} + \varphi_{3,1} = \Delta^2 f_{3,2} - \varphi_{4,2}$$

$$(1,3): \quad \varphi_{2,3} - 4\varphi_{1,3} + \varphi_{1,2} = \Delta^2 f_{1,3} - \varphi_{0,3} - \varphi_{1,4}$$

$$(2,3): \quad \varphi_{3,3} - 4\varphi_{2,3} + \varphi_{1,3} + \varphi_{2,2} = \Delta^2 f_{2,3} - \varphi_{2,4}$$



Step 6: Matrix Form (only for illustration, never code!)

- Example:

- use Dirichlet boundary conditions
- $M=N=4 \Rightarrow 3 \times 3$ interior points

➔ the solution vector φ covers 2D space
 \Rightarrow need to define a sorting

$$(1,1): \quad \varphi_{2,1} - 4\varphi_{1,1} + \varphi_{1,2} = \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0}$$

$$(2,1): \quad \varphi_{3,1} - 4\varphi_{2,1} + \varphi_{1,1} + \varphi_{2,2} = \Delta^2 f_{2,1} - \varphi_{2,0}$$

$$(3,1): \quad -4\varphi_{3,1} + \varphi_{2,1} + \varphi_{3,2} = \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0}$$

$$(1,2): \quad \varphi_{2,2} - 4\varphi_{1,2} + \varphi_{1,3} + \varphi_{1,1} = \Delta^2 f_{1,2} - \varphi_{0,2}$$


$$(2,2): \quad \varphi_{3,2} - 4\varphi_{2,2} + \varphi_{1,2} + \varphi_{2,3} + \varphi_{2,1} = \Delta^2 f_{2,2}$$

$$(3,2): \quad -4\varphi_{3,2} + \varphi_{2,2} + \varphi_{3,3} + \varphi_{3,1} = \Delta^2 f_{3,2} - \varphi_{4,2}$$

$$(1,3): \quad \varphi_{2,3} - 4\varphi_{1,3} + \varphi_{1,2} = \Delta^2 f_{1,3} - \varphi_{0,3} - \varphi_{1,4}$$

$$(2,3): \quad \varphi_{3,3} - 4\varphi_{2,3} + \varphi_{1,3} + \varphi_{2,2} = \Delta^2 f_{2,3} - \varphi_{2,4}$$

$$(3,3): \quad -4\varphi_{3,3} + \varphi_{2,3} + \varphi_{3,2} = \Delta^2 f_{3,3} - \varphi_{4,3} - \varphi_{3,4}$$



Step 6: Matrix Form (only for illustration, never code!)

• Example:

$$\begin{bmatrix}
 -4 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
 1 & -4 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\
 0 & 1 & -4 & 0 & 0 & 1 & 0 & 0 & 0 \\
 1 & 0 & 0 & -4 & 1 & 0 & 1 & 0 & 0 \\
 0 & 1 & 0 & 1 & -4 & 1 & 0 & 1 & 0 \\
 0 & 0 & 1 & 0 & 1 & -4 & 0 & 0 & 1 \\
 0 & 0 & 0 & 1 & 0 & 0 & -4 & 1 & 0 \\
 0 & 0 & 0 & 0 & 1 & 0 & 1 & -4 & 1 \\
 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & -4
 \end{bmatrix}
 \begin{bmatrix}
 \varphi_{1,1} \\
 \varphi_{2,1} \\
 \varphi_{3,1} \\
 \varphi_{1,2} \\
 \varphi_{2,2} \\
 \varphi_{3,2} \\
 \varphi_{1,3} \\
 \varphi_{2,3} \\
 \varphi_{3,3}
 \end{bmatrix}
 =
 \begin{bmatrix}
 \Delta^2 f_{1,1} - \varphi_{0,1} - \varphi_{1,0} \\
 \Delta^2 f_{2,1} - \varphi_{2,0} \\
 \Delta^2 f_{3,1} - \varphi_{4,1} - \varphi_{3,0} \\
 \Delta^2 f_{1,2} - \varphi_{0,2} \\
 \Delta^2 f_{2,2} \\
 \Delta^2 f_{3,2} - \varphi_{4,2} \\
 \Delta^2 f_{1,3} - \varphi_{0,3} - \varphi_{1,4} \\
 \Delta^2 f_{2,3} - \varphi_{2,4} \\
 \Delta^2 f_{3,3} - \varphi_{4,3} - \varphi_{3,4}
 \end{bmatrix}$$

➔ Symmetric block-tridiagonal matrix!

$$A = \begin{bmatrix} B & C & 0 \\ C & B & C \\ 0 & C & B \end{bmatrix} \quad B = \begin{bmatrix} -4 & 1 & 0 \\ 1 & -4 & 1 \\ 0 & 1 & -4 \end{bmatrix} \quad C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$


➔ next: we need to solve $A\vec{\varphi} = \vec{b}$

Step 7: Solve $A\vec{\phi} = \vec{b}$

- Two options:
 - 1) Direct method: Gaussian elimination
 - 2) Iterative methods
 - Gaussian elimination usually not efficient
 - operation count is $O(N^3)$
 - solution only available after $O(N^3)$ operations at the very end!
- ➡ Iterative methods are typically preferable

Step 7: Solve $A\vec{\varphi} = \vec{b}$: Iterative Methods

- Idea: Split A into $A = A_1 - A_2$

$$A\vec{\varphi} = \vec{b} \quad \Leftrightarrow \quad (A_1 - A_2)\vec{\varphi} = \vec{b} \quad \Leftrightarrow \quad A_1\vec{\varphi} = A_2\vec{\varphi} + \vec{b}$$


➡ to find the solution, iteratively solve this

$$A_1\vec{\varphi}^{(k+1)} = A_2\vec{\varphi}^{(k)} + \vec{b}$$


iteration counter/index

- This works, if

1) A_1 is easily invertible

$$\vec{\varphi}^{(k+1)} = A_1^{-1} A_2 \vec{\varphi}^{(k)} + A_1^{-1} \vec{b}$$

2) Iterations converge to true solution

$$\lim_{k \rightarrow \infty} \vec{\varphi}^{(k)} = \vec{\varphi}$$

How can we ensure this? let's define the error:

$$\vec{\varepsilon}^{(k)} = \vec{\varphi} - \vec{\varphi}^{(k)} \quad \Rightarrow \quad \lim_{k \rightarrow \infty} \vec{\varepsilon}^{(k)} = \vec{0}$$

$$A_1 \vec{\varphi} = A_2 \vec{\varphi} + \vec{b}$$

$$\vec{\varepsilon}^{(k)} = \vec{\varphi} - \vec{\varphi}^{(k)}$$

$$\text{ - } A_1 \vec{\varphi}^{(k+1)} = A_2 \vec{\varphi}^{(k)} + \vec{b}$$

$$A_1 \left(\vec{\varphi} - \vec{\varphi}^{(k+1)} \right) = A_2 \left(\vec{\varphi} - \vec{\varphi}^{(k)} \right)$$

$$A_1 \vec{\varepsilon}^{(k+1)} = A_2 \vec{\varepsilon}^{(k)}$$

$$\vec{\varepsilon}^{(k+1)} = A_1^{-1} A_2 \vec{\varepsilon}^{(k)}$$

now use linear algebra

$$\vec{\varepsilon}^{(k)} = \left(A_1^{-1} A_2 \right)^k \vec{\varepsilon}^{(0)}$$

from linear algebra we also know:

$$\lim_{k \rightarrow \infty} \vec{\varepsilon}^{(k)} = 0 \quad \text{if} \quad \rho \left(A_1^{-1} A_2 \right) = |\lambda_i|_{max} < 1$$

ρ : spectral radius
 λ_i : eigenvalues of $(A_1^{-1} A_2)$

\Rightarrow the smaller the spectral radius ρ , the faster the convergence!

• We thus need

- split of A into $A = A_1 - A_2$ with
 - A_1 easily invertible
 - $\max |\lambda_i| < 1$

Step 7: Solve $A\vec{\varphi} = \vec{b}$: Iterative Methods

Point Jacobi Method

- Let $A_I = D$: diagonal entries of A
 - in our example:

$$A_1 = -4I$$

- A_1^{-1} is easy: $A_1^{-1} = -\frac{1}{4}I$

we had: $\vec{\varphi}^{(k+1)} = A_1^{-1} A_2 \vec{\varphi}^{(k)} + A_1^{-1} \vec{b}$

$$\Rightarrow \vec{\varphi}^{(k+1)} = -\frac{1}{4} A_2 \vec{\varphi}^{(k)} - \frac{1}{4} \vec{b} \quad \text{with} \quad A_2 = A_1 - A = -4I - A$$

$$\Rightarrow \varphi_{i,j}^{(k+1)} = \frac{1}{4} \left(\varphi_{i-1,j}^{(k)} + \varphi_{i+1,j}^{(k)} + \varphi_{i,j-1}^{(k)} + \varphi_{i,j+1}^{(k)} \right) - \frac{1}{4} b_{i,j}$$

$$A = \begin{bmatrix} -4 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & -4 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & -4 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & -4 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & -4 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & -4 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & -4 & 1 & 0 \\ 0 & -1 & 0 & 0 & -1 & 0 & 0 & -4 & 0 \\ -1 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & 0 \\ 0 & -1 & 0 & -1 & 0 & -1 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 \end{bmatrix}$$

$$A_2 = \begin{bmatrix} 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 & -1 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & 0 \\ 0 & -1 & 0 & -1 & 0 & -1 & 0 & -1 & 0 \\ 0 & 0 & -1 & 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & -1 & 0 \end{bmatrix}$$

Step 7: Solve $A\vec{\phi} = \vec{b}$: Iterative Methods

Point Jacobi Method

- Let $A_I = D$: diagonal entries of A
 - What are the eigenvalues of $(A_1^{-1} A_2)$?

$$\lambda_{m,n} = \frac{1}{2} \left[\cos \left(\frac{m\pi}{M} \right) + \cos \left(\frac{n\pi}{N} \right) \right]$$

$$m = 1, \dots, M-1$$

$$n = 1, \dots, N-1$$

- expand cosines [...]:

$$|\lambda|_{\max} = 1 - \frac{1}{4} \left(\frac{\pi^2}{M^2} + \frac{\pi^2}{N^2} \right) + \dots$$

$$\text{in 1D: } |\lambda|_{\max} = \cos \left(\frac{\pi}{N} \right)$$

- ➔ if $M, N \uparrow \Rightarrow \rho \rightarrow 1$ (but always $\rho < 1$)
- ➔ slow convergence \Rightarrow Point Jacobi seldom used in practice

Step 7: Solve $A\vec{\varphi} = \vec{b}$: Iterative Methods

Point Jacobi Method

- How to code?
 - use 2 storage vectors, one for $\vec{\varphi}^{(k+1)}$ and one for $\vec{\varphi}^{(k)}$

```
while not converged (or loop from k=1 to #requested iterations)
  loop from j = 1 to N-1
    loop from i = 1 to M-1
      phinew(i,j) = 0.25*(phi(i-1,j)+phi(i+1,j)+...
    end loop i
  end loop j
  phi = phinew
end while
```

$$\varphi_{i,j}^{(k+1)} = \frac{1}{4} \left(\varphi_{i-1,j}^{(k)} + \varphi_{i+1,j}^{(k)} + \varphi_{i,j-1}^{(k)} + \varphi_{i,j+1}^{(k)} \right) - \frac{1}{4} b_{i,j}$$


Step 7: Solve $A\vec{\varphi} = \vec{b}$: Iterative Methods

Gauss-Seidel Method

$$\text{PJ: } \varphi_{i,j}^{(k+1)} = \frac{1}{4} \left(\varphi_{i-1,j}^{(k)} + \varphi_{i+1,j}^{(k)} + \varphi_{i,j-1}^{(k)} + \varphi_{i,j+1}^{(k)} \right) - \frac{1}{4} b_{i,j}$$

- In Point-Jacobi, we use only $\vec{\varphi}^{(k)}$ to calculate $\vec{\varphi}^{(k+1)}$
- Idea: Why don't we use $\vec{\varphi}^{(k+1)}$ instead of $\vec{\varphi}^{(k)}$ for nodes we have already calculated?

for our sorting, for $\vec{\varphi}_{i,j}^{(k+1)}$ this would be
 $\vec{\varphi}_{i,j-1}^{(k+1)}$ and $\vec{\varphi}_{i-1,j}^{(k+1)}$



$$\Rightarrow \varphi_{i,j}^{(k+1)} = \frac{1}{4} \left(\varphi_{i-1,j}^{(k+1)} + \varphi_{i+1,j}^{(k)} + \varphi_{i,j-1}^{(k+1)} + \varphi_{i,j+1}^{(k)} \right) - \frac{1}{4} b_{i,j}$$

➡ to code: use just one storage vector!

Step 7: Solve $A\vec{\varphi} = \vec{b}$: Iterative Methods

Gauss-Seidel Method

- How to code?
 - use 1 storage vector that gets overwritten

```
while not converged (or loop from k=1 to #requested iterations)
  loop from j = 1 to N-1
    loop from i = 1 to M-1
      phi(i,j) = 0.25*(phi(i-1,j)+phi(i+1,j)+...
    end loop i
  end loop j
end while
```

$$\varphi_{i,j}^{(k+1)} = \frac{1}{4} \left(\varphi_{i-1,j}^{(k+1)} + \varphi_{i+1,j}^{(k)} + \varphi_{i,j-1}^{(k+1)} + \varphi_{i,j+1}^{(k)} \right) - \frac{1}{4} b_{i,j}$$

Step 7: Solve $A\vec{\phi} = \vec{b}$: Iterative Methods

Gauss-Seidel Method

- in matrix notation: $A_1 = D - L$ and $A_2 = U$
 - What are the eigenvalues of $(A_1^{-1}A_2)$?

$$\lambda_{m,n} = \frac{1}{4} \left[\cos\left(\frac{m\pi}{M}\right) + \cos\left(\frac{n\pi}{N}\right) \right]$$

L : - lower triangular part of A
 U : - upper triangular part of A
Note: $A \neq LU$

$$m = 1, \dots, M - 1$$

$$n = 1, \dots, N - 1$$

- factor 2 faster than Point Jacobi (and need only half the storage!)

$$\text{PJ: } \lambda_{m,n} = \frac{1}{2} \left[\cos\left(\frac{m\pi}{M}\right) + \cos\left(\frac{n\pi}{N}\right) \right]$$