# EAVA: An R package for Expert Algorithm Verbal Autopsy (EAVA) cause of death assignment

30 May 2025

#### Summary

EAVA (Expert Algorithm Verbal Autopsy) is an R package for determining cause of death (COD) using verbal autopsy (VA) interview questionnaire data, a predominant tool to determine cause of death in under-resourced settings (Nichols et al. 2018). The EAVA R package translates the methodology created and validated by Kalter and colleagues (Kalter, Perin, and Black 2016) to diagnose a cause of death from VA records of neonates and children (1 to 59 months of age). EAVA inputs data from the 2016 version of the WHO Verbal Autopsy (VA) questionnaire and outputs a cause of death (COD) for each VA record based on a deterministic hierarchy of causes. The assignment approach utilizes the decedent's age to decide which of two separate hierarchies will determine a cause of death.

#### Statement of need

Most deaths occur outside of a medical setting and as a result, the causes of these deaths are not captured. Verbal autopsies conducted in the community, usually at the homes of children who died, are used to understand the most common causes of death in settings where civil registration systems need to be strengthened (WHO 2016). Physicians assess signs and symptoms, which were reported to be present at the time of death by a decedent's caregiver during the Verbal Autopsy, and assign a cause of death deemed most likely. To account for bias, physician-coded verbal autopsy (PCVA) requires that verbal autopsy questionnaire data are read by multiple physicians. PCVA is time-intensive for physicians in resource-limited settings and repeatability can be low in childhood deaths (Chandramohan et al. 1998).

Over the last decade, there has been increased adoption of algorithmic cause-ascertainment from VA. These algorithms, termed as Computer-Coded Verbal Autopsy (CCVA), are considerably less time- and resource-intensive than PCVA, facilitating scalability of obtaining COD for large (national- or sub-national-level) VA databases. There now exists a suite of CCVA algorithms for COD ascertainment from VA data – EAVA, InterVA4, InterVA5, InSilicoVA, Tariff, and Naïve Bayes Classifier (Li et al. 2022). Each algorithm differs in implementation details and underlying methodology. However, except EAVA, most of these are primarily data-driven, estimating some form of a conditional probability symptom-given-cause matrix. These estimates are primarily derived from the Population Health Metrics Research Consortium (PHMRC) study, conducted in 2011, which contains both VA records and validated 'gold-standard' causes (Murray et al. 2011). Reliance on PHMRC limits generalizability in newer VA studies. It has been shown that most CCVA algorithms misclassify the cause for a substantial proportion of deaths (Datta et al. 2021; Fiksel et al. 2022).

EAVA addresses some of the drawbacks of physician coding and is not reliant on PHMRC data. It automates and replicates the decision trees of human physician coders as it assesses signs and symptoms of common causes of death, arriving at a single diagnosis using ICD-10 classifications in the hierarchy. If diagnostic criteria are not met for any cause, the neonate or 1-to-59-month-old child is assigned a COD of "unspecified" (Appendices 1 and 2). EAVA has been shown to yield comparable accuracy to the other CCVA algorithms

(Fiksel et al. 2023; Gilbert et al. 2023). The COD outputs from EAVA can also be used in the VA calibration algorithm which combines COD ascertainment from multiple CCVA algorithms and adjusts for their biases to produce a calibrated estimate of population-level cause-specific mortality fractions (Datta et al. 2021; Fiksel et al. 2022; Pramanik et al. 2023). Inclusion of CCVA algorithms with different cause-ascertainment logic ensures robustness of results for VA calibration. Hence, due to the fundamentally different decision-making nature of EAVA compared to other CCVA algorithms, it is now a central component of the VA calibration algorithm and has been used in VA-calibration to produce bias-corrected estimates of CSMF for child (1-59 months) and neonatal deaths in Mozambique (Fiksel et al. 2023; Gilbert et al. 2023; Macicame et al. 2023)

The EAVA R package takes EAVA analytical scripts originally compiled in SAS and R and makes the methodology publicly available in CRAN, which expands the potential for research use, ongoing development, and future integration into VA pipelines and toolkits.

#### State of the field

Many of the aforementioned CCVA algorithms (EAVA, InterVA4, InterVA5, InSilicoVA, Tariff, and Naïve Bayes Classifier) are implemented in the openVA R package (Zehang et al. 2024). The CrossVA package converts 2016 WHO VA questionnaire data to a standard input format for use in openVA (Thomas et al. 2021). There is also standalone software for some CCVA algorithms, for example, the InterVA algorithm (Byass 2020) and the SmartVA algorithm (Flaxman 2025). There has been no publicly available version of EAVA prior to this R package.

#### The EAVA package


The EAVA R package comprises two functions (Wilson et al. 2025). The first function is odk2EAVA which builds on the CrossVA package to convert interview responses from the 2016 WHO Verbal Autopsy questionnaire into standardized inputs for use in codEAVA. The second function, codEAVA, evaluates whether reported symptoms meet diagnostic criteria of common causes of death and assigns a main cause based on a hierarchy of causes. The algorithm utilizes age-group specific ascertainment logic due to significant differences between the common causes of death of neonates 0-27 days (Appendix 1) and children aged 1-to-59-months (Appendix 2).

### Acknowledgements


This work was supported by The Gates Foundation Grant INV-03484. We would like to thank the families who participated in VA interviews for the Countrywide Mortality Surveillance for Action project.


### Appendix 1: deterministic hierarchical algorithm to reach a single cause of death in neonates


age 3-27 days at death and convulsions or spasms and either (able to suckle normally duringthe first day of life and neonatal tetanus stopped being able to sucklenormally) or (cried within 5 minutes afterbirth and stopped being able to cry)) head size very small at birth or YES congenital malformation head size very large at birth or mass defect on back of head or spine abnormality ↓ ио (NRD [neonatal respiratory depression]: Did not cry within 5 minutes after birth or did not breathe immediately after birth) + YES intrapartum related events ((NE [neonatal encephalopathy]: not able to suckle normally in the first day of life or convulsions or spasms or lethargy) or 0 days old at death); Bruises or signs of injury on the baby's body at birth J NO Pregnancy <9 months and fast breathing YES → preterm starting on day 0 and no fever and no cold to touch): Pregnancy <8 months Fever and (bulging fontanelle or YES → meningitis convulsions) and (lethargic or unresponsive/unconscious); √ мо More frequent loose or liquid stools than usual and >4 stools on the day diarrhea was most frequent; ↓ NO (fast breathing that lasted 1 day or more or difficult breathing that lasted 1 day or more and lasted until death) and pneumonia 2 or more of: (chest indrawing or grunting or never cried or: stopped crying) 3 NO



## Appendix 2: deterministic hierarchical algorithm to reach a single cause of death in children 1-to-59-months-of-age







#### References

- Byass, Peter. 2020. "InterVA." http://www.byass.uk/interva.
- Chandramohan, Daniel, Gillian H. Maude, Laura C. Rodrigues, and Richard J. Hayes. 1998. "Verbal Autopsies for Adult Deaths: Their Development and Validation in a Multicenter Study." *Tropical Medicine and International Health* 3 (6). https://doi.org/10.1046/j.1365-3156.1998.00255.x.
- Datta, Abhirup, Jacob Fiksel, Agbessi Amouzou, and Scott L. Zeger. 2021. "Regularized Bayesian Transfer Learning for Population-Level Etiological Distributions." *Biostatistics (Oxford, England)* 22 (4): 836–57. https://doi.org/10.1093/biostatistics/kxaa001.
- Fiksel, Jacob, Abhirup Datta, Agbessi Amouzou, and Scott Zeger. 2022. "Generalized Bayes Quantification Learning Under Dataset Shift." *Journal of the American Statistical Association* 117 (540): 2163–81. https://doi.org/10.1080/01621459.2021.1909599.
- Fiksel, Jacob, Brian Gilbert, Emily Wilson, Henry Kalter, Almamy Kante, Aveika Akum, Dianna Blau, et al. 2023. "Correcting for Verbal Autopsy Misclassification Bias in Cause-Specific Mortality Estimates." *The American Journal of Tropical Medicine and Hygiene* 108 (5\_Suppl): 66–77. https://doi.org/10.4269/ajtmh.22-0318.
- Flaxman, Abraham. 2025. "SmartVA-Analyze." https://github.com/ihmeuw/SmartVA-Analyze/releases/tag/v3.0.0.
- Gilbert, Brian, Jacob Fiksel, Emily Wilson, Henry Kalter, Almamy Kante, Aveika Akum, Dianna Blau, et al. 2023. "Multi-Cause Calibration of Verbal Autopsy–Based Cause-Specific Mortality Estimates of Children and Neonates in Mozambique." The American Journal of Tropical Medicine and Hygiene 108 (5 Suppl): 78–89. https://doi.org/10.4269/ajtmh.22-0319.
- Kalter, Henry D., Jamie Perin, and Robert E. Black. 2016. "Validating Hierarchical Verbal Autopsy Expert Algorithms in a Large Data Set with Known Causes of Death." *Journal of Global Health* 6 (1): 010601. https://doi.org/10.7189/jogh.06.010601.
- Li, Zehang Richard, Jason Thomas, Eungang Choi, Tyler H. McCormick, and Samuel J Clark. 2022. "The openVA Toolkit for Verbal Autopsies." *The R Journal* 14 (4): 316–34. https://doi.org/10.32614/RJ-2023-020.
- Macicame, Ivalda, Almamy M. Kante, Emily Wilson, Brian Gilbert, Alain Koffi, Sheila Nhachungue, Celso Monjane, et al. 2023. "Countrywide Mortality Surveillance for Action in Mozambique: Results from a National Sample-Based Vital Statistics System for Mortality and Cause of Death." The American Journal of Tropical Medicine and Hygiene 108 (5\_Suppl): 5–16. https://doi.org/10.4269/ajtmh.22-0367.
- Murray, Christopher JL, Alan D. Lopez, Robert Black, Ramesh Ahuja, Said Mohd Ali, Abdullah Baqui, Lalit Dandona, et al. 2011. "Population Health Metrics Research Consortium Gold Standard Verbal Autopsy Validation Study: Design, Implementation, and Development of Analysis Datasets." *Population Health Metrics* 9 (1): 27–27. https://doi.org/10.1186/1478-7954-9-27.
- Nichols, Erin K., Peter Byass, Daniel Chandramohan, Samuel J. Clark, Abraham D. Flaxman, Robert Jakob, Jordana Leitao, et al. 2018. "The WHO 2016 Verbal Autopsy Instrument: An International Standard Suitable for Automated Analysis by InterVA, InSilicoVA, and Tariff 2.0." *PLOS Medicine* 15 (1): e1–1002486. https://doi.org/10.1371/journal.pmed.1002486.
- Pramanik, Sandipan, Scott Zeger, Dianna Blau, and Abhirup Datta. 2023. "Modeling Structure and Country-Specific Heterogeneity in Misclassification Matrices of Verbal Autopsy-Based Cause of Death Classifiers." https://doi.org/10.48550/arxiv.2312.03192.
- Thomas, Jason, Eungang Choi, Zehang Li, Nicolas Marie, Tyler McCormick, Peter Byass, and Samuel Clark. 2021. "CrossVA: Verbal Autopsy Data Transformation for InSilicoVA and InterVA5 Algorithms." Https://CRAN.R-Project.org/Package=CrossVA. https://doi.org/10.32614/cran.package.crossva.
- WHO. 2016. "Verbal Autopsy Standards: The 2016 WHO Verbal Autopsy Instrument Manual." https://www.who.int/publications/m/item/verbal-autopsy-standards-the-2016-who-verbal-autopsy-instrument.
- Wilson, Emily B., Henry Kalter, Abhi Datta, Sandipan Pramanik, and Robert E. Black. 2025. "EAVA: Deterministic Verbal Autopsy Coding with Expert Algorithm Verbal Autopsy." *Https://CRAN.R-Project.org/Package=EAVA*. https://doi.org/10.32614/cran.package.eava.
- Zehang, Richard Li, Jason Thomas, Tyler H. McCormick, and Samuel J. Clark. 2024. "openVA: Automated Method for Verbal Autopsy." *Https://CRAN.R-Project.org/Package=openVA*. https://doi.org/10.32614/CRAN.package.openVA.