

1 Trigonometric Identities

1.1 Double-Angle Formulas

You can use the formula for $\cos(2x)$ with the identity $\sin^2 x + \cos^2 x = 1$ to produce other useful formulas.

$$\begin{aligned}\sin(2x) &= 2 \sin(x) \cos(x) \\ \cos(2x) &= \cos^2(x) - \sin^2(x) && ** \\ &= \cos^2(x) - (1 - \cos^2(x)) \\ &= 2 \cos^2(x) - 1 && ** \\ \cos(2x) &= \cos^2(x) - \sin^2(x) \\ &= (1 - \sin^2(x)) - \sin^2(x) \\ &= 1 - 2 \sin^2(x) && **\end{aligned}$$

1.2 Half-Angle Formulas

You can then use the double-angle formulas to derive the following.

These are the ones most useful for integral calculus.

Memorizing the original double-angle formulas allows one to derive these easily.

$$\begin{aligned}\cos^2(x) &= \frac{1 + \cos(2x)}{2} \\ \sin^2(x) &= \frac{1 - \cos(2x)}{2}\end{aligned}$$

2 Limits

2.1 e

The function e is defined as a continuous, differentiable function $f(x)$ that satisfies $f'(x) = f(x)$ for all x and $f(0) = 1$.

$$e = \lim_{n \rightarrow 0} (1 + n)^{\frac{1}{n}}$$

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

3 Integrals

3.1 Improper Integral Summary

Integral	$p \leq 1$	$p > 1$	Value
----------	------------	---------	-------

$\int_0^1 \frac{1}{x^p}$	divergent	convergent	$\frac{1}{1-p}$
--------------------------	-----------	------------	-----------------

$\int_1^\infty \frac{1}{x^p}$	divergent	convergent	$\frac{1}{p-1}$
-------------------------------	-----------	------------	-----------------

3.2 Comparison Theorem

If f and g are continuous and $f(x) \geq g(x) \geq 0$ for $x \geq a$ (there is some a where f is now always larger than g) then,

If $\int_a^\infty f(x)dx$ is convergent then the “smaller” integral $\int_a^\infty g(x)dx$ must be convergent too.

If $\int_a^\infty g(x)dx$ is divergent then the “larger” integral $\int_a^\infty f(x)dx$ must be divergent too.

4 Sequences

4.1 Precise Limit Definition

Say we have an arbitrary number $\epsilon > 0$ as a “tolerance band” from the limit L . Assume the sequence converges. There will be some integer N where every $n > N$ holds $|a_n - L| < \epsilon$.

This allows subsequent terms in the sequence to oscillate around the limit L , so long as they remain in our tolerance band ϵ .

4.2 Convergence

A sequence is convergent if:

- Its limit exists.
- We can make a_n closer and closer to L by increasing n .

4.3 Limit Theorems

- If $\lim_{x \rightarrow \infty} f(x) = L$ and $f(n) = a_n$ when n is an integer, then the sequence has the same limit. Essentially, if a function has the same value as the sequence for every integer, then its limit is the same.
- Given an arbitrary value, there will be a number N where every $a_n, n > N$ is larger than the arbitrary value, if the sequence diverges to infinity.
- $\lim_{n \rightarrow \infty} a_n^p = [\lim_{n \rightarrow \infty} a_n]^p$ if $p > 0$ and $a_n > 0$
- $\lim_{n \rightarrow \infty} |a_n| = 0$ then $\lim_{n \rightarrow \infty} a_n = 0$. If the limit of the absolute terms of the sequence is 0, then the limit of the terms is 0.
- If the terms of a convergent sequence ($\lim a_n = L$) are applied to a continuous function, then the result is convergent too.

$$\lim_{n \rightarrow \infty} f(a_n) = f(L)$$

4.4 Squeeze Theorem

If $a_n \leq b_n \leq c_n$ for $n \geq n_0$ and

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$$

then

$$\lim_{n \rightarrow \infty} b_n = L$$

4.5 r^n sequences

Sequences defined as r^n are convergent if $-1 < r \leq 1$.

$$\lim_{n \rightarrow \infty} r^n = \begin{cases} 0 & \text{if } -1 < r < 1 \\ 1 & \text{if } r = 1 \end{cases}$$

4.6 Monotonic Sequence Theorem

Every bounded, monotonic sequence is convergent.

5 Series

5.1 Definition

A series is simply the sum of terms in a sequence.

An infinite series (often simply just called a “sum” or “series”) is what we get when we sum an infinite number of terms in a sequence.

A partial sum is what we get when we sum a finite number of terms in a sequence, s_3 for a_n is the sum of a_1, a_2 , and a_3 .

A series s_n forms its own sequence s_n .

As we increase n in s_n we get closer and closer to the limit—the infinite sum—of the series.

$$s_n = \sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

$$s = \sum_{n=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_n + \dots$$

Suppose $\sum a_n = 3$ and s_n is the n th partial sum of the series.

Since $\sum a_n$ converges on 3, $\lim_{n \rightarrow \infty} a_n = 0$ must be true.

This is a fundamental theorem, if a series is convergent then the limit of its sequence is 0.

However, this does not mean that if the limit of a sequence is 0 that its series is convergent.

Since $\sum a_n$ converges on 3, $\lim_{n \rightarrow \infty} s_n = 3$.

5.2 Example

Take the sequence $a_n = \frac{1}{2^n}$.

This gives us $a_1 = \frac{1}{2}, a_2 = \frac{1}{4}$, etc....

s_2 would then be $a_1 + a_2 = \frac{3}{4}$.

This forms a sequence from the series, $\frac{1}{2}, \frac{3}{4}, \frac{7}{8}, \dots$

This sequence converges on 1 the more terms we add.

This is the infinite sum.

$$\sum_{n=1}^{\infty} \frac{1}{2^n} = \frac{1}{2} + \frac{3}{4} + \cdots + \frac{1}{2^n} + \cdots = 1$$

The sum of a series is $s = \lim_{n \rightarrow \infty} s_n$.

The series will be divergent—and not have a sum—if the sequence s_n diverges.

5.3 Geometric Series

A geometric series occurs when each term of the sequence is multiplied by the preceding one by a common ratio.

$$a \neq 0 \quad \sum_{n=1}^{\infty} ar^{n-1} = a + ar + ar^2 + ar^3 + \cdots + ar^{n-1} + \dots$$

This series is convergent if $|r| < 1$.

The partial sum is defined by the following.

$$s_n = \frac{a(1 - r^n)}{1 - r}$$

The sum of a convergent geometric series is defined by the following.

$$\sum_{n=1}^{\infty} ar^{n-1} = \frac{a}{1-r}$$

5.4 Sum and Difference Theorem

The following are all convergent series if $\sum a_n$ and $\sum b_n$ are both convergent.

- $\sum ca_n = c \sum a_n$
- $\sum(a_n + b_n) = \sum a_n + \sum b_n$
- $\sum(a_n - b_n) = \sum a_n - \sum b_n$

If $\sum a_n$ was convergent and $\sum b_n$ was divergent, you could use this theorem to show the divergence of $\sum(a_n + b_n)$. By the theorem, $\sum(a_n + b_n) - a_n$ would be convergent if both $\sum a_n$ and $\sum b_n$ were. This is equal to just $\sum b_n$. However, since we know $\sum b_n$ is divergent, $\sum(a_n + b_n)$ must also be divergent.

If we were given a power series $\sum(c_n + d_n)x^n$ and told that $\sum c_n x^n$ has a radius of convergence 2 and that $\sum d_n x^n$ has a radius of convergence 3, we can determine the radius of convergence for the whole series.

$\sum c_n x^n$ only converges if $|x| < 2$ and $\sum d_n x^n$ only converges if $|x| < 3$. We know from this theorem that both series must converge for the sum to be convergent. Thus $\sum(c_n + d_n)x^n$ is convergent for $|x| < 2$ and has a radius of convergence 2.

5.5 Test for Divergence

If the series is convergent then the limit of a_n will be 0.

However, we cannot conclude that a series is convergent just because a_n has a limit of 0.

$$\text{If } \sum a_n \text{ converges, then } \lim_{n \rightarrow \infty} a_n = 0$$

Even though we cannot make conclusions about the series being convergent, we can check if the series is divergent.

$$\sum a_n \text{ divergent if } \lim_{n \rightarrow \infty} a_n \neq 0 \text{ or the limit does not exist.}$$

5.6 Integral Test

If f is continuous, positive, and decreasing on $[1, \infty)$ then let $a_n = f(n)$.

$$\text{If } \int_1^{\infty} f(x)dx \text{ is convergent, then } \sum_{n=1}^{\infty} a_n \text{ is convergent.}$$

If $\int_1^\infty f(x)dx$ is divergent, then $\sum_{n=1}^\infty a_n$ is divergent.

5.7 p-series Test

$\sum_{n=1}^\infty \frac{1}{n^p}$ is convergent if $p > 1$ and divergent if $p \leq 1$.

5.8 Error with the Integral Test

Estimating an infinite sum with a finite number of terms yields a remainder $R_n = s - s_n$. This remainder is our error.

$$\int_{n+1}^\infty f(x)dx \leq R_n \leq \int_n^\infty f(x)dx$$

$$s_n + \int_{n+1}^\infty f(x)dx \leq s \leq s_n + \int_n^\infty f(x)dx$$

5.9 Comparison Tests

If $a_n \leq b_n$ for all n , then $\sum a_n$ will be convergent if $\sum b_n$ is convergent.

If $a_n \geq b_n$ for all n , then $\sum a_n$ will be divergent if $\sum b_n$ is divergent.

Essentially, if a bigger series is convergent, then smaller series must be as well.

If a smaller series is divergent, then larger series must be as well.

5.10 Limit Comparison Test ($c > 0$)

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c > 0$$

If this holds ($c > 0$) then either both $\sum a_n$ and $\sum b_n$ converge, or they both diverge.

5.11 Alternating Series Test

An alternating series is one defined by the following.

$$\sum (-1)^{n-1} b_n \quad b_n > 0$$

If the following are satisfied then the series will be convergent.

- $b_{n+1} \leq b_n$ for all n
- $\lim_{n \rightarrow \infty} b_n = 0$

5.12 Alternating Series Estimation Theorem

If the Alternating Series Test is satisfied then the following holds.

$$|R_n| = |s - s_n| \leq b_{n+1}$$

This lets us find a desired error by simply computing a value in the sequence.

How many terms are needed to estimate the sum $\sum \frac{(-1)^n}{n^6}$ such that $|\text{error}| < 0.00005$?

$$|\text{error}| \leq b_{n+1} < \frac{1}{20000}$$

$$(n+1)^6 > 20000$$

$$(5+1)^6 > 20000$$

$$6^6 > 20000$$

$$b_6 > 20000$$

The sixth term is less than the desired error.

Adding the sixth term will not yield more than our desired accuracy.

This means we need five terms to yield a sum with the desired accuracy.

5.13 Error

$$0.00005 = \frac{1}{20,000}$$

$$0.00001 = \frac{1}{100,000}$$

$$0.01 = \frac{1}{100}$$

$$0.02 = \frac{1}{50}$$

$$0.05 = \frac{1}{20}$$

5.14 Absolute and Conditional Convergence

A series is absolutely convergent if $\sum |a_n|$ is convergent.

A series is convergent if it is absolutely convergent.

A series is conditionally convergent if $\sum a_n$ is convergent but not $\sum |a_n|$.

5.15 Ratio and Ratio Test

Take a series $\sum a_n$.

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = L \quad \text{or} \quad \lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L$$

If $L < 1$ then the series is absolutely convergent.

If $L > 1$ or $L = \infty$ then the series is divergent.

If $L = 1$ then the nothing about the series can be concluded.

5.16 Product of Convergent Series

If two series are divergent, their product is not necessarily divergent.

$$a_n = b_n = \frac{(-1)^n}{\sqrt{n}}$$
$$a_n b_n = \frac{1}{n}$$

$\sum a_n$ and $\sum b_n$ are convergent by the alternating series test. $\sum a_n b_n$ is divergent.