


Journal Pre-proof

Ibuprofen use and clinical outcomes in COVID-19 patients

Ehud Rinott, BSc, Eran Kozler, MD, Yonatan Shapira, MD, Adina Bar-Haim, PhD, Ilan Youngster, MD, MMSc



PII: S1198-743X(20)30343-8

DOI: <https://doi.org/10.1016/j.cmi.2020.06.003>

Reference: CMI 2092

To appear in: *Clinical Microbiology and Infection*

Received Date: 12 May 2020

Revised Date: 3 June 2020

Accepted Date: 6 June 2020

Please cite this article as: Rinott E, Kozler E, Shapira Y, Bar-Haim A, Youngster I, Ibuprofen use and clinical outcomes in COVID-19 patients, *Clinical Microbiology and Infection*, <https://doi.org/10.1016/j.cmi.2020.06.003>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

Ibuprofen Use and Clinical Outcomes in COVID-19 Patients

Ehud Rinott, BSc¹, Eran Kozer MD^{2,6}, Yonatan Shapira MD^{3,6}, Adina Bar-Haim PhD⁴
Ilan Youngster MD, MMSc^{5,6}.

¹Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer-Sheva, Israel. ²Pediatric Emergency Unit, ³Division of Internal Medicine, ⁴Molecular Laboratory, ⁵Pediatric Infectious Diseases, Shamir (Assaf Harofeh) Medical Center, Zerifin, Israel. ⁶Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel.

Corresponding author: Ilan Youngster, MD, MMSc. Pediatric Infectious Diseases Unit and the Center for Microbiome Research, Shamir Medical Center, Zerifin 70300, Israel. Phone: +972-506301191 Fax: +972-89779136
Email: youngsteri@shamir.gov.il

Keywords: Ibuprofen; COVID-19; Antipyretics.

Word count: 1200

ABSTRACT

Objective: It was recently suggested that Ibuprofen might increase the risk for severe and fatal COVID-19 disease and should therefore be avoided in this patient population. We aimed to evaluate whether ibuprofen use in patients with COVID-19 was associated with more severe disease, compared to patients using paracetamol or no antipyretics.

Methods: In a retrospective cohort study of patients with COVID-19 from Shamir Medical Center, Israel, we monitored any use of ibuprofen from a week prior to diagnosis of COVID-19 throughout the disease. Primary outcomes were mortality and the need for respiratory support, including oxygen administration and mechanical ventilation.

Results: The study included 403 confirmed cases of COVID-19, with a median age of 45 years. Of the entire cohort, 44 patients (11%) needed respiratory support and 12 (3%) patients died. One hundred and seventy-nine (44%) patients had fever, with 32% using paracetamol and 22% using ibuprofen, for symptom-relief.

In the Ibuprofen group, 3 (3.4%) patients died, while in the non-Ibuprofen group 9 (2.8%) patients died ($P=0.95$). Nine (10.3%) patients from the Ibuprofen group needed respiratory support, as compared with 35 (11%) from the non-Ibuprofen group ($P=1$). When compared to exclusive paracetamol users, no differences were observed in mortality rates or the need for respiratory support among patients using ibuprofen.

Conclusions: In this cohort of COVID-19 patients, Ibuprofen use was not associated with worse clinical outcomes, compared to paracetamol or no antipyretic.

INTRODUCTION

The current outbreak of coronavirus disease 2019 (COVID-19), caused by the novel coronavirus SARS-CoV-2, has rapidly spread throughout the world, and has become a worldwide pandemic with more than two million confirmed cases by mid-April 2020. Though COVID-19 is primarily a respiratory disease, its' insidious and nonspecific clinical course can result in a delayed diagnosis and a prolonged period of self-administered symptomatic treatment. As fever is one of the most common symptoms of COVID-19, antipyretic medications, including ibuprofen, play an important role in controlling patients' symptoms

On March 14, 2020, during the emergence of the COVID-19 outbreak, the French Minister of health published a recommendation to avoid the use of anti-inflammatory medications like ibuprofen or cortisone, claiming it could aggravate infections [1]. Though no reference was provided, the statement may stem from a report published by the French Agency for the Safety of Health Products, concerning 400 cases of severe infections that were linked temporally to Ibuprofen administration [2]. Adding to these concerns, recent in-vitro work has supported the hypothesis that pathogenic coronaviruses have a high affinity to the angiotensin converting enzyme 2 (ACE2) receptor, and that ACE2 production as well as the ACE 2 receptor expression, can be increased by ibuprofen [3,4].

The public statement prompted a global debate about the safety of Ibuprofen as antipyretic treatment for patients with confirmed or even suspected COVID-19 during the current pandemic [1,5]. Several national health agencies worldwide, including the World Health Organization, have responded with official statements emphasizing that no evidence supports these claims [6–8]. Nevertheless some authorities, erring

on the side of caution while awaiting clinical data, have suggested paracetamol should be considered the recommended first-line antipyretic, with ibuprofen reserved for patients that are unable to tolerate paracetamol [9].

In the current study, we aimed to evaluate whether Ibuprofen administration in patients with COVID-19 was associated with worse clinical outcomes, compared to paracetamol or no antipyretic.

METHODS

We conducted a retrospective cohort study of SARS-Cov-2 positive patients tested at Shamir Medical Center, Israel, between March 15-April 15 2020. At that time the policy of the Israeli ministry of health was to test every patient with symptoms that could be compatible with COVID-19 (cough, dyspnea, fever). Nasopharyngeal swabs were tested by real-time reverse transcriptase polymerase chain reaction in the hospital's molecular laboratory that is approved by the Israeli ministry of health. Patients were contacted by phone (mean 13 days after the diagnosis, range 2-30 days), and a short structured questionnaire was administered. Information collected included age, gender, chronic diseases and medications, date of diagnosis, symptoms and factors relating to the clinical course of disease (admission to the hospital and the intensive care unit, need for oxygen or ventilation). For our purpose, severe disease was defined as needing any respiratory support (supplemental oxygen administration or ventilation), ICU admission or death. Patients were asked about the use of any medication containing ibuprofen (a complete list of local commercial names was supplied), paracetamol or dipyrrone starting a week prior to diagnosis of COVID-19. In case of mortality, information was obtained from medical charts and referral letters. Chi-square test of independence, Wilcoxon rank sum-test and Fischer's exact test were applied as needed (significance threshold, $P < .05$).

Statistical analyses were performed using R software, version 3.5.3. The study was approved by the Shamir Medical Center Institutional Review Board.

RESULTS

The study population included 403 confirmed cases of COVID-19 of 437 positive patients during the inclusion period (92%). Median age was 45 years, 220 (55%) were male, and 101 (25%) had chronic diseases.

Of the entire cohort, 47% were discharged to home isolation following their diagnosis, with the remainder being admitted to the hospital (45%) or a nursing facility (8.4%). The vast majority of patients (369, 89%) had only mild symptoms with no need for any respiratory support, however 32 (7.9%) patients received oxygen support, 18 (4.5%) were admitted to the intensive care unit, 17 (4.2%) were ventilated and 12 (3%) patients died. 179 (44%) patients had fever during their disease. The most commonly used antipyretic was paracetamol (32% of the population), with 22% reporting using ibuprofen and 3.7% using dipyron (Table 1).

Ibuprofen intake, mortality and respiratory support in COVID-19 patients

Patients who took ibuprofen at any time during the course of their disease had similar characteristics as those who did not receive ibuprofen except for the fact that 79% percent of patients using Ibuprofen had fever, while only 35% of patients that avoided the use Ibuprofen reported fever ($P<0.001$).

In the Ibuprofen group, 3 (3.4%) of patients died, while in the non-Ibuprofen group 9 (2.8%) of patients died ($P=0.95$). Nine (10%) patients from the Ibuprofen group needed respiratory support, as compared with 35 (11%) from the non-Ibuprofen group ($P=1$) (Table 1).

Ibuprofen vs paracetamol use in febrile COVID-19 patients

Patients with fever that used paracetamol or ibuprofen exclusively had similar age and gender.

No difference in mortality rates was observed (0 vs 3, $P=0.3$). One (2%) of the exclusive ibuprofen users vs 11 (12.9%) of the exclusive paracetamol users needed any respiratory support during their disease course ($P=0.06$).

DISCUSSION

In this retrospective cohort study of 403 COVID-19 cases admitted to Shamir Medical Center in central Israel, we did not observe an increased risk for mortality or the need for respiratory support in patients treated with Ibuprofen. Among patients with fever, no excess mortality or need for respiratory support were observed in patients who chose to use ibuprofen exclusively. In fact, the need for respiratory support was higher in the paracetamol group with borderline significance. One possible explanation for this could be that elderly patients with more severe chronic illnesses are more likely to be treated with paracetamol due to concern about ibuprofen causing renal injury.

Our study has several limitations. First, as with any retrospective study, recall-bias is a concern. However, given the relatively short timeframe included in the study we expect this to be of minor concern. Furthermore, this possible limitation should not favor one treatment over the other. Second, information from deceased patients could only be collected from medical chart review. However, medications administered prior to hospitalization are noted in the charts. Thirdly, our sample size was not sufficient to allow multivariable analyses. Lastly, the policy of the Israeli Ministry of Health was to test only patients with symptoms that are suggestive of COVID-19. Thus, we have no information about ibuprofen in asymptomatic carriers,

and whether antipyretics can influence their clinical course. A major strength of this study is the high response rate and nearly complete follow-up of patients admitted to a tertiary hospital. The balanced representation of the different severities of the disease, compatible with the distribution of disease in the Israeli population, is another strength.

The claim that Ibuprofen is unsafe for use in patients with COVID-19 symptoms was raised in the early stages of the COVID-19 outbreak, following the observation that SARS-CoV-2 bind to its target cell through angiotensin-converting enzyme 2 (ACE2) in the lung [10]. This led to the hypothesis that ACE2-stimulating drugs, such as ACE inhibitors and Ibuprofen, might increase the risk for severe and fatal COVID-19 disease [3]. Regarding ibuprofen, this hypothesis was not supported in our cohort. Larger prospective studies are needed to validate our results.

The authors report no conflict of interest

Funding: No funding was received

Contribution: EK and IY conceptualized the study, IY, YS and ABH collected the data, IY and ER conducted the data analyses, ER wrote the first draft of the manuscript, IY and EK revised the manuscript, all authors approved the final version.

Table 1: Characteristics of study participants

Characteristic	Overall	Without Ibuprofen intake	With Ibuprofen intake	P value	Proportion difference in % (95% confidence interval)
Subjects – no.	403	316	87		
Gender Female/Male (%)	183/220 (45.4/54.6)	142/174 (44.9/55.1)	41/46 (47.1/52.9)	0.81	0.8 (-14.8 to 10.4)
Age (median, interquartile range)	45.00 (25.00, 62.00)	46.00 (25.00, 61.00)	40.00 (24.50, 64.00)	0.96	
Chronic Diseases (%)	101 (25.1)	79 (25.0)	22 (25.3)	1	0.3 (-10.3 to 10.9)
Cardiovascular Disease (%)	52 (12.9)	40 (12.6)	12 (13.7)	0.85	1.1 (-8.2 to 9.6)
Diabetes (%)	38 (9.4)	28 (8.8)	10 (11.4)	0.53	2.6 (-13.7 to 17.6)
Admitted				0.16	
No (%)	188 (46.7)	145 (45.9)	43 (49.4)		
Hospital (%)	181 (44.9)	148 (46.8)	33 (37.9)		
Nursing facility (%)	34 (8.4)	23 (7.3)	11 (12.6)		
Fever (%)	179 (44.4)	110 (34.8)	69 (79.3)	<0.001	44.5 (33.8 to 55.2)
Respiratory Symptoms (%)	194 (48.1)	160 (50.6)	34 (39.1)	0.074	-11.5 (-23.9 to 0.8)
Observed clinical outcomes					
Received supplemental oxygen (%)	32 (7.9)	27 (8.5)	5 (5.7)	0.53	-2.8 (-9.3 to 3.7)
Mechanically ventilated (%)	17 (4.2)	13 (4.1)	4 (4.6)	>0.95	0.5 (-4.9 to 5.9)
Admitted to the Intensive Care Unit (%)	18 (4.5)	13 (4.1)	5 (5.7)	0.72	1.6 (-4.5 to 7.7)
Administration of respiratory support (%)	44 (10.9)	35 (11.1)	9 (10.3)	>0.95	-0.8 (-8.7 to 7.3)
Died (%)	12 (3.0)	9 (2.8)	3 (3.4)	>0.95	0.6 (-4.3 to 5.5)

Statistics presented: n(%), median (IQR)

Statistical test performed: chi-square of independence, Wilcoxon rank-sum test; Fisher's exact test

Table 2: Febrile COVID-19 patients that exclusively used ibuprofen or paracetamol.

Characteristic	Overall	Exclusive paracetamol intake	Exclusive ibuprofen intake	P value	Proportion difference in % (95% confidence interval)
Subjects – no.	134	85	49		
Gender Female/Male	60/74 (44.8/55.2)	41/44 (48.2/51.8)	19/30 (38.8/61.2)	0.4	9.4 (-9.4 to 28.3)
Age	46.00 (24.00, 65.00)	48.00 (27.25, 67.00)	34.50 (23.25, 56.50)	0.09	
Administration of respiratory support	12 (9.0)	11 (12.9)	1 (2.0)	0.06	10.9 (0.3 to 103)
Died	3 (2.2)	3 (3.5)	0 (0.0)	0.3	-3.5 (-9.1 to 2)

Statistics presented: n(%), median (IQR)

Statistical test performed: chi-square of independence, Wilcoxon rank-sum test; Fisher's exact test

REFERENCES

- [1] Day M. Covid-19: ibuprofen should not be used for managing symptoms, say doctors and scientists. BMJ 2020;368:m1086.
<https://doi.org/10.1136/bmj.m1086>.
- [2] Anti-inflammatoires non stéroïdiens (AINS) et complications infectieuses graves - Point d'Information - ANSM : Agence nationale de sécurité du médicament et des produits de santé n.d. https://www.ansm.sante.fr/S-informer/Points-d-information-Points-d-information/Anti-inflammatoires-non-steroidiens-AINS-et-complications-infectieuses-graves-Point-d-Information#_ftn1 (accessed April 16, 2020).

- [3] Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* 2020;8:e21. [https://doi.org/10.1016/S2213-2600\(20\)30116-8](https://doi.org/10.1016/S2213-2600(20)30116-8).
- [4] Shang J, Ye G, Shi K, Wan Y, Luo C, Aihara H, et al. Structural basis of receptor recognition by SARS-CoV-2. *Nature* 2020;581:1–4. <https://doi.org/10.1038/s41586-020-2179-y>.
- [5] Little P. Non-steroidal anti-inflammatory drugs and covid-19. *BMJ* 2020;368:m1185. <https://doi.org/10.1136/bmj.m1185>.
- [6] FDA advises patients on use of non-steroidal anti-inflammatory drugs (NSAIDs) for COVID-19 | FDA n.d. <https://www.fda.gov/drugs/drug-safety-and-availability/fda-advises-patients-use-non-steroidal-anti-inflammatory-drugs-nsaids-covid-19> (accessed April 20, 2020).
- [7] Commission on Human Medicines advice on ibuprofen and coronavirus (COVID-19) - GOV.UK n.d. <https://www.gov.uk/government/news/commission-on-human-medicines-advice-on-ibuprofen-and-coronavirus-covid-19> (accessed April 22, 2020).
- [8] Could Ibuprofen worsen disease for people with COVID19? World Heal Organ 2020:Twitter. <https://twitter.com/WHO/status/1240409217997189128/photo/1> (accessed April 23, 2020).
- [9] EMA advice on the use of NSAIDs for Covid-19 2020. <https://doi.org/10.1016/s2213>.
- [10] Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor Recognition by the Novel Coronavirus from Wuhan: an Analysis Based on Decade-Long Structural Studies of SARS Coronavirus. *J Virol* 2020;94. <https://doi.org/10.1128/JVI.00127-20>.