

Wormholes and the Possibility of Time Travel

Emily Tieu

March 23, 2026

Abstract

By analyzing solutions to Einstein's field equations, it is evident that spacetime admits various nontrivial geometries, including structures that connect two distant regions. Beginning with the Schwarzschild solution, the extension to the Einstein–Rosen bridge demonstrated how Einstein's equations naturally permitted such connections. However, these early models were restricted as non-traversable, limiting their physical applicability. The later development of traversable wormhole models introduced metrics explicitly allowing passage, though at the cost of violating fundamental energy conditions. These spacetime geometries thereby lend themselves to structures called closed timelike curves, creating a model theoretically capable of time travel. Despite their mathematical validity, such solutions require unphysical conditions, presenting limitations to what is permitted by general relativity and what may actually be realized in nature.

1 Introduction

Wormholes can be defined as a multiply-connected spatial geometry with non-trivial topologies that connect two distant regions, either in the same universe or two different universes. Structurally, wormholes are a tube-like passage between two regions in four-dimensional spacetime, where their entrances are referred to as the mouths, while the asymptotically flat tunnel between them is called the wormhole's throat [7]. To motivate the possibility of time travel, wormholes are believed to be a shortcut through space; while the distance between the mouths may be infinite, the distance through the wormhole, by contrast, may be arbitrarily small [11]. However, time travel

requires the wormhole to be traversable, which only exists in the presence of exotic matter, but this violates classical energy conditions [7]. Nonetheless, wormholes effectively allow superluminal travel and generate closed timelike curves, where one mouth of a traversable wormhole may experience a different passage of time than the other [5]. That is, this configuration could hypothetically permit exiting out of the wormhole at an earlier time, enabling backwards time travel. While they may remain as a speculation, these models raise profound questions about the nature of spacetime, causality, and the limits of physical law.

The theory of wormholes emerged from the early solutions to Einstein’s field equations of general relativity. One of the most notable results was Schwarzschild’s solution to the vacuum Einstein equations, which describes the spacetime geometry of static, spherically symmetric mass [6]. Schwarzschild discovered that Einstein’s equations predicted the existence of black holes containing singularities at their centres, a point in which all laws of physics collapse [1]. In response to the anomaly of Schwarzschild singularities, Einstein and Rosen rewrote Schwarzschild’s solution using a coordinate transformation to eliminate the point of infinite density altogether; this approach instead revealed that the singularity would become a bridge connecting two separate regions of spacetime, now known as the Einstein–Rosen bridge [1]. Modern theoretical developments in wormhole models have since expanded upon these early ideas on the basis that general relativity permits exotic spacetime structures, allowing implications of time travel, though physical constraints limit their realization.

2 Evolution of Wormhole Models

2.1 Einstein–Rosen Bridge

To begin, it is important to recall the Schwarzschild solution given by metric

$$ds^2 = - \left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi)^2 \quad (1)$$

where $r = 2M$ is referred to as the Schwarzschild radius [10]. The region containing this hypersurface, known as the event horizon, marks the point of no return. That is, an object within this horizon would be required to travel

faster than the speed of light to escape, which is physically impossible [1]. However, Einstein and Rosen realized that the Schwarzschild radius was not an intrinsic breakdown of spacetime but merely a coordinate singularity. Hence, they introduced the coordinate transformation $u^2 = r - 2M$ to derive

$$ds^2 = -4(u^2 + 2m)du^2 - (u^2 + 2m)^2 d\Omega^2 + \frac{u^2}{u^2 + 2m} dt^2, \quad (2)$$

where $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$ [13]. Thus, the Schwarzschild solution becomes free of singularities for all values of $u \in (-\infty, +\infty)$, and r ranging from 2 to $+\infty$. The line element for the Einstein–Rosen bridge 2 represents a space of two congruent, asymptotically flat sheets corresponding to the Schwarzschild solutions $u > 0$ and $u < 0$ [13]. These disconnected sheets are then connected by a bridge at $u = 0$ (or the $r = 2M$ hypersurface), solidifying itself as the first theoretical wormhole model [13] [3].

However, due to the existence of the event horizons, a second event horizon must appear on the other end of this bridge, effectively blocking the exit [1]. Hence, having event horizons on both ends of the bridge prevents any particle or signal from passing completely through. Furthermore, attaching the two spacetime sheets would reduce the space to a single sheet, posing a “topological gluing problem” [14]. Physically speaking, an observer falling into one mouth of the Einstein–Rosen bridge would be unable to exit the other, as the bridge collapses before traversal can occur. Consequently, the Einstein–Rosen bridge has been ruled out as traversable, though it inspired further studies of human-traversable wormhole models to be explored.

2.2 Morris–Thorne Model

Decades later, Morris and Thorne revolutionized wormhole physics with a new proposed class of traversable wormholes solutions specifically designed to allow passage through the wormhole. In the Schwarzschild coordinates (t, r, θ, ϕ) 1, the geometry of a static, spherically symmetric wormhole can be represented by the following Morris–Thorne metric:

$$ds^2 = -e^{2\varphi(r)} dt^2 + \frac{dr^2}{1 - \frac{b(r)}{r}} + r^2(d\theta^2 + \sin^2 \theta d\phi^2), \quad (3)$$

where $\varphi(r)$ is the gravitational redshift function and $b(r)$ is the shape function.

Morris and Thorne sought to remove the existence of the event horizon to address the limitation of Einstein and Rosen's model 2. To achieve this, $\varphi(r)$ must remain finite everywhere in the domain, ensuring that the metric coefficient $g_{tt} = -e^{2\varphi(r)}$ never vanishes [12]. If $\varphi(r)$ were to diverge, an event horizon would form, preventing two-way travel and rendering the wormhole non-traversable. Thus, the finiteness of $\varphi(r)$ guarantees that signals and observers can pass freely through the wormhole without encountering infinite gravitational redshift. [12]

The radial coordinate r must also range from r_0 , the wormhole's throat radius, to $+\infty$. Unlike the Schwarzschild geometry 1, where r decreases monotonically toward a singularity, the coordinate r reaches a minimum value of r_0 and then increases again, indicating that spacetime opens up into another region. As such, this behaviour reflects the nontrivial topology of the wormhole. Furthermore, the shape function $b(r)$, which determines the spatial geometry of a wormhole, must satisfy certain conditions in order to be traversable [12] [8]. At the throat $r = r_0$, the function follows

$$b(r_0) = r_0, \tag{4}$$

and for $r > r_0$, the inequality

$$\frac{b(r)}{r} < 1 \tag{5}$$

must hold, ensuring that the wormhole remains open and the radial coordinate is non-monotonically decreasing to a minimum at r_0 [8]. Additionally, the radial derivative at the throat must satisfy

$$b'(r_0) < 1, \tag{6}$$

enforcing that the wormhole geometry flares outward at the throat, a necessary condition for traversability [12]. Finally, spacetime should reach asymptotic flatness far from the throat, which requires

$$\lim_{r \rightarrow \infty} \frac{b(r)}{r} = 0, \tag{7}$$

as shown in [12].

A key consequence of these geometric requirements emerges when computing the stress-energy tensor components $T_{\mu\nu}$ for the Morris–Thorne metric 3. In particular, it is evident that the flaring-out condition 6 forces $\rho + p_r$ to be negative, where ρ is the energy density and p_r is the radial pressure. This is a direct violation to the null energy condition (NEC), which requires $\rho + p_r$ to be non-negative. This result implies that the radial tension required to keep the wormhole open must exceed the local energy density. Consequently, Morris and Thorne found that violating the NEC suggests that some form of exotic matter with negative energy density must exist at the throat to maintain the flaring geometry [2].

Unlike earlier solutions, Morris–Thorne wormholes demonstrate that the obstruction to traversability is not purely geometric but also arises from the interaction between geometry and matter, as dictated by the field equations. Thus, despite the dependence of traversable wormholes on exotic matter presenting a major physical limitation, they demonstrate how general relativity permits the mathematical possibility of traversable connections under the right conditions.

3 Implications

Since Gödel’s discovery of rotating universes in 1949, closed timelike curves (CTCs) have been well-established to be permitted in such universes as solutions to Einstein’s field equations [14] [9]. One approach to form CTCs was to modify the Einstein–Rosen bridge 2 by using Eddington–Finkelstein coordinates, resolving both the horizon problem and topological problem, allowing it to be traversable in finite time [14]. This Einstein–Rosen model was then extended to introduce another one-directional bridge, connecting two regions with two distinct bridges to become a closed curve [14]. For a particle following a geodesic parameterized by proper time τ along trajectory $\gamma(\tau)$,

$$g_{\mu\nu} \frac{d\gamma^\mu}{d\tau} \frac{d\gamma^\nu}{d\tau} < 0, \forall \tau \in [\tau_1, \tau_2], \quad (8)$$

where $\tau_1 < \tau_2$ and $\gamma(\tau_1) = \gamma(\tau_2)$ defines a closed timelike curve.

While coordinate time t is continuous across the wormhole throat, its derivative with respect to proper time τ undergoes a sign change. That

is, prior to crossing the throat, the evolution satisfies $\frac{dt}{d\tau} > 0$, indicating a forward progression in coordinate time. After traversing the throat, the relation becomes $\frac{dt}{d\tau} < 0$, corresponding to an apparent reversal toward the past. Furthermore, at the throat itself ($t = t_0$), one obtains $\frac{dt}{d\tau}|_{t_0} = 0$, indicating a time inversion. Physically speaking, this does not imply that time flows backwards; rather, it reflects a global property of spacetime in which the specified coordinate system no longer aligns with the universal notion of time [14]. Despite this reversal in $t(\tau)$, proper time continues to evolve monotonically, ensuring that local causality along $\gamma(\tau)$ is preserved.

As a result, CTCs represent a special closed loop that enables a particle to return to its original spacetime point, allowing it to essentially revisit its own past [9]. Wormholes provide an intuitive mechanism for constructing such curves, effectively acting as shortcuts between different temporal regions. CTCs thereby induces a time loop that warps spacetime, providing a theoretical framework for a time machine [1]. As such, one would only need to obtain a rocket capable of travelling close to the speed of light to transport through this time warp, and the special relativistic effect of time dilation will handle the rest [1].

As closed timelike curves allow for time travel, the concern for upholding causality arises. Specifically, time travel to the past introduces paradoxes, reflecting solutions to Einstein's field equations that do not admit a globally consistent time ordering [9]. However, these paradoxes may be resolved in the quantum regime, leaving CTCs consistent with relativity [9]. Consequently, the study of such systems lends itself to topics of causal structures and nonlinearities in spacetime, crucial components of quantum gravity [9]. While CTC solutions are mathematically valid, their implications of time travel and causality violation highlights the tension between geometric possibility and physical plausibility.

4 Limitations

Although wormhole solutions naturally arise from the Einstein field equations, their physical realization is subject to significant physical constraints. These limitations predominantly stem from the properties of stress-energy tensors, stability considerations, and lack of empirical support. While such

models proposed by Morris and Thorne [3] were successfully able to add traversability to a wormhole solution, an improvement to the original Einstein–Rosen bridge, their solution poses a direct violation of the NEC at the wormhole throat. As the weakest of the energy conditions, a violation of the NEC implies a violation of the others [4]. Such exotic matter have no known classical analogue, thereby posing one of the most challenging obstacles to the physical realization of traversable wormholes.

Moreover, even if exotic matter were available, wormhole solutions are dynamically unstable. Any small perturbation in the metric functions $\varphi(r)$ and $b(r)$ failing to meet their conditions 4–7 may cause singular behaviour or the formation of an event horizon, destroying traversability [8]. This sensitivity accentuates the need for precise and continuous control over matter distributions within a wormhole. Finally, there does not yet exist experimental data supporting the presence of exotic matter satisfying all of the conditions to sustain a traversable wormhole. As there is no empirical evidence supporting the existence of wormholes or exotic matter, the general consensus among theoretical physicists is that wormholes are not plausible in nature [1].

5 Conclusion

The study of wormholes provides insight into the depth and flexibility of general relativity as a mathematical framework. An analysis of these models reveals a consistent pattern: while the geometry required for a wormhole may be depicted through metric functions and curvature conditions, the corresponding stress–energy tensor must exhibit properties that do not exist in classical physics. These results thereby support the central thesis that general relativity permits exotic spacetime structures of mathematical solutions, though physical constraints significantly hinder their realization. Besides from piquing the interest of many, wormholes occupy a unique position in expanding the scope of general relativity by testing the limits of geometry, matter, and causality. Moving forward, future research may look into determining solutions that do not violate energy conditions, or whether these violations may exist in a physically meaningful way through quantum effects. In particular, advances in the study of quantum fields in curved spacetime may provide further insight into the viability of these structures. As such, the investigation of wormholes offers a deeper understanding of the relationship between

spacetime geometry and the fundamental nature of matter and energy.

References

- [1] J. Al-Khalili, *Black holes, Wormholes, and Time Machines*, 2nd ed. Boca Raton, FL: Taylor and Francis, 2012.
- [2] A. Anand, K. Jusufi, and M. Latifi, “Exploring a Novel Einstein–Rosen BTZ wormhole,” *The European Physical Journal. C, Particles and Fields*, Vol. 84, No. 11, Art. No. 1211, Nov. 2024.
- [3] C. Bambi and D. Stojkovic, “Astrophysical Wormholes,” *Universe (Basel)*, vol. 7, no. 5, p. 136, May 2021.
- [4] B. B. Hazarika, “Introducing a General Time Machine Solution and Analysis of a Vacuum Spacetime Generating Closed Timelike Curves,” *European Physical Journal Plus*, vol. 137, no. 1, Art. no. 13, Jan. 2022.
- [5] F. S. N. Lobo, “From the Flamm–Einstein–Rosen bridge to the modern renaissance of traversable wormholes,” *International Journal of Modern Physics. D, Gravitation, astrophysics, cosmology*, vol. 25, no. 7, p. 1630017, June 2016.
- [6] D. R. Mayerson, A. M. Charles, and J. E. Golec, *Relativity: A Journey Through Warped Space and Time*, 1st ed. Cham: Springer International Publishing, 2019.
- [7] P. H. R. S. Moraes, R. A. C. Correa, and R. V. Lobato, “Analytical General Solutions for Static Wormholes in $f(R, T)$ Gravity,” *Journal of Cosmology and Astroparticle Physics*, vol. 2017, no. 7, pp. 29–29, July 2017.
- [8] P. H. R. S. Moraes and P. K. Sahoo, “Modeling wormholes in $f(R, T)$ gravity,” *Physical review. D*, vol. 96, no. 4, Art. no. 044038, Aug. 2017.
- [9] M. Ringbauer, M. A. Broome, C. R. Myers, A. G. White, and T. C. Ralph, “Experimental simulation of closed timelike curves,” *Nature communications*, vol. 5, no. 1, Art. no. 4145, June 2014.

- [10] B. F. Schutz, *A first course in general relativity*, 2nd ed. Cambridge, UK; Cambridge University Press, 2009.
- [11] L. Susskind, “Wormholes and Time Travel? Not Likely,” Mar. 2005.
- [12] V. Venkatesha, N. S. Kavya, and P. K. Sahoo, “Geometric structures of Morris-Thorne wormhole metric in $f(L, m)$ gravity and energy conditions,” *Physical scripta*, vol. 98, no. 6, p. 65020, June 2023.
- [13] G. Weinstein, “The Einstein–Rosen Bridge and the Einstein–Podolsky–Rosen Argument: Singularities and Separability,” in *Thinking About Space and Time*, vol. 15, C. Wüthrich, T. Sauer, and C. Beisbart, Eds., Switzerland: *Springer International Publishing AG*, 2020, pp. 91–109.
- [14] H. Zejli, “Closed timelike curves in $f(R)$ -symmetric wormholes,” *International journal of modern physics. D, Gravitation, astrophysics, cosmology*, vol. 34, no. 11, Aug. 2025.