

MATH3195/M5195 EXERCISE SHEET 3

DUE: MARCH 11, 2024

- Problem 1.** (a) Let $R = \mathbb{Q}[[x, y]]$ and let $J = \langle xy + y^3, x + x^2y, xy + 3y, x^4 - 5y^2 + x^2y \rangle$ be an ideal in R . Show that J is minimally generated by two elements in R .
- (b) Let $R = K[t]$ and consider $M = K[t, t^{-1}]$ as R -module and let $I = tR$ be an ideal in R . Show that $M = IM$ but $M \neq 0$. Why does this example not contradict Nakayama's lemma?

Problem 2. Prove the isomorphism theorems for modules.

Problem 3. Prove the 3×3 -lemma: Let R be a ring. Assume that

$$\begin{array}{ccccccc}
 & 0 & & 0 & & 0 & \\
 & \downarrow & & \downarrow & & \downarrow & \\
 0 & \longrightarrow & A_1 & \xrightarrow{\alpha} & A_2 & \xrightarrow{\alpha'} & A_3 \longrightarrow 0 \\
 & & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 \\
 0 & \longrightarrow & B_1 & \xrightarrow{\beta} & B_2 & \xrightarrow{\beta'} & B_3 \longrightarrow 0 \\
 & & \downarrow g_1 & & \downarrow g_2 & & \downarrow g_3 \\
 0 & \longrightarrow & C_1 & \xrightarrow{\gamma} & C_2 & \xrightarrow{\gamma'} & C_3 \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

is a commutative diagram of R -modules and all columns and the middle row is exact. Show that the top row is exact if and only if the bottom row is exact.

Problem 4. (Localisation of a module) Let R be a ring and $A \subset R$ be multiplicatively closed. Let M be an R -module. Assume we know that $(m, a) \sim (n, b)$ if and only if $mbc = nac$ for some $c \in A$ defines an equivalence relation on $M \times A$. (Note: recall the definition of an equivalence relation.)

- (a) Writing $A^{-1}M$ for the set of equivalence classes of \sim , and $\frac{m}{a}$ for the class containing (m, a) , show that the operation

$$\frac{m}{a} + \frac{n}{b} = \frac{bm + an}{ab}$$

is well defined and hence that $A^{-1}M$ is an abelian group.

- (b) By defining an appropriate multiplication rule, show that $A^{-1}M$ has the structure of an $A^{-1}R$ -module.

Problem 5. Let R be a ring and $A \subset R$ be multiplicatively closed.

- (a) Suppose that $\phi : M \rightarrow N$ is a homomorphism of R modules. Show ϕ induces an $A^{-1}R$ -homomorphism $A^{-1}M \rightarrow A^{-1}N$.
- (b) Suppose $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is an exact sequence of R -modules. Show that $0 \rightarrow A^{-1}L \rightarrow A^{-1}M \rightarrow A^{-1}N \rightarrow 0$, with the induced maps from (i), is an exact sequence of $A^{-1}R$ -modules. (*Remark:* This means that localization is an exact functor from the category of R -modules to the category of $A^{-1}R$ -modules.)