

1. What is the mean of the dataset $D = \{1, 2, 3\}$?

1 / 1 punto

Do the exercises using pen and paper.

☐ 3

☐ 6

☒ 2



Correcto

That's it. Good job!

2. Compute the mean of the following dataset:

1 / 1 punto

$$D = \left\{ \begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ 8 \end{bmatrix}, \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix} \right\}$$

Do the exercises using pen and paper.

☐ $\begin{bmatrix} 6 \\ 15 \\ 24 \end{bmatrix}$

☐ $\begin{bmatrix} -2 \\ -5 \\ -8 \end{bmatrix}$

☒ $\begin{bmatrix} 2 \\ 5 \\ 8 \end{bmatrix}$



Correcto

Well done!

3. What is the mean of the following dataset, **after** multiplying each sample in the dataset by 2?

1 / 1 punto

$$D = \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}, \begin{bmatrix} 5 \\ 3 \\ 1 \end{bmatrix} \right\}$$

☐ $\begin{bmatrix} 3 \\ 3 \\ 3 \end{bmatrix}$

☐ $\begin{bmatrix} 18 \\ 18 \\ 18 \end{bmatrix}$

☒ $\begin{bmatrix} 6 \\ 6 \\ 6 \end{bmatrix}$

✓ **Correcto**
Well done!

4.

What is the mean of the following dataset, **after** adding $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ to each sample in the following dataset?

1 / 1 punto

$$D = \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}, \begin{bmatrix} 5 \\ 3 \\ 1 \end{bmatrix} \right\}$$

☒ $\begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$

☐ $\begin{bmatrix} 3 \\ 3 \\ 3 \end{bmatrix}$

☐ $\begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$

✓ **Correcto**
Well done!

5. Assuming that we know the mean \bar{x}_{n-1} of a dataset D_{n-1} with $n - 1$ data points. Now, suppose that we collect another data point, which we denote by x_* . Select the correct formula that computes the correct new mean \bar{x}_n of the full data set $D_n = D_{n-1} \cup \{x_*\}$, i.e., we add x_* to the dataset D .

1 / 1 punto

☐ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n-1}(x_* - \bar{x}_{n-1})$

☐ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n+1}(x_* - \bar{x}_{n-1})$

☐ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n+1}(\bar{x}_{n-1} - x_*)$

☒ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n}(x_* - \bar{x}_{n-1})$

✓ **Correcto**
Excellent!

6. Assuming you are given an image as a two dimensional array of shape 28 x 28. Write a small piece of python code to reshape this image to a vector of length 784 (=28 x 28).

1 / 1 punto

Hint: This can be a one-liner.

```
1 import numpy as np
2
3 def reshape(x):
4     """return x_resaped as a flattened vector of the multi-dimensional array x"""
5     x_resaped = np.reshape(x, -1)
6     return x_resaped
7
```

Ejecutar

✓ **Correcto**

Good job!