

Schur Rings Over Numerical Semigroups

Thomas Reese
Southern Utah University

February 27, 2026

1. Numerical Semigroups
2. Schur Rings Over Numerical Semigroups
3. Research Objectives and Results
4. New Paradigm

Numerical Semigroups

Definition

A Numerical Semigroup is a subset of the natural numbers which is closed under addition.

Example

- $\mathbb{Z}^+$
- $n\mathbb{Z}^+, \forall n \in \mathbb{Z}^+$
- $\{3, 5, 6, 8, 9, 10, 11, \rightarrow\}$

Properties of Numerical Semigroups

Any numerical semigroup can be reduced to a finite generating set.

Example

- $\mathbb{Z}^+ = \langle 1 \rangle$
- $n\mathbb{Z}^+ = \langle n \rangle$
- $\{3, 5, 6, 8, 9, 10, 11, \rightarrow\} = \langle 3, 5 \rangle$

It can be assumed the gcd of a generating set is 1.

Properties of Numerical Semigroups, Cont.

A numerical semigroup with at least two generators greater than 0 has a Frobenius number and a conductor.

Definition

The Frobenius number is the greatest integer not in the semigroup, e.g. $F(\langle 3, 5 \rangle) = 7$

Definition

The Conductor is the smallest element of the numerical semigroup such that all greater natural numbers are in the semigroup, e.g. $C(\langle 3, 5 \rangle) = F(\langle 3, 5 \rangle) + 1 = 8$

Note that $\langle a_0, a_1, \dots, a_n \rangle$ can be expressed as $\langle x^{a_0}, x^{a_1}, \dots, x^{a_n} \rangle$, as the algebraic structure is identical with polynomial multiplication. This notation is helpful when discussing Schur rings.

Definition

The coarsest Schur ring is the partition such that if $a, b \in S$ always have the same coefficient in $\mathbb{Z}[S]$, then a, b are combined into the same element of the partition.

Schur Rings Cont.

Let $S = \langle x^3, x^5 \rangle$.

Start by combining the generators, $x^3 + x^5$.

Then check higher orders.

$$(x^3 + x^5)^2 = x^6 + 2x^8 + x^{10}$$

There is no other way to generate x^6 or x^{10} , so they can be combined.

Similarly, there is no other way to generate x^8 , so it is a singleton.

Schur Rings Cont.

$$(x^3 + x^5)^3 = x^9 + 3x^{11} + 3x^{13} + x^{15}$$

$x^{15} \in S^5, x^9 \notin S^5$, thus they are both singletons.

x^{11}, x^{13} ? They both can only be generated by $x^8(x^3 + x^5)$, so they can be combined.

No further combining is possible.

Final Coarsest Schur Ring: $x^3 + x^5, x^6 + x^{10}, x^8, x^9, x^{11} + x^{13}, x^{12}, \rightarrow$

Finer Schur rings are possible.

Primary Research Question

Conjecture

All Schur Rings over numerical semigroups are eventually discrete.

Theorems

Theorem 1

For $S = \langle p, q \rangle$ where $p < q$ and $\gcd(p, q) = 1$, no element of S greater than $(p + q)(p - 1)$ can be combined in a Schur ring.

Theorem 2

For any numerical semigroup of the form $S = \langle m, m + 1, m + 2, \dots, 2m - 1 \rangle$, any Schur ring over S is eventually discrete.

Theorem 3

For the numerical semigroup, $S = \langle a_0, a_1, \dots, a_k \rangle$, if $\gcd(a_i, a_j) = 1$ for all $0 \leq i, j \leq k$ with $i \neq j$, then any Schur ring over S is eventually discrete.

Sketch of Proof of Theorem 1

Theorem 1

For $S = \langle p, q \rangle$ where $p < q$ and $\gcd(p, q) = 1$, no element of S greater than $(p + q)(p - 1)$ can be combined in a Schur ring.

Proof

For $s \in S$, there exists a unique maximal decomposition which contains more generators than all other decompositions.

Two elements of S can only have a maximal decomposition of the same order if they are both less than $(p + q)(p - 1)$

A similar theorem for 3 generators was recently developed.

Sketch of Proof of Theorem 2

Theorem 2

For any numerical semigroup of the form $S = \langle m, m+1, m+2, \dots, 2m-1 \rangle$, any Schur ring over S is eventually discrete.

Proof

$$\begin{aligned}(x^m + x^{m+1} + \dots + x^{2m-1})^2 &= x^{2m} + 2x^{2m+1} + \dots + bx^{3m-2} + ax^{3m-1} + bx^{3m} + \dots + x^{4m-2} \\ (x^m + x^{m+1} + \dots + x^{2m-1})^3 &= x^{3m} + \dots\end{aligned}$$

Thus $3m-1$ and $3m$ are both singletons. Note also $\gcd(3m-1, 3m) = 1$. This generates a numerical semigroup within the Schur ring, which is eventually discrete.

Sketch of Proof of Theorem 3

Theorem 3

For a numerical semigroup, $S = \langle a_0, a_1, \dots, a_k \rangle$, if $\gcd(a_i, a_j) = 1$ for all $0 \leq i, j \leq k$ with $i \neq j$, then any Schur ring over S is eventually discrete.

Proof

The case for 2 and 3 generators are covered by other theorems. For 4 or more, two coprime singletons can be guaranteed within the Schur ring.

New Paradigm

Definition

"Freshman Exponentiation" for a polynomial $X = a_0 + a_1x + \dots + a_nx^n$, denoted $X^{(m)}$ is defined as $X^{(m)} = a_0^m + (a_1x)^m + \dots + (a_nx^n)^m$.

Definition

The Hadamard product of two polynomials

$X_a = a_0 + a_1x + \dots + a_nx^n$, $X_b = b_0 + b_1x + \dots + b_mx^m$, denoted $X_a \circ X_b$ is defined as $X_a \circ X_b = \sum_{i=0}^k a_i b_i x^i$ where $k = \max(n, m)$.

A Schur ring is closed under both operations.

Two Generator Case

Let $S = \langle a, b \rangle$. Thus $x^a + x^b$ is an element of the Schur Ring over S .

Consider $(x^a + x^b)^2 = x^{2a} + 2x^{a+b} + x^{2b}$. No two terms of this polynomial can be combined, as $a \neq b$, thus x^{a+b} is a singleton in the partition for the Schur Ring.

Consider $(x^a + x^b)^{(a)} \circ (x^a + x^b)^{(b)}$.

$$\begin{aligned} & (x^a + x^b)^{(a)} \circ (x^a + x^b)^{(b)} \\ &= (x^{a^2} + x^{ab}) \circ (x^{ab} + x^{b^2}) \\ &= x^{ab} \text{ because } a \neq b \end{aligned}$$

Thus x^{ab} is a singleton in the Schur Ring.

Two Generator Case Cont.

Note $\gcd(a + b, ab) = 1$, as $\gcd(a, b) = 1$.

Therefore x^{a+b}, x^{ab} form a numerical semigroup in the Schur Ring, meaning the Schur Ring is eventually discrete.

Note the argument can be extended to 3 and 4 generators.

A modified form of this argument can be inducted to any number of generators.

Abdallah Assi, Pedro A. Garcia-Sanchez
Numerical Semigroups and Applications

The End