

Quantifying comprehensive marine vessel emissions using data fusion

emLab, UCSB and GFW

Last Updated on July 28, 2026

Table of contents

1	In-line manuscript statistics	2
1.1	Abstract	2
1.2	Introduction	2
1.3	Results	3
1.4	Discussion	4
1.5	Methods	5
2	Results	6
2.1	Figure 1	6
2.2	Figure 2	7
2.3	Figure 3	8
2.4	Figure 4	9
3	Methods	10
3.1	Figure 6	10
3.2	Table 1	12
3.3	Figure 7	12
3.4	Figure 8	13
3.5	Figure 9	14
3.6	Figure 10	15
3.7	Table 2	16
3.8	Figure 11	18
3.9	Figure 12	19
3.10	Figure 13	20
3.11	Figure 14	21
3.12	Figure 15	22

3.13	Table 3	24
3.14	Table 4	24
3.15	Figure 16	24
3.16	Figure 17	26
4	Supplementary materials: Supplementary figures	27
4.1	Figure S1	27
4.2	Figure S2	28
4.3	Figure S3	29
5	Supplementary materials: Supplementary tables	30
5.1	Table S1	30
5.2	Table S2	30
5.3	Table S3	31
5.4	Table S4	31
5.5	Table S5	31
6	Supplementary materials: Supplementary figures: Comparing our emission estimates to other inventories	32
6.1	Figure S4	32
6.2	Table S6	33
6.3	Figure S5	33
6.4	Figure S6	34
6.5	Figure S7	35

1 In-line manuscript statistics

1.1 Abstract

- We found that vessels emitted approximately 1.53 billion metric tonnes of CO₂ in 2025. In 2024, the most recent year for which EDGAR data are available, marine vessels accounted for 16% of global fossil fuel emissions from the transportation sector.
- Current AIS-only approaches underestimate the magnitude of 2025 emissions by 22%. Meanwhile, because AIS coverage has generally been increasing over time, current AIS-only approaches overestimate the increasing emissions trend by 15% (from 2017 to 2025).

1.2 Introduction

- Climate change mitigation requires accurate accounting across sectors; while the transportation sector as a whole was responsible for 21% of all global CO₂ emissions in 2024, fundamental gaps remain in understanding the contribution by marine vessels.

- For over 780,000 AIS-broadcasting vessels, we quantify emissions for each vessel using precise latitude/longitude/timestamp information for over 119 billion individual AIS messages.

1.3 Results

- Using this data fusion approach, we find that total global CO₂ emissions at sea in 2025 was 1.53 billion MT, with an increase of 38% from 2017 to 2025. If only considering emissions from vessels that broadcast AIS, in line with currently accepted state-of-the-art models, global CO₂ emissions in 2025 would be only 1.19 billion MT, with a more pronounced increase from 2017 to 2025 of 44%.
- Thus using only AIS data from broadcasting vessels underestimates current global CO₂ emissions by 22%, and overestimates the increasing trend for 2017–2025 by 15%.
- Across the entire 2017 to 2025 time series, 96.4% of non-broadcasting emissions occurred inside the S1 footprint and in pixel-months that were imaged; 3.5% occurred inside the S1 footprint but in pixel-months that weren't imaged; and 0.079% occurred outside the S1 footprint entirely.
- In 2025, 36% of CO₂ emissions from fishing vessels were from non-broadcasting vessels, and 21% of CO₂ emissions from non-fishing vessels were from non-broadcasting vessels. For both types of vessels, this fraction of emissions from non-broadcasting vessels has been declining steadily since 2020.
- Across the entire 2017 to 2025 time series, almost all non-broadcasting emissions (96.4%) occurred inside the S1 footprint and in pixel-months that were imaged by S1.
- Our machine learning model for predicting these emissions directly from observed S1 detection density achieves a high out-of-sample predictive performance (traditional R² = 0.94 and total percent error of -8.6% for CO₂).
- Only 3.5% of non-broadcasting emissions occurred inside the S1 footprint but in pixel-months that weren't imaged by S1, requiring us to use a two-stage hurdle framework using both a classification model (F1 Score = 0.72) and a regression model (traditional R² = 0.73) to predict S1 detection density before predicting emissions (traditional R² = 0.94).
- Only 0.079% of non-broadcasting emissions occurred outside the S1 footprint entirely, since most vessels operating in these open ocean areas use AIS.
- Most emissions for non-fishing vessels occur during international trips, while most emissions for fishing vessels occur during domestic trips.
- China, Japan, and United States had the most CO₂ emissions during within-country domestic trips in 2025.

- China, Malaysia, and United States had the most CO₂ emissions during international trips that started or ended in these countries in 2025 (where the emissions from each trip are partitioned equally to its departure and arrival country).
- China, United States, and Indonesia had the most CO₂ emissions during port stays.
- Of the roughly 775,000 AIS-broadcasting vessels in our analysis, only 93,400 (12%) are listed with the IMO, while 45,700 (6%) are listed on some other public registry; this leaves over 636,000 vessels (82%) that have no public registry information and must rely completely on modeled vessel characteristics (main engine power and design speed).
- Because many large vessels are already listed with the IMO, the majority of our total CO₂ emissions come from these vessels (39%); 19% come from vessels listed on other registries; and 42% come from vessels with no public registry information and which rely entirely on modeled vessel characteristics.
- Although vessel activity in the Southern Ocean represents a very small fractions of the world’s marine vessel emissions (0.04%), it represents the part of the world experiencing the highest growth rate of marine vessel emissions (267%).
- These polar estimates are primarily driven by AIS-broadcasting vessels, with only a small fraction coming from non-broadcasting vessels (1.02% in the Southern Ocean).
- Polar regions have extremely limited S1 coverage (only 0.06% of pixel-months in the Southern Ocean were imaged by S1), meaning that trends there may be more susceptible to AIS adoption biases.
- Using the latest data available from the EU’s Emissions Database for Global Atmospheric Research (EDGAR), we estimate that in 2024 marine vessels accounted for 16% of all transportation-related CO₂ emissions. Marine vessel CO₂ emissions increased by 32% between 2017 and 2024, 10 times faster than the growth rate of other transportation emissions, and 5 faster than the growth rate of total global emissions across all sectors.

1.4 Discussion

- For non-fishing vessels, the total social cost of CO₂, CH₄, and N₂O emissions in 2021 was 344 billion USD. This represents about 5% of the total value of all goods transported by maritime trade (using 6.63 trillion USD as the 2021 value of maritime-traded goods).
- For fishing vessels, the total social cost of CO₂, CH₄, and N₂O emissions in 2022 was 24 billion USD. This represents about 15% of the total value of all wild capture fisheries (using 159 billion USD as the 2022 value of wild capture fisheries).

1.5 Methods

- The time between pings is typically small (the mean value is 0.0881 hours and the median value is 0.0447 hours, or about 5 minutes and 3 minutes respectively), while the maximum value for this across our dataset is 47 hours (meaning this is the most amount of time over which we will estimate emissions for a single message).
- In all, we analyze AIS data for over 780,000 AIS-broadcasting vessels, across 119 billion individual timestamped AIS messages which are assigned to 14.4 million international trips, 112 million domestic trips, and 96.4 million port visits that occurred between 2017 and 2025.
- As a result, we used a dataset containing 31.9 million vessel detections from 2017 to 2025, of which 14.9 million detections were not matched to a broadcasting vessel ('unmatched'). These detections come from across more than 924,000 individual S1 scenes.
- We were able to use this direct estimation approach for 96.4% of the total estimated non-broadcasting CO₂ emissions .
- Of the total predicted non-broadcasting CO₂ emissions, 3.5% corresponded to indirect estimations within the S1 footprint, while only 0.079% occurred entirely outside the footprint

2 Results

2.1 Figure 1




Figure 1: (A) Annual marine vessel CO₂ emissions, with each line colored based on the data source. (B) Relative change in annual marine vessel CO₂ emissions from a 2017 baseline, with each line colored based on the data source. (C) Spatial map of CO₂ emissions in 2024 (top) and change in CO₂ emissions between 2017 and 2024 (bottom), aggregated across AIS-broadcasting vessels and non-broadcasting vessels, and shown at a 1x1 degree spatial resolution; emissions are area-normalized for each pixel (metric tonnes per square kilometer, shown on a pseudolog-10 scale).

2.2 Figure 2




Figure 2: (A) Time series of monthly global CO₂ emissions, shown for both AIS-broadcasting and S1-detected non-broadcasting vessels, and for both fishing and non-fishing vessels. (B) Spatial distribution of cumulative 2017-2024 CO₂ emissions from both AIS-broadcasting and S1-detected non-broadcasting vessels, for both fishing and non-fishing vessels, and shown at a 1x1 degree spatial resolution; emissions are area-normalized for each pixel (metric tonnes per square kilometer, shown on a pseudolog-10 scale).

2.3 Figure 3




Figure 3: Richness of AIS-broadcasting vessel emissions data, showing: (A) CO₂ emissions by vessel type for AIS-broadcasting vessels during different activity types (international trips, port stays, and domestic trips), aggregated for activities ending in 2024. (B) CO₂ emissions by country for AIS-broadcasting vessels during different activity types (international trips, port stays, and domestic trips), aggregated for activities ending in 2024. For both panels, emissions for each international trip are partitioned equally to its departure and arrival country). Countries with no values are shown in black.

2.4 Figure 4




Figure 4: (A) Trend in CO₂ emissions by ocean (normalized to 2017 baseline), use our fused AIS+S1 marine vessel data. (B) Comparison in trend of CO₂ emissions data (normalized to 2017 baseline) between marine transportation emissions from GFW, other transportation emissions from EDGAR, and all sector emissions from EDGAR. On the right-hand side of the figure, 2024 absolute values (in millions of metric tonnes) are shown as text and color-coded for each line.

3 Methods

3.1 Figure 6




Figure 5: Comparison of daily vessel-level CO_2 emissions estimated using our model with registered emissions from a proprietary vessel database (based on main engine fuel consumption over 24 hours). Estimates are shown using main engine power and design speed from IMO (left), our machine learning random forest models (center), and the same registered dataset (right). The solid red line shows a linear regression fit, while the dashed line represents the 1:1 line. Performance values for traditional R^2 are shown for each set of vessel characteristics. Over 15,000 vessels are represented in the figure, and blue shading indicates point density. All data are rule-of-three compliant, meaning each cell (20x20 mt) displayed contains at least three vessels.

Table 1: Performance summary across different thresholds for the hours at sea tolerance between datasets. Results are shown using the R^2 (measuring linear association between observed and predicted values) and traditional R^2 (proportion of variance explained compared to the mean-only baseline).

Tolerance	Number of vessels	R^2	Traditional R^2
0.05	921	0.78	0.77
0.10	1,594	0.79	0.78
0.15	2,274	0.80	0.80
0.20	2,910	0.77	0.76
0.25	3,598	0.78	0.77
0.30	4,373	0.71	0.67
0.35	5,254	0.72	0.64
0.40	6,298	0.72	0.58

3.2 Table 1

3.3 Figure 7




Figure 6: Comparison of annual vessel-level CO₂ emissions estimated using our model with published emissions from the EU MRV database. The sample includes 2,276 observations from yearly aggregated vessel trips where hours at sea differ by no more than 15% between our data and the MRV dataset. The solid red line shows the linear regression fit, and the dashed line represents the 1:1 line.

3.4 Figure 8




Figure 7: Map showing the percentage of months from 2017 through 2024 that were imaged by S1 for each 1x1 degree pixel.

3.5 Figure 9




Figure 8: Distributions showing the number of unique AIS vessels and number of S1 detections for each length size bin, disaggregated by fishing and non-fishing vessels.

3.6 Figure 10




Figure 9: Relationship between main engine power (killowatts) and length (meters) for all AIS-broadcasting vessels, disaggregated by non-fishing and fishing vessels.

3.7 Table 2

Table 2: Data sources for all included model features, including which random forest model uses the features (M1: detections classification; M2: detections regression; and/or M3: emissions regression), the units, feature type, whether it varies spatially, whether it varies temporally, and the source name and reference citation.

Model feature	Category	M1	M2	M3	Units	Type	Spatial	Temporal	Source	Ref.
Presence/absence of unmatched S1 detections	Dependent variable	Yes	No	No	-	Boolean	Yes	Yes	GFW	[?]
Unmatched S1 detections per km ² imaged	Dependent variable	No	Yes	No	detects/km ²	Numeric	Yes	Yes	GFW	[?]
Log(AIS emissions)	Dependent variable	No	No	Yes	log(MT)	Numeric	Yes	Yes	GFW	-
Vessel type (fishing or non-fishing)	Dependent variable	Yes	Yes	Yes	-	Boolean	Yes	Yes	GFW	[?]
Lower threshold of vessel length bin	S1 detection features	Yes	Yes	No	m	Numeric	No	No	GFW	[?]
Vessel length bin (4 categories)	S1 detection features	Yes	Yes	No	-	Factor	Yes	Yes	GFW	[?]
S1 detections per km ² of area for each length bin	S1 detection features	No	No	Yes	detects/km ²	Numeric	Yes	Yes	GFW	[?]
Days since start of study	Temporal features	Yes	Yes	Yes	days	Numeric	No	Yes	GFW	[?]
Post-2020 IMO Sulphur policy	Temporal features	Yes	Yes	Yes	-	Boolean	No	Yes	IMO	[?]
Ocean basin	Geographic features	Yes	Yes	Yes	-	Factor	Yes	No	GFW	[?]
Pixel water area	Geographic features	Yes	Yes	Yes	km ²	Numeric	Yes	No	GFW	[?]
Mean ocean depth	Geographic features	Yes	Yes	Yes	m	Numeric	Yes	No	GFW	[?]
Mean distance from shore	Geographic features	Yes	Yes	Yes	m	Numeric	Yes	No	GFW	[?]
Mean distance from nearest port	Geographic features	Yes	Yes	Yes	m	Numeric	Yes	No	GFW	[?]
Distance to top 20 ports	Geographic features	Yes	Yes	Yes	m	Numeric	Yes	No	Lloyd's List	[?]
Positions reported by AIS type a	AIS reception quality	Yes	Yes	Yes	-	Numeric	Yes	No	GFW	[?]
Positions reported by AIS type b	AIS reception quality	Yes	Yes	Yes	-	Numeric	Yes	No	GFW	[?]
Proportion of messages, dynamic AIS receivers	AIS reception quality	Yes	Yes	Yes	-	Numeric	Yes	Yes	GFW	[?]
Proportion of messages, terrestrial AIS receivers	AIS reception quality	Yes	Yes	Yes	-	Numeric	Yes	Yes	GFW	[?]
Proportion of messages, satellite AIS receivers	AIS reception quality	Yes	Yes	Yes	-	Numeric	Yes	Yes	GFW	[?]
Mean wind speed	Weather features	Yes	Yes	Yes	m/s	Numeric	Yes	Yes	ERA5	[?]
Mean primary wave height	Weather features	Yes	Yes	Yes	m	Numeric	Yes	Yes	ERA5	[?]
Mean primary wave period	Weather features	Yes	Yes	Yes	s	Numeric	Yes	Yes	ERA5	[?]

3.8 Figure 11




Figure 10: Monthly statistics of global S1 coverage including total number of S1 scenes, total imaged area across all S1 scenes, total number of vessel detections, total number of vessel detections that could not be matched to an AIS-broadcasting vessels, and fraction of the total number of vessel detections that could not be matched to an AIS-broadcasting vessels.

3.9 Figure 12




Figure 11: Maps of pixels that were used in the training and testing datasets for the simulated outside S1 footprint performance tests for each model. Pixels $\leq 100\text{km}$ from the shore were used for training; pixels $> 100\text{km}$ from shore were used for testing.

3.10 Figure 13




Figure 12: Precision-recall curves of the detection presence/absence classification model, shown for both the inside S1 footprint test and simulated outside S1 footprint test.

3.11 Figure 14




Figure 13: Receiver Operating Characteristic (ROC) curves of the detection classification model, shown for both the inside S1 footprint test and simulated outside S1 footprint test.

3.12 Figure 15




Figure 14: Confusion matrices of the detection classification model, shown for both the inside S1 footprint test and simulated outside S1 footprint test.

Table 3: Summary of dark fleet model performance metrics for the: 1) emissions regression model; 2) detections classification mode; and 3) detections regression model. Performance estimates for each metric are shown for both the inside S1 footprint and outside S1 footprint tests. All metrics are calculated at the unit of observation except for 'Total percent error', which is the percentage error in the total sum of all predictions relative to the total sum of all truth values.

Model	Pollutant	Metric	Inside footprint	Outside footprint
Emissions regression	CO2	Traditional R ²	0.94	0.73
Emissions regression	CH4	Traditional R ²	0.90	0.63
Emissions regression	N2O	Traditional R ²	0.94	0.71
Emissions regression	CO	Traditional R ²	0.92	0.63
Emissions regression	NOX	Traditional R ²	0.87	0.57
Emissions regression	SOX	Traditional R ²	0.95	0.73
Emissions regression	PM2.5	Traditional R ²	0.94	0.65
Emissions regression	PM10	Traditional R ²	0.94	0.65
Emissions regression	VOCs	Traditional R ²	0.88	0.60
Emissions regression	CO2	Total percent error	-8.61	-10.08
Emissions regression	CH4	Total percent error	-20.26	-30.01
Emissions regression	N2O	Total percent error	-10.70	-15.51
Emissions regression	CO	Total percent error	-19.10	-29.29
Emissions regression	NOX	Total percent error	-23.68	-36.44
Emissions regression	SOX	Total percent error	-8.23	-9.01
Emissions regression	PM2.5	Total percent error	-16.21	-25.52
Emissions regression	PM10	Total percent error	-16.21	-25.52
Emissions regression	VOCs	Total percent error	-22.23	-32.45
Detections classification	-	Accuracy	0.92	0.95
Detections classification	-	F1 score	0.72	0.48
Detections classification	-	Precision	0.72	0.52
Detections classification	-	Precision-recall AUC	0.81	0.49
Detections classification	-	ROC AUC	0.95	0.92
Detections classification	-	Sensitivity	0.73	0.44
Detections classification	-	Specificity	0.96	0.98
Detections classification	-	recall	0.73	0.44
Detections regression	-	Traditional R ²	0.73	0.46
Detections regression	-	Total percent error	1.18	28.46


Table 4: Linear regression fit statistics for the final models trained to estimate emissions of non-CO₂ gases from CO₂ emissions. The model for each gas has 2 fit terms.

Pollutant	Term	Coefficient estimate	Standard error	T-statistic	p.value
CH4	emissionsco2mt:log(meandistancefromshorem + 1)	0.000	3.29e-09	379.875	0
CH4	emissionsco2mt:post2020FALSE:fishingFALSE	0.000	3.17e-08	-38.042	0
CH4	emissionsco2mt:post2020TRUE:fishingFALSE	0.000	3.14e-08	-61.349	0
CH4	emissionsco2mt:post2020FALSE:fishingTRUE	0.000	8.42e-08	43.904	0
CH4	emissionsco2mt:post2020TRUE:fishingTRUE	0.000	4.52e-08	82.642	0
N2O	emissionsco2mt:log(meandistancefromshorem + 1)	0.000	2.32e-09	307.957	0
N2O	emissionsco2mt:post2020FALSE:fishingFALSE	0.000	2.23e-08	1869.423	0
N2O	emissionsco2mt:post2020TRUE:fishingFALSE	0.000	2.21e-08	1865.728	0
N2O	emissionsco2mt:post2020FALSE:fishingTRUE	0.000	5.92e-08	727.041	0
N2O	emissionsco2mt:post2020TRUE:fishingTRUE	0.000	3.18e-08	1379.919	0
CO	emissionsco2mt:log(meandistancefromshorem + 1)	0.000	1.57e-07	382.287	0
CO	emissionsco2mt:post2020FALSE:fishingFALSE	0.000	1.51e-06	19.236	0
CO	emissionsco2mt:post2020TRUE:fishingFALSE	0.000	1.50e-06	-3.662	0
CO	emissionsco2mt:post2020FALSE:fishingTRUE	0.000	4.02e-06	67.869	0
CO	emissionsco2mt:post2020TRUE:fishingTRUE	0.000	2.16e-06	128.069	0
NOX	emissionsco2mt:log(meandistancefromshorem + 1)	0.002	4.33e-06	466.681	0
NOX	emissionsco2mt:post2020FALSE:fishingFALSE	-0.007	4.00e-05	-176.274	0
NOX	emissionsco2mt:post2020TRUE:fishingFALSE	-0.008	4.00e-05	-205.243	0
NOX	emissionsco2mt:post2020FALSE:fishingTRUE	-0.001	1.10e-04	-8.817	0
NOX	emissionsco2mt:post2020TRUE:fishingTRUE	-0.002	6.00e-05	-36.129	0
SOX	emissionsco2mt:log(meandistancefromshorem + 1)	0.000	7.15e-08	-173.062	0
SOX	emissionsco2mt:post2020FALSE:fishingFALSE	0.006	6.88e-07	9164.189	0
SOX	emissionsco2mt:post2020TRUE:fishingFALSE	0.002	6.81e-07	2441.103	0
SOX	emissionsco2mt:post2020FALSE:fishingTRUE	0.006	1.83e-06	3486.129	0
SOX	emissionsco2mt:post2020TRUE:fishingTRUE	0.002	9.81e-07	1705.575	0
PM2.5	emissionsco2mt:log(meandistancefromshorem + 1)	0.000	9.11e-08	400.992	0
PM2.5	emissionsco2mt:post2020FALSE:fishingFALSE	0.000	8.76e-07	472.536	0
PM2.5	emissionsco2mt:post2020TRUE:fishingFALSE	0.000	8.67e-07	58.147	0
PM2.5	emissionsco2mt:post2020FALSE:fishingTRUE	0.001	2.33e-06	228.924	0
PM2.5	emissionsco2mt:post2020TRUE:fishingTRUE	0.000	1.25e-06	168.405	0
PM10	emissionsco2mt:log(meandistancefromshorem + 1)	0.000	9.90e-08	400.992	0
PM10	emissionsco2mt:post2020FALSE:fishingFALSE	0.000	9.52e-07	472.536	0
PM10	emissionsco2mt:post2020TRUE:fishingFALSE	0.000	9.43e-07	58.147	0
PM10	emissionsco2mt:post2020FALSE:fishingTRUE	0.001	2.53e-06	228.924	0
PM10	emissionsco2mt:post2020TRUE:fishingTRUE	0.000	1.36e-06	168.405	0
VOCs	emissionsco2mt:log(meandistancefromshorem + 1)	0.000	2.02e-07	495.650	0
VOCs	emissionsco2mt:post2020FALSE:fishingFALSE	0.000	1.94e-06	-203.024	0
VOCs	emissionsco2mt:post2020TRUE:fishingFALSE	0.000	1.92e-06	-227.991	0
VOCs	emissionsco2mt:post2020FALSE:fishingTRUE	0.000	5.15e-06	-39.044	0
VOCs	emissionsco2mt:post2020TRUE:fishingTRUE	0.000	2.77e-06	-75.686	0

3.13 Table 3

3.14 Table 4

3.15 Figure 16



3.16 Figure 17




Figure 16: Spatial coverage of the non-broadcasting emissions model. Areas are categorized according to whether they fall inside or outside the S1 footprint and whether they have been imaged by S1 at any point in time. A given pixel may be imaged in some months but not in others. Pixels outside the footprint have never been imaged. The percentage of total CO₂ emissions from non-broadcasting vessels attributed to each category is shown.

4 Supplementary materials: Supplementary figures

4.1 Figure S1




Figure 17: Spatial maps 2024 emissions for each GHG and pollutant, aggregated across AIS-broadcasting vessels and non-broadcasting vessels, and shown at a 1x1 degree spatial resolution. Emissions are area-normalized for each pixel (metric tonnes per square kilometer), and shown using a log-10 scale.

4.2 Figure S2




Figure 18: Time series of monthly global emissions for each GHG and pollutant, aggregated across AIS-broadcasting vessels and non-broadcasting vessels for which we only have Sentinel-1 data.

4.3 Figure S3




Figure 19: Annual total CO₂ emissions by ocean, disaggregated by data source (AIS-broadcasting vessels, and non-broadcasting vessels detected by S1).

5 Supplementary materials: Supplementary tables

5.1 Table S1

Table 5: Summary of 2017 and 2025 emissions, data source and pollutant.

Source	Pollutant	2017 emissions, MT	2025 emissions, MT	Percent 2025 total	2017 to 2025 change
AIS	CO2	8.24e+08	1.19e+09	78%	44%
S1	CO2	2.83e+08	3.43e+08	22%	21%
AIS + S1	CO2	1.11e+09	1.53e+09	100%	38%
AIS	CH4	11,500	15,800	80%	37%
S1	CH4	3,420	3,990	20%	17%
AIS + S1	CH4	14,900	19,700	100%	33%
AIS	N2O	41,800	59,500	78%	42%
S1	N2O	13,900	16,800	22%	21%
AIS + S1	N2O	55,700	76,300	100%	37%
AIS	CO	628,000	865,000	80%	38%
S1	CO	189,000	222,000	20%	17%
AIS + S1	CO	817,000	1,090,000	100%	33%
AIS	NOX	14,100,000	18,500,000	80%	31%
S1	NOX	3,960,000	4,490,000	20%	14%
AIS + S1	NOX	18,100,000	2.3e+07	100%	27%
AIS	SOX	5,030,000	1,830,000	78%	-64%
S1	SOX	1,750,000	527,000	22%	-70%
AIS + S1	SOX	6,770,000	2,360,000	100%	-65%
AIS	PM2.5	701,000	590,000	79%	-16%
S1	PM2.5	226,000	153,000	21%	-32%
AIS + S1	PM2.5	927,000	743,000	100%	-20%
AIS	PM10	762,000	641,000	79%	-16%
S1	PM10	246,000	166,000	21%	-32%
AIS + S1	PM10	1,010,000	807,000	100%	-20%
AIS	VOCs	641,000	868,000	80%	35%
S1	VOCs	185,000	213,000	20%	15%
AIS + S1	VOCs	825,000	1,080,000	100%	31%

5.2 Table S2

Table 6: Underestimation of 2025 emissions using only AIS data, and overestimation of 2017 to 2025 emissions trend using only AIS data, by pollutant.

Pollutant	Underestimation of 2025 emissions	Overestimation of 2017-2025 emissions trend
CO2	22%	15%
CH4	20%	15%
N2O	22%	15%
CO	20%	14%
NOX	20%	14%
SOX	22%	-2%
PM2.5	21%	-20%
PM10	21%	-20%
VOCs	20%	15%

Table 7: For each ocean, summary of: 2025 total (AIS+S1) CO₂ emissions (metric tonnes); percentage of global total (AIS+S1) 2025 CO₂ emissions from across all oceans; percent of 2025 total (AIS+S1) CO₂ emissions by non-broadcasting vessels; percent change in total (AIS+S1) CO₂ emissions from 2017 to 2025; and the percentage of all observations in the dataset (pixel-months) from 2017-2025 that were imaged by S1.

Summary statistic	Atlantic	Southern	Pacific	Indian	Arctic
2025 total CO2 emissions (million MT)	449	0.567	457	222	4.13
Percent of total CO2 2025 emissions	39.37%	0.04%	41.28%	18.93%	0.37%
Percent of total CO2 emissions by non-broadcasting	22.64%	1.02%	24.81%	20.58%	24.50%
Percent change in total CO2 emissions (2017 to 2025)	34.31%	267.19%	48.27%	32.49%	3.74%
Percent observations imaged by S1 (across 2017-2025)	35.94%	0.06%	17.74%	16.99%	13.55%

5.3 Table S3

5.4 Table S4

Table 8: For each data source, summary of: total 2025 CO₂ emissions (metric tonnes); and percent change in CO₂ emissions from 2017 to 2025.

Data source	2024 emissions, billion MT	Change in emissions, 2017 to 2024
Other transportation sectors (EDGAR)	7.44	3.35%
All sectors (EDGAR)	39.6	7.06%
Marine vessels (AIS + S1)	1.47	32.49%

5.5 Table S5

Table 9: For both non-fishing vessels and fishing vessels, we summarize: the total emissions (metric tonnes) of each GHG gas; the social cost associated with each GHG (USD/metric tonne); the social cost of each (USD); the total social cost summed across all three gasses (USD). We use 2021 for non-fishing vessels since it is the year for which we have the total value of all traded goods at sea. We use 2022 for fishing vessels since it is the year for which we have the total value of wild capture fisheries.

Year	Vessel type	GHG	Emissions (MT)	SC (USD/MT)	SC (billion USD)
2021	Non-fishing	CH4	15,296	2,900	0.044
2021	Non-fishing	CO2	1,205,568,569	283	341.176
2021	Non-fishing	N2O	60,152	49,600	2.984
Total	Non-fishing	-	-	-	344.204
2022	Fishing	CH4	1,373	2,900	0.004
2022	Fishing	CO2	82,662,425	283	23.393
2022	Fishing	N2O	4,211	49,600	0.209
Total	Fishing	-	-	-	23.606

6 Supplementary materials: Supplementary figures: Comparing our emission estimates to other inventories

6.1 Figure S4




Figure 20: Comparison of annual marine vessel CO₂ emissions inventory estimates.

Table 10: Comparison of annual marine vessel CO₂ emissions inventory estimates (billions of metric tonnes).

Year	GFW (AIS + S1)	GFW (AIS)	EDGAR	OECD	IMO
2017	1.11	0.824	0.881	-	1.06
2018	1.14	0.861	0.887	-	1.06
2019	1.17	0.884	0.875	0.887	-
2020	1.2	0.879	0.798	0.852	-
2021	1.27	0.941	0.833	0.891	-
2022	1.34	1	0.837	0.909	-
2023	1.38	1.05	0.89	0.917	-
2024	1.47	1.14	0.86	0.97	-
2025	1.53	1.19	-	-	-

6.2 Table S6

6.3 Figure S5




Figure 21: Annual total CO₂ emissions (metric tonnes) by AIS-broadcasting vessels, AIS pings (i.e., individual vessel timestamped location messages), and number of unique AIS-broadcasting vessels. Trends are disaggregated by AIS receiver type (satellite, terrestrial, and dynamic), and also shown as a total aggregated across receiver types. A dashed vertical line is shown for 2022, the first year in which dynamic AIS was widely adopted.

6.4 Figure S6




Figure 22: Annual total CO₂ emissions (metric tonnes) by AIS-broadcasting vessels, disaggregated by AIS receiver type (satellite, terrestrial, and dynamic) and total aggregated across receiver types, for the top 10 flags with the most emissions in 2024. A dashed vertical line is shown for 2022, the first year in which dynamic AIS was widely adopted.

6.5 Figure S7




Figure 23: Spatial change in CO₂ emissions by AIS-broadcasting vessels between 2017 and 2024 (metric tonnes) by receiver type (satellite, terrestrial, and dynamic) and total across receiver types, at a 1x1 degree spatial resolution; emissions are area-normalized for each pixel (metric tonnes per square kilometer, shown on a pseudolog-10 scale).