

# **Toward sustainable and resilient fisheries management for the Humboldt squid deep-water fishery**

emLab, UCSB

Last Updated on September 25, 2024

# Table of contents

<b>Introduction</b>	<b>3</b>
Background and Motivation . . . . .	3
Approach . . . . .	4
Deliverables . . . . .	5
<b>1 Methods</b>	<b>6</b>
1.1 Data sources . . . . .	6
1.1.1 AIS-based Fishing effort . . . . .	6
1.1.2 Vessel info . . . . .	6
1.1.3 Sea surface temperature (SST) . . . . .	7
1.1.4 Night light detections using VIIRS (Visible Infrared Imaging Radiometer Suite) . . . . .	9
1.1.5 Joined dataset . . . . .	10
<b>2 Results</b>	<b>13</b>
2.1 Fishery summary . . . . .	13
2.1.1 Fleet description . . . . .	13
2.2 Fishing effort . . . . .	15
2.2.1 VIIRS detections . . . . .	25
<b>References</b>	<b>27</b>

# Introduction

## Background and Motivation

The Humboldt squid (*Dosidicus gigas*) is one of the most important deep-water fisheries (DWF) globally, playing a crucial role in both the economy and food security. These extraordinarily large squid are particularly sensitive to environmental changes, including fluctuations in sea surface temperature (SST), driven by the El Niño-Southern Oscillation (“ENSO”), that climate change increasingly intensifies. SST plays a pivotal role in determining where squid populations thrive, influencing their migration patterns and spawning grounds (@yu2018ocean). As these temperatures shift, so do the squid, leading to changes in fishing grounds and market demand (Powell, Levine, and Ordonez-Gauger (2022)). Understanding these dynamics is essential for sustainable fisheries management in an era of rapid climate change.

The urgency of this issue is highlighted by recent trends. From 2017 to 2020, the global squid fleet’s fishing effort increased by 68%, with most activity concentrated beyond a nation’s Exclusive Economic Zone (EEZ) particularly from international vessels (Seto et al. (2023)). This has led to significant geopolitical and environmental concerns, particularly around South America where the Humboldt squid is abundant (Montecalvo et al. (2023)). Although some regional fisheries management organizations, such as the South Pacific Regional Fisheries Management Organization (SPRFMO), have attempted to introduce regulations to moderate squid fishing, most squid stocks on the high seas remain largely unregulated. This lack of regulation, coupled with the sensitivity of squid populations to environmental changes, underscores the need for comprehensive research and effective management strategies.

To address these challenges and seize opportunities for sustainable management, it is crucial to understand how sea surface temperature (SST) changes will affect the Humboldt squid. While some research has begun to explore the impact of sea surface temperature (SST) on the Humboldt squid, few studies have connected these findings to policy implications (Yu and Chen (2018)). We propose to investigate the sensitivity of this fishery to climate change, with a particular focus on the implications for policy interventions. Our goal is to provide policymakers with the necessary data and insights to make informed, sustainable management decisions. Specifically, we will address the following research questions:

- How do fluctuations in sea surface temperature influence the distribution and abundance of Humboldt squid?
- What are the subsequent shifts in fishing demand and potential conflicts among fisheries?

- What specific challenges and opportunities for management enhancement exist in light of changing SST?
- What policy recommendations and fisheries management strategies can be developed to manage Humboldt squid fisheries sustainably in the context of climate change?
- How the relevant RFMO and DWF fleets or adjacent countries can collectively contribute to the sustainable management of jumbo flying squid in the context of climate change?

## **Approach**

The project will be conducted in three phases:

### **Phase One: Literature review**

We will begin with a review of scientific literature and existing studies on biological and ecological characteristics of Humboldt squid, including their life cycles, breeding patterns, and feeding behaviors. We will also compile information on how changes in SST have historically affected squid populations and identify key indicators of climate resilience. In addition, we will review socioeconomic studies that highlight the economic importance of Humboldt squid fisheries, their role in local economies, and the impact of environmental changes on these aspects.

### **Phase Two: Structural model development**

In the second phase, we will develop a data-driven fishery model to simulate future SST scenarios and predict Humboldt squid responses. This model will integrate historical and current data from Global Fishing Watch and SPRFMO, as well as bioeconomic parameters in the existing study of this fishery. We will use this model to analyze the migration patterns of these squid in response to changing SST to predict shifts in fishing demand and the potential for conflict among fisheries.

### **Phase Three: Policy review and recommendations**

Finally, we will assess various fisheries management strategies specific to individual country and regions, such as voluntary seasonal closures and enforced fishing moratoriums, to identify their effectiveness in addressing the challenges posed by climate change and overfishing. This will help us develop evidence-based policy recommendations aimed at ensuring the future profitability, resiliency, and sustainability of the Humboldt squid fishery.

## **Deliverables**

- **Kickoff meeting:** Initial meeting with EDF to align on project priorities.
- **Literature review:** In-depth review of existing knowledge on Humboldt squid ecology, distribution, and response to SST fluctuations.
- **Model development:** A model predicting Humboldt squid responses to various climate change scenarios, integrating empirical data on fishing effort and catch.
- **Draft white paper:** A report on the impact of climate change on Humboldt squid fisheries, including policy recommendations supporting the sustainable management of the sector.

# 1 Methods

## 1.1 Data sources

### 1.1.1 AIS-based Fishing effort

We use satellite-based individual vessel monitoring AIS data processed by Global Fishing Watch (Kroodsma et al. (2018)). We use the V3 pipeline table `pipe_ais_v3_published.messages`. Variables of interest within this table include the following (descriptions are taken directly from the schema for `pipe_ais_v3_published.messages`):

- `ssvid`: source specific vessel id; MMSI for AIS
- `hours`: time since the previous position in the segment
- `timestamp`: timestamp for position
- `lon`: longitude
- `lat`: latitude
- `night_loitering`: 1 if the `seg_id` of every message of a `squid_jigger` that is at night and not moving, 0 if not.

In order to minimize noisy data, we only include AIS messages that have a `clean_segs` boolean (i.e., all messages must have `good_seg` boolean and must not have an `overlapping_and_short` boolean). We filter to just those messages where `night_loitering` = 1. For squid jigging vessels, GFW uses the heuristic of night loitering to identify when they are fishing. Therefore, any `hours` where `night_loitering` = 1 can be classified as `fishing_hours`

We take the raw high-resolution AIS data and aggregate `fishing_hours` spatially (by 0.5x0.5 degree pixels, which are roughly 55.5km x 55.5km at the equator), temporally by month, and by flag. We currently process data from 2016-01-01 through 2024-08-31.

### 1.1.2 Vessel info

Vessel characteristics data processed are by Global Fishing Watch (Park et al. (2023)). We use the V3 pipeline table `pipe_ais_v3_published.vi_ssvid_v20240601`. Variables of interest within this table include the following (descriptions are taken directly from the schema for `pipe_ais_v3_published.vi_ssvid_v20240301`):

- `ssvid`: source specific vessel id; MMSI for AIS

- `best.flag`: best flag state (ISO3) for the vessel
- `best.best_vessel_class`: best vessel class for the vessel (using official registry information where available, or the GFW vessel characteristics algorithm where not available)
- `best.best_engine_power_kw`: best engine power (kilowatts) for the vessel (using official registry information where available, or the GFW characteristics algorithm where not available)
- `activity.active_hours`: hours the vessel was broadcasting AIS and moving more than 0.1 knots
- `activity.offsetting`: true if this vessel has been seen with an offset position at some point between 2012 and 2019
- `activity.overlap_hours_multinames`: the total numbers of hours of overlap between two segments where, over the time period of the two segments that overlap (including the non-overlapping time of the segments), the vessel broadcast two or more normalized name, where each normalized name was broadcast at least 10 or more times. That is a bit complicated, but the goal is to identify overlapping segments where there were likely more than one identity. (this should be 0; if it is > 0, it can be used as a filter to remove potentially erroneous/noisy vessels)

We filter to just those vessels where `best.best_vessel_class = squid_jigger`. Additionally, to reduce noise, we filter out vessels that broadcast exceedingly infrequently (i.e., `activity.active_hours < 24`) or are noisy/spoofing/offsetting vessels (i.e., NOT `activity.offsetting OR activity.overlap_hours_multinames > 0.`) They are simply not reliable and will not provide good effort estimates. This leaves us with 1,561 squid vessels for our analysis.

### 1.1.3 Sea surface temperature (SST)

Sea surface temperature (SST) data come from [NOAA's Optimum Interpolation Sea Surface Temperature \(OISST\) version 2.1](#) (Huang et al. (2021)), which were downloaded from their Coast Watch ERDDAP server. The raw data are provided globally at 0.25x0.25 degree daily resolution. We aggregate the data to 0.5x0.5 degree monthly resolution by calculating the mean, standard deviation, minimum, and maximum SST for each 0.5x0.5 degree pixel and month.

We can look at a map of SST, using August 2024 as an example (Figure 1.1).




Figure 1.1: Map of mean sea surface temperature in August 2024, using 0.5x0.5 degree pixels.

Aggregating across the mean sea surface temperatures of each pixel, we can calculate the mean global sea surface temperature over time (Figure 1.2). This allows us to see both seasonal trends, and what appears to be a generally increasing trend over time.




Figure 1.2: Time series of global mean sea surface temperature.

#### 1.1.4 Night light detections using VIIRS (Visible Infrared Imaging Radiometer Suite)

We use the NOAA VIIRS (Visible Infrared Imaging Radiometer Suite) data product to detect night light emissions from vessels. When the appropriate radiance threshold is applied, these can generally be assumed to represent light-luring squid vessels. Since use only the single most accurate VIIRS detection measurements each day, these can also be thought of as representing a fishing effort metric of vessel-days. We use the VIIRS data as an alternative fishing effort metric to AIS-based fishing effort since AIS is not used on all vessels, and since it can also be disabled on the vessels that do use it.

We leverage the VIIRS boat detection dataset developed by Elvidge et al. (2015). We then apply similar processing as Seto et al. (2023), which includes the following steps:

- Reduce false detections near South America caused by the the South Atlantic Anomaly (an abundance of high-energy particles in the atmosphere).
- Use a same radiance threshold of  $10 \text{ nW cm}^{-2} \text{ sr}^{-1}$  used by Seto et al. (2023) (and established by Park et al. (2020)) in order to filter detections to those likely engaged in pelagic light-luring activity.

- To eliminate double-counting when there may be multiple satellite overpasses on a single night, only count detections from the overpass with the smallest satellite zenith angle (smaller zenith angles are more accurate)

We then finally aggregate the total detections for each 0.5x0.5 degree pixel and month.

We can compare AIS-based fishing effort with VIIRS detections by aggregating AIS effort across flags for each month and pixel, and then matching this with VIIRS detections by month and pixel. There appears to be a generally strong relationship between the two, although there are many outliers that appear to have very low AIS-based fishing effort but actually have a very high number of VIIRS detections.




Figure 1.3: Comparison of AIS-based fishing effort (x-axis) and VIIRS detections (y-axis) for each pixel and month.

### 1.1.5 Joined dataset

The final dataset we use for our analysis is a combination of the gridded AIS-based fishing effort data, the gridded SST data, and the gridded VIIRS detections. We inner join the AIS-based effort and SST datasets by 0.5x0.5 degree pixel and month, since both datasets cover the entire time series. We then left join the VIIRS detection dataset by pixel and month, since this dataset only covers January 2017 through December 2021. Note that the joined dataset does not contain flag-pixel-months with zero AIS-based fishing effort (i.e., the data are conditional

on there being some effort for any given flag-pixel-month). If desired, once we have a spatial scope with which to restrict the analysis, we could construct a dataset for that bounding box that includes zero fishing effort flag-pixel-month rows.

The joined dataset can be loaded in R using the command `targets::tar_load(joined_dataset)`. The dataset has the following columns:

- **month**: Month (first day of month) (date)
- **lon\_bin**: 0.5 degree longitude bin (degrees) (numeric)
- **lat\_bin**: 0.5 degree latitude bin (degrees) (numeric) \* **flag**: Fishing flag (character)
- **mean\_sst**: Mean sea surface temperature, aggregated from the raw daily 0.25x0.25 degree data (degrees C) (numeric)
- **sd\_sst**: Standard deviation of sea surface temperature, aggregated from the raw daily 0.25x0.25 degree data (degrees C) (numeric)
- **min\_sst**: Minimum sea surface temperature (degrees C), aggregated from the raw daily 0.25x0.25 degree data (numeric)
- **max\_sst**: Maximum sea surface temperature (degrees C), aggregated from the raw daily 0.25x0.25 degree data (numeric)
- **fishing\_hours**: Total fishing effort across vessels (hours) (numeric)
- **fishing\_kw\_hours**: Total fishing effort across vessels (kW-hours) (numeric)
- **viirs\_detections**: Night light detections from the VIIRS dataset (numeric)

Here we summarize these data (Table 1.1):

Table 1.1: Summary statistics for joined dataset that includes, gridded SST, AIS-based fishing effort, and VIIRS detections

(a) Data summary

Name	joined_dataset
Number of rows	1610486
Number of columns	11
Column type frequency:	
character	1
Date	1
numeric	9
Group variables	None

**Variable type: character**

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
flag	179745	0.89	3	3	0	45	0

**Variable type: Date**

skim_variable	n_missing	complete_rate	min	max	median	n_unique
month	0	1	2016-01-01	2024-08-01	2021-02-01	104

**Variable type: numeric**

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
mean_sst	0	1.00	20.04	6.81	-1.65	14.31	20.41	26.08	35.46	
sd_sst	302	1.00	0.17	0.19	0.00	0.05	0.11	0.22	0.45	
min_sst	0	1.00	19.85	6.90	-1.68	13.95	20.24	25.98	35.45	
max_sst	0	1.00	20.23	6.73	-1.63	14.66	20.59	26.17	35.51	
lon_bin	0	1.00	16.89	110.83	-	-	-0.50	132.50	179.50	
					180.00	82.00				
lat_bin	0	1.00	4.22	30.56	-	-	0.50	36.00	80.50	
					54.50	16.50				
fishing_hours	0	1.00	134.28	456.21	0.00	8.03	20.90	77.28	21005.80	
fishing_kw_hours	0	1.00	164088.1636584.244.73	8351.4823717.340379.8031939325.02						
viirs_detections	0.41	79.22	209.98	1.00	4.00	18.00	69.00	4608.00		

## 2 Results

### 2.1 Fishery summary

#### 2.1.1 Fleet description

Our analysis includes 1,561 squid vessels spread across 52 flags. For each flag, we look at the total engine power (kW) across the fleet and total number of unique vessels and (Figure 2.1). China dominates the fleet, with Taiwan a fairly distant second.



Figure 2.1: Squid jigger vessel summary by flag, showing the total engine power (kW) and total number of unique vessels.

Next we look at total global fishing effort from 2016 through August 2024, by flag (Figure 2.2).

The top 10 flags are shown individually, with all other flags aggregated into “Other”. Again, China dominates the fishing effort, with Taiwan a fairly distant second.




Figure 2.2: Total fishing effort, by flag, from 2016 through August 2024. The top 10 flags are shown, with all other flags aggregated into “Other”.

## 2.2 Fishing effort

Next we look at a time series of total monthly global effort by fishing flag over time (Figure 2.3). The top top 10 flags are shown individually, with all other flags aggregated into “Other”.




Figure 2.3: Monthly fishing effort by flag from 2016 through August 2024. The top 10 flags are shown individually, with all other flags aggregated into “Other”.

Next we can look at a map of squid fishing effort (Figure 2.4), aggregating effort across

effort and flags and time for each pixel across the entire processed time series (2016-01-01 through 2024-08-31). EEZ boundaries from [Marine Regions v12](#) are shown in orange (Institute (2023)).








Figure 2.4: Map of squid jigger fishing effort from 2016 through August 2024, aggregating effort across effort, flags, and time for each pixel. EEZ boundaries from Marine Regions V12 are shown in orange.

We can also look at these maps, broken apart by flag, and still aggregating squid jigger fishing effort from 2016 through August 2024, aggregating effort across time for each pixel. The top 10 flags are shown individually, with all other flags aggregated into “Other”.


```
[[1]]
```




[[2]]




[[3]]




Fishing effort (kW·h·hours)




[[4]]





[[5]]





[[6]]




[[7]]




[[8]]





[[9]]





Fishing effort (kwW-hours)



[[10]]



[[11]]



### 2.2.1 VIIRS detections

First we look at a time series of monthly VIIRS detections globally (Figure 2.5). These can generally be thought of as representing light-luring squid vessels. Again we can see a clear seasonal pattern.




Figure 2.5: Monthly VIIRS detections.

Next we can look at a map of these VIIRS detections (Figure 2.6), aggregating across time for each pixel across the entire processed time series (January 2017 through December 2021). The general spatial patterns seem to be consistent with those seen using AIS-based fishing effort data (Figure 2.4).




Figure 2.6: Map of VIIRS detections, aggregating across time for each pixel. EEZ boundaries from Marine Regions V12 are shown in orange.

## References

- Elvidge, CD, M Zhizhin, K Baugh, and FC Hsu. 2015. "Automatic Boat Identification System for VIIRS Low Light Imaging Data," *Remote Sens.*, 7, 3020–3036."
- Huang, Boyin, Chunying Liu, Viva Banzon, Eric Freeman, Garrett Graham, Bill Hankins, Tom Smith, and Huai-Min Zhang. 2021. "Improvements of the Daily Optimum Interpolation Sea Surface Temperature (DOISST) Version 2.1." *Journal of Climate* 34 (8): 2923–39.
- Institute, Flanders Marine. 2023. "Maritime Boundaries Geodatabase, Version 12." Available Online at <Https://Www.marineregions.org/>. <Https://Doi.org/10.14284/628>.
- Kroodsma, David A, Juan Mayorga, Timothy Hochberg, Nathan A Miller, Kristina Boerder, Francesco Ferretti, Alex Wilson, et al. 2018. "Tracking the Global Footprint of Fisheries." *Science* 359 (6378): 904–8.
- Montecalvo, Isabella, Philippe Le Billon, Chris Arsenault, and Milko Schwartzman. 2023. "Ocean Predators: Squids, Chinese Fleets and the Geopolitics of High Seas Fishing." *Marine Policy* 152: 105584.
- Park, Jaeyoon, Jungsam Lee, Katherine Seto, Timothy Hochberg, Brian A Wong, Nathan A Miller, Kenji Takasaki, et al. 2020. "Illuminating Dark Fishing Fleets in North Korea." *Science Advances* 6 (30): eabb1197.
- Park, Jaeyoon, Jennifer Van Osdel, Joanna Turner, Courtney M Farthing, Nathan A Miller, Hannah L Linder, Guillermo Ortúño Crespo, Gabrielle Carmine, and David A Kroodsma. 2023. "Tracking Elusive and Shifting Identities of the Global Fishing Fleet." *Science Advances* 9 (3): eabp8200.
- Powell, Farrah, Arielle Levine, and Lucia Ordóñez-Gauger. 2022. "Climate Adaptation in the Market Squid Fishery: Fishermen Responses to Past Variability Associated with El Niño Southern Oscillation Cycles Inform Our Understanding of Adaptive Capacity in the Face of Future Climate Change." *Climatic Change* 173 (1): 1.
- Seto, Katherine L, Nathan A Miller, David Kroodsma, Quentin Hanich, Masanori Miyahara, Rui Saito, Kristina Boerder, Masaki Tsuda, Yoshioki Oozeki, and Osvaldo Urrutia S. 2023. "Fishing Through the Cracks: The Unregulated Nature of Global Squid Fisheries." *Science Advances* 9 (10): eadd8125.
- Yu, Wei, and Xinjun Chen. 2018. "Ocean Warming-Induced Range-Shifting of Potential Habitat for Jumbo Flying Squid *Dosidicus gigas* in the Southeast Pacific Ocean Off Peru." *Fisheries Research* 204: 137–46.