Displacement of fishing effort by Large Scale Marine Protected Areas

Updated on 2018-10-12

Juan Carlos Villaseñor-Derbez[1] John Lynham[2]

[1]*Bren School of Environmental Science and Management, University of California Santa Barbara, Santa Barbara, CA*

[2]*Department of Economics, University of Hawaii at Manoa, Honolulu, HI*

Abstract

Large-scale Marine Protected Areas (LSMPAs) have seen a significant increase over the last years. Fishing effort is effectively eliminated within these protected areas upon implementation. The benefits of reducing effort have been largely studied, and include increases in abundance, biomass, and diversity within the bounded regions. These no-take zones may produce spillover effects, which provide fish for outside areas. However, the economic and ecological implications of displacing fishing effort are not yet fully understood. Novel data products that track fishing effort at the vessel-level allow us to identify changes in fleet- and vessel-level behavior upon the implementation of protected areas, as well as how these redistribute. This papers evaluates the implications of implementing LSMPA, by evaluating changes in fishing hours, showing that vessels in the effected region reduce fishing effort after the implementation of PIPA. Our results are robust to a set of specifications. We also track the relative spatial allocation of fishing events thorugh time, and identify that areas closer to PIPA show an increase in relative fishing hourse due to the displacement of PIPA-fishing vessels. Our results not only provide an impact evaluation of the effect of LSMPAs on fishing activity, but provide insights into vessel redistribution dynamics, which may have ecological and economic implications.

1 Introduction

Marine Protected Areas (MPAs) are intended to safeguard parts of the ocean from fishing and other extractive activities. Current international goals aim to "effectively protect 10% of the ocean environments by 2020". In an effort to meet this target, the world has seen a rapid increase in MPA coverage [Wood et al., 2008, Sala et al., 2018a]. A significant part of this rapid increase can be attributed to the designation of a small number of Large Scale Marine Protected Areas [LSMPAs; [Gray et al., 2017]]. These are defined as MPAS with at least 250,000 Km^2 in extension (Fig. 1), and are often implemented in the pelagic environment, where the dominant human activity is industrial fishing [Toonen et al., 2013, Gray et al., 2017, Kroodsma et al., 2018].

Due to weak property rights, limited habitat transformation, and potentially lower management costs, pelagic MPAs provide an opportunity to safeguard the oceans Game et al. [2009]. A growing body of literature has shown that closing the high seas to all fishing -effectively, turning them into a LSMPA- could increase fishery yields and profitability of fisheries, with negligible costs to food security [White and Costello, 2014, Sumaila et al., 2015, Sala et al., 2018b, Schiller et al., 2018]. However, as with customary MPAs, it is important that we understand the socioeconomic implications of management interventions.

Given the relatively recent establishment of most LSMPAs, very little is known about their human dimensions and implication for fisheries [Gray et al., 2017]. LSMPAs were erroneously assumed to have little social implications due to their remoteness [Agardy et al., 2011, Gray et al., 2017]. However, the anticipation of a LSMPA can lead to preemptive overfishing, which can erode or delay the expected benefits of the intervention [McDermott et al., 2018]. LSMPAs have received great attention in terms of governance and enforcement, but are yet to be the focus of economic analyses [Gray et al., 2017].

Research on LSMPAs have been the focus of analysing X, Y, and Z. However, no studies look into displacement of fishing effort. Redistribution of fishing effort has been evaluated in ssf by X, Y ,and Z. Cabral et al. [2017] analyse the redistribution of fishing and non-fishing vessels following the implementation of MPAs in California, and find that responses are idiosyncratic; commercial dive boats follow a fishing-the-line pattern, while some fishing boats follow an ideal free distribution. The way in which fishers react to a spatial closure can have major implications in its outcome [Smith and Wilen, 2003, Hilborn et al., 2006]. This highlights the need to understand how fishers react to the implementation of a LSMPA, and fishing effort changes and is spatially redistributed.

The main objective of this paper is to identify how fishers change their behavior due to the implementation of LSMPAs. We combine novel vessel tracking technologies and causal inference techniques to identify behavioral changes of fishing vessels due to the implementation of PIPA. We focus on fishing hours and distance traveled as outcome variables that fishers might adjust following implementation of a LSMPA in an impact-evaluation fashion. Additionally, we evaluate the spatial redistribution of fishing effort that existed within PIPA before its implementation. This work provides novel empirical insights into fisher's responses to the implementation of LSMPAs, and can help guide future interventions. Our work is novel in the sense that it provides empirical evidence of the effect of large-acale Marine Protected Areas in fishing behavior and distribution.

The next sections are as follows: Section 2 provides an overview of the Nauru Agreement and associated countries, a description of the fleet that operates in the region, and a brief history of PIPA. Section 3 describes our data and identification strategy. Section 4 presents our results, section 5 provides an extension of our results to other LSMPAs and discusses our results.



Figure 1:

2 Background

The Nauru Agreement was established in 1982 by Pacific island nations to manage their important tuna resources. PNA Members include Federated States of Micronesia, Kiribati, Marshall Islands, Nauru, Palau, Papua New Guinea, Solomon Islands, and Tuvalu.The Nauru Agreement regulated access of foreign vessels (*i.e.* those from non-PNA countries). Holding ~80% of the historical purse seining grounds within their Exclusive Economic Zones, PNA countries gained bargaining power when providing access to foreign fleets [Havice, 2010].

The cooperation that emerged thanks to the PNA allowed for subsequent agreements that strengthened fisheries management, like the Palau Agreement, which limited the number of purse seiners at 205 vessels from 1995-2007[1]. However, the most notable regulation is their approach to manage fishing effort: a Vessel Day Scheme (VDS) implemented in 2007 [Havice, 2013]. This effectively modified how fishing effort was managed, from number of vessels under the Palau Agreement to fishing hours. The VDS works as follows: Each year, scientific advisors recomend a total number of fishing vessel-days per year. Hours are allocated to each PNA country based on catch history, and

[1]See Havice [2010] for a detailed description of the Nauru, Palau, and Federal States of Micronesia agreements, their objectives, and outcomes.

they then sell fishing rights to other non-PNA countries [Aqorau et al., 2018]. While the effectiveness of this scheme has been debated in terms of meeting their fishery management and conservation objectives, the licensing significantly contributes to the economy of these island nations [Havice, 2010].



Figure 2:

The main tuna species caught in the region are skipjack (*Katsuwonus pelamis*), yellowfin (*Thunnus albacares*), albacore (*Thunnus alalunga*) and bigeye (*Thunnus obesus*). From these, the first two are amongst the top-10 species represented in global fisheries production statistics, with catches increasing relative to the 2005-2014 average [Food and of the United Nations, 2018]. The pacific region has historically accounted for a large portion of tuna species production [Aqorau and Bergin, 1997]. Today, the PNA controls close to 50% of the global skipjack tuna production [PNA, 2018]. A large portion of these catches derive from purse seine vessels licensed under the VDS. Fishing vessels from Australia, New Zeland, China, France, Korea, Japan, the Philippines, Taiwan and the United States participate in the purse-seining vessel day schemes.

For example, McDermott et al. [2018] show that fishing effort within the Phoenix Islands Marine Protected Area (PIPA) is effectively reduced after implementation, and describe changes in fishing behavior pre-implementation.

3 Methods

This section is divided into two main parts. First, we provide a general description of AIS data and the process of identification of vessel-level fishing events done by Global Fishing Watch[2]. Alongside, we describe the subset of data that we use for these analyses. When relevant, we also point out possible shortcomings in the data, or factors that must be considered in the later analyses. We then move on to explain our empirical strategy for the identification of the behavioral changes and redistribution of fishing effort.

3.1 Data

Automatic Identification Systems are on-board devices intended to provide at-sea safety and prevent ship collisions by broadcasting vessel position, course, and activities to surrounding vessels. These broadcasted messages can be received by satellites and land-based antennas. GFW uses a neural network to infer vessel characteristics and whether each broadcasted position represents a fishing event, thus allowing us to estimate near real-time fishing events globally since 2012 [Kroodsma et al., 2018]. Our data contain information for 2012 - 2017. The recent addition of satellites that can receive AIS signals causes an apparent increase in the number of broadcasted AIS messages (*i.e.* points), and therefore number of vessels and fishing hours. The variability in AIS data and ocean conditions require that temporal trends be taken into account. We do that by obtaining a subset of data that meet a BACI design, which gives us the full tracks for vessels affected and unaffected by the implementation of PIPA.

Our data contain over 45 million individual AIS messages (*i.e.* positions) for 371 purse seiners and longliners. A total of 233 vessels have fished within PIPA waters; 217 did so at least once before 2015. However, not all vessels continued to fish elsewhere after PIPA implementation: 34 vessels have no recorded AIS messages after 2015[3], leaving us with 176 vessels that fished inside PIPA before its implementation, and continued to fish elsewhere afterward. New vessels might have also entered the fishery after PIPA closure, and were likely not exposed to the policy intervention in the pre-treatment period. To account for this, we identify a subset of vessels which we track since before the implementation of PIPA, and categorize them as treated or control vessels. Our treatment and control groups are defined as follows.

The treatment group contains all vessels (n = 176) that fished within PIPA at least once before the closure, and that continued to fish elsewhere afterwards. Vessels in the control group meet all three of the following conditions: i) vessels never fished within PIPA waters, ii) vessels belong to other PNA countries, and iii) vessels have fished in surrounding areas (*i.e.* PNA-countries' EEZ) before and after PIPA closure. For each vessel meeting these characteristics, we calculate their total monthly fishing hours. Figure 3 provides a visual representation of the vessel-level fishing events that make up each group through time. Table 1 shows the number of vessels following a BACI design, as well as the fishing hours, before and after PIPA.

[2] Global Fishing Watch: globalfishingwatch.org

[3] The 34 missing vessels might have exited the fishery, been decommissioned or sold (therefore changing their AIS and mmsi), or turned off their AIS transmitters. In either case, we are not able to observe these.



Figure 3: Stream of fishing events by vessels through time. Each line represents a vessel, with dots indicating months with fishing activity and colors indicating the pre and post periods.

Table 1: Number of fishing vessels and fishing hours by gear and treatment group before and after PIPA.

Gear	Treatment	n	Before	After	Change (A / B)
drifting_longlines	FALSE	77	39.03181	37.79044	0.9681958
drifting_longlines	TRUE	115	41.62659	40.43857	0.9714600
purse_seines	FALSE	26	11.73804	20.13247	1.7151474
purse_seines	TRUE	61	10.66939	18.36289	1.7210820

3.2 Analysis

The first analysis focuses on identifying the response of fishing vessels to PIPA closure. Our variables of interests are fishing effort, indicated by total fishing hours per month, and distance traveled (Km) on every fishing trip. We compare fishing hours[4] before and after the implementation of PIPA using a Difference-in-Differences approach, where we track the variable of interest for vessels that used to fish inside PIPA and vessels that never fished inside PIPA, before and after PIPA implementation. Our specification is the following:

$$y_{i,t} = \alpha + \beta_1 Post_t + \beta_2 Treat_i + \beta_3 Post_t \times Treat_i + \mu_1 Y_t + \mu_2 Y_t^2 + \phi_t + \gamma_i + \epsilon_{i,t}$$

Where $y_{i,t}$ is the variable of interest for vessel i in time period t. A dummy variable $Post_t$ takes the value of 0 for all dates prior to PIPA implementation and a value of 1 for all dates including and following PIPA implementation. $Treat_i$ is a dummy variable indicating whether a vessel belongs to the control ($Treat_i = 0$) or treatment ($Treat_i = 1$) group. α is the standard intercept, β_1 captures the temporal trend change, β_2 captures the difference between treated and control groups, and β_3 is our parameter of interest: de DiD estimate capturing the treatment effect. Finally, μ_1 and μ_2 are coefficients for a second order polynomial for years (Y_t), while ϕ_t and γ_i represent month-, and flag-level dummies that account for seasonality or country-level management interventions[5].

Our second part of the analyses focuses on the redistribution of fishing effort. In other words, identifying where do vessels that used to fish inside PIPA go after its establishment. We calculate the monthly relative distribution of fishing hours by all treated vessels across all fished EEZs and the high seas. These trends are shown in Figure 5, and the relative temporal change is presented in Table **??**. EEZs that had sporadic fishing events were pooled into a group of "others", leaving us with a total of n = 23 and n = 0[6] spatially defined regions (*i.e.* EEZs, High Seas, "other EEZs", and PIPA) for purse seiners and longliners, respectively.

To evaluate this change in effort allocation, we regress our variable of interest (*i.e.*fishing hours) on the interaction between a dummy variable indicating the policy intervention and a dummy variable for countries. This gives us the by-country change in proportional allocation of fishing effort:

$$y_{i,t} = \alpha + \beta_1 Post_t + \beta_{2,i} Country_i + \beta_{3,i} Post_t \times Country_i + \epsilon_{i,t}$$

Our variable of interest, $y_{i,t}$ represents the proportion of fishing hours that country i receives at time t. *Post* also represents a policy dummy that takes the value of 0 for all dates before implementation of PIPA, and 1 otherwise. *Country* is a dummy variable for countries, interpreted as individual EEZs, the high seas, and a group of "other EEZs". Our parameter of interest is $\beta_{3,i}$, which captures the country-level change in proportional fishing effort.

All regression coefficients were estimated via ordinary least squares, and heteroskedastic-robust standard errors were calculated. All analyses were performed in R version 3.5.1 [R Core Team, 2018]. Raw data and code used in this work are available on github.

[4] And soon, distance

[5] An earlier specification included years as a dummy variable. Such results are included in the appendix, but are similar to the ones found under current specification.

[6] This number is likely to change upon finalizing the spatial analysis of longliners, which is currently running.

4 Results

Our data suggest that purse seiners and longliners have different responses to the implementation of a Large-Scale Marine Protected Areas. Fig. 4 shows that mean fishing hours for purse seiners have an abrupt increase, just before January 1st, 2015. This trend is observed for both treated and control groups. The pattern closely corresponds with the increase in total hours, but the total number of vessels doesn't entirely follow this pattern. The increase in fishing hours might be caused by the increased number of satellites[7]. Longliners, however, show no apparent trend with a clear seasonality [Ortuño-Crespo et al., 2018]. The number of mmsi codes also increases slightly through time, but becomes stable after 2015. For both gears and across all measures, the treatment and control vessels follow similar patterns, confirming our claim that the control group provides a plausible counterfactual.

Figure 4: Fishing hours and number of vessels by month for all vessels. Vertical dashed line indicates PIPA closure.

Our DiD analysis shows that treated purse seiners reduce their fishing effort after PIPA implementation in the order of 16 hours per month. This result is robust and significant ($p < 0.05$) for

[7]Need to check this. Not sure any satellites were incorporated during 2014. It is also possible that PNA countries started enforcing the requirement of having an AIS unit. On either case, both treatment and control groups seem to be affected equally.

all model specifications, with the effect varying between $\beta_3 = -16.457$ and $\beta_3 = -18.709$. Model specifications that include the year polynomial show lower values for the β_1 coefficient associated to the $Post_t$ policy dummy, and show positive and negative values for μ_1 and μ_2, the linear and quadratic terms for Y_t, respectively. These effectively represent the patterns observed in Figure 4.

Longliners show a similar pattern of effort reduction. However, the magnitude of the β_3 coefficient is smaller (ranging from $\beta_3 = -9.851$ to $\beta_3 = -14.850$) and not significant across all model specifications. This, along with higher standard error values suggest that longliners have a smaller and more variable response to the implementation of LSMPAs.

Regressions coefficients for each gear type are shown in Tables **??** and 4. Column (1) presents the DiD regression with no fixed effects, column (2) includes month fixed effects, column (3) includes month and the second degree polynomial for years, and column (4) includes all of the above and country-level fixed effects.

Recall that to evaluate the redistribution of fishing effort we only track fishing vessels that belong to the treated group. In this case, we calculate the proportion of fishing effort allocated every month to each spatially explicit region outlined by EEZs and the high seas. For purse seiners, these represent 9 main EEZs, PIPA, the high seas, and a group of other EEZs. Figure 5 shows the monthly relative fishing hours that each region received by all 176 treated vessels. The top-left panel shows the change in fishing effort inside PIPA, including the preemptive fishing and immediate reduction previously reported [McDermott et al., 2018].

The change in the relative allocation of fishing effort by purse seiners increases in eight of the 12 regions after PIPA implementation (Table 2). The largest increase is observed for the I-Kiribati EEZ, with an average increase of 0.11 ($p < 0.001$). In other words, the redistribution of treated vessels caused a 10% increase in the *relative* allocation of fishing effort within I-Kiribati waters. The only decrease is observed for Papua New Guinea, but the coefficient is not significant. Figure 6 provides a spatial representation of these changes. It is evident that the increase in relative fishing effort is greater for for regions closer to PIPA.



Figure 5: Monthly relative allocation of fishing effort by PIPA-fishing vessels before and after PIPA for 9 EEZs, PIPA, the high seas and 'other EEZs'.



Figure 6: Spatial representation of the mean change in the monthly allocation of fishing effort for purse seiners.

Table 2: Change in the relative allocation of fishing hours by purse seiners for each region. Asterisks indicate significance levels. Numbers in parenthesis represent heteroskedastic-robust standard errors.

	Dependent variable:
	h_prop
post	0.005*** (0.001)
countryEEZ FSM 1	0.052*** (0.001)
countryEEZ KIR 1	0.067*** (0.002)
countryEEZ KIR 2	0.020*** (0.001)
countryEEZ KIR 3	0.250*** (0.003)
countryEEZ MHL 1	0.022*** (0.001)
countryEEZ NRU 1	0.049*** (0.001)
countryEEZ PNG 2	0.184*** (0.004)
countryEEZ SLB 1	0.062*** (0.002)
countryEEZ TUV 1	0.059*** (0.001)
countryHS	0.073*** (0.001)
countryHS COK 1	0.001*** (0.0004)
countryHS FSM 1	−0.004*** (0.0003)
countryHS KIR 1	0.015*** (0.001)
countryHS KIR 2	−0.001* (0.0003)
countryHS KIR 3	0.002*** (0.0004)
countryHS MHL 1	−0.004*** (0.0003)
countryHS MUS 1	−0.004*** (0.0003)
countryHS NRU 1	−0.003*** (0.001)
countryHS PNG 2	−0.001* (0.001)
countryHS SLB 1	−0.003*** (0.0003)
countryHS TUV 1	−0.003*** (0.0003)
countryPIPA PIPA 1	0.083*** (0.002)
post:countryEEZ FSM 1	0.003 (0.002)
post:countryEEZ KIR 1	−0.007*** (0.002)
post:countryEEZ KIR 2	0.040*** (0.003)
post:countryEEZ KIR 3	−0.033*** (0.005)
post:countryEEZ MHL 1	−0.010*** (0.002)
post:countryEEZ NRU 1	−0.005*** (0.002)
post:countryEEZ PNG 2	−0.096*** (0.004)
post:countryEEZ SLB 1	0.026*** (0.004)
post:countryEEZ TUV 1	0.007*** (0.002)
post:countryHS	0.034*** (0.002)
post:countryHS COK 1	−0.005*** (0.001)
post:countryHS FSM 1	−0.005*** (0.001)
post:countryHS KIR 1	0.024*** (0.001)
post:countryHS KIR 2	0.005*** (0.001)
post:countryHS KIR 3	0.020*** (0.001)
post:countryHS MHL 1	−0.004*** (0.001)
post:countryHS MUS 1	−0.005*** (0.001)
post:countryHS NRU 1	−0.006*** (0.001)
post:countryHS PNG 2	−0.007*** (0.001)
post:countryHS SLB 1	−0.006*** (0.001)
post:countryHS TUV 1	−0.005*** (0.001)
post:countryPIPA PIPA 1	−0.088*** (0.002)
Constant	0.004*** (0.0003)
Observations	69,048

5 Discussion

Our findings provide interesting insights into the effect that LSMPAs can have on vessel behavior and the redistribution of fishing effort. These collection of results shows that the implementation of PIPA caused treated vessels to reduce their fishing hours, and that this effect is greater for purse seiners than longliners. Even though treated vessels fish less, their relative allocation of fishing hours increased for all other fishing grounds. This does not imply that there is more fishing effort exerted by treated vessels, but rather that each region receives a greater portion of the post-PIPA fishing effort of these same vessels, which is lower than pre-PIPA levels. In this section we discuss the implications of vessel-level reductions in fishing effort and the increase in relative allocation of the remaining effort through space. We also provide plausible explanations as to why purse seiners seem to be more reactive to the spatial closure.

A major shortcoming of our analyses is that we do not observe catches or revenues, which ultimately are the factors that guide the decision making process of profit-maximizing agents. Therefore, it is difficult to know whether the reduction in fishing effort represents a positive or negative impact. A decrease in fishing effort is associated to an increase in catches (and therefore greater CPUE) only when the entire fleet does it, and if previous levels of effort were greater than F_{MEY} (*i.e.* the effort that would yield the maximum economic yield). Therefore, it is plausible that the reduction of fishing hours is not done by choice, but rather results from fishers having to increase search time. Upon being relocated, fishers may not identify the best fishing grounds as easily as before, and therefore invest a greater proportion of their time searching for their catch. Further analysis of temporal trends in non-fishing hours, as well as distance traveled should provide us with insights as to why fishers reduced fishing hours.

Previous studies on insular environments suggest that vessels move to distant places, which might be translated as increased costs [Stevenson et al., 2013]. Nevertheless, they do not use counterfactuals that could help account for system- or fleet-level changes that occur through time. Others have used similar satellite-tracking systems to show that fishing effort accumulates near the edges of spatial closures, yielding greater catches [Murawski et al., 2005]. Yet, these vessel tracks do not cover the pre-reserve period, making it difficult identify the contribution of spatial closures to the observed spatial distribution of fishing vessels. Recent work by Elahi et al. [2018] identified that total fishing effort in a focal region where a short-term MPA was implemented showed little change, likely indicating that fishers redistributed fishing effort to compensate for the reduction in available space. Our data is assembled in a similar way, with fishing positions before and after the implementation of PIPA and vessels grouped into treated and control groups. Our BACI design, along with our difference-in-differences analysis allows us to make causal inferences about the effect that large scale marine protected areas have on fishing effort.

The different responses observed between purse seiners and longliners might have two possible explanations. It is likely that PIPA did not contain habitat that longliners would consider optimal. Therefore, the sporadic fishing events that occurred there are of little importance to the fleet, and it is unlikely that the implementation of PIPA has an effect on them. Alternatively, the differences may be due to the nature of each fishing gear. Purse seiners are often constrained by seafloor and termocline depth, and have a smaller spatial footprint [Kroodsma et al., 2018]. Tuna purse seiners are known to have greater proportion of null sets (*i.e.* where purse seines effectively cast their nets, but no catch is obtained) during El Niño years, where the termocline deepens in the Eastern Pacific [Dreyfus-Leon, 2015]. On the other hand, longliners may be more flexible as to where they can deploy their longlines. Ortuño-Crespo et al. [2018] evaluated the ecological niche of the pelagic longline fleet, and suggest that the fleet may be under-utilizing the ocean, meaning that they can easily redistribute elsewhere.

Our work suggests that the implementation of LSMPAs can have important implications for purse seiners, and less so for longliners. We also show that fishing effort is redistributed to areas close by. Future management interventions that aim to close large portion of the oceans should consider how fishing effort will change in space and through time, and the ecological implications of this redistribution to ensure that fishing effort is not just displaced elsewhere, leading to overfishing in adjacent waters.

6 Appendix

6.1 Other model specifications

6.1.1 Year as dummy variables

Table 3: Fishing hours from GFW for purse seiners (n = 106; 38 control, 68 treatment). Asterisks indicate significance levels. Numbers in parenthesis represent heteroskedastic-robuste standard errors.

	Dependent variable:			
	hours			
	(1)	(2)	(3)	(4)
post	8.394***	9.334***	14.934***	17.284***
	(0.267)	(0.254)	(0.312)	(0.403)
treated	−1.069***	−0.770***	−0.993***	0.259
	(0.249)	(0.234)	(0.219)	(0.269)
post:treated	−0.701**	−0.899***	−0.530*	−0.580**
	(0.308)	(0.295)	(0.283)	(0.292)
Constant	11.738***	11.281***	7.574***	10.077***
	(0.220)	(0.292)	(0.318)	(0.422)
Month FE	No	Yes	Yes	Yes
Year FE	No	No	Yes	Yes
Flag FE	No	No	No	Yes
Observations	37,840	37,840	37,840	37,840
R^2	0.087	0.136	0.179	0.192
Note:				*p<0.1; **p<0.05; ***p<0.01

```
##
## % Table created by stargazer v.5.2 by Marek Hlavac, Harvard University. E-mail: hlavac at fa
## % Date and time: vie., oct. 12, 2018 - 06:16:13 p. m.
## \begin{table}[!htbp] \centering
##   \caption{}
##   \label{}
## \begin{tabular}{@{\extracolsep{5pt}}lccc}
## \\[-1.8ex]\hline
## \hline \\[-1.8ex]
##  & \multicolumn{3}{c}{\textit{Dependent variable:}} \\
## \cline{2-4}
## \\[-1.8ex] & \multicolumn{3}{c}{hours} \\
## \\[-1.8ex] & (1) & (2) & (3)\\
```

Table 4: Fishing hours from GFW for longliners (n = 203; 88 control, 115 treatment).. Asterisks indicate significance levels. Numbers in parenthesis represent heteroskedastic-robuste standard errors.

	Dependent variable:			
	hours			
	(1)	(2)	(3)	(4)
post	-1.241^{***}	-1.026^{***}	-0.944^{***}	13.292^{***}
	(0.116)	(0.116)	(0.157)	(0.330)
treated	2.595^{***}	2.708^{***}	2.639^{***}	1.493^{***}
	(0.101)	(0.101)	(0.102)	(0.116)
post:treated	0.053	-0.069	-0.029	-0.060
	(0.133)	(0.133)	(0.134)	(0.135)
Constant	39.032^{***}	38.105^{***}	38.388^{***}	36.993^{***}
	(0.089)	(0.131)	(0.150)	(0.178)
Month FE	No	Yes	Yes	Yes
Year FE	No	No	Yes	Yes
Flag FE	No	No	No	Yes
Observations	227,873	227,873	227,873	227,873
R^2	0.010	0.015	0.016	0.030
Note:	*p<0.1; **p<0.05; ***p<0.01			

```
## \hline \\[-1.8ex]
##  post & 8.394$^{***}$ & 9.334$^{***}$ & 9.139$^{***}$ \\
##   & (0.271) & (0.265) & (0.300) \\
##   & & & \\
##  treated & $-$1.069$^{***}$ & $-$0.770$^{***}$ & 0.041 \\
##   & (0.259) & (0.253) & (0.317) \\
##   & & & \\
##  month\_c10 &  & 2.885$^{***}$ & 2.639$^{***}$ \\
##   &  & (0.302) & (0.335) \\
##   & & & \\
##  month\_c11 &  & 3.687$^{***}$ & 3.556$^{***}$ \\
##   &  & (0.304) & (0.335) \\
##   & & & \\
##  month\_c12 &  & 3.001$^{***}$ & 2.594$^{***}$ \\
##   &  & (0.309) & (0.339) \\
##   & & & \\
##  month\_c2 &  & $-$0.989$^{***}$ & $-$1.062$^{***}$ \\
##   &  & (0.308) & (0.334) \\
##   & & & \\
##  month\_c3 &  & $-$2.238$^{***}$ & $-$2.412$^{***}$ \\
##   &  & (0.313) & (0.340) \\
##   & & & \\
##  month\_c4 &  & $-$3.643$^{***}$ & $-$4.008$^{***}$ \\
##   &  & (0.317) & (0.345) \\
##   & & & \\
##  month\_c5 &  & $-$4.590$^{***}$ & $-$4.914$^{***}$ \\
##   &  & (0.316) & (0.349) \\
##   & & & \\
##  month\_c6 &  & $-$4.420$^{***}$ & $-$4.977$^{***}$ \\
##   &  & (0.332) & (0.368) \\
##   & & & \\
##  month\_c7 &  & $-$3.580$^{***}$ & $-$3.351$^{***}$ \\
##   &  & (0.336) & (0.366) \\
##   & & & \\
##  month\_c8 &  & 0.317 & 0.204 \\
##   &  & (0.311) & (0.346) \\
##   & & & \\
##  month\_c9 &  & 2.865$^{***}$ & 2.305$^{***}$ \\
##   &  & (0.309) & (0.341) \\
##   & & & \\
##  flagECU &  &  & $-$6.188$^{***}$ \\
##   &  &  & (0.434) \\
##   & & & \\
##  flagESP &  &  & $-$2.949$^{***}$ \\
##   &  &  & (0.497) \\
##   & & & \\
##  flagFSM &  &  & $-$4.729$^{***}$ \\
##   &  &  & (0.470) \\
```

```
##   & & & \\
##  flagKOR &  &  & $-$3.708$^{***}$ \\
##   &  &  & (0.289) \\
##   & & & \\
##  flagMHL &  &  & $-$1.879$^{***}$ \\
##   &  &  & (0.310) \\
##   & & & \\
##  flagOTH &  &  & $-$2.244$^{***}$ \\
##   &  &  & (0.290) \\
##   & & & \\
##  flagTWN &  &  & $-$2.493$^{***}$ \\
##   &  &  & (0.332) \\
##   & & & \\
##  post:treated & $-$0.701$^{**}$ & $-$0.899$^{***}$ & $-$0.908$^{***}$ \\
##   & (0.314) & (0.306) & (0.344) \\
##   & & & \\
##  Constant & 11.738$^{***}$ & 11.281$^{***}$ & 13.942$^{***}$ \\
##   & (0.229) & (0.313) & (0.422) \\
##   & & & \\
## \hline \\[-1.8ex]
## Observations & 37,840 & 37,840 & 30,359 \\
## R$^{2}$ & 0.087 & 0.136 & 0.151 \\
## Adjusted R$^{2}$ & 0.087 & 0.136 & 0.150 \\
## Residual Std. Error & 12.752 (df = 37836) & 12.405 (df = 37825) & 12.257 (df = 30337) \\
## F Statistic & 1,203.929$^{***}$ (df = 3; 37836) & 427.020$^{***}$ (df = 14; 37825) & 256.286
## \hline
## \hline \\[-1.8ex]
## \textit{Note:}  & \multicolumn{3}{r}{$^{*}$p$<$0.1; $^{**}$p$<$0.05; $^{***}$p$<$0.01} \\
## \end{tabular}
## \end{table}

##
## % Table created by stargazer v.5.2 by Marek Hlavac, Harvard University. E-mail: hlavac at fa
## % Date and time: vie., oct. 12, 2018 - 06:16:13 p. m.
## \begin{table}[!htbp] \centering
##   \caption{}
##   \label{}
## \begin{tabular}{@{\extracolsep{5pt}}lccc}
## \\[-1.8ex]\hline
## \hline \\[-1.8ex]
##  & \multicolumn{3}{c}{\textit{Dependent variable:}} \\
## \cline{2-4}
## \\[-1.8ex] & \multicolumn{3}{c}{hours} \\
## \\[-1.8ex] & (1) & (2) & (3)\\
## \hline \\[-1.8ex]
##  post & 8.222$^{***}$ & 8.665$^{***}$ & 8.923$^{***}$ \\
##   & (0.444) & (0.431) & (0.429) \\
##   & & & \\
```

```
##  treated & $-$0.015 & 0.059 & 2.910$^{***}$ \\
##   & (0.475) & (0.462) & (0.602) \\
##   & & & \\
##  month\_c10 &  & 3.140$^{***}$ & 3.489$^{***}$ \\
##   &  & (0.711) & (0.877) \\
##   & & & \\
##  month\_c11 &  & 4.203$^{***}$ & 4.775$^{***}$ \\
##   &  & (0.695) & (0.828) \\
##   & & & \\
##  month\_c12 &  & 2.878$^{***}$ & 3.087$^{***}$ \\
##   &  & (0.699) & (0.837) \\
##   & & & \\
##  month\_c2 &  & $-$0.880 & $-$1.082 \\
##   &  & (0.739) & (0.885) \\
##   & & & \\
##  month\_c3 &  & $-$2.181$^{***}$ & $-$1.776$^{**}$ \\
##   &  & (0.740) & (0.891) \\
##   & & & \\
##  month\_c4 &  & $-$4.056$^{***}$ & $-$4.186$^{***}$ \\
##   &  & (0.745) & (0.893) \\
##   & & & \\
##  month\_c5 &  & $-$4.023$^{***}$ & $-$4.108$^{***}$ \\
##   &  & (0.730) & (0.891) \\
##   & & & \\
##  month\_c6 &  & $-$4.743$^{***}$ & $-$5.336$^{***}$ \\
##   &  & (0.779) & (0.968) \\
##   & & & \\
##  month\_c7 &  & $-$4.457$^{***}$ & $-$4.115$^{***}$ \\
##   &  & (0.815) & (1.022) \\
##   & & & \\
##  month\_c8 &  & 0.972 & 1.249 \\
##   &  & (0.724) & (0.895) \\
##   & & & \\
##  month\_c9 &  & 4.947$^{***}$ & 4.322$^{***}$ \\
##   &  & (0.714) & (0.878) \\
##   & & & \\
##  flagMHL &  &  & 2.969$^{***}$ \\
##   &  &  & (0.467) \\
##   & & & \\
##  post:treated & $-$2.014$^{***}$ & $-$2.047$^{***}$ & $-$2.984$^{***}$ \\
##   & (0.610) & (0.592) & (0.766) \\
##   & & & \\
##  Constant & 11.415$^{***}$ & 10.972$^{***}$ & 8.160$^{***}$ \\
##   & (0.342) & (0.616) & (0.835) \\
##   & & & \\
## \hline \\[-1.8ex]
## Observations & 7,756 & 7,756 & 5,288 \\
## R$^{2}$ & 0.069 & 0.132 & 0.145 \\
```

```
## Adjusted R$^{2}$ & 0.069 & 0.131 & 0.142 \\
## Residual Std. Error & 13.095 (df = 7752) & 12.652 (df = 7741) & 12.518 (df = 5272) \\
## F Statistic & 191.591$^{***}$ (df = 3; 7752) & 84.244$^{***}$ (df = 14; 7741) & 59.406$^{**
## \hline
## \hline \\[-1.8ex]
## \textit{Note:}  & \multicolumn{3}{r}{$^{*}$p$<$0.1; $^{**}$p$<$0.05; $^{***}$p$<$0.01} \\
## \end{tabular}
## \end{table}

##
## % Table created by stargazer v.5.2 by Marek Hlavac, Harvard University. E-mail: hlavac at fa
## % Date and time: vie., oct. 12, 2018 - 06:16:14 p. m.
## \begin{table}[!htbp] \centering
##   \caption{}
##   \label{}
## \begin{tabular}{@{\extracolsep{5pt}}lccc}
## \\[-1.8ex]\hline
## \hline \\[-1.8ex]
##  & \multicolumn{3}{c}{\textit{Dependent variable:}} \\
## \cline{2-4}
## \\[-1.8ex] & \multicolumn{3}{c}{hours} \\
## \\[-1.8ex] & (1) & (2) & (3)\\
## \hline \\[-1.8ex]
##  post & 9.018$^{***}$ & 10.574$^{***}$ & 10.492$^{***}$ \\
##   & (0.746) & (0.727) & (0.722) \\
##   & & & \\
##  treated & $-$0.530 & $-$0.039 & 0.430 \\
##   & (0.629) & (0.612) & (0.683) \\
##   & & & \\
##  month\_c10 &  & 2.437$^{***}$ & 2.224$^{***}$ \\
##   &  & (0.347) & (0.386) \\
##   & & & \\
##  month\_c11 &  & 3.376$^{***}$ & 3.150$^{***}$ \\
##   &  & (0.350) & (0.389) \\
##   & & & \\
##  month\_c12 &  & 2.625$^{***}$ & 2.072$^{***}$ \\
##   &  & (0.357) & (0.394) \\
##   & & & \\
##  month\_c2 &  & $-$1.197$^{***}$ & $-$1.256$^{***}$ \\
##   &  & (0.354) & (0.386) \\
##   & & & \\
##  month\_c3 &  & $-$2.779$^{***}$ & $-$2.850$^{***}$ \\
##   &  & (0.360) & (0.395) \\
##   & & & \\
##  month\_c4 &  & $-$3.767$^{***}$ & $-$4.013$^{***}$ \\
##   &  & (0.361) & (0.397) \\
##   & & & \\
##  month\_c5 &  & $-$4.653$^{***}$ & $-$5.032$^{***}$ \\
```

```
##   &  & (0.362) & (0.402) \\
##   & & & \\
##  month\_c6 &  & $-$4.797$^{***}$ & $-$5.323$^{***}$ \\
##   &  & (0.381) & (0.426) \\
##   & & & \\
##  month\_c7 &  & $-$3.646$^{***}$ & $-$3.487$^{***}$ \\
##   &  & (0.381) & (0.417) \\
##   & & & \\
##  month\_c8 &  & 0.153 & $-$0.202 \\
##   &  & (0.358) & (0.399) \\
##   & & & \\
##  month\_c9 &  & 3.151$^{***}$ & 2.287$^{***}$ \\
##   &  & (0.354) & (0.393) \\
##   & & & \\
##  flagECU &  &  & $-$5.462$^{***}$ \\
##   &  &  & (0.585) \\
##   & & & \\
##  flagESP &  &  & $-$3.357$^{***}$ \\
##   &  &  & (0.557) \\
##   & & & \\
##  flagFSM &  &  & $-$1.449$^{**}$ \\
##   &  &  & (0.675) \\
##   & & & \\
##  flagKOR &  &  & $-$4.298$^{***}$ \\
##   &  &  & (0.379) \\
##   & & & \\
##  flagMHL &  &  & $-$2.361$^{***}$ \\
##   &  &  & (0.501) \\
##   & & & \\
##  flagOTH &  &  & $-$3.311$^{***}$ \\
##   &  &  & (0.402) \\
##   & & & \\
##  flagTWN &  &  & $-$3.266$^{***}$ \\
##   &  &  & (0.473) \\
##   & & & \\
##  post:treated & $-$1.325$^{*}$ & $-$2.136$^{***}$ & $-$2.253$^{***}$ \\
##   & (0.762) & (0.742) & (0.742) \\
##   & & & \\
##  Constant & 11.200$^{***}$ & 10.748$^{***}$ & 14.339$^{***}$ \\
##   & (0.617) & (0.654) & (0.812) \\
##   & & & \\
## \hline \\[-1.8ex]
## Observations & 28,315 & 28,315 & 22,766 \\
## R$^{2}$ & 0.083 & 0.134 & 0.144 \\
## Adjusted R$^{2}$ & 0.083 & 0.133 & 0.143 \\
## Residual Std. Error & 12.672 (df = 28311) & 12.322 (df = 28300) & 12.231 (df = 22744) \\
## F Statistic & 857.512$^{***}$ (df = 3; 28311) & 311.587$^{***}$ (df = 14; 28300) & 182.169$
## \hline
```

```
## \hline \\[-1.8ex]
## \textit{Note:}  & \multicolumn{3}{r}{$^{*}$p$<$0.1; $^{**}$p$<$0.05; $^{***}$p$<$0.01} \\
## \end{tabular}
## \end{table}
##
## % Table created by stargazer v.5.2 by Marek Hlavac, Harvard University. E-mail: hlavac at fa
## % Date and time: vie., oct. 12, 2018 - 06:16:14 p. m.
## \begin{table}[!htbp] \centering
##   \caption{}
##   \label{}
## \begin{tabular}{@{\extracolsep{5pt}}lccc}
## \\[-1.8ex]\hline
## \hline \\[-1.8ex]
##  & \multicolumn{3}{c}{\textit{Dependent variable:}} \\
## \cline{2-4}
## \\[-1.8ex] & \multicolumn{3}{c}{hours} \\
## \\[-1.8ex] & (1) & (2) & (3)\\
## \hline \\[-1.8ex]
##  post & 7.875$^{***}$ & 8.778$^{***}$ & 8.770$^{***}$ \\
##   & (0.303) & (0.296) & (0.336) \\
##   & & & \\
##  treated & $-$0.751$^{***}$ & $-$0.445$^{*}$ & 0.557 \\
##   & (0.269) & (0.262) & (0.345) \\
##   & & & \\
##  month\_c10 &  & 2.848$^{***}$ & 2.445$^{***}$ \\
##   &  & (0.317) & (0.356) \\
##   & & & \\
##  month\_c11 &  & 3.647$^{***}$ & 3.320$^{***}$ \\
##   &  & (0.319) & (0.355) \\
##   & & & \\
##  month\_c12 &  & 3.011$^{***}$ & 2.457$^{***}$ \\
##   &  & (0.324) & (0.358) \\
##   & & & \\
##  month\_c2 &  & $-$1.143$^{***}$ & $-$1.308$^{***}$ \\
##   &  & (0.326) & (0.357) \\
##   & & & \\
##  month\_c3 &  & $-$2.430$^{***}$ & $-$2.707$^{***}$ \\
##   &  & (0.331) & (0.364) \\
##   & & & \\
##  month\_c4 &  & $-$3.542$^{***}$ & $-$4.025$^{***}$ \\
##   &  & (0.332) & (0.365) \\
##   & & & \\
##  month\_c5 &  & $-$4.442$^{***}$ & $-$4.842$^{***}$ \\
##   &  & (0.332) & (0.371) \\
##   & & & \\
##  month\_c6 &  & $-$4.524$^{***}$ & $-$5.275$^{***}$ \\
##   &  & (0.349) & (0.392) \\
```

```
##   & & & \\
##  month\_c7 &  & $-$3.670$^{***}$ & $-$3.545$^{***}$ \\
##   &  & (0.354) & (0.389) \\
##   & & & \\
##  month\_c8 &  & 0.228 & 0.002 \\
##   &  & (0.328) & (0.368) \\
##   & & & \\
##  month\_c9 &  & 3.002$^{***}$ & 2.320$^{***}$ \\
##   &  & (0.324) & (0.362) \\
##   & & & \\
##  flagECU &  &  & $-$6.396$^{***}$ \\
##   &  &  & (0.515) \\
##   & & & \\
##  flagESP &  &  & $-$3.432$^{***}$ \\
##   &  &  & (0.552) \\
##   & & & \\
##  flagFSM &  &  & $-$4.834$^{***}$ \\
##   &  &  & (0.553) \\
##   & & & \\
##  flagKOR &  &  & $-$4.212$^{***}$ \\
##   &  &  & (0.378) \\
##   & & & \\
##  flagMHL &  &  & $-$1.837$^{***}$ \\
##   &  &  & (0.442) \\
##   & & & \\
##  flagOTH &  &  & $-$3.284$^{***}$ \\
##   &  &  & (0.401) \\
##   & & & \\
##  flagTWN &  &  & $-$2.638$^{***}$ \\
##   &  &  & (0.439) \\
##   & & & \\
##  post:treated & $-$0.182 & $-$0.331 & $-$0.455 \\
##   & (0.342) & (0.333) & (0.379) \\
##   & & & \\
##  Constant & 11.420$^{***}$ & 10.972$^{***}$ & 14.036$^{***}$ \\
##   & (0.239) & (0.329) & (0.548) \\
##   & & & \\
## \hline \\[-1.8ex]
## Observations & 34,489 & 34,489 & 27,008 \\
## R$^{2}$ & 0.082 & 0.133 & 0.147 \\
## Adjusted R$^{2}$ & 0.082 & 0.132 & 0.147 \\
## Residual Std. Error & 12.712 (df = 34485) & 12.359 (df = 34474) & 12.188 (df = 26986) \\
## F Statistic & 1,031.434$^{***}$ (df = 3; 34485) & 377.251$^{***}$ (df = 14; 34474) & 222.04
## \hline
## \hline \\[-1.8ex]
## \textit{Note:}  & \multicolumn{3}{r}{$^{*}$p$<$0.1; $^{**}$p$<$0.05; $^{***}$p$<$0.01} \\
## \end{tabular}
## \end{table}
```



608


5
0
−5
−10
estimate
2012
2013
2014
2015
2016
year
model
1
2


estimate
20
10
0
2013
2014
2015
2016
year
model
1
2


0
−2
−4
−6
estimate
2012
2013
2014
2015
2016
year
model
1
2



612



613



614



615

References

Tundi Agardy, Giuseppe Notarbartolo di Sciara, and Patrick Christie. Mind the gap: Addressing the shortcomings of marine protected areas through large scale marine spatial planning. *Marine Policy*, 35(2):226–232, mar 2011. ISSN 0308597X. doi: 10.1016/j.marpol.2010.10.006. URL http://linkinghub.elsevier.com/retrieve/pii/{S0308597X10001740}.

Transform Aqorau and Anthony Bergin. Ocean governance in the western pacific purse seine fishery - palau arrangement. *Marine Policy*, 21(2):173–186, mar 1997. ISSN 0308597X. doi: 10.1016/S0308-{597X}(96)00054-1. URL http://linkinghub.elsevier.com/retrieve/pii/{S0308597X96000541}.

Transform Aqorau, Johann Bell, and John N. Kittinger. Good governance for migratory species. *Science*, 361(6408):1208.2–1209, sep 2018. ISSN 0036-8075. doi: 10.1126/science.aav2051. URL http://www.sciencemag.org/lookup/doi/10.1126/science.aav2051.

Reniel B Cabral, Steven D Gaines, Brett A Johnson, Tom W Bell, and Crow White. Drivers of redistribution of fishing and non-fishing effort after the implementation of a marine protected area network. *Ecol Appl*, 27(2):416–428, mar 2017. ISSN 10510761. doi: 10.1002/eap.1446. URL http://doi.wiley.com/10.1002/eap.1446.

Michel J Dreyfus-Leon. Analysis of null sets (zero catch) made by the mexican tuna purse seine fleet (2000–2013). *Cienc Mar*, 41(2):85–92, jun 2015. ISSN 01853880. doi: 10.7773/cm.v41i2.2471. URL http://www.cienciasmarinas.com.mx/index.php/cmarinas/article/view/2471/1552.

Robin Elahi, Francesco Ferretti, Azzurra Bastari, Carlo Cerrano, Francesco Colloca, Jonathan Kowalik, Mary Ruckelshaus, Andreas Struck, and Fiorenza Micheli. Leveraging vessel traffic data and a temporary fishing closure to inform marine management. *Front Ecol Environ*, aug 2018. ISSN 15409295. doi: 10.1002/fee.1936. URL http://doi.wiley.com/10.1002/fee.1936.

Food and Agriculture Organization of the United Nations. *The state of world fisheries and aquaculture 2018: meeting the sustainable development goals*. The state of world fisheries and aquaculture. UN, jul 2018. ISBN 9789210472340. doi: 10.18356/8d6ea4b6-en. URL https://www.un-ilibrary.org/agriculture-rural-development-and-forestry/the-state-of-world-fisheries-and-aquaculture-2018_8d6ea4b6-en.

Edward T Game, Hedley S Grantham, Alistair J Hobday, Robert L Pressey, Amanda T Lombard, Lynnath E Beckley, Kristina Gjerde, Rodrigo Bustamante, Hugh P Possingham, and Anthony J Richardson. Pelagic protected areas: the missing dimension in ocean conservation. *Trends Ecol Evol (Amst)*, 24(7):360–369, jul 2009. doi: 10.1016/j.tree.2009.01.011. URL http://dx.doi.org/10.1016/j.tree.2009.01.011.

Noella J. Gray, Nathan J. Bennett, Jon C. Day, Rebecca L. Gruby, T. 'Aulani Wilhelm, and Patrick Christie. Human dimensions of large-scale marine protected areas: Advancing research and practice. *Coastal Management*, pages 1–9, oct 2017. ISSN 0892-0753. doi: 10.1080/08920753.2017.1373448. URL https://www.tandfonline.com/doi/full/10.1080/08920753.2017.1373448.

Elizabeth Havice. The structure of tuna access agreements in the western and central pacific ocean: Lessons for vessel day scheme planning. *Marine Policy*, 34(5):979–987, sep 2010. ISSN 0308597X. doi: 10.1016/j.marpol.2010.02.004. URL http://linkinghub.elsevier.com/retrieve/pii/{S0308597X10000461}.

Elizabeth Havice. Rights-based management in the western and central pacific ocean tuna fishery: Economic and environmental change under the vessel day scheme. *Marine Policy*, 42:259–267, nov 2013. ISSN 0308597X. doi: 10.1016/j.marpol.2013.03.003. URL http://linkinghub.elsevier.com/retrieve/pii/{S0308597X13000717}.

Ray Hilborn, Fiorenza Micheli, and Giulio A De Leo. Integrating marine protected areas with catch regulation. *Can. J. Fish. Aquat. Sci.*, 63(3):642–649, mar 2006. ISSN 0706-652X. doi: 10.1139/f05-243. URL http://www.nrcresearchpress.com/doi/abs/10.1139/f05-243.

David A Kroodsma, Juan Mayorga, Timothy Hochberg, Nathan A Miller, Kristina Boerder, Francesco Ferretti, Alex Wilson, Bjorn Bergman, Timothy D White, Barbara A Block, Paul Woods, Brian Sullivan, Christopher Costello, and Boris Worm. Tracking the global footprint of fisheries. *Science*, 359(6378):904–908, feb 2018. ISSN 0036-8075. doi: 10.1126/science.aao5646. URL http://www.sciencemag.org/lookup/doi/10.1126/science.aao5646.

Grant R McDermott, Kyle C Meng, Gavin G McDonald, and Christopher J Costello. The blue paradox: Preemptive overfishing in marine reserves. *Proc Natl Acad Sci USA*, aug 2018. doi: 10.1073/pnas.1802862115. URL http://dx.doi.org/10.1073/pnas.1802862115.

S Murawski, S Wigley, M Fogarty, P Rago, and D Mountain. Effort distribution and catch patterns adjacent to temperate MPAs. *ICES Journal of Marine Science*, jul 2005. ISSN 10543139. doi: 10.1016/j.icesjms.2005.04.005. URL https://academic.oup.com/icesjms/article-lookup/doi/10.1016/j.icesjms.2005.04.005.

Guillermo Ortuño-Crespo, Daniel C. Dunn, Gabriel Reygondeau, Kristina Boerder, Boris Worm, William Cheung, Derek P. Tittensor, and Patrick N. Halpin. The environmental niche of the global high seas pelagic longline fleet. *Sci. Adv.*, 4(8):eaat3681, aug 2018. ISSN 2375-2548. doi: 10.1126/sciadv.aat3681. URL http://advances.sciencemag.org/lookup/doi/10.1126/sciadv.aat3681.

PNA. Parties to the nauru agreement, 2018. URL https://www.pnatuna.com/.

R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2018. URL https://www.R-project.org/.

Enric Sala, Jane Lubchenco, Kirsten Grorud-Colvert, Catherine Novelli, Callum Roberts, and U. Rashid Sumaila. Assessing real progress towards effective ocean protection. *Marine Policy*, 91 (2):11–13, may 2018a. ISSN 0308597X. doi: 10.1016/j.marpol.2018.02.004. URL http://linkinghub.elsevier.com/retrieve/pii/{S0308597X17307686}.

Enric Sala, Juan Mayorga, Christopher Costello, David Kroodsma, Maria L D Palomares, Daniel Pauly, U Rashid Sumaila, and Dirk Zeller. The economics of fishing the high seas. *Sci Adv*, 4 (6):eaat2504, jun 2018b. ISSN 2375-2548. doi: 10.1126/sciadv.aat2504. URL http://advances.sciencemag.org/lookup/doi/10.1126/sciadv.aat2504.

Laurenne Schiller, Megan Bailey, Jennifer Jacquet, and Enric Sala. High seas fisheries play a negligible role in addressing global food security. *Sci Adv*, 4(8):eaat8351, aug 2018. ISSN 2375-2548. doi: 10.1126/sciadv.aat8351. URL http://advances.sciencemag.org/lookup/doi/10.1126/sciadv.aat8351.

Martin D. Smith and James E. Wilen. Economic impacts of marine reserves: the importance of spatial behavior. *Journal of Environmental Economics and Management*, 46(2):183–206, sep 2003. ISSN 00950696. doi: 10.1016/S0095-0696(03)00024-X. URL http://linkinghub.elsevier.com/retrieve/pii/{S009506960300024X}.

Todd C. Stevenson, Brian N. Tissot, and William J. Walsh. Socioeconomic consequences of fishing displacement from marine protected areas in hawaii. *Biological Conservation*, 160:50–58, apr 2013. ISSN 00063207. doi: 10.1016/j.biocon.2012.11.031. URL http://linkinghub.elsevier.com/retrieve/pii/S0006320712005277.

U. Rashid Sumaila, Vicky W. Y. Lam, Dana D. Miller, Louise Teh, Reg A. Watson, Dirk Zeller, William W. L. Cheung, Isabelle M. Côté, Alex D. Rogers, Callum Roberts, Enric Sala, and Daniel Pauly. Winners and losers in a world where the high seas is closed to fishing. *Sci Rep*, 5(1):8481, jul 2015. ISSN 2045-2322. doi: 10.1038/srep08481. URL http://www.nature.com/articles/srep08481.

Robert J Toonen, T 'Aulani Wilhelm, Sara M Maxwell, Daniel Wagner, Brian W Bowen, Charles R C Sheppard, Sue M Taei, Tukabu Teroroko, Russell Moffitt, Carlos F Gaymer, Lance Morgan, Nai'a Lewis, Anne L S Sheppard, John Parks, Alan M Friedlander, and Big Ocean Think Tank. One size does not fit all: the emerging frontier in large-scale marine conservation. *Mar Pollut Bull*, 77(1-2):7–10, dec 2013. doi: 10.1016/j.marpolbul.2013.10.039. URL http://dx.doi.org/10.1016/j.marpolbul.2013.10.039.

Crow White and Christopher Costello. Close the high seas to fishing? *PLoS Biol*, 12(3):e1001826, mar 2014. ISSN 1545-7885. doi: 10.1371/journal.pbio.1001826. URL http://dx.plos.org/10.1371/journal.pbio.1001826.

Louisa J. Wood, Lucy Fish, Josh Laughren, and Daniel Pauly. Assessing progress towards global marine protection targets: shortfalls in information and action. *Oryx*, 42(03), jul 2008. ISSN 0030-6053. doi: 10.1017/{S003060530800046X}. URL http://www.journals.cambridge.org/{abstract_S003060530800046X}.