

# EMMA'S ATLAS OF NUMBERS AND SHAPES

- A NOT-SO-BRIEF GUIDE TO ALL THE MATH I KNOW -

EMMA BACH

2026-09-12

# Contents

|            |                                               |           |
|------------|-----------------------------------------------|-----------|
| <b>I</b>   | <b>Foundations</b>                            | <b>6</b>  |
| 1          | An Introduction to Mathematical Reasoning     | 8         |
| 2          | Sets                                          | 9         |
| 2.1        | Infinity: $\emptyset$ and $\mathbb{N}$        | 10        |
| 2.2        | Extensionality                                | 10        |
| 2.3        | Specification                                 | 11        |
| 2.4        | Pairing, Union, and Power Sets                | 11        |
| 2.5        | Relations and Functions                       | 11        |
| 2.6        | Cartesian Products                            | 11        |
| 2.7        | Replacement                                   | 12        |
| 2.8        | Regularity and Well-Foundedness               | 12        |
| 2.9        | Choice                                        | 12        |
| 2.10       | Cardinality                                   | 12        |
| <b>II</b>  | <b>Algebraic Structures</b>                   | <b>13</b> |
| 1          | $(\mathbb{N}, +)$ is a Monoid                 | 15        |
| 2          | $(\mathbb{Z}, +, -)$ is a Group               | 16        |
| 2.1        | Group Homomorphisms                           | 16        |
| 2.2        | Permutations and the Symmetric Groups         | 17        |
| 2.3        | Product Groups                                | 18        |
| 2.4        | Group Actions                                 | 19        |
| 3          | $(\mathbb{Z}, +, -, \cdot)$ is a Ring         | 21        |
| 4          | $(\mathbb{Q}, +, -, \cdot, ^{-1})$ is a Field | 22        |
| <b>III</b> | <b>Foundations of Analysis</b>                | <b>23</b> |
| 1          | The Real Numbers $\mathbb{R}$                 | 25        |
| 1.1        | Ordered Algebraic Structures                  | 25        |
| 1.2        | Dedekind Cuts                                 | 25        |
| 1.3        | Uniqueness of $\mathbb{R}$                    | 25        |
| 1.4        | Cardinality of $\mathbb{R}$                   | 25        |
| 2          | The Concept of Distance                       | 26        |
| 2.1        | Metric Spaces                                 | 26        |
| 2.2        | Topologies                                    | 29        |
| 2.3        | Convergence                                   | 33        |
| 2.4        | Continuity                                    | 34        |
| 2.5        | Compactness                                   | 35        |

|           |                                                               |           |
|-----------|---------------------------------------------------------------|-----------|
| 2.6       | Connectedness.....                                            | 38        |
| 2.7       | Function Spaces.....                                          | 40        |
| 2.8       | Banach's Fixed Point Theorem .....                            | 43        |
| <b>3</b>  | <b>Infinite Series</b> .....                                  | <b>44</b> |
| 3.1       | Achilles and the Turtle.....                                  | 44        |
| 3.2       | Convergence Tests.....                                        | 44        |
| 3.3       | Euler's Number $e$ .....                                      | 44        |
| 3.4       | The Exponential Function.....                                 | 44        |
| <b>4</b>  | <b>Euclidean Geometry</b> .....                               | <b>45</b> |
| 4.1       | The Complex Numbers $\mathbb{C}$ .....                        | 45        |
| 4.2       | Circles, Angles and Trigonometry .....                        | 45        |
| <b>IV</b> | <b>Calculus in one Variable</b> .....                         | <b>46</b> |
| <b>1</b>  | <b>Derivatives</b> .....                                      | <b>47</b> |
| <b>2</b>  | <b>Integrals</b> .....                                        | <b>48</b> |
| 2.1       | The Gamma Function.....                                       | 48        |
| <b>V</b>  | <b>Linear Algebra</b> .....                                   | <b>51</b> |
| <b>1</b>  | <b>Analytic Geometry</b> .....                                | <b>54</b> |
| 1.1       | Arithmetic in $n$ dimensions.....                             | 54        |
| 1.2       | The Inner Product: Angles and Lengths in $\mathbb{R}^n$ ..... | 56        |
| 1.3       | The Cross Product.....                                        | 60        |
| 1.4       | The Quaternions $\mathbb{H}$ .....                            | 63        |
| 1.5       | Quaternions and $\mathbb{R}^3$ .....                          | 67        |
| <b>2</b>  | <b>Linear Maps</b> .....                                      | <b>68</b> |
| 2.1       | Vector Spaces .....                                           | 68        |
| 2.2       | Subspaces and Quotient Spaces.....                            | 69        |
| 2.3       | Bases and Dimension .....                                     | 69        |
| 2.4       | Matrices .....                                                | 69        |
| 2.5       | Systems of Linear Equations .....                             | 69        |
| 2.6       | The Method of Least Squares .....                             | 69        |
| <b>3</b>  | <b>Affine Spaces</b> .....                                    | <b>70</b> |
| <b>4</b>  | <b>Determinants</b> .....                                     | <b>74</b> |
| 4.1       | Defining Volume .....                                         | 74        |
| 4.2       | Determinants of Endomorphisms .....                           | 74        |
| 4.3       | Orientation.....                                              | 74        |
| <b>5</b>  | <b>Eigenvectors and Eigenvalues</b> .....                     | <b>75</b> |
| 5.1       | Characteristic Polynomials .....                              | 77        |
| 5.2       | Algebras and the Cayley-Hamilton Theorem .....                | 77        |
| 5.3       | Minimal Polynomials .....                                     | 78        |
| <b>6</b>  | <b>Normal Forms of Endomorphisms</b> .....                    | <b>79</b> |
| 6.1       | Decomposing Vector Spaces using Polynomials .....             | 79        |
| 6.2       | The Jordan Normal Form.....                                   | 82        |
| 6.3       | The Weierstrass Normal Form .....                             | 84        |
| <b>7</b>  | <b>Inner Product Spaces</b> .....                             | <b>85</b> |
| 7.1       | Inner Products.....                                           | 85        |

|             |                                                     |            |
|-------------|-----------------------------------------------------|------------|
| 7.2         | Orthogonality .....                                 | 87         |
| 7.3         | Matrix Representations of Inner Products .....      | 88         |
| 7.4         | Dual Spaces.....                                    | 88         |
| <b>8</b>    | <b>Normal Endomorphisms</b>                         | <b>92</b>  |
| 8.1         | Linear Isometries.....                              | 94         |
| 8.2         | Affine Isometries .....                             | 94         |
| <b>9</b>    | <b>Revisiting Synthetic Geometry</b>                | <b>96</b>  |
| 9.1         | Incidence Geometries .....                          | 96         |
| 9.2         | Projection Maps.....                                | 98         |
| 9.3         | Projectivization.....                               | 99         |
| <b>10</b>   | <b>Hyperbolic Geometry</b>                          | <b>101</b> |
| 10.1        | Circle Inversions.....                              | 101        |
| 10.2        | Möbius Transformations.....                         | 102        |
| 10.3        | Complex Möbius Geometry .....                       | 103        |
| <b>VI</b>   | <b>Group Theory</b>                                 | <b>106</b> |
| <b>1</b>    | <b>Discrete Symmetry Groups</b>                     | <b>107</b> |
| 1.1         | Finite subgroups of the orthogonal groups .....     | 107        |
| 1.2         | Lattice Groups .....                                | 108        |
| 1.3         | The Two-Dimensional Point Groups.....               | 108        |
| 1.4         | The Three-Dimensional Point Groups .....            | 109        |
| 1.5         | Classification of Lattices .....                    | 110        |
| 1.6         | Wallpaper Groups.....                               | 110        |
| 1.7         | Three-Dimensional Space Groups.....                 | 110        |
| <b>VII</b>  | <b>Multivariable Calculus</b>                       | <b>111</b> |
| <b>1</b>    | <b>Differentiation in <math>\mathbb{R}^n</math></b> | <b>112</b> |
| 1.1         | Differentials and Partial Derivatives .....         | 112        |
| 1.2         | Rules of Differentiation .....                      | 118        |
| 1.3         | Higher Partial Derivatives .....                    | 122        |
| 1.4         | Differential Operators.....                         | 123        |
| <b>2</b>    | <b>The Two Queens of Multivariable Calculus</b>     | <b>124</b> |
| 2.1         | The Inverse Function Theorem .....                  | 124        |
| 2.2         | The Implicit Function Theorem .....                 | 125        |
| <b>3</b>    | <b>Multiple Integrals</b>                           | <b>126</b> |
| <b>4</b>    | <b>Curves</b>                                       | <b>128</b> |
| <b>5</b>    | <b>Surfaces</b>                                     | <b>129</b> |
| <b>VIII</b> | <b>Point-Set Topology</b>                           | <b>130</b> |
| <b>1</b>    | <b>Topological Spaces</b>                           | <b>131</b> |
| 1.1         | Topologies .....                                    | 131        |
| 1.2         | Basic Separation Properties .....                   | 132        |
| 1.3         | Bases and Countability .....                        | 134        |
| <b>2</b>    | <b>New Spaces from Old</b>                          | <b>136</b> |
| 2.1         | The Subspace Topology.....                          | 136        |

|           |                                                              |            |
|-----------|--------------------------------------------------------------|------------|
| 2.2       | Product Spaces .....                                         | 139        |
| 2.3       | Disjoint Union Spaces.....                                   | 140        |
| 2.4       | Quotient Spaces.....                                         | 141        |
| 2.5       | Adjunction Spaces.....                                       | 142        |
| <b>3</b>  | <b>Topological Manifolds</b>                                 | <b>143</b> |
| 3.1       | Manifolds from Embeddings .....                              | 146        |
| <b>4</b>  | <b>Topologies on Algebraic Structures</b>                    | <b>149</b> |
| 4.1       | Topological Groups.....                                      | 149        |
| <b>5</b>  | <b>Connectedness</b>                                         | <b>150</b> |
| <b>6</b>  | <b>Compactness</b>                                           | <b>151</b> |
| <b>7</b>  | <b>Filters</b>                                               | <b>154</b> |
| 7.1       | Filter Convergence .....                                     | 154        |
| 7.2       | Tychonoff's Theorem .....                                    | 154        |
| <b>8</b>  | <b>Higher Separation Properties</b>                          | <b>157</b> |
| 8.1       | Regular Spaces.....                                          | 157        |
| 8.2       | Normal Spaces.....                                           | 160        |
| 8.3       | Urysohn's Metrization Theorem.....                           | 162        |
| <b>9</b>  | <b>Lattice Theory</b>                                        | <b>163</b> |
| 9.1       | Lattices.....                                                | 163        |
| 9.2       | Value Quantaes .....                                         | 164        |
| 9.3       | Generalized metrics metrize every topological space .....    | 164        |
| <br>      |                                                              |            |
| <b>IX</b> | <b>Algebraic Topology</b>                                    | <b>165</b> |
| <b>1</b>  | <b>Homotopy</b>                                              | <b>166</b> |
| 1.1       | The Fundamental Group.....                                   | 168        |
| <b>2</b>  | <b>Fibrations and Coverings</b>                              | <b>172</b> |
| <br>      |                                                              |            |
| <b>X</b>  | <b>Measure Theory</b>                                        | <b>175</b> |
| <b>1</b>  | <b>Measure Theory</b>                                        | <b>177</b> |
| 1.1       | The Measure Problem .....                                    | 177        |
| 1.2       | Lattices and Boolean Algebras .....                          | 180        |
| 1.3       | Set Algebras .....                                           | 183        |
| 1.4       | $\sigma$ -Algebras.....                                      | 185        |
| 1.5       | Dynkin Systems.....                                          | 188        |
| 1.6       | Monotone Classes .....                                       | 189        |
| 1.7       | Measures .....                                               | 191        |
| 1.8       | Carathéodory Extension .....                                 | 196        |
| 1.9       | Measures on $\mathbb{R}^n$ .....                             | 197        |
| <b>2</b>  | <b>Measurable Functions</b>                                  | <b>202</b> |
| <b>3</b>  | <b>Lebesgue Integration</b>                                  | <b>206</b> |
| 3.1       | Step Functions .....                                         | 206        |
| 3.2       | Defining the Lebesgue Integral.....                          | 208        |
| 3.3       | Comparing Riemann Integration and Lebesgue Integration ..... | 212        |
| 3.4       | Convergence Theorems .....                                   | 213        |
| 3.5       | $L^p$ -Spaces.....                                           | 215        |
| 3.6       | Density Functions .....                                      | 221        |

|          |                                                                           |            |
|----------|---------------------------------------------------------------------------|------------|
| <b>4</b> | <b>Integration in <math>\mathbb{R}^n</math></b>                           | <b>223</b> |
| 4.1      | Products of Measure Spaces.....                                           | 223        |
| 4.2      | Product Measures and Fubini's Theorem .....                               | 230        |
| 4.3      | Change of Variables .....                                                 | 238        |
| 4.4      | Alternative coordinate systems on $\mathbb{R}^2$ and $\mathbb{R}^3$ ..... | 241        |
| 4.5      | The Surface Area Measure.....                                             | 250        |
| 4.6      | The Divergence Theorem.....                                               | 256        |
| <b>5</b> | <b>Convolutions</b>                                                       | <b>259</b> |

**Part I**

**Foundations**

# Contents

|          |                                                  |          |
|----------|--------------------------------------------------|----------|
| <b>1</b> | <b>An Introduction to Mathematical Reasoning</b> | <b>8</b> |
| <b>2</b> | <b>Sets</b>                                      | <b>9</b> |
| 2.1      | Infinity: $\emptyset$ and $\mathbb{N}$ .....     | 10       |
| 2.2      | Extensionality .....                             | 10       |
| 2.3      | Specification .....                              | 11       |
| 2.4      | Pairing, Union, and Power Sets .....             | 11       |
| 2.5      | Relations and Functions .....                    | 11       |
| 2.6      | Cartesian Products .....                         | 11       |
| 2.7      | Replacement .....                                | 12       |
| 2.8      | Regularity and Well-Foundedness .....            | 12       |
| 2.9      | Choice .....                                     | 12       |
| 2.10     | Cardinality .....                                | 12       |

## **Chapter 1**

# **An Introduction to Mathematical Reasoning**

# Chapter 2

## Sets

Almost all of modern mathematics can be put in the context of the study of *sets* and their *elements*. Generally, given a collection of objects  $a, b, c, d, \dots$ , they can be taken together as a set, generally written in curly brackets and represented by a capital letter:

$$M := \{a, b, c, d, e, \dots\}$$

However, an approach where any collection of mathematical objects forms a set quickly runs into trouble - famously, *Russel's paradox* asks:

*"Let  $R$  be the set of all sets that don't contain themselves. Does  $R$  contain itself?"*

If  $R$  doesn't contain itself, then by definition it must contain itself, but if it contains itself, then it cannot contain itself! If  $R \in R$  is true, then it must be false! We arrive at a classic paradox. The conclusion, then, is that the set of all sets that don't contain themselves cannot exist - we cannot define sets to be any collection of things defined by any property, and must instead establish ground rules about which sets exist and exactly how we can use sets we already know to construct new ones.

These ground rules are given in the form of *axioms*, which are statements that mathematicians take for granted, which can then be used as a starting point to then derive "the rest of mathematics" from. The following chapters contain a brief and hopefully easily understandable introduction to the ZFC axiom system, which was developed in the early 20th century by Ernst Zermelo and Abraham Fraenkel and is now the most widely accepted axiomatization of mathematics. Informally stated, the ZFC axioms are:

1. The empty set is a set, and the natural numbers are a set.
2. Sets are uniquely determined by their elements.
3. Any definable subset of a set is a set.
4. Given two sets, there exists a set whose elements are those two sets.
5. Given two sets, there exists a set containing the elements of both sets.
6. Given a set, there exists a set containing all subsets of that set as elements.

7. The image of a set under a definable function is a set.
8. There are no infinitely nested sets.
9. Given any set of nonempty sets, there exists a function mapping each sets to one of its elements.

## 2.1 Infinity: $\emptyset$ and $\mathbb{N}$

## 2.2 Extensionality

To say “ $x$  is an element of  $M$ ”, or equivalently “ $x$  contains  $M$ ”, we simply write:

$$x \in M$$

One of the most basic axioms of set theory is the *Axiom of Extensionality*, which states:

### Axiom 2.1: Axiom of Extensionality

Two sets  $A$  and  $B$  are equal if and only if they contain the same elements.

The Axiom of Extensionality establishes that sets are uniquely defined by the objects they contain and have no additional structure. In particular:

1. The order of elements in a set doesn’t matter.  $\{1, 2\}$  and  $\{2, 1\}$  are the same set.
2. The number of times an element appears within the same set doesn’t matter.  $\{1\}$ ,  $\{1, 1\}$ , and  $\{1, 1, 1\}$  are the same set.

### Definition 2.2: Subsets

We call a set  $A$  a **subset** of a set  $B$  if and only if every element of  $A$  is contained in  $B$ . We write “ $A$  is a subset of  $B$ ” as  $A \subseteq B$ .

The demand that “every element of  $A$  is contained in  $B$ ” can be stated more formally as “ $x \in A$  implies  $x \in B$ ”. Importantly, this means that every set is a subset of itself - we always have  $A \subseteq A$ .

This definition gives us another way of phrasing the Axiom of Extensionality - two sets are equal if and only if each is a subset of the other:

$$A = B \Leftrightarrow A \subseteq B \text{ and } B \subseteq A$$

This idea is incredibly important for writing out proofs in practice - if a problem asks you to establish that two sets  $A$  and  $B$  are the same, then the easiest approach is almost always to show that any element of  $A$  must be contained in  $B$ , and to then show that any arbitrary element of  $B$  must be contained in  $A$ .

The subset relation is frequently written as  $A \subset B$ . However, other authors reserve  $A \subset B$  to mean “ $A$  is a subset of  $B$  that isn’t equal to  $B$ ” - we call such a

set a *proper subset* of  $B$ . I will therefore attempt to only use the symbols  $\subseteq$  and  $\subsetneq$ , which clear up this ambiguity.

## 2.3 Specification

## 2.4 Pairing, Union, and Power Sets

## 2.5 Relations and Functions

## 2.6 Cartesian Products

We can generalize the cartesian product to infinite products as follows:

### Definition 2.3: Cartesian Product of sets

Let  $I$  be any set. Let  $(X_i)_{i \in I}$  be a Family of nonempty sets indexed by  $I$ , i.e. there exists a function assigning to each element  $i \in I$  an element of  $(X_i)_{i \in I}$ . Then the cartesian product of  $(X_i)_{i \in I}$  is the unique set:

$$\prod_{i \in I} X_i := \left\{ x : I \rightarrow \bigcup_{i \in I} X_i \mid x_i := x(i) \in X_i \text{ for all } i \in I \right\}$$

Written more simply: Each element of  $\prod_{i \in I} X_i$  is a function assigning to each  $i \in I$  an element  $x(i) \in X_i$ , where we generally write the argument in the index ( $x_i := x(i)$ ), to stay consistent with the established notation in the countable case.

This definition is simply a reinterpretation of the definition of finite cartesian products: It amounts to viewing, for example, the tuple  $T := (x, y, z)$  as a map:

$$T : \{1, 2, 3\} \rightarrow \{x, y, z\}$$

such that  $T(1) = x$ ,  $T(2) = y$ , and  $T(3) = z$ .

### Definition 2.4: Projections

Let  $J \subset I$ . Then we define the projection of  $I$  to  $J$  as the map

$$\begin{aligned} \pi_J &:= \pi_J^I : \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_j \\ x &\mapsto \left( x|_J : J \rightarrow \bigcup_{j \in J} X_j \right) \end{aligned}$$

which restricts the domain of a given element  $x$  of the product to the subset  $J$ .

In particular, if  $J$  is a set consisting of a single element  $j$ , we have:

$$\begin{aligned} \pi_j &:= \pi_{\{j\}}^I : \prod_{i \in I} X_i \rightarrow X_j \\ x &\mapsto x_j \end{aligned}$$

## **2.7 Replacement**

## **2.8 Regularity and Well-Foundedness**

## **2.9 Choice**

## **2.10 Cardinality**

## **Part II**

# **Algebraic Structures**

# Contents

|          |                                                                 |           |
|----------|-----------------------------------------------------------------|-----------|
| <b>1</b> | <b><math>(\mathbb{N}, +)</math> is a Monoid</b>                 | <b>15</b> |
| <b>2</b> | <b><math>(\mathbb{Z}, +, -)</math> is a Group</b>               | <b>16</b> |
| 2.1      | Group Homomorphisms .....                                       | 16        |
| 2.2      | Permutations and the Symmetric Groups.....                      | 17        |
| 2.3      | Product Groups.....                                             | 18        |
| 2.4      | Group Actions .....                                             | 19        |
| <b>3</b> | <b><math>(\mathbb{Z}, +, -, \cdot)</math> is a Ring</b>         | <b>21</b> |
| <b>4</b> | <b><math>(\mathbb{Q}, +, -, \cdot, ^{-1})</math> is a Field</b> | <b>22</b> |

## **Chapter 1**

### **$(\mathbb{N}, +)$ is a Monoid**

## Chapter 2

# $(\mathbb{Z}, +, -)$ is a Group

### 2.1 Group Homomorphisms

**Theorem 2.1.** Let  $\varphi : (G, \circ_G) \rightarrow (H, \circ_H)$  be a map that respects group operations, i.e:

$$\varphi(g_1 \circ_G h_2) = \varphi(g_1) \circ_H \varphi(g_2)$$

Then  $\varphi$  also respects the neutral element and inverses, i.e:

$$\begin{aligned}\varphi(e_G) &= e_H \\ \varphi(g^{-1}) &= \varphi(g)^{-1}\end{aligned}$$

*Proof.* 1. For any  $h \in H$ , we have:

$$\begin{aligned}e_H &= e_H \circ_H e_H \\ &= (\varphi(e_G)^{-1} \circ_H \varphi(e_G)) \circ_H e_H \\ &= \varphi(e_G)^{-1} \circ_H \varphi(e_G) \\ &= \varphi(e_G)^{-1} \circ_H \varphi(e_G \circ_G e_G) \\ &= \varphi(e_G)^{-1} \circ_H \varphi(e_G) \circ_H \varphi(e_G) \\ &= e_H \circ_H \varphi(e_G) \\ &= \varphi(e_G)\end{aligned}$$

So  $\varphi$  respects the neutral element.

2. Since  $\varphi$  respects the neutral element, we get:

$$\begin{aligned}\varphi(g) \circ_H \varphi(g^{-1}) &= \varphi(g \circ_G g^{-1}) \\ &= \varphi(e_G) \\ &= e_H,\end{aligned}$$

i.e.  $\varphi(g^{-1}) = \varphi(g)^{-1}$ .

□

## 2.2 Permutations and the Symmetric Groups

## 2.3 Product Groups

### Theorem 2.2

Let  $G = G_1 \times G_2$  be a product group. Let  $H \subseteq G$  be a subgroup. Then  $H$  isn't necessarily isomorphic to a product  $H_1 \times H_2$  of subgroups of  $G_1$  and  $G_2$ .

*Proof.* Let  $G = S_3 \times S_3$ . Let  $H$  be subgroup of elements  $(\sigma_1, \sigma_2)$  with  $\text{sgn}(\sigma_2\sigma_1) = 1$ . We have  $|G| = 36$ , giving us  $|H| = 18$ .

The subgroups of  $S_3$  are  $S_3$  itself,  $C_1$ ,  $A_3 \cong C_3$  and  $C_2$ . If  $H$  was a product of subgroups, we would thus need to have  $H \cong A_3 \times S_3$ , since every other combination has the wrong order.

Now, notice that  $A_3 \times S_3$  contains elements of order 6, such as  $((1, 2), (1, 2, 3))$ . Since  $S_3$  isn't cyclic, it doesn't contain elements of order 6, so an element of order 6 in  $H$  must be a pair consisting of an element  $\sigma_2$  of order two (i.e. a transposition, which has odd order) and an element  $\sigma_3$  of order three (i.e. a three-cycle, which has even order). But then  $\text{sgn}(\sigma_2\sigma_1) = -1$ , so no such element can be contained in  $H$ .  $\square$

## 2.4 Group Actions

### Definition 2.3

Let  $(G, \circ, e)$  be a group. Let  $X$  be a set. Then a map  $\cdot : G \times X \rightarrow X$  is called a **group action** of  $G$  on  $X$  iff:

1. It respects the neutral element:

$$\begin{aligned} \forall x \in X : \\ e.x = x \end{aligned}$$

2. It associates with the group operation:

$$\begin{aligned} \forall g, h \in G, x \in X : \\ (g \circ h).x = g.(h.x) \end{aligned}$$

**Exercise 2.4.** Show that a map  $(g, x) \rightarrow g.x$  is a group operation if and only if the map  $g \rightarrow (x \rightarrow g.x)$  is a group homomorphism from  $G$  into the symmetric group over  $X$ .

**Definition 2.5.** Let  $\cdot : G \times X \rightarrow X$  be an action of a group  $G$  on a set  $X$ . Let  $x \in X$ . Then we call the set

$$Gx := \{g.x \mid g \in G\}$$

of all points that  $x$  can be taken to by an element of  $G$  the **orbit** of  $x$  under  $G$ .

**Exercise 2.6.** Let  $X = \mathbb{R}^2$  be the plane. Let  $G$  be the group of rotations around the origin. Sketch the orbits of different points  $x \in X$ .

**Definition 2.7.** Let  $\cdot : G \times X \rightarrow X$  be an action of a group  $G$  on a set  $X$ . Then we call the set

$$X/G := \{Gx \mid x \in X\}$$

of all orbits under  $G$  the **orbit space** of  $G$  in  $X$ .

**Exercise 2.8.** Show that the relation

$$x \sim y \Leftrightarrow \exists g \in G : g.x = y$$

is an equivalence relation. Show that for any  $x \in X$ , the equivalence class of  $x$  is exactly the orbit of  $x$ .

**Definition 2.9.** Let  $\cdot : G \times X \rightarrow X$  be an action of a group  $G$  on a set  $X$ . Then we call the action **transitive** if any point  $x \in X$  can be taken to any point  $y \in X$  by an element of  $G$ :

$$\begin{aligned} \forall x, y \in X : \\ \exists g \in G : \\ g.x = y \end{aligned}$$

**Exercise 2.10.** Show that a group action is transitive if and only if  $Gx = X$  for all  $x \in X$ .

**Definition 2.11.** Let  $\cdot : G \times X \rightarrow X$  be an action of a group  $G$  on a set  $X$ . Let  $x \in X$ . Then we call the set

$$G_x := \{g \in G \mid g.x = x\}$$

of all elements of  $G$  leaving  $x$  unchanged the *isotropy group* of  $x$ .

**Exercise 2.12.** Show that the isotropy group  $G_x$  is always a subgroup of  $G$ .

**Exercise 2.13.** Show that if the action  $(g, x) \mapsto g.x$  is transitive, the map

$$\begin{aligned} \varphi_x : G/G_x &\rightarrow X \\ [g] &\mapsto g.x, \end{aligned}$$

where  $G/G_x$  is the quotient group of  $G$  by  $G_x$ , is well-defined and bijective.

We can also define the isotropy groups of points:

**Definition 2.14.** Let  $\cdot : G \times X \rightarrow X$  be an action of a group  $G$  on a set  $X$ . Let  $A \subseteq X$ . Then we define

$$g.A := \{g.a \mid a \in A\}$$

and call the set

$$G_A := \{g \in G \mid g.A = A\}$$

of all elements of  $G$  mapping elements of  $A$  to elements of  $A$  the *isotropy group* of  $A$ .

## Chapter 3

**$(\mathbb{Z}, +, -, \cdot)$  is a Ring**

## Chapter 4

**$(\mathbb{Q}, +, -, \cdot, ^{-1})$  is a Field**

## **Part III**

# **Foundations of Analysis**

# Contents

|          |                                                 |           |
|----------|-------------------------------------------------|-----------|
| <b>1</b> | <b>The Real Numbers <math>\mathbb{R}</math></b> | <b>25</b> |
| 1.1      | Ordered Algebraic Structures .....              | 25        |
| 1.2      | Dedekind Cuts.....                              | 25        |
| 1.3      | Uniqueness of $\mathbb{R}$ .....                | 25        |
| 1.4      | Cardinality of $\mathbb{R}$ .....               | 25        |
| <b>2</b> | <b>The Concept of Distance</b>                  | <b>26</b> |
| 2.1      | Metric Spaces.....                              | 26        |
| 2.2      | Topologies .....                                | 29        |
| 2.3      | Convergence .....                               | 33        |
| 2.4      | Continuity .....                                | 34        |
| 2.5      | Compactness.....                                | 35        |
| 2.6      | Connectedness.....                              | 38        |
| 2.7      | Function Spaces.....                            | 40        |
| 2.8      | Banach's Fixed Point Theorem .....              | 43        |
| <b>3</b> | <b>Infinite Series</b>                          | <b>44</b> |
| 3.1      | Achilles and the Turtle.....                    | 44        |
| 3.2      | Convergence Tests.....                          | 44        |
| 3.3      | Euler's Number $e$ .....                        | 44        |
| 3.4      | The Exponential Function.....                   | 44        |
| <b>4</b> | <b>Euclidean Geometry</b>                       | <b>45</b> |
| 4.1      | The Complex Numbers $\mathbb{C}$ .....          | 45        |
| 4.2      | Circles, Angles and Trigonometry .....          | 45        |

# Chapter 1

## The Real Numbers $\mathbb{R}$

### 1.1 Ordered Algebraic Structures

### 1.2 Dedekind Cuts

### 1.3 Uniqueness of $\mathbb{R}$

### 1.4 Cardinality of $\mathbb{R}$

## Chapter 2

# The Concept of Distance

### 2.1 Metric Spaces

A metric space is a set, alongside with a *distance function*, more commonly known as a *metric*,  $d$  which takes two points and outputs a distance between the points. Naturally, we want this "distance" to actually give us some usable structure on the set.

Now, what pieces of geometric intuition should we bake into our notion of distance? I would argue that the most fundamental use of "distance" is to be able to *compare distances between any two pairs of points*. Therefore, our metric should have as its output a *totally ordered set*.

We also want to be able to add distances together as a way of comparing them - logically, the distance between two points  $x$  and  $y$  plus the distance between  $y$  and  $z$  gives us an upper bound to the distance between  $x$  and  $z$ , since we can always take this path to get from  $x$  to  $z$ , but there might be a shorter path we haven't found yet. As a formula, this is the *triangle inequality*

$$d(x, z) \leq d(x, y) + d(y, z)$$

Finally, if we have an ordered group structure, we would really like for our order to be archimedean - the distance between two distinct points should be greater than zero, and it shouldn't be infinitesimal.

This leaves  $\mathbb{R}_{\geq 0}$  as the obvious candidate for the domain of our distance function, since any archimedean ordered group is a subgroup of  $\mathbb{R}_{\geq 0}$  anyways. This shouldn't be too much of a surprise, even without this logical background - I assume that when most people read "distance between two points", they intuitively already assume that these distances will be given by positive real numbers.

### Definition 2.1: Metric Spaces

A **metric space** is a set  $M$  together with a function  $d : M \times M \rightarrow \mathbb{R}_{\geq 0}$ , such that:

1.  $d$  is **positive definite**, i.e.  $d(x, y) = 0$  if and only if  $x = y$ .
2.  $d$  is **symmetric**, i.e.  $d(x, y) = d(y, x)$
3.  $d$  fulfills the **triangle inequality**

$$d(x, z) \leq d(x, y) + d(y, z)$$

**Theorem 2.2.** Let  $M^n := \prod_{i=1}^n M$  be a finite cartesian product of metric spaces. Then the **Euclidean distance function**

$$d_\infty(x, y) := \sqrt{\sum_{i=1}^n (x_i - y_i)^2}$$

gives a metric on  $M^n$ .

**Theorem 2.3.** Let  $M^n := \prod_{i=1}^n M$  be a finite cartesian product of metric spaces. Then the **Chebyshev distance function**

$$d_\infty(x, y) := \max_{i \in [1..n]} d(x_i - y_i)$$

gives a metric on  $M^n$ .

### Definition 2.4: Balls

Let  $(M, d)$  be a metric space. Then the **open ball of radius  $r$  around a point  $x$**  is the set

$$B_r(x) := \{y \in M : d(x, y) < r\},$$

and the **closed ball of radius  $r$  around  $x$**  is the set

$$\overline{B}_r(x) := \{y \in M : d(x, y) \leq r\},$$

### Definition 2.5: Open and Closed Sets

Let  $(M, d)$  be a metric space. Then we call a subset  $U \subseteq M$  **open** (in  $M$ ) if it contains an open ball (of any radius) around all of its points, i.e.

$$\begin{aligned} \forall x \in U : \\ \exists r \in \mathbb{R}_{\geq 0} : \\ B_r(x) \subseteq U. \end{aligned}$$

We call  $U$  **closed** if its complement  $M \setminus U$  is open.

Intuitively, a set is open if it contains "a little bit of wiggle room" around any point contained in it. This means that sets with a "clear border", such as our closed balls, aren't open. For example, the interval  $[0, 1] \subset \mathbb{R}$  doesn't leave any wiggle room around 0 and 1.

Importantly, while it may seem unintuitive, there are in fact sets that are both closed and open at the same time. We call such sets *clopen*.

**Exercise 2.6.** *There are exactly two clopen subsets of  $\mathbb{R}$ . What are they?*

**Definition 2.7.** Let  $(M, d)$  be a metric space. We call a subset  $N \subseteq M$  *dense in  $M$*  if every nonempty open subset of  $M$  contains an element of  $N$ .

**Lemma 2.8.** Every nonempty open subset of  $\mathbb{R}$  is uncountable.

*Proof.* 1. Since every nonempty open subset of  $\mathbb{R}$  contains an open interval  $(a, b)$ , it suffices to show that open intervals are uncountable.

2. The map

$$\begin{aligned}\varphi_1 : (a, b) &\rightarrow \left(-\frac{b-a}{2}, \frac{b-a}{2}\right) \\ x &\mapsto x - \frac{b+a}{2}\end{aligned}$$

is a bijection.

3. The map

$$\begin{aligned}\varphi_2 : \left(-\frac{b-a}{2}, \frac{b-a}{2}\right) &\rightarrow (-1, 1) \\ x &\mapsto \frac{2x}{b-a}\end{aligned}$$

is also a bijection.

4. Finally, the map

$$\begin{aligned}\varphi_3 : (-1, 1) &\rightarrow \mathbb{R} \\ x &\mapsto \frac{x}{1-|x|}\end{aligned}$$

is a bijection (the inverse is the map we get by replacing the  $-$  with a  $+$ ).

5. Therefore,  $\varphi_3 \circ \varphi_2 \circ \varphi_1$  is a bijection  $(a, b) \cong \mathbb{R}$ , so any open interval is uncountable, so any open subset of  $\mathbb{R}$  is uncountable.  $\square$

### Theorem 2.9

The set of rational numbers  $\mathbb{Q}$  is dense in  $\mathbb{R}$ . The set of irrational numbers  $\mathbb{R} \setminus \mathbb{Q}$  is also dense in  $\mathbb{R}$ .

*Proof.* 1. Since any nonempty open subset  $U \subseteq \mathbb{R}$  contains an open interval  $(a, b)$ , it suffices to show that any open interval contains a rational number. We assume without loss of generality that  $0 \leq$

$a < b$ , since  $a < 0 < b$  already implies  $0 \in (a, b)$ , while  $a < b \leq 0$  implies that  $(a, b)$  contains a rational number if and only if  $(-b, -a)$  contains a rational number.

Now, since the ordering on the reals is archimedean, we can find a natural number  $n \in \mathbb{N}$  such that  $\frac{1}{n} < b - a$ . Now, the set

$$L = \left\{ k \in \mathbb{N} \mid \frac{k}{n} > a \right\}$$

is nonempty - if  $a = 0$ , then  $1 \in M$ , and if  $a \neq 0$ , then the archimedean property tells us that there exists a  $k \in \mathbb{N}$  such that

$$\frac{1}{k} < \frac{1}{na},$$

which implies  $\frac{k}{n} > a$ . Since  $M$  is a nonempty subset of  $\mathbb{N}$ , there must be a smallest element  $m$ . Since  $m$  is minimal, we have  $m - 1 \notin M$ , which implies

$$\frac{m-1}{n} \leq a,$$

which in turn implies

$$\frac{m}{n} = \frac{m-1}{n} + \frac{1}{n} < a + (b-a) = b$$

Therefore,  $\frac{m}{n}$  is a rational number contained in  $(a, b)$ , so  $\mathbb{Q}$  is dense in  $\mathbb{R}$ .

2. Since any open subset  $U \subseteq \mathbb{R}$  is uncountable, while the rationals are countable,  $U$  must contain an uncountable number of irrationals. Therefore,  $\mathbb{R} \setminus \mathbb{Q}$  is dense in  $\mathbb{R}$ .

□

### Corollary 2.10

The set of rational numbers  $\mathbb{Q}$  is neither closed nor open in  $\mathbb{R}$ .

*Proof.* Since both sets are dense in  $\mathbb{R}$ , every open ball containing a rational number must also contain an irrational number, and vice versa. □

## 2.2 Topologies

Topological spaces generalize metric spaces. By inheriting the concept of open and closed sets from metric spaces, they are able to still capture notions of closeness without putting a numerical value on exactly how close any given pair of points are. The notion of a topology is based on the following insight:

**Exercise 2.11.** Let  $(M, d)$  be a metric space. Then:

1. *The empty set is open in  $M$ .*
2. *Any union of sets that are open in  $M$  is open in  $M$ .*
3. *Any finite intersection of sets that are open in  $M$  is open in  $M$ .*
4.  *$M$  is open in  $M$ .*

We take this idea and make it a definition:

**Definition 2.12**

Let  $X$  be a set. Then a **topology on  $X$**  is a set  $\mathcal{O}$  of subsets of  $X$ , called **open sets**, such that:

1. The empty set is open.
2. Any union of open sets is open.
3. Any finite intersection of open sets is open.
4.  $X$  is open.

Most of the topological spaces we care about in practice are already metric spaces, but many proofs flow more elegantly when phrased in the generalized language of topological spaces, and there *are* topological spaces that aren't metric spaces.

**Theorem 2.13: Extremal Topologies**

Let  $X$  be an arbitrary set. Then the following two sets of subsets are always topologies on  $X$ :

1. The **indiscrete topology**, where the only open sets are  $X$  and the empty set.
2. The **discrete topology**, where every subset is open.

*Proof Sketch.* Both topologies fulfill all requirements trivially. :) "□"

**Definition 2.14.** We say that a metric  $d$  on a space  $M$  **generates** a given topology  $\mathcal{O}$  on  $M$  if they give the same open sets, i.e. the open sets in  $\mathcal{O}$  are exactly the open sets in  $(M, d)$ .

**Exercise 2.15.**

1. Show that the discrete metric generates the discrete topology.
2. Show that there is no metric generating the indiscrete topology.

Two different metrics can generate the same topology:

**Exercise 2.16.** Show that the euclidean metric also generates the discrete topology on  $\mathbb{N}$ .

**Exercise 2.17.** Let  $d$  be a metric on a set  $M$ . Let  $c \in \mathbb{R}_{>0}$ .

1. Show that

$$d_c(x, y) = c \cdot d(x, y)$$

is a metric on  $M$ .

2. Show that  $d$  and  $d_c$  generate the same topologies on  $M$ .

This fact can be seen as a formalization of the intuition that changing the specific unit we choose to measure space with doesn't change its underlying structure.

#### Definition 2.18: Interior, Exterior and Boundary

Let  $X$  be a topological space. Let  $A \subseteq X$ . Then:

1. The **interior**  $\text{int } A$  of  $A$  is the set of all points that have an open neighbourhood entirely contained in  $A$ .
2. The **exterior**  $\text{ext } A$  of  $A$  is the interior of the complement  $X \setminus A$ , i.e. the set of all points that have an open neighbourhood entirely contained in  $X \setminus A$ .
3. The **boundary**  $\partial A$  of  $A$  is the set of all points  $x \in X$  lying neither in  $\text{int } A$ , nor in  $\text{ext } A$ .

The interior of  $A$  is occasionally written as  $\overset{\circ}{A}$ . Intuitively, the interior is the largest open subset of  $A$ , and the exterior is the largest open set not containing any point of  $A$ .

**Exercise 2.19.** Show that:

1. The interior of any set is open.
2. The exterior of any set is open.
3. The interior of  $A$  is the union of all open subsets of  $A$ .
4. The exterior of  $A$  is the union of all open subsets of  $X \setminus A$ .
5. The boundary of  $A$  is the set of all points  $x \in X$  such that every neighbourhood of  $x$  contains both points in  $A$  and points in  $X \setminus A$ .

#### Definition 2.20: Closure

The closure of  $A$  is the set  $\overline{A} := A \cup \partial A$  containing  $A$  and all of its boundary points.

**Exercise 2.21.**

1. Show that  $\overline{A}$  is the set of all points  $x \in X$  such that every neighbourhood of  $x$  contains points in  $A$ .
2. Show that  $\overline{A}$  is closed.
3. Show  $\overline{A}$  is the intersection of all closed sets containing  $A$ .

**Theorem 2.22.** Let  $B \subseteq \mathbb{R}$  be a bounded subset of  $\mathbb{R}$ . Then  $\inf B$  and  $\sup B$  are contained in  $\overline{B}$ .

*Proof.* Since  $B$  is bounded, completeness of the real numbers tells us that  $\inf B$  and  $\sup B$  are finite real numbers. If  $\inf B \notin B$ , then there would exist an open neighbourhood  $U_{\inf}$  of  $\inf B$  not containing any points of  $B$ . Since any neighbourhood of a real number  $x$  must contain numbers greater than  $x$ , this would give us a lower bound of  $B$  greater than  $\inf B$ , which contradicts the definition of  $\inf B$ . Similarly, if  $\sup B \notin \overline{B}$ , we could get a lesser upper bound of  $B$ , which is again a contradiction.  $\square$

## 2.3 Convergence

### Theorem 2.23: Detecting openness and closedness using sequences

Let  $M$  be a metric space. Let  $A \subseteq M$ . Let  $x \in M$ . Then:

1.  $x \in \text{int } A$  if and only if every sequence in  $M$  converging to  $x$  is eventually in  $A$ .
2.  $x \in \overline{A}$  if and only if there exists a sequence in  $A$  converging to  $x$ .

*Proof.* 1. If  $x \in \text{int } A$ , then there exists an  $\varepsilon > 0$  such that  $B_\varepsilon(x) \subseteq A$ . If a sequence converges to  $x$ , it must thus be eventually in  $B_\varepsilon(x)$ , and thus eventually in  $A$ .

Vice versa, if  $x \notin \text{int } A$ , then every open ball  $B_n := B_{\frac{1}{n}}(x)$  around  $A$  contains at least one point not in  $A$ , and so we can construct a sequence converging to  $x$  that isn't eventually in  $A$  by picking these points.

2. If there exists sequence of points in  $A$  converging to  $x$ , then every neighbourhood of  $x$  must contain at least one point in  $A$ , so  $x \in \overline{A}$ .

Vice versa, if  $x \in \overline{A}$ , then every open ball around  $x$  must contain a point from  $A$ , so we can construct a sequence in  $A$  converging to  $x$  by taking the sequence  $B_n := B_{\frac{1}{n}}(x)$  of open balls and picking a point  $x_n \in A \cap B_n$  for each  $n \in \mathbb{N}$ .

□

### Corollary 2.24.

1.  $A$  is closed in  $X$  if and only if it contains every limit of every convergent sequence of points in  $A$ .
2.  $A$  is open in  $X$  if and only if every sequence in  $X$  converging to a point of  $A$  is eventually in  $A$ .

## 2.4 Continuity

**Definition 2.25.** Let  $f : M \rightarrow N$  be a function between metric spaces. Then we call  $f$  *uniformly continuous* if it is continuous and fulfills the additional requirement that, given an  $\varepsilon$ , we can pick the same  $\delta$  for all points  $x \in M$ , i.e:

$$\forall \varepsilon \in \mathbb{R}_{>0} :$$

$$\exists \delta \in \mathbb{R}_{>0} :$$

$$\forall x, y \in M :$$

$$d(x, y) < \delta \implies d(f(x), f(y)) < \varepsilon$$

## 2.5 Compactness

**Exercise 2.26.** Any closed subset of a compact set is compact.

**Exercise 2.27.** A compact subset of a metric space is always closed.

### Theorem 2.28: Compactness in Metric Spaces

1. A subset of a metric space is compact if and only if it is sequentially compact.
2. A compact subset of a metric space is closed and bounded.

*Proof.* 1.

2. Let  $K \subseteq M$  be compact. Since  $M$  is a metric space,  $K$  is closed. If we cover  $K$  using open balls, that cover must have a finite subcover. Thus,  $K$  must be contained in a finite union of open balls, which means  $K$  must be bounded.

□

**Theorem 2.29.** Any closed interval  $I = [a, b] \subset \mathbb{R}$  is compact.

**Theorem 2.30.** Let  $f : M \rightarrow N$  be a continuous function between metric spaces. Let  $K \subseteq M$  be compact. Then  $f(K)$  is compact.

*Proof.* Let  $K \subseteq M$  be compact. Let  $\mathcal{U}$  be an open cover of  $f(K)$ . Since  $f$  is continuous,  $f^{-1}[\mathcal{U}] := \{f^{-1}[U] \mid U \in \mathcal{U}\}$  is an open cover of  $K$ , which must have a finite subcover  $\mathcal{V} \subseteq f^{-1}[\mathcal{U}]$ . Since  $f : K \rightarrow f(K)$  is surjective by definition,  $f[\mathcal{V}]$  must be a finite cover of  $f(K)$ . □

### Corollary 2.31: Maximum-Minimum-Theorem

Let  $f : K \rightarrow \mathbb{R}$  be a function from a compact subset of a metric space into the real numbers. Then  $f$  attains a maximum and a minimum.

*Proof.* By the previous theorem,  $f(K)$  is a compact subset of  $\mathbb{R}$ , i.e. closed and bounded by Theorem 2.28. But then Theorem 2.22 tells us that  $\inf f(K)$  and  $\sup f(K)$  are already contained in  $\overline{f(K)} = f(K)$ . □

**Lemma 2.32.** Let  $M, N$  be metric spaces. Let  $K \subseteq M$  be compact. Let  $f : K \rightarrow N$  be continuous. Then the preimage of any compact subset of  $N$  is a compact subset of  $K$ .

*Proof.* Since  $N$  is a metric space, compact subsets are closed. Since  $f$  is continuous, the preimage of a closed subset is closed. Since  $K$  is compact, any closed subset of  $K$  is compact.  $\square$

### Theorem 2.33: Countable Metric Tychonoffs Theorem

Let  $M_1, M_2$  be metric spaces. Let  $K_1 \subseteq M_1, K_2 \subseteq M_2$  be compact. Then  $K_1 \times K_2 \subseteq M_1 \times M_2$  is compact.

*Proof.* Using whichever bijection  $\mathbb{N}^2 \cong \mathbb{N}$  you prefer, we can unroll a sequence  $s_n$  in  $K_1 \times K_2$  into a sequence  $x_n$  in  $K_1$  and a sequence  $y_n$  in  $K_2$ , such that the elements of  $s_n$  are in one-to-one correspondence with pairs  $(x_i, y_j)$  of elements of  $x_n$  and  $y_n$ . Since  $K_1$  and  $K_2$  are compact,  $x_n$  and  $y_n$  have convergent subsequences. Using our bijection, these can be reassembled into a subsequence of  $s_n$ , and since both components of this subsequence converge, the subsequence itself converges. Thus, every sequence in  $K_1 \times K_2$  has a convergent subsequence. Since sequential compactness and compactness are equivalent in metric spaces by Theorem 2.28,  $K_1 \times K_2$  is compact.  $\square$

### Theorem 2.34: Heine-Borel Theorem

Every closed and bounded subset of  $\mathbb{R}^n$  is compact.

*Proof.* A bounded set is a subset of a cube  $C^n = \prod_{i=1}^n [-r, r]$ . Such a cube is a product of closed intervals, which are compact by Theorem 2.29. Every set that is closed and bounded is thus a closed subset of a compact set, making them compact by Exercise 2.26.  $\square$

**Theorem 2.35.** Let  $M, N$  be metric spaces. Let  $K \subseteq M$  be compact. Then every continuous function  $f : K \rightarrow N$  is uniformly continuous.

*Proof.* Assume that  $f : K \rightarrow N$  isn't uniformly continuous. Then there exists an  $\varepsilon \in \mathbb{R}_{>0}$  such that for every  $n \in \mathbb{N}$ , we can find points  $x_n, y_n \in K$  such that  $d(x_n, y_n) < 1/n$ , but  $d(f(x_n), f(y_n)) \geq \varepsilon$ . Since  $K$  is compact, it is sequentially compact by Theorem 2.28, so there exists a subsequence  $(x_k) \subset (x_n)$  converging to a point  $p$ . Since  $d(x_n, y_n) < 1/n$ , there also exists a subsequence  $x_k \subset y_n$  converging to  $p$ . If  $f$  was continuous, we would have

$$\lim_{k \rightarrow \infty} f(x_k) = \lim_{k \rightarrow \infty} f(y_k),$$

contradicting the fact that we have  $d(f(x_k) - f(y_k)) \geq \varepsilon$  for all  $k \in \mathbb{N}$ .  $\square$

## 2.6 Connectedness

### Definition 2.36

A metric space is called **connected** if it can't be written as the disjoint union of two nonempty open subsets.

**Exercise 2.37.** 1. Show that  $\mathbb{N}$  is not connected.

2. Show that  $\mathbb{Q}$  is not connected.

**Exercise 2.38.** Show that a metric space  $M$  is connected if and only if its only clopen subsets are  $M$  and  $\emptyset$ .

**Theorem 2.39.** The connected subsets of  $\mathbb{R}$  are the intervals  $I \subseteq \mathbb{R}$ .

*Proof.* 1. Let  $I \subseteq \mathbb{R}$  be an interval. Assume we have a decomposition

$$I = I_1 \cup I_2$$

of  $I$  into disjoint nonempty open subsets. Without loss of generality, assume that there exists  $x_1 \in I_1$  and  $x_2 \in I_2$  such that  $x_1 < x_2$  (otherwise swap the two subsets). Now, consider the set

$$S = \{x \in I_1 \mid x \leq x_2\}$$

$S$  is nonempty, since  $x_1 \in S$ , and  $S$  has  $x_2$  as an upper bound. Now, let  $s = \sup S$ . We have  $x_1 \leq s \leq x_2$ , so  $s \in I$ . By Theorem 2.22, we have  $s \in \bar{S} \subseteq \bar{I}_1$ .

Since we have  $I_1 = I \setminus I_2$ , we know that  $I_1$  is closed in  $I$ , giving us  $s \in \bar{I}_1 = I_1$ .

Since  $I_1$  is by definition also open in  $I$ , and  $s \leq x_2$  cannot be an upper bound of  $I$ , there exists  $\varepsilon \in \mathbb{R}_{>0}$  such that  $s + \varepsilon \in I_1$ , which contradicts  $s = \sup I_1$ . Therefore, all intervals must be connected.

2. Let  $C \subseteq \mathbb{R}$  be connected. If  $C$  is empty or only contains one point, it vacuously qualifies as an interval  $(x, x)$  or  $[x, x]$ . Now, assume that  $C$  contains at least two points, but isn't an interval. Then there exist  $c_1 \neq c_2 \in C$  and  $c \in [c_1, c_2]$  such that  $c \notin C$ . But then

$$C_1 = (-\infty, c) \cap C$$

$$C_2 = (c, \infty) \cap C$$

is a decomposition of  $C$  into disjoint nonempty open subsets.

□

**Theorem 2.40.** Let  $f : M \rightarrow N$  be a continuous map between metric spaces. Let  $M$  be connected. Then  $f(M)$  is connected.

*Proof.* Assume  $f(M)$  wasn't connected. Then we would have some decomposition  $f(M) = U \cup V$  of  $f(M)$  into disjoint open sets. Since  $f$  is continuous,  $f^{-1}(U)$  and  $f^{-1}(V)$  are open. But then  $M = f^{-1}(U) \cup f^{-1}(V)$  can't be connected.  $\square$

**Corollary 2.41: Intermediate Value Theorem**

Let  $f : [a, b] \rightarrow \mathbb{R}$  be a continuous map from a closed real interval into the real numbers. Let  $y$  be some number between  $f(a)$  and  $f(b)$ <sup>1</sup>. Then there exists  $x \in [a, b]$  such that  $f(x) = y$ .

<sup>1</sup> i.e.  $f(a) \leq y \leq f(b)$  or  $f(b) \leq y \leq f(a)$

*Proof.* Since  $f$  is continuous,  $f([a, b])$  is a connected subset of  $\mathbb{R}$ , i.e. an interval. Since  $f([a, b])$  contains  $f(a)$  and  $f(b)$ , it must therefore also contain all numbers in between the two.  $\square$

**Corollary 2.42: Bolzano's Theorem**

If a continuous function  $f : M \rightarrow \mathbb{R}$  has a point  $x \in M$  with  $f(x) < 0$  and a point  $y \in M$  with  $f(y) > 0$ , there exists a root  $z \in M : f(z) = 0$  between  $x$  and  $y$ .

- Exercise 2.43.**
1. Assuming that temperature varies continuously, show that there are always two points at a given latitude on earth at which the temperature is the same.
  2. Assuming there are two points on earth that are a given distance apart from each other, are there always two points on earth that are that distance apart from each other and have the same temperature?

## 2.7 Function Spaces

Since we know that a sequence in a finite cartesian product converges if and only if all components converge, it seems natural to generalize this notion to arbitrary, potentially infinite cartesian products, i.e. arbitrary functions:

### Definition 2.44: Pointwise Convergence

We say that a sequence of functions  $f_n : M \rightarrow N$  converges *pointwise* to a given function  $f : M \rightarrow N$  if and only if all components  $f_n(x)$  converge in the usual sense, i.e:

$$\begin{aligned} \forall x \in M : \\ \forall \varepsilon \in \mathbb{R}_{>0} : \\ \exists N \in \mathbb{N} : \\ \forall n \geq N : \\ d(f_n(x) - f(x)) < \varepsilon \end{aligned}$$

We can't put a metric on  $N^M$  in which this form of converges matches regular metric convergence, but the intuition that this is the most natural form of convergence in  $N^M$  will be vindicated once we study abstract product spaces in point-set topology, at which point we confirm that in the natural product topology, an infinite cartesian product still converges if and only if every component converges individually.

However, there *is* a natural metric to put on  $N^M$ , which leads to a stronger form of convergence:

**Definition 2.45.** An *extended metric space* is a metric space where  $d$  can also take on the value  $\infty$ .

### Theorem 2.46: Supremum Metric

The set  $N^M$  of functions from a metric space  $M$  to a metric space  $N$  can be metrized by the extended metric

$$d_\infty(f, g) := \sup_{x \in M} |f(x) - g(x)|.$$

Additionally, this function is a regular metric on the set  $B(M, N) \subset N^M$  of bounded functions  $M \rightarrow N$ .

Note that is simply a direct generalization of the Chebyshev metric 2.3 to infinite cartesian products!

**Theorem 2.47: Uniform Convergence**

In  $(N^M, d_\infty)$ , a sequence of functions  $f_n$  converges to a given function  $f$  if and only if

$$\begin{aligned} &\forall \varepsilon \in \mathbb{R}_{>0} : \\ &\exists N \in \mathbb{N} : \\ &\forall n \geq N : \\ &\forall x \in M : \\ &\quad d(f_n(x) - f(x)) < \varepsilon, \end{aligned}$$

i.e. if the speed at which  $f_n(x)$  converges to  $f(x)$  doesn't depend on the specific  $x$  we choose. We call this form of convergence of a sequence of functions *uniform convergence*.

**Theorem 2.48.** Let  $(f_n)_{n \in \mathbb{N}}$  be a sequence of continuous functions  $f_n : M \rightarrow N$  converging uniformly to a function  $f : M \rightarrow N$ . Then  $f$  is continuous.

*Proof.* Choose some  $\varepsilon \in \mathbb{R}_{>0}$ . Since  $(f_n)_{n \in \mathbb{N}}$  converges to  $f$  uniformly, there exists an  $N \in \mathbb{N}$  such that

$$\begin{aligned} &\forall x \in M : \\ &\quad n \geq N \implies d(f(x), f_n(x)) < \varepsilon/3 \end{aligned}$$

Let  $x_0 \in M$  be arbitrary. Since  $f_N$  is continuous, there exists a  $\delta \in \mathbb{R}_{>0}$  such that

$$\begin{aligned} &\forall x \in M : \\ &\quad d(x, x_0) < \delta \implies d(f_N(x), f_N(x_0)) < \varepsilon/3 \end{aligned}$$

Applying the triangle inequality twice gives

$$\begin{aligned} &d(f(x), f(x_0)) \\ &\leq d(f(x), f_N(x)) + d(f_N(x), f_N(x_0)) + d(f_N(x_0), f(x_0)) \end{aligned}$$

to which we can apply the first inequality, then the second, and then the first again, to get:

$$\begin{aligned} &d(f(x), f(x_0)) \\ &\leq d(f(x), f_N(x)) + d(f_N(x), f_N(x_0)) + d(f_N(x_0), f(x_0)) \\ &< \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon \end{aligned}$$

for all  $x \in M$  with  $d(x, x_0) < \delta$ . □

**Theorem 2.49.** Let  $U \subseteq (X, d)$  be open and path-connected. Then there exists a piecewise linear path  $\gamma_n : [0, 1] \rightarrow U$  between any two  $p, q \in U$ .

*Proof Sketch.* 1. Since  $U$  is path connected, there exists a curve  $\gamma : [0, 1] \rightarrow U$  connecting  $p$  and  $q$ .

2. We can approximate  $\gamma$  via a piecewise linear curve  $\gamma_n$  by setting

$$\gamma_n\left(\frac{x}{n}\right) := \gamma\left(\frac{x}{n}\right)$$

3. Since  $\gamma$  is continuous and defined on a compact set, it is uniformly continuous by Theorem 2.35. From this, it can be shown that the  $\gamma_n$  converge to  $\gamma$  uniformly.

4. Since  $\gamma([0, 1]) \subseteq U$  is compact and  $X \setminus U$  is closed and disjoint from  $U$ , we have  $\varepsilon := d(\gamma([0, 1]), X \setminus U) > 0$ .

5. Therefore, uniform convergence guarantees there is an  $n$  such that  $d(\gamma_n([0, 1]), \gamma([0, 1])) < \varepsilon$ , which implies  $\gamma_n([0, 1]) \subseteq U$ .

”□”

## **2.8 Banach's Fixed Point Theorem**

## Chapter 3

# Infinite Series

3.1 Achilles and the Turtle

3.2 Convergence Tests

3.3 Euler's Number  $e$

3.4 The Exponential Function

## Chapter 4

# Euclidean Geometry

### 4.1 The Complex Numbers $\mathbb{C}$

### 4.2 Circles, Angles and Trigonometry

## **Part IV**

# **Calculus in one Variable**

## **Chapter 1**

# **Derivatives**

## Chapter 2

# Integrals

### 2.1 The Gamma Function

In this section, we discover some basic properties of the so-called gamma function. We will find that it is the canonical extension of the factorial function to positive real numbers. With some knowledge of complex analysis, it can easily be extended to arbitrary complex numbers, but that is beyond the scope of these notes.

**Definition 2.1: Gamma function**

For any  $x \in \mathbb{R}_{>0}$ , the gamma function is defined as the improper integral

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt$$

**Theorem 2.2.** The gamma function obeys the recurrence relation

$$\Gamma(x + 1) = x \cdot \Gamma(x)$$

*Proof.* Integration by parts gives:

$$\begin{aligned} \Gamma(x + 1) &= \int_0^{\infty} t^x e^{-t} dt \\ &= \left[ -t^x e^{-t} \right]_0^{\infty} + \int_0^{\infty} x t^{x-1} e^{-t} dt \\ &= 0 - \lim_{n \rightarrow \infty} (-t^x e^{-t}) + x \int_0^{\infty} t^{x-1} e^{-t} dt \\ &= x \int_0^{\infty} t^{x-1} e^{-t} dt \\ &= x \cdot \Gamma(x) \end{aligned}$$

□

**Corollary 2.3.** For any  $n \in \mathbb{N}$ , we have

$$\Gamma(n + 1) = n!$$

*Proof.* Our recurrence relation gives

$$\Gamma(n + 1) = n! \cdot \Gamma(1)$$

and we have

$$\begin{aligned} \Gamma(1) &= \int_0^\infty t^{1-1} e^{-t} dt \\ &= \int_0^\infty e^{-t} dt \\ &= [-e^{-t}]_0^\infty \\ &= \left( \lim_{n \rightarrow \infty} -e^{-n} \right) - (-e^0) \\ &= 0 - (-1) \\ &= 1 \end{aligned}$$

□

**Lemma 2.4.** We have:

$$\Gamma(x) = 2c^x \int_0^\infty t^{2x-1} e^{-ct^2} dt$$

*Proof.* Substituting  $t := cu^2$ , we get:

$$\begin{aligned} \Gamma(x) &= \int_0^\infty t^{x-1} e^{-t} dt \\ &= \int_0^\infty (cu^2)^{x-1} e^{-cu^2} \cdot 2cu du \\ &= \int_0^\infty c^{x-1} u^{2(x-1)} e^{-cu^2} \cdot 2cu du \\ &= \int_0^\infty c^x u^{2x-1} e^{-cu^2} du \\ &= c^x \cdot \int_0^\infty u^{2x-1} e^{-cu^2} du \\ &= c^x \cdot \int_0^\infty t^{2x-1} e^{-ct^2} dt \end{aligned}$$

□

**Proposition 2.5.** We have  $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$

*Proof Sketch.* By the last lemma, we have:

$$\begin{aligned}\Gamma\left(\frac{1}{2}\right) &= 2 \int_0^{\infty} t^{2 \cdot \frac{1}{2} - 1} e^{-t^2} dt \\ &= 2 \int_0^{\infty} e^{-t^2} dt \\ &= \int_{-\infty}^{\infty} e^{-t^2} dt\end{aligned}$$

The fact that  $\int_{-\infty}^{\infty} e^{-t^2} dt = \sqrt{\pi}$  might be familiar if you have a basic knowledge of probability theory, since this is exactly the reason why  $\pi$  appears in the formula for a normal distribution:

$$\mathcal{N}(\mu, \sigma^2)(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{\left(-\frac{(t-\mu)^2}{2\sigma^2}\right)}$$

Setting  $\mu = 0$  and  $\sigma^2 = \frac{1}{2}$ , which was Gauss' original definition for the standard normal distribution, we get:

$$\mathcal{N}(0, 1)(t) = \frac{1}{\sqrt{\pi}} e^{-t^2},$$

where the factor  $\frac{1}{\sqrt{\pi}}$  appears to normalize the integral of the distribution to 1, since it wouldn't be a probability distribution otherwise.

We will formally prove this fact much later, when we introduce the substitution formula for multidimensional Lebesgue integration. ”□”

**Part V**

**Linear Algebra**

# Contents

|          |                                                               |           |
|----------|---------------------------------------------------------------|-----------|
| <b>1</b> | <b>Analytic Geometry</b>                                      | <b>54</b> |
| 1.1      | Arithmetic in $n$ dimensions.....                             | 54        |
| 1.2      | The Inner Product: Angles and Lengths in $\mathbb{R}^n$ ..... | 56        |
| 1.3      | The Cross Product.....                                        | 60        |
| 1.4      | The Quaternions $\mathbb{H}$ .....                            | 63        |
| 1.5      | Quaternions and $\mathbb{R}^3$ .....                          | 67        |
| <b>2</b> | <b>Linear Maps</b>                                            | <b>68</b> |
| 2.1      | Vector Spaces .....                                           | 68        |
| 2.2      | Subspaces and Quotient Spaces.....                            | 69        |
| 2.3      | Bases and Dimension .....                                     | 69        |
| 2.4      | Matrices .....                                                | 69        |
| 2.5      | Systems of Linear Equations .....                             | 69        |
| 2.6      | The Method of Least Squares .....                             | 69        |
| <b>3</b> | <b>Affine Spaces</b>                                          | <b>70</b> |
| <b>4</b> | <b>Determinants</b>                                           | <b>74</b> |
| 4.1      | Defining Volume .....                                         | 74        |
| 4.2      | Determinants of Endomorphisms .....                           | 74        |
| 4.3      | Orientation.....                                              | 74        |
| <b>5</b> | <b>Eigenvectors and Eigenvalues</b>                           | <b>75</b> |
| 5.1      | Characteristic Polynomials .....                              | 77        |
| 5.2      | Algebras and the Cayley-Hamilton Theorem .....                | 77        |
| 5.3      | Minimal Polynomials .....                                     | 78        |
| <b>6</b> | <b>Normal Forms of Endomorphisms</b>                          | <b>79</b> |
| 6.1      | Decomposing Vector Spaces using Polynomials .....             | 79        |
| 6.2      | The Jordan Normal Form.....                                   | 82        |
| 6.3      | The Weierstrass Normal Form .....                             | 84        |
| <b>7</b> | <b>Inner Product Spaces</b>                                   | <b>85</b> |
| 7.1      | Inner Products .....                                          | 85        |
| 7.2      | Orthogonality .....                                           | 87        |
| 7.3      | Matrix Representations of Inner Products .....                | 88        |
| 7.4      | Dual Spaces.....                                              | 88        |
| <b>8</b> | <b>Normal Endomorphisms</b>                                   | <b>92</b> |
| 8.1      | Linear Isometries.....                                        | 94        |
| 8.2      | Affine Isometries .....                                       | 94        |
| <b>9</b> | <b>Revisiting Synthetic Geometry</b>                          | <b>96</b> |
| 9.1      | Incidence Geometries .....                                    | 96        |

|           |                               |            |
|-----------|-------------------------------|------------|
| 9.2       | Projection Maps.....          | 98         |
| 9.3       | Projectivization.....         | 99         |
| <b>10</b> | <b>Hyperbolic Geometry</b>    | <b>101</b> |
| 10.1      | Circle Inversions.....        | 101        |
| 10.2      | Möbius Transformations.....   | 102        |
| 10.3      | Complex Möbius Geometry ..... | 103        |

# Chapter 1

## Analytic Geometry

### 1.1 Arithmetic in $n$ dimensions

We now focus our attention on  $n$ -dimensional euclidean space, which is modelled by the cartesian product  $\mathbb{R}^n$ :

$$\mathbb{R}^n = \left\{ \vec{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \mid x_1, \dots, x_n \in \mathbb{R} \right\} = \{ \vec{x} : \{1, \dots, n\} \rightarrow \mathbb{R} \}$$

Elements of  $\mathbb{R}^n$  are known as *vectors*<sup>1</sup>. In the context of linear algebra, they are usually written vertically, as *column vectors*.

Vectors simultaneously represent both *positions* and *directions* in space - in the same way that the number 5 is both the point that lies 5 units away from 0 on the real number line *and* the distance between the numbers 3 and 8 on that same number line, the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  represents both

- The point on the cartesian plane that can be reached by going one unit in the direction of one coordinate axis and then two units in the direction of the other axis, *and*
- the distance between two points with coordinates  $\begin{pmatrix} 2 \\ 3 \end{pmatrix}$  and  $\begin{pmatrix} 3 \\ 5 \end{pmatrix}$ , or any other pair of points that are one unit apart in one coordinate direction and two in the other.

As is expected for algebraic structures defined on cartesian products, addition is defined **componentwise**:

$$\begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} + \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} x_1 + y_1 \\ \vdots \\ x_n + y_n \end{pmatrix}$$

Given a real number  $c$ , we can scale elements of  $\mathbb{R}^n$  by that number:

$$c \cdot \vec{x} = c \cdot \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} c \cdot x_1 \\ \vdots \\ c \cdot x_n \end{pmatrix}$$

<sup>1</sup> We will see in the next chapter that  $\mathbb{R}^n$  is an example of a *vector space*, including the case  $n = 1$  (and  $n = 0$ , if you want to get a bit pathological). Therefore, formally speaking and only in certain contexts, regular old real (or even rational) numbers can be vectors too. However, when encountering the term "vector" in the wild, it is generally implied that  $n \geq 2$ .

The factor  $c$  is known as a *scalar*, and the operator  $\cdot : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  as *scalar multiplication*. For the sake of readability, we differentiate scalars from vectors by writing a little arrow on top of a vector.

Arguably, it is these two operations - vector addition and scalar multiplication - that define linear algebra - next chapter, we will abstractly define *vector spaces* to be those structures where addition and scalar multiplication are well-behaved and the scalars come from a field, and *modules* to be their cousins whose scalars come from a ring.

In general, there is no well-behaved<sup>2</sup> vector multiplication that works for any  $n$ . Later on, we will prove this observation in the form of *Frobenius' Theorem*, which roughly states:

**Theorem 1.1: Frobenius' Theorem**

The only dimensions in which an associative multiplication exists on  $\mathbb{R}^n$  are  $\mathbb{R}^1$ , i.e.  $\mathbb{R}$  itself,  $\mathbb{R}^2$ , where multiplication is given by treating two-dimensional vectors as complex numbers, and  $\mathbb{R}^4$ , where multiplication is given treating four-dimensional vectors as *quaternions*.

We will take a closer look at the complex numbers  $\mathbb{C} \cong \mathbb{R}^2$  and the quaternions  $\mathbb{H} \cong \mathbb{R}^4$  in the following sections. In particular, we will see that quaternion multiplication isn't commutative, so if we require multiplication to be commutative, our options shrink even further, leaving just  $\mathbb{R}$  and  $\mathbb{C} \cong \mathbb{R}^2$ .

<sup>2</sup> In this case, our criterion for multiplication being well-behaved is that multiplicative inverses should exist, i.e. that division should be possible.

## 1.2 The Inner Product: Angles and Lengths in $\mathbb{R}^n$

Another operation that is of central importance to euclidean geometry is the *inner product*, also known as the *dot product* or *scalar product*.

### Definition 1.2: Euclidean Inner Product

Given two vectors  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , their *Euclidean inner product*, written  $\langle \vec{x}, \vec{y} \rangle$  or simply  $\vec{x} \cdot \vec{y}$ , is the sum of the products of the components of  $\vec{x}$  with the components of  $\vec{y}$ :

$$\langle \vec{x}, \vec{y} \rangle = \sum_{i=1}^n x_i \cdot y_i$$

### Theorem 1.3: Properties of Inner Products

The Euclidean inner product has the following properties:

1. **(Positive definiteness):** For every  $\vec{x} \in \mathbb{R}^n$ , we have  $\langle \vec{x}, \vec{x} \rangle = 0$  if and only if  $\vec{x} = \vec{0}$ .
2. **(Linearity in the first argument):** For every  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$  and  $a, b \in \mathbb{R}$ , we have:

$$\langle a\vec{x} + b\vec{y}, \vec{z} \rangle = a \langle \vec{x}, \vec{z} \rangle + b \langle \vec{y}, \vec{z} \rangle$$

3. **(Symmetry):** For every  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , we have:

$$\langle \vec{x}, \vec{y} \rangle = \langle \vec{y}, \vec{x} \rangle$$

**Exercise 1.4.** Show that the Euclidean inner product fulfills the following version of the binomial formula:

$$\begin{aligned} \langle \vec{x} + \vec{y}, \vec{x} + \vec{y} \rangle &= \langle \vec{x}, \vec{x} \rangle + 2 \langle \vec{x}, \vec{y} \rangle + \langle \vec{y}, \vec{y} \rangle \\ \langle \vec{x} - \vec{y}, \vec{x} - \vec{y} \rangle &= \langle \vec{x}, \vec{x} \rangle - 2 \langle \vec{x}, \vec{y} \rangle + \langle \vec{y}, \vec{y} \rangle \end{aligned}$$

The importance of inner products comes from the fact they let us define *lengths* and *angles*.

### Definition 1.5: Euclidean Norm

The length of a vector  $\vec{x} \in \mathbb{R}^n$ , also known as its *Euclidean norm*, written as  $\|\vec{x}\|_2$ ,  $\|\vec{x}\|$ , or just  $|x|$ , is given by:

$$\|\vec{x}\|_2 = \sqrt{\langle \vec{x}, \vec{x} \rangle} = \sqrt{x_1^2 + \dots + x_n^2}$$

Note that for  $n = 1$ , we simply get  $\|x\|_2 = \sqrt{x^2} = |x|$ , so any result concerning the Euclidean norm on  $\mathbb{R}^n$  also holds for the absolute value function.

**Theorem 1.6: Some properties of Norms**

The Euclidean norm has the following properties:

1. **(Non-negativity):** For every  $\vec{x} \in \mathbb{R}^n$ ,  $\|\vec{x}\|_2 \geq 0$ .
2. **(Positive definiteness):** For every  $\vec{x} \in \mathbb{R}^n$ ,  $\|\vec{x}\|_2 = 0$  if and only if  $\vec{x} = \vec{0}$ .
3. **(Absolute homogeneity):** For every  $c \in \mathbb{R}$  and  $\vec{x} \in \mathbb{R}^n$ , we have

$$\|c \cdot \vec{x}\|_2 = |c| \cdot \|\vec{x}\|_2$$

The little subscript 2 in  $\|\vec{x}\|_2$  differentiates the Euclidean norm from the many other norms we will meet over time - in general, for any  $p \in \mathbb{N}_1$ ,  $\|\vec{x}\|_p$  denotes the function

$$\|\vec{x}\|_p = \sqrt[p]{|x_1|^p + \dots + |x_n|^p} \quad \left( = \sqrt[p]{\sum_{i=1}^n |x_i|^p} \right),$$

known as the  **$p$ -norm**. For now, we only really care about the Euclidean norm, but other values of  $p$  will start coming in handy once we start applying tools from analysis to vector spaces. The limiting case  $p \rightarrow \infty$  is particularly useful: We will find that

$$\|\vec{x}\|_\infty = \max |x_i|.$$

**Theorem 1.7: Cauchy-Schwartz Inequality**

Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$ . Then we have

$$|\langle \vec{x}, \vec{y} \rangle| \leq \|\vec{x}\|_2 \cdot \|\vec{y}\|_2$$

*Proof.* 1. Assume  $\|\vec{x}\|_2 = 0$ . Then we immediately get

$$\langle \vec{x}, \vec{y} \rangle = 0 = \|\vec{x}\|_2 \cdot \|\vec{y}\|_2.$$

2. Assume  $\|\vec{x}\|_2 = \|\vec{y}\|_2 = 1$ . Then we have  $\|\vec{x}\|_2^2 + \|\vec{y}\|_2^2 = 2$ , and thus:

$$\begin{aligned} 1 + \langle \vec{x}, \vec{y} \rangle &= \frac{1}{2}(\|\vec{x}\|_2^2 + \|\vec{y}\|_2^2) + \langle \vec{x}, \vec{y} \rangle \\ &= \frac{1}{2}(\|\vec{x}\|_2^2 + 2\langle \vec{x}, \vec{y} \rangle + \|\vec{y}\|_2^2) \\ &= \frac{1}{2}\|\vec{x} + \vec{y}\|_2^2 \\ &\geq 0 \end{aligned}$$

Thus we get  $\langle \vec{x}, \vec{y} \rangle \geq -1$ . Applying the same reasoning to  $1 - \langle \vec{x}, \vec{y} \rangle$ , we get

$$1 - \langle \vec{x}, \vec{y} \rangle = \frac{1}{2}\|\vec{x} - \vec{y}\|_2^2 \geq 0,$$

i.e.  $\langle \vec{x}, \vec{y} \rangle \leq 1$ . We thus have

$$|\langle \vec{x}, \vec{y} \rangle| \leq 1 = \|\vec{x}\|_2 \cdot \|\vec{y}\|_2,$$

as desired.

3. Now, let  $\vec{x}, \vec{y} \in \mathbb{R}^n \setminus \{\vec{0}\}$  be arbitrary. Then  $\frac{\vec{x}}{\|\vec{x}\|_2}$  and  $\frac{\vec{y}}{\|\vec{y}\|_2}$  both have norm 1, letting us simply use our previous result:

$$\begin{aligned} |\langle \vec{x}, \vec{y} \rangle| &= \left| \|\vec{x}\| \cdot \|\vec{y}\| \cdot \left\langle \frac{\vec{x}}{\|\vec{x}\|}, \frac{\vec{y}}{\|\vec{y}\|} \right\rangle \right| \\ &\leq \left| \|\vec{x}\| \cdot \|\vec{y}\| \cdot \left\| \frac{\vec{x}}{\|\vec{x}\|} \right\| \cdot \left\| \frac{\vec{y}}{\|\vec{y}\|} \right\| \right| \\ &= \|\vec{x}\| \cdot \|\vec{y}\| \end{aligned}$$

□

### Corollary 1.8: Triangle Inequality

For every  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , we have

$$\|\vec{x} + \vec{y}\|_2 \leq \|\vec{x}\|_2 + \|\vec{y}\|_2$$

We had previously formalized the concept of an ‘angle between two lines’ via the complex exponential function, and then defined the trigonometric functions  $\sin x$  and  $\cos x$  as the complex and real parts of  $e^{ix}$ , respectively. If we take the cosine function as a given, then scalar products provide a convenient alternate way of defining angles, which works in any dimension:

### Definition 1.9: Inner angle between two vectors

Given two vectors  $\vec{x}, \vec{y} \in \mathbb{R}^n \setminus \{\vec{0}\}$ , we define the *inner angle* between them to be:

$$\angle(\vec{x}, \vec{y}) = \arccos \left( \frac{\langle \vec{x}, \vec{y} \rangle}{\|\vec{x}\|_2 \cdot \|\vec{y}\|_2} \right)$$

Note that Cauchy-Schwartz gives us  $\frac{\langle \vec{x}, \vec{y} \rangle}{\|\vec{x}\|_2 \cdot \|\vec{y}\|_2} \in [-1, 1]$ , showing that this expression actually is well-defined.

**Exercise 1.10.** Verify that we have:

1.  $\angle(\vec{x}, \vec{y}) = 0$  if and only if there exists  $c \in \mathbb{R}_{>0}$  such that  $\vec{x} = c\vec{y}$ .
2.  $\angle(\vec{x}, \vec{y}) = \frac{\pi}{2}$  if and only if  $\langle \vec{x}, \vec{y} \rangle = 0$ .
3.  $\angle(\vec{x}, \vec{y}) = \pi$  if and only if there exists  $c \in \mathbb{R}_{>0}$  such that  $\vec{x} = -c\vec{y}$ .

Notably, this suggests that our definition is sensible even in the case  $n = 1$ , since any pair of non-zero vectors fulfills either condition 1 or 2, implying that the angle between two vectors on the real number line is either 0 or  $\pi$ .

These inner angles match our earlier definition of angles in terms of complex numbers, as long as we are content with not being able to meaningfully talk about negative angles or angles greater than  $\pi$  anymore:

**Exercise 1.11.** Earlier in this book, we defined angles in terms of complex numbers. Given a complex number  $y = re^{i\varphi}$  we saw that the corresponding real vector is given by  $\vec{y} = \begin{pmatrix} r \cos \varphi \\ r \sin \varphi \end{pmatrix}$ .

Show that our two definitions match as long as  $\varphi \in [0, \pi]$ , i.e. that given a vector  $\vec{x} = \begin{pmatrix} x \\ 0 \end{pmatrix}$  lying on the  $x$ -axis, we have

$$\angle(\vec{x}, \vec{y}) = \varphi$$

(Hint: Remember  $\cos^2 \varphi + \sin^2 \varphi = 1$ )

We have to be content with this restricted interval  $[0, \pi]$ , since the cosine function isn't injective on larger intervals, meaning the arccosine would no longer be well-defined.

More generally, any definition of an 'angle between two arbitrary vectors' will suffer from this - if we only specify two vectors, it's left ambiguous which side we should measure the angle on.

This problem can be solved by finding a way to take a tuple  $(\vec{x}_1, \dots, \vec{x}_n)$  of vectors and recognize whether it is 'positively ordered' or 'negatively ordered' - in  $\mathbb{R}^2$ , this would be asking whether the second vector lies 'clockwise' or 'counterclockwise' from the first, while in  $\mathbb{R}^3$  the question is whether the vectors are 'left-handed' or 'right-handed'. A map that gives us this information is known as an **orientation**. We will formalize this idea later, when we have introduced the necessary tools, those being vector spaces, their **bases**, and **determinant functions**.

In the complex case, we could dodge this issue because one of our vectors, call it  $\vec{x}$ , was always part of the  $x$ -axis, and thus we could obtain a canonical orientation by declaring that the angle between  $\vec{x}$  and  $\vec{y}$  was positive if  $y_2 > 0$  and negative if  $y_2 < 0$ .

**Exercise 1.12.** Prove the following theorems of classical Euclidean geometry:

1. **The Law of Cosines:**

$$\|\vec{x} - \vec{y}\|_2^2 = \|\vec{x}\|_2^2 + \|\vec{y}\|_2^2 - 2\|\vec{x}\|_2\|\vec{y}\|_2 \cdot \cos(\angle(\vec{x}, \vec{y}))$$

2. **The Pythagorean Theorem:**

$$\angle(\vec{x}, \vec{y}) = \frac{\pi}{2} \implies \|\vec{x} - \vec{y}\|_2^2 = \|\vec{x}\|_2^2 + \|\vec{y}\|_2^2$$

## 1.3 The Cross Product

The *cross product* is an operation  $\times : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ , such that:

1.  $\vec{u} \times \vec{v}$  is orthogonal to both  $\vec{u}$  and  $\vec{v}$ ,
2. the length of  $\vec{u} \times \vec{v}$  is the area of the parallelogram spanned by  $\vec{u}$  and  $\vec{v}$ :

$$\|\vec{u} \times \vec{v}\|_2 = \|\vec{u}\|_2 \|\vec{v}\|_2 \sin(\angle(\vec{u}, \vec{v}))$$

We also declare that our cross product should be *anticommutative*: given two vectors, we want to be able to change the sign of the cross product by switching the order in which we multiply:

$$\vec{v} \times \vec{w} := -\vec{w} \times \vec{v}$$

Notice how this immediately implies

$$\begin{aligned} \vec{v} \times \vec{v} &= -(\vec{v} \times \vec{v}) \\ \implies \vec{v} \times \vec{v} &= 0 \end{aligned}$$

Finally, notice that our geometric intuition immediately tells us that our product should be *linear* and *distributive over vector addition*.

We can now derive an explicit formula for this operation on our own. First, note that the standard basis

$$\vec{e}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \vec{e}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \vec{e}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix},$$

consists of orthogonal vectors of length one, so we can start by declaring

$$\begin{aligned} \vec{e}_1 \times \vec{e}_2 &:= \vec{e}_3 \\ \vec{e}_2 \times \vec{e}_3 &:= \vec{e}_1 \\ \vec{e}_3 \times \vec{e}_1 &:= \vec{e}_2 \end{aligned}$$

Note that once again, the direction in which these vectors point depends on a somewhat arbitrary choice of orientation - we could invert all of the directions and still end up with a valid definition.

These observations are already enough to calculate the cross product of two

arbitrary vectors: Using distributivity, we get:

$$\begin{aligned}
 \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \times \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} &= (v_1 \vec{e}_1 + v_2 \vec{e}_2 + v_3 \vec{e}_3) \times (w_1 \vec{e}_1 + w_2 \vec{e}_2 + w_3 \vec{e}_3) \\
 &= v_1 w_1 (\vec{e}_1 \times \vec{e}_1) \\
 &\quad + v_1 w_2 (\vec{e}_1 \times \vec{e}_2) \\
 &\quad + v_1 w_3 (\vec{e}_1 \times \vec{e}_3) \\
 &\quad + v_2 w_1 (\vec{e}_2 \times \vec{e}_1) \\
 &\quad + v_2 w_2 (\vec{e}_2 \times \vec{e}_2) \\
 &\quad + v_2 w_3 (\vec{e}_2 \times \vec{e}_3) \\
 &\quad + v_3 w_1 (\vec{e}_3 \times \vec{e}_1) \\
 &\quad + v_3 w_2 (\vec{e}_3 \times \vec{e}_2) \\
 &\quad + v_3 w_3 (\vec{e}_3 \times \vec{e}_3)
 \end{aligned}$$

To which we can apply anticommutativity and our formulas for multiplying the basis vectors to get:

$$\begin{aligned}
 \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \times \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} &= v_1 w_1 \cdot \vec{0} \\
 &\quad + v_1 w_2 \cdot \vec{e}_3 \\
 &\quad + v_1 w_3 \cdot (-\vec{e}_2) \\
 &\quad + v_2 w_1 \cdot (-\vec{e}_3) \\
 &\quad + v_2 w_2 \cdot \vec{0} \\
 &\quad + v_2 w_3 \cdot \vec{e}_1 \\
 &\quad + v_3 w_1 \cdot \vec{e}_2 \\
 &\quad + v_3 w_2 \cdot (-\vec{e}_1) \\
 &\quad + v_3 w_3 \cdot \vec{0} \\
 &= \begin{pmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - w_1 v_3 \\ v_1 w_2 - v_2 w_1 \end{pmatrix}
 \end{aligned}$$

**Corollary 1.13: Cross Product**

The cross product  $\vec{v} \times \vec{w}$  of two vectors  $\vec{v}, \vec{w} \in \mathbb{R}^3$  returns a vector whose direction is orthogonal to both  $\vec{v}$  and  $\vec{w}$  and whose length equals the area of the parallelogram spanned between them. Given coordinate representations of  $\vec{v}$  and  $\vec{w}$ , the cross product can be calculated via the following formula:

$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \times \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - w_1 v_3 \\ v_1 w_2 - v_2 w_1 \end{pmatrix}$$

**Proposition 1.14.** We have

$$\langle \vec{u} \times \vec{v}, \vec{w} \times \vec{t} \rangle = \langle \vec{u}, \vec{w} \rangle \langle \vec{v}, \vec{t} \rangle - \langle \vec{u}, \vec{t} \rangle \langle \vec{v}, \vec{w} \rangle.$$

**Proposition 1.15.** We have

$$\langle \vec{u} \times \vec{v}, \vec{w} \rangle = \langle \vec{v} \times \vec{w}, \vec{u} \rangle = \langle \vec{w} \times \vec{u}, \vec{v} \rangle$$

This expression is known as the **box product** of  $\vec{u}, \vec{v}, \vec{w}$ . Its absolute value corresponds to the volume of the parallelepiped spanned by the three vectors.

**Proposition 1.16.** We have

$$(\vec{u} \times \vec{v}) \times \vec{w} = \langle u, w \rangle \cdot v - \langle v, w \rangle \cdot u = w \times (v \times u).$$

This result is known as **Grassmann's Identity**.

**Proposition 1.17.** We have

$$(\vec{u} \times \vec{v}) \times \vec{w} + (\vec{v} \times \vec{w}) \times \vec{u} + (\vec{w} \times \vec{u}) \times \vec{v} = 0$$

This result is known as **Jacobi's Identity**.

## 1.4 The Quaternions $\mathbb{H}$

### Definition 1.18: Quaternions

The Quaternions  $\mathbb{H}$  consist of the set  $\mathbb{R}^4$ , with the following special notation:

$$\vec{1} := (1, 0, 0, 0)$$

$$i := (0, 1, 0, 0)$$

$$j := (0, 0, 1, 0)$$

$$k := (0, 0, 0, 1)$$

Such that in addition to scalar multiplication and vector addition, which work as usual, we have an additional vector multiplication operation  $\cdot : \mathbb{R}^4 \times \mathbb{R}^4 \rightarrow \mathbb{R}^4$  that behaves as follows:

1.  $\vec{1}$  is a neutral element of vector multiplication:

$$\forall x \in \mathbb{H} :$$

$$\vec{1} \cdot x = x \cdot \vec{1} = x$$

2. Vector multiplication is associative:

$$\vec{x}(\vec{y}\vec{z}) = (\vec{x}\vec{y})\vec{z}$$

3. Multiplicative inverses exist:

$$\forall \vec{x} \in \mathbb{H}_{\neq 0} :$$

$$\exists \vec{x}^{-1} \in \mathbb{H} :$$

$$\vec{x} \cdot \vec{x}^{-1} = \vec{x}^{-1} \cdot \vec{x} = \vec{1}$$

4. Vector multiplication distributes over vector addition, from both sides:

$$\forall \vec{x}, \vec{y}, \vec{z} \in \mathbb{H} :$$

$$\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x}\vec{y} + \vec{x}\vec{z}$$

$$(\vec{x} + \vec{y}) \cdot \vec{z} = \vec{x}\vec{z} + \vec{y}\vec{z}$$

5. Vector multiplication is compatible with scalar multiplication:

$$\forall a, b \in \mathbb{R}, \vec{x}, \vec{y} \in \mathbb{H} :$$

$$a\vec{x} \cdot b\vec{y} = (ab)(\vec{x}\vec{y})$$

6. The unit vectors fulfill the *Quaternion multiplication formula*:

$$i^2 = j^2 = k^2 = ijk = -\vec{1}$$

Apart from the Quaternion multiplication formula, which is the defining feature of the Quaternions, all of our other demands together state that  $\mathbb{H}$  should

be an *associative real division algebra over  $\mathbb{R}^4$* .

One very important consequence of this definition is that vectors of the form  $\vec{a} := (a, 0, 0, 0)$  behave exactly like real numbers - let  $\vec{x} \in \mathbb{H}$ , then we have:

$$\vec{a}\vec{x} = (a \cdot \vec{1}) \cdot \vec{x} = a \cdot (\vec{1} \cdot \vec{x}) = a\vec{x}$$

For this reason, it is common to write  $\vec{1}$  simply as 1, and to then proceed as we did for the complex numbers and write:

$$(a, b, c, d) = a + bi + cj + dk$$

We can now already get most of the way to deriving multiplication formula for quaternions, by simply applying distributivity:

$$\begin{aligned} (v_1 + v_2i + v_3j + v_4k) \cdot (u_1 + u_2i + u_3j + u_4k) \\ = v_1u_1 + (v_1u_2)i + (v_1u_3)j + (v_1u_4)k \\ + (v_2u_1)i + (v_2u_2)i^2 + (v_2u_3)ij + (v_2u_4)ik \\ + (v_3u_1)j + (v_3u_2)ji + (v_3u_3)j^2 + (v_3u_4)jk \\ + (v_4u_1)k + (v_4u_2)ki + (v_4u_3)kj + (v_4u_4)k^2 \end{aligned}$$

Now, using  $i^2 + j^2 + k^2 = ijk = -1$ , we can simplify slightly:

$$\begin{aligned} (v_1 + v_2i + v_3j + v_4k) \cdot (u_1 + u_2i + u_3j + u_4k) \\ = v_1u_1 + (v_1u_2)i + (v_1u_3)j + (v_1u_4)k \\ + (v_2u_1)i + (v_2u_2)i^2 + (v_2u_3)ij + (v_2u_4)ik \\ + (v_3u_1)j + (v_3u_2)ji + (v_3u_3)j^2 + (v_3u_4)jk \\ + (v_4u_1)k + (v_4u_2)ki + (v_4u_3)kj + (v_4u_4)k^2 \\ = v_1u_1 + (v_1u_2)i + (v_1u_3)j + (v_1u_4)k \\ + (v_2u_1)i - (v_2u_2) + (v_2u_3)ij + (v_2u_4)ik \\ + (v_3u_1)j + (v_3u_2)ji - (v_3u_3) + (v_3u_4)jk \\ + (v_4u_1)k + (v_4u_2)ki + (v_4u_3)kj - (v_4u_4) \end{aligned}$$

All that's left is to figure out what the products of  $i, j$  and  $k$  should be.

We can derive  $ij$  and  $ji$  as follows:

1. Start with  $ijk = -1$ .
2. Multiply  $k$  from the right to get  $ijk^2 = -k$ , i.e.  $-ij = -k$ , i.e.  $ij = k$ .
3. Multiply  $ij$  from the left to get  $ijij = ijk = -1$ .
4. Multiply  $i$  from the left and  $j$  from the right to get  $i(ijij)j = -ij$ .
5. Simplify the left side to get  $i(ijij)j = (ii)(ji)(jj) = ji$ , i.e.:

$$ij = -ji = k$$

**Exercise 1.19.** Show that we also have  $jk = -kj = i$  and  $ki = -ik = j$ .

This means that Quaternion multiplication isn't commutative!

Our final multiplication table for  $1, i, j$  and  $k$  is thus:

| $\cdot$ | 1   | $i$  | $j$  | $k$  |
|---------|-----|------|------|------|
| 1       | 1   | $i$  | $j$  | $k$  |
| $i$     | $i$ | -1   | $k$  | $-j$ |
| $j$     | $j$ | $-k$ | -1   | $i$  |
| $k$     | $k$ | $j$  | $-i$ | -1   |

**Exercise 1.20.** Verify that this multiplication table doesn't violate associativity.

Entering these new results into our partial multiplication formula, we get:

$$\begin{aligned}
 (v_1 + v_2i + v_3j + v_4k) \cdot (u_1 + u_2i + u_3j + u_4k) \\
 &= v_1u_1 + (v_1u_2)i + (v_1u_3)j + (v_1u_4)k \\
 &+ (v_2u_1)i - (v_2u_2) + (v_2u_3)ij + (v_2u_4)ik \\
 &+ (v_3u_1)j + (v_3u_2)ji - (v_3u_3) + (v_3u_4)jk \\
 &+ (v_4u_1)k + (v_4u_2)ki + (v_4u_3)kj - (v_4u_4) \\
 &= v_1u_1 + (v_1u_2)i + (v_1u_3)j + (v_1u_4)k \\
 &+ (v_2u_1)i - (v_2u_2) + (v_2u_3)k - (v_2u_4)j \\
 &+ (v_3u_1)j - (v_3u_2)k - (v_3u_3) + (v_3u_4)i \\
 &+ (v_4u_1)k + (v_4u_2)j - (v_4u_3)i - (v_4u_4)
 \end{aligned}$$

which we can finally simplify to get:

$$\begin{aligned}
 (v_1 + v_2i + v_3j + v_4k) \cdot (u_1 + u_2i + u_3j + u_4k) \\
 &= (v_1u_1 - v_2u_2 - v_3u_3 - v_4u_4) \\
 &+ (v_1u_2 + v_2u_1 + v_3u_4 - v_4u_3)i \\
 &+ (v_1u_3 - v_2u_4 + v_3u_1 + v_4u_2)j \\
 &+ (v_1u_4 + v_2u_3 - v_3u_2 + v_4u_1)k
 \end{aligned}$$

Now, this is obviously a long and unreasonably complicated formula - is there any way to think of it differently? The complex numbers had a very nice and intuitive geometric definition, maybe the quaternions do too?

Well, the key trick here is to split the underlying set  $\mathbb{R}^4$  into  $\mathbb{R} \times \mathbb{R}^3$ , thinking of each Quaternion as consisting of a **real part**  $a \in \mathbb{R}$  and an **imaginary part**  $\vec{v} \in \mathbb{R}^3$ . Our multiplication formula now reads:

$$\begin{aligned}
 (a, \vec{v}) \cdot (b, \vec{u}) \\
 &= (ab - v_1u_1 - v_2u_2 - v_3u_3) \\
 &+ (au_1 + bv_1 + v_2u_3 - v_3u_2)i \\
 &+ (au_2 - v_1u_3 + bv_2 + v_3u_1)j \\
 &+ (au_3 + v_1u_2 - v_2u_1 + bv_3)k
 \end{aligned}$$

which you might notice consists exclusively of familiar vector operations:

$$1. \quad ab - v_1u_1 - v_2u_2 - v_3u_3 = ab - \langle \vec{v}, \vec{u} \rangle$$

$$2. \quad \begin{pmatrix} au_1 \\ au_2 \\ au_3 \end{pmatrix} = a\vec{u}, \begin{pmatrix} bv_1 \\ bv_2 \\ bv_3 \end{pmatrix} = b\vec{v}$$

$$3. \quad \begin{pmatrix} v_2u_3 - v_3u_2 \\ v_1u_3 - v_3u_1 \\ v_1u_2 - v_2u_1 \end{pmatrix} = \vec{v} \times \vec{u}$$

Putting everything together, we can now write out our multiplication formula in a much nicer form:

$$(a, \vec{v}) \cdot (b, \vec{u}) = (ab - \langle \vec{v}, \vec{u} \rangle, a\vec{u} + b\vec{v} + \vec{v} \times \vec{u}),$$

**Exercise 1.21.** Verify that quaternion multiplication is associative and distributes over vector addition.

**Exercise 1.22.** Let  $p = (a, \vec{v}) \in \mathbb{H}$ . Then  $pq = qp$  holds for all  $q \in \mathbb{H}$  if and only if  $\vec{v} \neq 0$ .

Similar to complex numbers, we can define *quaternion conjugation*:

**Definition 1.23: Quaternion Conjugation**

Just like for  $\mathbb{C}$ , we define the *conjugate* of a quaternion  $(a, \vec{v})$  to be the quaternion

$$\overline{(a, \vec{v})} = (a, -\vec{v})$$

An immediate application of quaternion conjugation is finding a formula for the inverse of a quaternion, which ones again ends up working out very similarly to the complex case - We have

$$\begin{aligned} \overline{(a, \vec{v})} \cdot (a, \vec{v}) &= (a, -\vec{v}) \cdot (a, \vec{v}) \\ &= (a^2 - \langle -\vec{v}, \vec{v} \rangle, a\vec{v} - a\vec{v} + (-\vec{v} \times \vec{v})) \\ &= (a^2 + \|\vec{v}\|^2, \vec{0}) \in \mathbb{R} \end{aligned}$$

And thus we can simply divide by this value to get our inverse formula:

**Corollary 1.24: Inverse Quaternion**

Let  $(a, \vec{v}) \in \mathbb{H}_{\neq 0}$ . Then  $(a, \vec{v})$  has a multiplicative inverse, which is given by:

$$(a, \vec{v})^{-1} = \frac{(a, -\vec{v})}{a^2 + \|\vec{v}\|^2}.$$

**Definition 1.25.** Let  $q = (a, \vec{v}) \in \mathbb{H}$ . Then

$$|q| := \sqrt{\bar{q} \cdot q} = a^2 + v_1^2 + v_2^2 + v_3^2 = \|q\|_2$$

Note how much this once again resembles both the real case  $|x| = \sqrt{x^2} = \|x\|_2$  and the complex case  $|z| = \sqrt{\bar{z}z} = \|z\|_2$ .

## 1.5 Quaternions and $\mathbb{R}^3$

### Theorem 1.26: Quaternions encode rotations

Let  $\|\vec{v}\| = 1$  and

$$q = (\cos \varphi, \vec{v} \sin \varphi) \in \mathbb{H}$$

Then  $qw\bar{q} \in \mathbb{R}^3$  is the vector obtained by rotating  $w$  around the axis  $\{\lambda\vec{v} : \lambda \in \mathbb{R}\}$  by the angle  $2\varphi$ .

Since  $2\pi$  corresponds to a full rotation,  $\varphi$  and  $\varphi + \pi$  describe the same rotation. Therefore, there are two different quaternions  $q$  and  $-q$  associated with each rotation - in topological terms, we say that the quaternions form a *double cover* of the space of all rotations.

This phenomenon is actually closely related to the theory of *spin* in quantum physics. All known quantum particles that constitute ordinary matter, including protons, neutrons, electrons, neutrinos, and quarks, are *spin- $\frac{1}{2}$* , meaning that they exhibit a curious behavior where a rotation of  $360^\circ$  leaves them in a different state than they were before, from which they return to their original state only after being rotated by another  $360^\circ$  around the same axis. This lets us mathematically model particles as quaternions, such that a  $360^\circ$  rotation transforms a particle from a state  $q \in \mathbb{H}$  to the "rotationally equivalent" state  $-q \in \mathbb{H}$ .

**Corollary 1.27.** For every isometry  $F : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ , there exist  $u \in \mathbb{R}^3$  and  $q \in \mathbb{H}$  such that we have either

1.  $F(w) = u + qw\bar{q}$ , if  $F$  preserves orientation, or
2.  $F(w) = u + q\bar{w}q$ , if  $F$  doesn't preserve orientation.

This representation turns out to be a lot more computationally efficient than other representations of the same rotation. Other, more intuitively "trivial" representations of rotation also run into issues such as *gimbal lock*, where they may get "stuck" in certain orientations.

This makes quaternions surprisingly practical in many applied contexts, including game development and animation<sup>1</sup>, robotics, and three-dimensional image processing.

---

<sup>1</sup>For an introduction to quaternions in game development, I highly recommend this excellent talk by Freya Holmer.

## Chapter 2

# Linear Maps

### 2.1 Vector Spaces

Intuitively, a vector space is any algebraic structure  $V$  equipped with a field  $\mathbb{F}$ , such that elements of  $V$  can be added to elements of  $V$  or multiplied by elements of  $\mathbb{F}$ , and these operations behave how vector addition and scalar addition behave in  $\mathbb{R}^n$ . More precisely:

#### Definition 2.1: Vector Spaces

A *vector space*  $V$  over a field of scalars  $\mathbb{F}$  is a structure  $(V, \mathbb{F}, +, \cdot)$  consisting of a *vector addition operation*  $+: V \times V \rightarrow V$  and a *scalar multiplication operation*  $\cdot: \mathbb{F} \times V \rightarrow V$  such that:

1. vectors form a commutative group under vector addition, and
2. scalar multiplication is a homomorphism from  $\mathbb{F}$  into the endomorphism ring of  $(V, +)$ .

Spelling everything out, the condition that  $(V, +)$  is a commutative group tells us that:

1. Vector addition is associative:

$$\begin{aligned}\forall v, w, u \in V : \\ (v + w) + u &= v + (w + u)\end{aligned}$$

2. Vector addition is commutative:

$$\begin{aligned}\forall v, w \in V : \\ v + w &= w + v\end{aligned}$$

3. There exists a vector  $0_V \in V$  such that  $v + 0_V = v$  for all  $v \in V$ .

4. For every vector  $v \in V$ , there exists an additive inverse  $-v \in V$ :

$$\begin{aligned}\forall v \in V : \\ \exists -v \in V : \\ v + (-v) &= 0_V\end{aligned}$$

As usual, we write  $v - w := v + (-w)$  as an abbreviation.

The second condition might seem a bit more obtuse - however, all it tells us is that  $\cdot$  assigns to each scalar  $\lambda \in \mathbb{F}$  an endomorphism of  $V$ , such that  $1_{\mathbb{F}}$  is the identity map  $\text{id}_{\mathbb{F}}$ . Putting everything together, we get the following demands for scalar addition:

1. Scalar multiplication by any scalar respects vector addition, i.e. the operations distribute:

$$\begin{aligned}\forall \lambda \in \mathbb{F} : \\ \forall v, w \in V : \\ \lambda.(v + w) &= \lambda.v + \lambda.w\end{aligned}$$

## 2.2 Subspaces and Quotient Spaces

## 2.3 Bases and Dimension

## 2.4 Matrices

## 2.5 Systems of Linear Equations

## 2.6 The Method of Least Squares

## Chapter 3

# Affine Spaces

Affine spaces are structures similar to vector spaces, but where the notions of "points" and "directions" are separate.

### Definition 3.1: Affine Space

A  $\mathbb{k}$ -*affine space* is a tuple  $(A, \vec{A})$ , consisting of a set  $A$  of *points* and a  $\mathbb{k}$ -vector space  $\vec{A}$  of *translation vectors*, alongside a translation map  $+ : A \times \vec{A} \rightarrow A$ , such that:

1. Translation is associative:

$$\begin{aligned} \forall a \in A, \vec{v}, \vec{w} \in \vec{A} : \\ (a + \vec{v}) + \vec{w} = a + (\vec{v} + \vec{w}) \end{aligned}$$

2. For any pair of points, there exists a unique vector translating one point onto the other:

$$\begin{aligned} \forall a, b \in P_A : \\ \exists! \vec{v} \in \vec{A} : a + \vec{v} = b \end{aligned}$$

The dimension of an affine space is simply the dimension of its associated vector space.

Any vector space  $V$  can be canonically turned into an affine space by taking  $V$  as both the set of points and the set of vectors. On the flipside, we can turn any affine space  $(A, \vec{A})$  into a vector space by restricting our attention to translations of one "base point"  $b \in A$ , and then identifying any point  $p \in A$  with the vector  $\vec{v} \in \vec{A}$  translating  $b$  onto  $p$ .

For most practical purposes, it is therefore sufficient to work with a vector space as your base space for whatever math you are doing, but the non-canonicity of having to choose an arbitrary base point  $b \in A$  to turn an affine space into a vector space means that some people prefer affine spaces for philosophical reasons - after all, what is the "base point" of our physical universe?

The distinction becomes a lot more important when considering affine subspaces,

which are essentially vector subspaces translated by arbitrary points:

**Definition 3.2: Affine subspaces**

Let  $(A, \vec{A})$  be an affine space. Then an affine subspace of  $(A, \vec{A})$  is a tuple  $(B, \vec{B})$  such that:

1.  $B \subseteq A$ .
2.  $\vec{B}$  is a vector subspace of  $\vec{A}$ .
3.  $\vec{B}$  is the set of translations between points in  $B$ , i.e. for any  $b \in B$ ,  $\vec{v} \in \vec{A}$ , we have  $b + \vec{v} \in B$  if and only if  $\vec{v} \in \vec{B}$ .

Thus, an affine subspace can always be written as

$$B = b + \vec{B} := \{b + \vec{v} \mid \vec{v} \in \vec{B} \subseteq \vec{A}\}$$

for an arbitrary base point  $b \in B$ .

An affine subspace of dimension 1 is known as a *line*. An affine subspace of dimension 2 is known as a *plane*. An affine subspace whose dimension is one lower than the space it resides in is known as an *hyperplane*.

**Definition 3.3.** Two affine subspaces  $(A, \vec{A})$ ,  $(B, \vec{B})$  are called *parallel* if  $\vec{A} = \vec{B}$ .

**Definition 3.4: Affine Maps**

Let  $A, B$  be  $\mathbb{K}$ -affine spaces. Then a map  $F : A \rightarrow B$  is called *affine* if there exists a linear map  $\vec{F} : \vec{A} \rightarrow \vec{B}$  such that

$$\forall a \in P_A, \vec{v} \in \vec{A} : F(a + \vec{v}) = F(a) + \vec{F}(\vec{v})$$

As previously mentioned, if our affine space is a vector space, we can identify any point in  $P_A$  with the vector translating the origin onto that point, which lets us write:

$$F(\vec{v}) := F(0 + \vec{v}) = F(0) + \vec{F}(\vec{v}) := \vec{F}(\vec{v}) + \vec{w}$$

thus, an affine map between vector spaces is simply a linear map composed with a translation.

**Theorem 3.5: Intercept Theorem**

Let  $v_1, v_2 \in \mathbb{R}$  be linearly independent. Let  $L_1$  be an affine line through  $v_2$  and  $v_1$ . Then if we have  $\alpha, \beta \in \mathbb{R}_{\neq 0}$ , and  $L_2$  is a parallel line through  $\alpha v_2$  and  $\beta v_1$ , we must have  $\alpha = \beta$ .

*Proof.* Since  $L_1$  and  $L_2$  are parallel, their direction vectors must point in

the same direction, which means there must be a  $\mu \in \mathbb{R}_{\neq 0}$  such that

$$\begin{aligned}\mu(v_2 - v_1) &= \mu(\alpha v_2 - \beta v_1) \\ \implies \mu v_2 - \mu v_1 &= \alpha v_2 - \beta v_1 \\ \implies \mu v_2 - \alpha v_2 &= \mu v_1 - \beta v_1 \\ \implies (\mu - \alpha)v_2 &= (\mu - \beta)v_1\end{aligned}$$

Since  $v_2$  and  $v_1$  are linearly independent, this implies  $\mu - \alpha = \mu - \beta = 0$ , i.e.  $\alpha = \beta = \mu$ .  $\square$

**Theorem 3.6.** An injective map  $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is linear if and only if it maps  $\vec{0}$  to  $\vec{0}$  and lines to lines.

*Proof.*  $\Rightarrow$  Let  $\Phi$  be linear. Let  $g := \{\vec{p} + \lambda \cdot \vec{v}\}$  be a line. Then we have

$$\begin{aligned}\Phi(g) &= \{\Phi(\vec{p} + \lambda \cdot \vec{v})\} \\ &= \{\Phi(\vec{p}) + \lambda \cdot \Phi(\vec{v})\},\end{aligned}$$

i.e.  $\Phi(g)$  is also a line.

$\Leftarrow$  Since  $\Phi$  maps  $\vec{0}$  to  $\vec{0}$  and lines to lines, it must map lines through the origin to lines through the origin. Let

$$\begin{aligned}\Phi(e_1) &:= v_1 \\ \Phi[\lambda e_1] &:= L_1 = \{\lambda v_1\} \\ \Phi(e_2) &:= v_2 \\ \Phi[\lambda e_2] &:= L_2 = \{\lambda v_2\}\end{aligned}$$

Since  $\Phi$  is injective,  $L_1$  and  $L_2$  can only intersect at  $\{0\}$ , and thus  $v_1$  and  $v_2$  must be linearly independent.

Now, let's take a closer look at images of individual points. Since  $\Phi[\lambda e_1] = \{\lambda v_1\}$  and  $\Phi[\lambda e_2] = \{\lambda v_2\}$ , we must have

$$\begin{aligned}\Phi[\lambda e_1] &= \varphi_1(\lambda)v_1 \\ \Phi[\lambda e_2] &= \varphi_2(\lambda)v_2\end{aligned}$$

for some  $\varphi_1, \varphi_2 : \mathbb{R} \rightarrow \mathbb{R}$ .

Now, notice that the line through  $\lambda e_1$  and  $\lambda e_2$  is parallel to the line through  $e_1$  and  $e_2$ . Since  $\Phi$  is injective, it must preserve parallelism, and so the line through  $v_1$  and  $v_2$  must be parallel to the line through  $\varphi_1(\lambda)v_1$  and  $\varphi_2(\lambda)v_2$ . The intercept theorem now gives us  $\varphi_1(\lambda) = \varphi_2(\lambda) = \mu$ . We thus write  $\varphi(\lambda) := \varphi_1(\lambda) = \varphi_2(\lambda)$ .

Since  $\Phi$  is injective,  $\varphi$  must be as well, since otherwise we would have some  $v$  with  $\Phi(\lambda_1 v) = \Phi(\lambda_2 v)$  despite  $\lambda_1 v \neq \lambda_2 v$ . We have also have  $\varphi(0) = 0$  and  $\varphi(1) = 1$ , since  $\Phi(0) = 0$  and  $\Phi(e_1) = v_1$ .

Now, we want to show that  $\Phi(v + w) = \Phi(v) + \Phi(w)$  holds for linearly independent  $v, w$ . Since  $\Phi$  preserves parallelism, it must map parallelograms to parallelograms. Therefore, the parallelogram spanned by  $v$  and  $w$  must be mapped to the parallelogram spanned by  $\Phi(v)$  and  $\Phi(w)$ . This means that the upper corner  $v + w$  gets mapped to the upper corner  $\Phi(v) + \Phi(w)$  which gives us  $\Phi(v + w) = \Phi(v) + \Phi(w)$  as desired.

Putting our insights together, we now know that

$$\begin{aligned}\Phi(\lambda_1 e_1 + \lambda_2 e_2) &= \Phi(\lambda_1 e_1) + \Phi(\lambda_2 e_2) \\ &= \varphi(\lambda_1)\Phi(e_1) + \varphi(\lambda_2)\Phi(e_2)\end{aligned}$$

it only remains to be shown that  $\varphi$  is the identity. For this, consider that we have:

$$\begin{aligned}(\varphi(a) + \varphi(b))\Phi(e_1) &= \varphi(a)\Phi(e_1) + \varphi(b)\Phi(e_1) \\ &= \Phi(ae_1 + be_1) \\ &= \Phi((a + b)e_1) \\ &= \varphi(a + b)\Phi(e_1),\end{aligned}$$

which gives us  $\varphi(a) + \varphi(b) = \varphi(a + b)$ . For  $a \neq 0$ , the line through  $e_1$  and  $ae_2$  must point in the same direction as the line through  $be_1$  and  $bae_2$ , so the line through  $\Phi(e_1)$  and  $\varphi(a)\Phi(e_2)$  must be parallel to the line through  $\varphi(b)\Phi(e_1)$  and  $\varphi(ab)\Phi(e_2)$ , from which the intercept theorem gives us

$$\begin{aligned}\frac{\varphi(ab)}{\varphi(a)} &= \varphi(b) \\ \implies \varphi(ab) &= \varphi(a)\varphi(b)\end{aligned}$$

Since we have  $\varphi(0) = 0$ ,  $\varphi(1) = 1$ ,  $\varphi(a + b) = \varphi(a) + \varphi(b)$ ,  $\varphi(ab) = \varphi(a)\varphi(b)$ , our map  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  must be a field homomorphism, of which the identity is the only one. Thus,  $\Phi$  is linear.

□

**Exercise 3.7.** *An injective map between real affine spaces of dimension  $\geq 2$  is affine if and only if it maps lines to lines.*

## **Chapter 4**

# **Determinants**

### **4.1 Defining Volume**

### **4.2 Determinants of Endomorphisms**

### **4.3 Orientation**

## Chapter 5

# Eigenvectors and Eigenvalues

### Definition 5.1: Eigenvalues and Eigenvectors

Let  $k$  be a field,  $F : V \rightarrow V$  an Endomorphism, and  $\lambda \in k$ . Then the  $\lambda$ -*eigenspace* of  $F$  is the set

$$\begin{aligned}\text{Eig}(F, \lambda) &:= \{\vec{v} \in V : F(\vec{v}) = \lambda\vec{v}\} \\ &= \{\vec{v} \in V : (F - \lambda \text{id}_V)(\vec{v}) = 0\} \\ &= \ker(F - \lambda \text{id}_V)\end{aligned}$$

If  $F(\vec{v}) = \lambda\vec{v}$ , with  $\vec{v} \neq \vec{0}$ , we say that  $\vec{v}$  is an *eigenvector* of  $F$  with *eigenvalue*  $\lambda$ .

Note that we exclude  $\vec{0}$  from being an eigenvector, since  $F(\vec{0}) = \lambda\vec{0}$  trivially holds for every endomorphism. Thus every  $\lambda$  would be an eigenvalue, which would make the definition pointless. We do, however, allow 0 as an eigenvalue - the 0-eigenspace of  $F$  is simply its kernel. Sometimes it ends up being useful to rephrase our definition of eigenspaces in terms of kernels:

$$\begin{aligned}\text{Eig}(F, \lambda) &= \{\vec{v} \in V : F(\vec{v}) = \lambda\vec{v}\} \\ &= \{\vec{v} \in V : F(\vec{v}) - \lambda\vec{v} = 0\} \\ &= \{\vec{v} \in V : (F - \lambda \text{id}_V)(\vec{v}) = 0\} \\ &= \ker(F - \lambda \text{id}_V)\end{aligned}$$

This shows that  $\text{Eig}(F, \lambda)$  is a subspace of  $V$  - by definition, these subspaces must then be the maximal subspaces that are closed under application of  $F$ , i.e.  $F(\text{Eig}(F, \lambda)) = \text{Eig}(F, \lambda)$ .

**Exercise 5.2.** *In an informal, intuition-based way:*

1. Find the real eigenvalues and eigenspaces of a reflection of  $\mathbb{R}^2$  along an axis  $\{\lambda\vec{v} : \lambda \in \mathbb{R}\}$
2. Find a rotation of  $\mathbb{R}^2$  that has no real eigenvalues. Find two different rotations of  $\mathbb{R}^2$  that do have real eigenvalues. Show that any rotation of  $\mathbb{R}^2$  has complex eigenvalues.
3. Find the eigenvectors of the derivative map  $\frac{d}{dx}$

Our characterisation  $\text{Eig}(F, \lambda) = \ker(F - \lambda \text{id}_V)$  means that, if  $F$  is finite-dimensional, we can find all eigenvectors of  $F$  with a given eigenvalue  $\lambda$  by solving  $(F - \lambda \text{id}_V)\vec{v} = 0$ , which is simply a system of linear equations. Finding the correct eigenvalues ends up being more tricky - in the section after this one, we will work towards proving the following key result:

**Proposition 5.3.** The eigenvalues of  $F$  are given by the zeroes of the *characteristic polynomial*  $\chi_F(\lambda) := \det(F - \lambda \text{id}_V)$ .

However, before we do so, we want to familiarize ourselves with some of the more basic properties of eigenvectors and eigenvalues.

One important application is that finding eigenvectors lets us diagonalize endomorphisms, greatly easing computation afterwards:

**Theorem 5.4.** Let  $V$  be a vector space over a field  $k$ . Let  $B = (b_1, \dots, b_n)$  be a basis of  $V$ , and let  $F$  be an endomorphism represented in  $B$  by a matrix  $A$ . Then:

1. A basis vector  $b_i$  is an eigenvector of  $F$  iff. the  $i$ -th column of  $A$  has the form  $\lambda e_i$ .
2.  $F$  is diagonalizable iff. there exists a basis of  $V$  consisting of eigenvectors of  $F$ .

**Theorem 5.5.** Let  $V$  be a vector space over a field  $k$ . Let  $F : V \rightarrow V$  be an endomorphism. Then eigenvectors of  $F$  with different eigenvalues are linearly independent.

**Corollary 5.6.** Let  $V$  be an  $n$ -dimensional vector space over a field  $k$ . Let  $F : V \rightarrow V$  be an endomorphism. Then:

1.  $F$  has no more than  $n$  distinct eigenvalues,
2. If  $(\lambda_i)_{i \in I}$  are mutually distinct eigenvalues of  $F$ , the sum

$$\sum_{i \in I} \text{Eig}(F, \lambda_i) \subseteq V$$

is direct.

3.  $F$  is diagonalizable if and only if  $V$  is a direct sum of eigenspaces of  $F$ .
4. If  $F$  has  $n$  distinct eigenvalues, it is diagonalizable.

**Corollary 5.7.** Let  $V, W$  be  $n$ -dimensional vector spaces over a field  $k$ . Let  $F : V \rightarrow V, G : W \rightarrow W$  be endomorphisms. Then there exists a unique endomorphism  $P : V \rightarrow W$  such that  $P \circ F = G \circ P$  if and only if  $F$  and  $G$  have the same eigenvalues.

## 5.1 Characteristic Polynomials

### Theorem 5.8: Characteristic Polynomial

Let  $V$  be a finite-dimensional vector space over a field  $\mathbb{k}$ . Let  $F : V \rightarrow V$  be an endomorphism. Then  $\lambda \in \mathbb{k}$  is an eigenvalue of  $F$  if and only if

$$\chi_F(\lambda) = \det(\lambda \cdot \text{id}_V - F) = 0_{\mathbb{k}}$$

We call  $\chi_F$  the *characteristic polynomial of  $F$* .

*Proof.* We know that  $\lambda \in \mathbb{k}$  is an Eigenvalue of  $F$  if and only if

$$\ker(F - \lambda \text{id}_V) \neq \{0_V\},$$

i.e. if and only if  $F - \lambda \text{id}_V$  isn't injective.

We also know that  $F - \lambda \text{id}_V$  isn't injective if and only if it isn't surjective, i.e. if and only if it isn't invertible, i.e. if and only if

$$\det(\lambda \cdot \text{id}_V - F) = \chi_F(\lambda) = 0_{\mathbb{k}}.$$

□

**Exercise 5.9.** Show that every endomorphism  $F : V \rightarrow V$  of a finite-dimensional complex vector space  $V$  has at least one eigenvalue.

**Exercise 5.10.** Show that the constant term of a characteristic polynomial  $\chi_F(\lambda)$  is given by  $(-1)^n \cdot (\det F)$ .

(...)

## 5.2 Algebras and the Cayley-Hamilton Theorem

### Theorem 5.11: Evaluation map

Let  $R$  be a commutative unital ring, and let  $A$  be a commutative unital algebra over  $R$ . Then, for every  $a \in A$ , there exists a unique algebra homomorphism  $\text{ev}_a : R[X] \rightarrow A$  such that:

1.  $\text{ev}_a(r) = 1_A \cdot r$  for all constant polynomials  $r \in R[X]$ ,
2.  $\text{ev}_a(X) = a$ .

We call  $\text{ev}_a$  the *evaluation map of  $R[X]$  at  $a$* .

Informally, this means that we can take elements of an algebra  $A$  over  $R$  and plug them into polynomials that were originally only defined on  $R$ , and we won't run into issues. In particular, we can plug matrices into polynomials over an underlying ring! Since  $\text{ev}_a$  is unique, we will generally just write  $P(a) := \text{ev}_a(P)$  from now on.

**Theorem 5.12: Cayley-Hamilton**

Let  $R$  be a commutative unital ring. Let  $A \in R^{n \times n}$ . Then  $\chi_A(A) = 0$ .

## 5.3 Minimal Polynomials

**Theorem 5.13: Minimal polynomial**

Let  $F$  be an endomorphism of a vector space  $V$  over a field  $\mathbb{k}$ . Then there exists a unique polynomial  $\mu_F \in \mathbb{k}[X]$  such that:

1.  $\mu_F$  is either normalized or equal to  $0_{\mathbb{k}[X]}$
2.  $\mu_F$  is a divisor of every  $P \in \mathbb{k}[X]$  with  $P(F) = 0_{\text{End}_{\mathbb{k}}(V)}$

We call  $\mu_F$  the *minimal polynomial* of  $F$ .

## Chapter 6

# Normal Forms of Endomorphisms

### 6.1 Decomposing Vector Spaces using Polynomials

**Exercise 6.1.** Let  $G$  be an abelian group. Show that  $G$  becomes a  $\mathbb{Z}$ -module when equipped with the operation  $n.g = \underbrace{g + \dots + g}_{n \text{ times}}$ .

**Exercise 6.2.** Let  $V$  be a vector space over a field  $\mathbb{k}$ . Show that  $V$  becomes an  $\text{End}(V)$ -Module when equipped with the operation  $F.v = F(v)$ .

**Theorem 6.3.** Any  $\mathbb{k}[X]$ -module is the same thing as a  $\mathbb{k}$ -vector space  $V$  equipped with a specified endomorphism  $F$ , and vice versa.

*Proof.* 1. Let  $V$  be a  $\mathbb{k}$ -vector space. We know from the previous exercise that  $V$  is an  $\text{End}(V)$ -module. Thus, the polynomial evaluation map  $\text{ev}_F : \mathbb{k}[X] \rightarrow \text{End}(V)$  turns  $V$  into  $\mathbb{k}$ -module via

$$P.v = \text{ev}_F(P)(v) = P(F)(v)$$

2. Now let  $M$  be a  $\mathbb{k}[X]$ -module. Then  $M$  becomes a  $\mathbb{k}$ -module, i.e. a  $\mathbb{k}$ -vector space, if we restrict scalar multiplication to only accept constant polynomials. Then it can easily be seen that  $G(v) = X.v$  is a linear transformation.

Now, note that if we start with a vector space  $(V, F)$ , then use our construction to get a module  $M$ , and then turn it back into a vector space, we will obtain our original  $F$  again, since

$$G(v) = X.Mv = X(F)(v) = F(v)$$

Conversely, if we start with a module  $M$ , then turn it into a vector space  $(V, F)$ , and then back into a module  $M$ , we reobtain our original scalar multiplication, since

$$P.Vv = P(G)(v) = P(X.M)(v) = P.Mv$$

Therefore, the two directions of our construction are inverse to each other, i.e. we actually have a bijection.  $\square$

Thus, given a  $\mathbb{K}[X]$ -module  $M$ , we can write  $M = (V, F)$

**Exercise 6.4.** Let  $M = (V, F)$  and  $N = (W, G)$  be  $\mathbb{K}[X]$ -Modules. Then a  $\mathbb{K}[X]$ -linear map  $H : M \rightarrow N$  is  $\mathbb{K}$ -linear and additionally fulfills  $H \circ F = G \circ H$ , and any  $\mathbb{K}$ -linear map  $H$  fulfilling these properties is  $\mathbb{K}[X]$ -linear.

Let  $M = (V, F)$  be a  $\mathbb{K}[X]$ -Module. Then, by definition, a subspace  $U \subseteq V$  is a  $\mathbb{K}[X]$ -submodule of  $M$  if and only if  $U$  is closed under scalar multiplication with elements of  $\mathbb{K}[X]$ .

**Exercise 6.5.** Show that  $U$  is closed under scalar multiplication with elements of  $\mathbb{K}[X]$  if and only if  $Xu \in U$ , which itself holds if and only if  $F(u) \in U$  holds for all  $u \in U$ , i.e.  $U$  is  **$F$ -invariant**.

**Exercise 6.6.** 1. Show that the scalar multiplication map  $P. = P(F)$  is linear.  
2. Conclude that  $\ker(P(F))$  and  $\text{im}(P(F))$  are submodules of  $M$ .  
3. Conclude that  $\ker(P(F))$  and  $\text{im}(P(F))$  are  $F$ -invariant.

**Theorem 6.7: Structure theorem for modules over a Euclidean ring**

Let  $M$  be a module over a euclidean domain  $R$ . Let  $a = bc \in R$  such that  $b$  and  $c$  share no divisors and  $a.m = 0$  for all  $m \in M$ . Then we have:

1.  $\ker(b.) = \text{im}(c.)$ ,
2.  $\text{im}(b.) = \ker(c.)$ ,
3.  $M = \ker(b.) \oplus \ker(c.)$ .

*Proof.* 1. Let  $c.m \in \text{im}(c.)$  for  $m \in M$ . Then we have  $b.(c.m) = a.m = 0$ , i.e.  $\text{im}(c.) \subseteq \ker(b.)$ . Now, applying the Euclidean algorithm gives us  $1 = k_1b + k_2c$ . If we now let  $m \in \ker(b.)$ , we get:

$$\begin{aligned} m &= 1m \\ &= (k_1b + k_2c)m \\ &= k_1 \underbrace{(bm)}_{=0} + k_2(cm) \\ &= c(k_2m) \in \text{im}(c.) \end{aligned}$$

Thus, we also have  $\ker(b.) \subseteq \text{im}(c.)$ , i.e.  $\ker(b.) = \text{im}(c.)$ .

2.  $\ker(c.) = \text{im}(b.)$  follows from the same argument by swapping the roles of  $b$  and  $c$ .

3. Now, let  $m \in M$ . Then, applying our identity from before, we get

$$\begin{aligned} m &= (k_1b + k_2c)m \\ &= b(k_1m) + c(k_2m) \\ &\in \text{im}(b.) + \text{im}(c.) \\ &= \ker(b.) + \ker(c.), \end{aligned}$$

which gives us  $M = \ker(b.) + \ker(c.)$  as desired.

If we have  $m \in \ker(b.) \cap \ker(c.)$ , we get

$$m = k_1(b(m)) + k_2(c(m)) = 0,$$

so  $\ker(b.) \cap \ker(c.) = \{0\}$ , so our sum is direct. □

**Corollary 6.8.** Let  $G$  be an abelian group. Let  $n = mk \in \mathbb{Z}$  such that  $m$  and  $k$  share no divisors and  $n \cdot g = 0$  for all  $g \in G$ . Then we have

$$G = \ker(m.) \oplus \ker(k.).$$

#### Corollary 6.9: Chinese Remainder Theorem

**Corollary 6.10.** Let  $V$  be a vector space over a field  $\mathbb{k}$  and  $F : V \rightarrow V$  be an Endomorphism. Let  $P = QR \in \mathbb{k}[X]$  such that  $Q$  and  $R$  share no divisors and  $P.v = P(F)(v) = 0$  for all  $v \in V$ . Then we have

$$V = \ker(Q(F)) \oplus \ker(R(F)).$$

#### Corollary 6.11: $P$ -Generalized Eigenspace Decomposition

Let  $V$  be a vector space over a field  $\mathbb{k}$  and  $F : V \rightarrow V$  be an Endomorphism. Let  $P \in \mathbb{k}[X]$  be a normed polynomial and let  $P = P_1^{n_1} \cdot \dots \cdot P_k^{n_k}$  be its prime factorization. Then we have

$$V = \bigoplus_{i=1}^k \ker(P_i^{n_i}(F))$$

## 6.2 The Jordan Normal Form

### Corollary 6.12: Generalized Eigenspace Decomposition

Let  $V$  be a vector space over a field  $\mathbb{K}$  and  $F : V \rightarrow V$  be an Endomorphism. Assume that its minimal polynomial  $\mu_F$  splits into linear factors over  $\mathbb{K}$ , i.e.

$$\mu_F(x) = \prod_{i=1}^k (x - \lambda_i)^{n_i}$$

with  $\lambda_i \in \mathbb{K}$ . Then we have

$$V = \bigoplus_{i=1}^k \ker((F - \lambda_i \text{id}_V)^{n_i})$$

We call  $\ker((F - \lambda_i \text{id}_V)^{n_i})$  the **generalized  $\lambda_i$ -Eigenspace of  $F$**  and denote it  $H(F, \lambda_i)$  (from the German "Hauptraum").

**Exercise 6.13.** Show that  $H(F, \lambda_i)$  contains all eigenvectors of  $F$  with eigenvalue  $\lambda_i$ .

**Exercise 6.14.** Show that  $H(F, \lambda_i)$  is an  $F$ -invariant subspace of  $V$ , i.e. a subspace such that  $F(H(F, \lambda_i)) \subseteq H(F, \lambda_i)$

**Exercise 6.15.** Let  $V = V_1 \oplus V_2$  be a decomposition of  $V$  into  $F$ -invariant subspaces. Let  $B_1$  be a basis of  $V_1$ , and let  $B_2$  be a basis of  $V_2$ . Show that:

1.  $B = B_1 \cup B_2$  is a basis of  $V$
2. If the restriction  $F|_{V_1}$  of  $F$  to  $V_1$  is represented in basis  $B_1$  as a matrix  $A_1$ , and the restriction  $F|_{V_2}$  of  $F$  to  $V_2$  is represented in basis  $B_2$  as a matrix  $A_2$ , then  $F$  is represented in basis  $B$  as

$$A = \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix}$$

**Exercise 6.16.** Let  $F : V \rightarrow W$  be a linear map. Let  $A$  be a basis of  $\ker F$ , and let  $B$  be a tuple of vectors such that  $F(B)$  is a basis of  $\text{im } F$ . Then  $A \cup B$  is a basis of  $V$ .

We now want to make use of this to construct a Jordan basis  $B$  of a nilpotent Endomorphism  $N$ :

**Corollary 6.17.** Let  $N : V \rightarrow V$  be a nilpotent endomorphism. Then there exists a basis  $B$  of  $V$  such that:

1.  $B \cup \{0\}$  is  $F$ -invariant, i.e. if  $b \in B$ , then we have  $F(b) \in B$  or  $F(b) = 0$ .
2. For every  $b \in B$ , there exists at most one element  $b' \in B$  with  $F(b') = b$ .

We call such a basis a **Jordan basis**.

*Proof.* We prove this theorem in three steps: We construct a basis  $B$ , then show it is  $F$ -invariant, then show that each element of  $B$  has at most one preimage in  $B$ .

1. We construct our basis by induction over  $n = \dim V$ .

If  $n = 0$ , then the empty set is a basis (since a sum over the empty set equals the additive identity, i.e. the zero vector) which (vacuously) fulfills our desired properties.

Now, since the kernel of  $N$  is non-trivial, we have  $\dim \operatorname{im} N < \dim V$ . Therefore, by induction, we can assume we already have a Jordan basis  $A$  of  $\operatorname{im} N$ . We split this basis  $A$  into two parts:

- (a)  $A_0 := \{a \in A \mid N(a) = 0\}$ ,
- (b)  $A_{\neq 0} := \{a \in A \mid N(a) \neq 0\}$ .

$A_0$  is by definition a set of linearly independent vectors contained in  $\ker A$ . By the Steinitz exchange lemma, we can add a set  $C$  of linearly independent vectors such that  $A_0 \cup C$  is a basis of  $\ker A$ .

$A_{\neq 0}$  is the set of all  $A$  such that  $N(A_{\neq 0}) \subseteq A$ . We add a set  $D$  containing preimages of all the remaining elements of  $A$ , which must exist since  $A \subseteq \operatorname{im} N$ . By construction, we now have  $F(A_{\neq 0} \cup D) = A$ .

Putting everything together,  $A_0 \cup C$  is a basis of  $\ker N$ , and  $F(A_{\neq 0} \cup D) = A$ , which is a basis of  $\operatorname{im} N$ . Therefore, by exercise 6.16,  $B = A \cup C \cup D$  is a basis of  $V$ .

2. Now,  $B$  must be  $F$ -invariant since  $A$  is  $F$ -invariant by definition,  $C$  is contained in the kernel of  $F$  and thus  $F(B) = \{0\}$ , and  $D$  consists of preimages of  $A$ , i.e.  $F(D) = A \subseteq B$ .
3. Finally, the preimages of  $A$  in  $B$  are contained in either  $A_{\neq 0}$  or  $D$ , which are disjoint by construction.

□

**Corollary 6.18.** Let  $F : V \rightarrow V$  be a matrix whose characteristic polynomial has  $n = \dim(V)$  roots. Then there exists a basis  $B$  of  $V$  such that  ${}_B F_B$  is a block diagonal matrix consisting of Jordan blocks.

*Proof.* Generalized Eigenspace decomposition 6.2 shows us that  $V$  is the direct sum of the generalized Eigenspaces of  $F$ . We know by exercise 6.14 that the generalized Eigenspaces of  $F$  are  $F$ -invariant. Therefore, exercise 6.15 tells us that if we find a basis  $B_i$  for each generalized Eigenspace  $H(F, \lambda_i)$  such that  $F|_{H(F, \lambda_i)}$  is represented by a matrix  $J_i$  in Jordan normal form, and we combine our bases into a basis  $B = \bigcup_{i=1}^n B_i$  of  $V$  as a whole, then the matrix representation of  $F$  in basis  $B$  will be a block diagonal matrix consisting of these  $J_i$ , and thus  $F$  will also be in Jordan normal

form.

Now, let  $F_i$  be the restriction of  $(F - \lambda_i \text{id}_V)$  to  $H(F, \lambda_i)$ . By definition of the generalized Eigenspace, we have

$$H(F, \lambda_i) = \ker((F - \lambda_i \text{id}_V)^{n_i})$$

therefore, by construction,  $F_i$  is nilpotent. Therefore, by theorem ??, there exists a basis  $B_i$  such that  ${}_{B_i}F_i{}_{B_i}$  is in Jordan normal form, with zeroes on the diagonal. But then, since we have

$$F|_{H(F, \lambda_i)} = F_i + \lambda_i \text{id}_{H(F, \lambda_i)}$$

the matrix representation of  $F|_{H(F, \lambda_i)}$  is also in Jordan form, since we simply have to add  $\lambda$  to the diagonal of  ${}_{B_i}F_i{}_{B_i}$ .

Therefore, the restriction to  $F$  to any of its generalized Eigenspaces is in Jordan normal form in basis  $B_i$ . As discussed at the beginning of this proof, this means  $F$  is in Jordan normal form in basis  $B = \bigcup_{i=1}^n B_i$ .  $\square$

**Exercise 6.19.** Let  $F$  be an endomorphism, and let  $J$  be its Jordan normal form. Then:

1. The algebraic multiplicity of each eigenvalue of  $F$  corresponds to the number of times that eigenvalue appears on the diagonal of  $J$ .
2. The geometric multiplicity of any eigenvalue  $\lambda$  of  $F$  corresponds to the number of  $\lambda$ -Jordan blocks with that eigenvalue in  $J$ .
3. The exponent of  $(X - \lambda)$  in the minimal polynomial of  $F$  corresponds to the size of the largest  $\lambda$ -Jordan block in  $J$ .

## 6.3 The Weierstrass Normal Form

# Chapter 7

## Inner Product Spaces

### 7.1 Inner Products

**Definition 7.1.** Let  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $V$  and  $W$  be vector spaces over  $\mathbb{k}$ . Then a map  $\varphi : V \rightarrow W$  is called ( $\mathbb{k}$ -) *semilinear* or *antilinear*, if we have

1. **Additivity:**  $\forall u, v \in V : \varphi(u + v) = \varphi(u) + \varphi(v)$
2. **Antihomogeneity:**  $\forall v \in V, \lambda \in \mathbb{k} : \varphi(\lambda \cdot v) = \overline{\lambda} \cdot \varphi(v)$ .

Notice that for  $\mathbb{k} = \mathbb{R}$ , there is no difference between linearity and antilinearity, since complex conjugation leaves real numbers unchanged.

**Definition 7.2.** Let  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $V$  be a  $\mathbb{k}$ -vector space. Then a map  $s : V \times V \rightarrow \mathbb{k}$  is called a *sesquilinear form* if it is antilinear in the first component and linear in the second component, i.e. the map  $s_w : v \rightarrow s(v, w)$  is antilinear and the map  $s_v : w \rightarrow s(v, w)$  is linear.

**Exercise 7.3.** Let  $A$  be a matrix with entries in  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Show that the map

$$s_A(v, w) := v^* A w$$

is sesquilinear. (Recall that  $v^*$  stands for the conjugated transpose of  $v$ .)

The latin prefix "sesqui-" means "one and a half" - a sesquilinear is a map that is linear in one component, and "sort of linear", i.e. "one-half-linear" in the other. Some sources require that a sesquilinear form is linear in the first component and antilinear in the second.

**Definition 7.4.** A sesquilinear form  $s : V \times V \rightarrow \mathbb{k}$  is called a *Hermitian form* if it is *conjugate symmetric*, i.e.  $s(v, w) = \overline{s(w, v)}$ .

Once again, notice that for  $\mathbb{k} = \mathbb{R}$ , this just means  $s(v, w) = s(w, v)$ .

**Exercise 7.5.** Show that a matrix is Hermitian (i.e. fulfills  $A = A^*$ ) if and only if its associated sesquilinear form  $s_A(v, w) = v^* A w$  is Hermitian.

**Definition 7.6.** A sesquilinear form  $s : V \times V \rightarrow \mathbb{k}$  is called *positive semidefinite* if  $s(v, v) \geq 0$  for all  $v \in V$ . It is called *positive definite* if it additionally fulfills  $s(v, v) = 0$  if and only if  $v = \vec{0}$ .

Positive definiteness is the primary reason we are working with sesquilinear forms instead of bilinear forms - we need to be able to compare  $s(v, v)$  to 0,

and there is no way to order all of  $\mathbb{C}$  that is compatible with its field structure. However, if we have a sesquilinear Hermitian map, we get  $s(v, v) = \overline{s(v, v)}$ , i.e.  $s(v, v) \in \mathbb{R}$ , which means we can start comparing things again.

**Definition 7.7.** A *scalar product* or *inner product* on a vector space  $V$  over  $\mathbb{R}$  or  $\mathbb{C}$  is a positive definite Hermitian form. We call a real vector space equipped with a scalar product *Euclidean*, and a complex vector space equipped with a scalar product *unitary*.

**Definition 7.8.** The standard scalar product on  $\mathbb{C}^n$  is given by

$$\langle v, w \rangle_{\mathbb{C}} = v^* w$$

**Exercise 7.9.** Show that  $\langle v, w \rangle_{\mathbb{C}}$  is actually a scalar product.

**Exercise 7.10.** Let  $V = C^\infty([0, 1], \mathbb{R})$  be the room of infinitely differentiable functions  $[0, 1] \rightarrow \mathbb{R}$ . Show that the  $L^2$  scalar product

$$\langle f, g \rangle_{L^2} = \int_0^1 f(t)g(t) dt$$

is actually a scalar product.

**Definition 7.11.** Let  $V$  be a vector space over  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ . Then a *Norm* is a map  $\|\cdot\| : V \rightarrow \mathbb{K}$  fulfilling the following three axioms:

1. **Positive definiteness:**  $\|v\| \geq 0$  and  $\|v\| = 0$  iff.  $v = \vec{0}$
2. **Homogeneity:**  $\|\lambda \cdot v\| = |\lambda| \cdot \|v\|$
3. **Triangle inequality:**  $\|v + w\| \leq \|v\| + \|w\|$

A norm is a way of assigning lengths to vectors. Once lengths of vectors are defined, one can easily define distances between vectors too:

**Exercise 7.12.** Let  $\|\cdot\|$  be a norm on a vector space  $V$  over a field  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ . Show that  $d(v, w) := \|v - w\|$  is a metric on  $V$ .

#### Theorem 7.13: Cauchy-Schwartz Inequality

Let  $s$  be a scalar product on a vector space  $V$  over a field  $\mathbb{K}$ . Then we have

$$|\langle v, w \rangle| \leq \sqrt{\langle v, v \rangle} \cdot \sqrt{\langle w, w \rangle}$$

One important property of scalar products is that they induce norms. Notice that the standard norm on  $\mathbb{R}^n$  is just the square root of the standard scalar product on  $\mathbb{R}^n$ . This property generalizes to all scalar products:

**Exercise 7.14.** Let  $s : V \times V \rightarrow \mathbb{K}$  be a scalar product on a vector space  $V$  over a field  $\mathbb{K}$ . Let  $\|v\|_s := \sqrt{s(v, v)}$ . Show that:

1.  $\|\cdot\|_s$  is a norm on  $V$ .
2. We have the **parallelogram identity**

$$\|v + w\|_s^2 + \|v - w\|_s^2 = 2\|v\|_s^2 + 2\|w\|_s^2$$

3. If  $\mathbb{K} = \mathbb{R}$ , we have the **real polarisation formula**

$$s(v, w) = \frac{1}{4} \left( \|v + w\|_s^2 - \|v - w\|_s^2 \right)$$

4. If  $\mathbb{k} = \mathbb{C}$ , we have the **complex polarisation formula**

$$s(v, w) = \frac{1}{4} \left( \|v + w\|_s^2 - \|v - w\|_s^2 - i\|v + i.w\|_s^2 + i\|v - i.w\|_s^2 \right)$$

**Exercise 7.15.** Find a norm on a vector space that cannot be constructed as the square root of a scalar product.

**Theorem 7.16.** Let  $V$  be a vector space over  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $\|\cdot\|$  be a norm on  $V$  fulfilling the parallelogram identity. Then there exists a corresponding scalar product  $s$  such that  $\|v\| = \sqrt{s(v, v)}$

## 7.2 Orthogonality

**Definition 7.17.** Let  $(V, s)$  be a vector space with scalar product over a field  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Then we call two vectors  $v, w$  **orthogonal** iff.

$$s(v, w) = 0$$

**Exercise 7.18.** Let  $B = (v_1, \dots, v_k)$  be a tuple of orthogonal vectors not containing the zero vector. Then  $B$  is linearly independent.

**Definition 7.19.** A basis consisting of pairwise orthogonal vectors is called an **orthogonal basis**. If the basis vectors are also normalized, i.e. have length 1, we call the basis an **orthonormal basis**.

**Exercise 7.20.** Let  $(V, s)$  be a vector space with scalar product. Show that  $B = (b_1, \dots, b_n) \subset V$  is an orthonormal basis iff.

$$s(b_i, b_j) = \delta_{ij}$$

(Recall that  $\delta_{ij}$  is defined as the function that is 1 if  $i = j$  and 0 otherwise).

**Definition 7.21.** Let  $(V, s)$  be a vector space with scalar product over a field  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $U \subseteq V$  be a subspace. Let  $B_U := \{b_1, \dots, b_m\}$  be a basis of  $U$ . Then the **orthogonal projection of  $V$  onto  $U$**  is the map

$$p : V \rightarrow U$$

$$v \mapsto \sum_{i=1}^m g(b_i, v) \cdot b_i$$

**Exercise 7.22.** 1. Show that the orthogonal projection of  $V$  onto  $U$  leaves vectors in  $u$  unchanged.

2. Show that for each  $v \in V$  and  $u \in U$ ,  $u$  is orthogonal to  $v - p(v)$ .

3. Show that for each  $v \in V$ ,  $p(v)$  is the unique point in  $U$  for which  $\|v - p(v)\|_s$  is minimal.

**Theorem 7.23.** Let  $(V, s)$  be a vector space with scalar product over a field  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $(v_1, \dots, v_n)$  be a basis of  $V$ . Then there are unique vectors  $(b_1, \dots, b_n) \subset V$  such that for all  $p \in (1, \dots, n)$ , we have:

1.  $(b_1, \dots, b_p)$  is an  $s$ -orthonormal Basis of  $\langle v_1, \dots, v_p \rangle \subseteq V$
2.  $s(b_p, v_p) \in \mathbb{R}_{>0}$

These vectors  $b_p$  can be constructed inductively via  $b_1 = \frac{v_1}{\|v_1\|}$  followed by projection onto the already constructed basis vectors, i.e.

$$w_p = v_p - \sum_{q=1}^{p-1} g(b_q, v_p) \cdot b_q$$

$$\hat{b}_p = \frac{w_p}{\|w_p\|}$$

**Corollary 7.24.** Every vector space with scalar product has an orthonormal basis.

## 7.3 Matrix Representations of Inner Products

**Definition 7.25.** Let  $s$  be a sesquilinear form on a finite dimensional vector space  $V$  over a field  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $B = (b_1, \dots, b_n)$  be a basis of  $V$ . Then the **Gram matrix of  $B$  by  $g$**  is defined as the matrix with entries

$$G_{ij} = s(b_i, b_j) \in \mathbb{k}^{n \times n}$$

**Exercise 7.26.** Show that the Gram matrix of the standard basis of  $\mathbb{R}^n$  by the scalar product is the identity matrix.

**Definition 7.27.** A matrix  $A \in \mathbb{k}^{n \times n}$  is called **positive (semi-)definite** if  $s_A(v, w) = v^* A w$  is positive (semi-)definite.

**Exercise 7.28.** Let  $s$  be a sesquilinear form on a finite-dimensional vector space  $V$  over a field  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Let  $G$  be its Gram matrix. Then we have  $s(v, w) = v^* G y$  in basis  $B$ . In particular,  $s$  is a scalar product if and only if its Gram matrix is Hermitian and positive definite.

**Theorem 7.29.** Let  $A \in \mathbb{k}^{n \times n}$ , where  $\mathbb{k} = \mathbb{R}$  or  $\mathbb{C}$ . Then the following are equivalent:

1.  $A$  is hermitian and positive definite.
2. **Cholesky Decomposition:** There exists an upper triangular Matrix  $B$  with positive real diagonal entries such that  $A = B^* B$ .
3. There exists an invertible Matrix  $B$  such that  $A = B^* B$ .
4. **Sylvester-Hurwitz Criterion:**  $A$  is Hermitian and all of its square upper left submatrices  $A_r$ , known as its **leading principal minors**, have strictly positive determinant.

## 7.4 Dual Spaces

### Definition 7.30: Dual Space

Let  $V$  be a vector space over a field  $\mathbb{k}$ . Then we define  $V^* = \text{Hom}_{\mathbb{k}}(V, \mathbb{k})$ . Elements of  $V^*$ , i.e. linear maps  $V \rightarrow \mathbb{k}$ , are known as **linear forms**.

**Theorem 7.31: Dual Basis**

iven a basis  $B = (b_1, \dots, b_n)$  of a vector space  $V$ , the set of linear forms  $E = (\varepsilon_1, \dots, \varepsilon_n)$  given by

$$\varepsilon_p(e_q) = \delta_{pq}$$

is a basis of  $V^*$ , known as the **dual basis** of  $B$ .

**Exercise 7.32.** Let  $C^n(\mathbb{K}, \mathbb{K})$ ,  $n \in 0, 1, \dots, \infty$  be the set of  $n$ -times continuously differentiable functions  $\mathbb{K} \rightarrow \mathbb{K}$ . Then for any  $x_0 \in \mathbb{K}$ , the map

$$\cdot(x_0) : f \rightarrow f^{(n)}(x_0)$$

is a linear form on  $C^n(\mathbb{K}, \mathbb{K})$  (where  $f^{(n)}$  denotes the  $n$ -th derivative of  $f$ ).

**Exercise 7.33.** The map mapping an integrable function  $f$  to  $\int_0^1 f(x) dx$  is a linear form. More generally, for any integrable  $g : [0, 1] \rightarrow \mathbb{K}$ , the map taking  $f$  to  $\int_0^1 \overline{g(x)} \cdot f(x) dx$  is a linear form.

**Theorem 7.34.** Let  $(V, g)$  be a  $\mathbb{K}$ -vector space with scalar product. Then  $g$  induces an injective antilinear map  $g_1 : V \rightarrow V^*$  via

$$g_1(v) = g(v, \cdot)$$

A linear form  $\alpha \in V^*$  is called **representable** by  $g$  if there exists a  $v \in V$  such that  $\alpha = g(v, \cdot)$ .

**Definition 7.35.** Let  $V$  be a  $\mathbb{K}$ -vector space. Then a **antilinear form** is an antilinear forms  $\gamma : V \rightarrow \mathbb{K}$ . The **antidual space**  $\overline{V}^*$  of  $V$  is the space of antilinear maps  $V \rightarrow \mathbb{K}$ .

**Theorem 7.36.** Let  $(V, g)$  be a finite-dimensional vector space over a field  $\mathbb{K}$ . Then the maps  $g : V \rightarrow V^*$  and  $\overline{g} : V \rightarrow \overline{V}^*$  are bijective.

**Corollary 7.37.** Let  $(V, g)$  be a finite-dimensional vector space. Then all linear and antilinear forms on  $V$  are representable by  $g$ .

**Definition 7.38.** Let  $(V, g)$  and  $(W, h)$  be  $\mathbb{K}$ -vector spaces with scalar products. Then a linear map  $F : V \rightarrow W$  is called **adjointable** if there exists a map  $G : W \rightarrow V$  such that

$$g(G(w), v) = h(w, F(v))$$

for all  $v \in V$  and  $w \in W$ . Such a map  $G$  is known as the **adjoint of  $F$** , and is written  $G = F^*$ . If  $F = F^*$ , we call  $F$  **self-adjoint**.

**Proposition 7.39.** If it exists, the adjoint of a given map is unique.

**Proposition 7.40.** Adjoint maps are linear.

**Proposition 7.41.** If  $G$  is the adjoint of  $F$ , then  $F$  is the adjoint of  $G$ , i.e. we have  $(F^*)^* = F$ .

**Proposition 7.42.** If  $A$  is a matrix representation of  $F$  by two orthonormal bases, then the conjugate transpose  $A^*$  is a matrix representation of  $F^*$ .

**Definition 7.43.** Let  $V$  and  $W$  be  $\mathbb{K}$ -vector spaces, with  $\mathbb{K}$  a field. Let  $F : V \rightarrow W$  be a linear map. Then the *dual* of  $F$  is the map

$$F^\vee : W^* \rightarrow V^* \\ \beta \mapsto \beta \circ F$$

i.e.

$$F^\vee(\beta)(\vec{v}) = \beta(F(\vec{v}))$$

for  $\beta \in W^*$ ,  $\vec{v} \in V$ .

We can construct a dual of any linear map this way, while linear maps involving infinite-dimensional vector spaces are generally not adjointable.

**Exercise 7.44.** Verify the following properties of the dual map:

1.  $(\text{id}_V)^\vee = \text{id}_{V^*}$
2. Let  $F : V \rightarrow W$  and  $G : U \rightarrow V$  be linear, then  $(F \circ G)^\vee = G^\vee \circ F^\vee$
3. The dual of a linear map is linear.
4. If  $F : V \rightarrow W$  is given by  $F(\vec{v}) = A \cdot \vec{v}$  for some matrix  $A$ , then  $F^\vee(\beta) = \beta \cdot A$ , where elements  $\beta \in W^*$  are represented as row vectors.

**Theorem 7.45.** Let  $U \subseteq V$  be vector spaces. Let  $\iota : U \rightarrow V$  be the inclusion map. Let  $q : V \rightarrow V/U$  be the quotient map. Then  $(V/U)^* \cong \text{im } q^\vee = \ker \iota^\vee$ .

*Proof.* Since  $q$  maps  $V$  to  $V/U$ , we have  $q^\vee : (V/U)^* \rightarrow \text{im } q^\vee \subseteq V^*$ . Since  $q$  is linear, so is  $q^\vee$ . By definition, any map surjectively maps onto its image. Therefore, all we need to show to obtain  $(V/U)^* \cong \text{im } q^\vee$  is injectivity.

So, let  $\beta \in \ker q^\vee \subseteq (V/U)^*$ , i.e. let  $q^\vee(\beta) = \beta(q)$  be the zero map. Since  $q$  is surjective,  $\beta(q(v)) = 0$  for all  $v \in V$  already implies  $\beta(w) = 0$  for all  $w \in V/U$ , so  $\beta$  must have already been the zero map.

Finally, we want to show  $\text{im } q^\vee = \ker \iota^\vee$ .

First off, recall that  $\iota^\vee : V^* \rightarrow U^*$ . By the universal property of the quotient map, for every  $\beta \in \ker \iota^\vee \subseteq V^*$ , there exists  $\bar{\beta} : V/U \rightarrow \mathbb{K}$  (i.e.  $\bar{\beta} \in (V/U)^*$ ) with  $\beta = \bar{\beta} \circ q = \bar{\beta}(q) \in \text{im } q^\vee$ , so we have  $\ker \iota^\vee \subseteq \text{im } q^\vee$ .

Conversely, for any  $\beta \in (V/U)^*$  and  $v \in (V/U)$ , we have

$$q^\vee(\beta)(v) = \beta(q(v)) = \beta(0) = 0,$$

so we also have  $\text{im } q^\vee \subseteq \ker \iota^\vee$ . □

**Theorem 7.46**

Let  $(V, g)$  and  $(W, h)$  be  $\mathbb{K}$ -vector spaces with scalar products. Let  $h_1 : W \rightarrow W^*$  and  $g_1 : V \rightarrow V^*$  be the antilinear maps induced by  $g$  and  $h$ , i.e.

$$\begin{aligned} h_1(\vec{w}) &= h(\vec{w}, \cdot) \\ g_1(\vec{v}) &= g(\vec{v}, \cdot) \end{aligned}$$

Let  $F : V \rightarrow W$  be linear. Then  $g_1 \circ F^* = F^\vee \circ h_1$ .

*Proof.* Let  $v \in V, w \in W$ , then we have:

$$\begin{aligned} g_1(F^*(w))(v) &= g(F^*(w), v) \\ &= h(w, F(v)) \\ &= h_1(w)(F(v)) \\ &= F^\vee(h_1(w))(v) \end{aligned}$$

□

**Corollary 7.47.** If  $V$  is finite-dimensional,  $F : V \rightarrow W$  is adjointable.

*Proof.* If  $V$  is finite-dimensional,  $g_1$  is invertible. We get  $F^* = g_1^{-1} \circ F^\vee \circ h_1$ . □

**Definition 7.48.** Let  $V$  and  $W$  be  $\mathbb{K}$ -vector spaces. Let  $F : V \rightarrow W$  be linear, and let  $s : W \times W \rightarrow \mathbb{K}$  be a sesquilinear form. Then we define the **pullback of  $s$  by  $F$**  to be the map  $F^*s : V \times V \rightarrow \mathbb{K}$  given by

$$(F^*s)(u, v) = s(F(u), F(v))$$

**Exercise 7.49.** Verify that:

1.  $F^*s$  is a sesquilinear form.
2. If  $s$  is Hermitian,  $F^*s$  is Hermitian.
3. If  $s$  is positive semidefinite,  $F^*s$  is positive semidefinite.
4. If  $s$  is positive definite and  $F$  is injective,  $F^*s$  positive definite.
5. If we fix bases of  $V$  and  $W$  such that  $F$  is represented by a matrix  $A$  and  $s$  is represented by a Gram matrix  $G$ , then  $F^*s$  is represented by  $A^*GA$ .
6. Given  $F : V \rightarrow W$  antilinear, we can achieve the same results by defining  $F^*s(u, v) = s(F(v), F(u))$ .

## Chapter 8

# Normal Endomorphisms

### Definition 8.1

Let  $(V, g)$  be a  $\mathbb{k}$ -vector space with scalar product. Let  $F : V \rightarrow V$  be an adjointable endomorphism. Then we call  $F$  **normal** if  $F^* \circ F = F \circ F^*$ .

**Exercise 8.2.** 1. Show that self-adjoint maps are normal.

2. A linear map with  $F^* = -F$  is called **skew**. Show that skew maps are normal.

3. A linear map represented in a given basis by a matrix of the form

$$\begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

is normal.

### Definition 8.3

Let  $(V, g)$  and  $(W, h)$  be  $\mathbb{k}$ -vector spaces with scalar products. Then a linear map  $F : V \rightarrow W$  is called an **isometric embedding** if  $g(v, w) = h(F(v), F(w))$  for all  $v, w \in V$ .

**Exercise 8.4.** Show that:

1. Isometric embeddings preserve lengths and distances.

2. Isometric embeddings preserve angles.

3. Isometric embeddings are injective.

4. If orthonormal bases are chosen for  $V$  and  $W$ , then the matrix representation  $A$  of  $F$  fulfills  $A^*A = E_n$ .

5. If  $\dim V = \dim W$ ,  $F$  is invertible with  $F^{-1} = F^*$ . An invertible isometric embedding is called an **isometry**.

6. Isometries are normal.

**Theorem 8.5: Fundamental Theorem of Normal Endomorphisms**

Let  $(V, g)$  be a finite-dimensional complex vector space with scalar product. Let  $F : V \rightarrow V$  be an endomorphism. Then there exists an orthonormal basis of  $(V, g)$  consisting of eigenvectors of  $F$  if and only if  $F$  is normal.

Note that this only works for complex numbers, not for reals:

**Exercise 8.6.** Recall that any endomorphism represented by a matrix of the form

$$A = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

is normal. Show that, if  $b \neq 0$ ,  $A$  has no real eigenvalues.

**Corollary 8.7: A Normal Form for Complex Normal Endomorphisms**

Let  $(V, g)$  be a finite-dimensional real vector space with scalar product. Let  $F : V \rightarrow V$  be an endomorphism. Then there exists an orthonormal basis of  $(V, g)$  in which  $F$  is represented by a block diagonal matrix consisting of  $1 \times 1$  blocks and  $2 \times 2$  blocks of the form

$$\begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

with  $b > 0$  if and only if  $F$  is normal. If such a representation exists, it is unique up to permutation of the blocks.

**Corollary 8.8: Spectral Theorem for Self-Adjoint Maps**

Let  $(V, g)$  be a finite dimensional real or complex vector space with scalar product. Let  $F : V \rightarrow V$  be an endomorphism. Then there exists an orthonormal basis of  $V$  in which  $F$  is represented by a diagonal matrix with real entries if and only if  $F$  is self-adjoint.

**Exercise 8.9.** Derive an analogous spectral theorem for skew maps.

**Corollary 8.10.** Let  $(V, g)$  be a finite-dimensional real or complex vector space with scalar product. Let  $s$  be a sesquilinear form on  $V$ . Then there exists a  $g$ -orthonormal Basis of  $V$  in which  $s$  is represented by a diagonal matrix with real entries if and only if  $s$  is hermitian.

**Corollary 8.11: Singular Value Decomposition**

Let  $(V, g)$  and  $(W, h)$  be finite-dimensional real or complex vector spaces with scalar products. Let  $F : V \rightarrow W$  be linear. Then there exist orthonormal bases of  $V$  and  $W$  in which  $F$  is represented by a diagonal matrix with uniquely determined real entries  $a_n$  such that  $a_1 \geq a_2 \geq \dots$  and  $a_n = 0$  if  $n \geq \text{rk } F$ . These entries are known as the *singular values* of  $F$ .

Singular value decomposition is fundamental to the JPG compression algorithm, which represents an image by a matrix, then calculates the singular

value decomposition of that matrix, and then sets all of the singular values below a given threshold to zero, leading to less data needing to be stored while still maintaining the singular values that ‘noticeably contribute’ to the resulting image.

## 8.1 Linear Isometries

### Corollary 8.12: Normal Form for Real Isometries

Let  $(V, g)$  be a finite-dimensional real vector space with scalar product  $g$ . Let  $F : V \rightarrow V$  be a linear isometry. Then there exists an orthonormal basis of  $(V, g)$  in which  $F$  is represented by a block diagonal matrix consisting of  $1 \times 1$  blocks and  $2 \times 2$  blocks of the form

$$\begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

with  $a^2 + b^2 = 1$ .

### Corollary 8.13: Normal Form for Complex Isometries

Let  $(V, g)$  be a finite-dimensional complex vector space with scalar product  $g$ . Let  $F : V \rightarrow V$  be a linear isometry. Then there exists an orthonormal basis of  $(V, g)$  in which  $F$  is represented by a diagonal matrix whose entries have absolute value 1.

### Corollary 8.14

Let  $(V, g)$  be a finite-dimensional complex vector space with scalar product  $g$ . Let  $F : V \rightarrow V$  be a linear isometry. Then  $V$  can be decomposed into a direct sum of subspaces  $U$  of dimension one or two, such that  $F$  acts on the one-dimensional subspaces as the identity or reflection, and on the two-dimensional subspaces as a rotation around the origin.

**Exercise 8.15.** A linear isometry is orientation-preserving if and only if the number of invariant subspaces on which it acts as a reflection is even.

## 8.2 Affine Isometries

**Theorem 8.16.** Let  $(A, \vec{A}, s), (B, \vec{B}, g)$  be finite-dimensional normed affine spaces of the same dimension over vector spaces with scalar products. Then there exists an affine isometry  $F : A \rightarrow B$ .

**Theorem 8.17.** Let  $(A, \vec{A}, g)$  be a finite-dimensional normed  $\mathbb{k}$ -affine space over a vector space with scalar product. Let  $F : A \rightarrow A$  be an affine isometry. Then there exists a point  $o \in A$  and a vector  $x \in \text{Eig}(\vec{F}, 1)$  such that

$$F(a) = o + x + \vec{F}(a - o)$$

$x$  is uniquely determined by  $A$ , while  $o$  is an arbitrary point chosen from an affine- $F$ -invariant subspace  $(B, \text{Eig}(\vec{F}, 1)) \subseteq (A, \vec{A})$ .

**Definition 8.18.** We call  $(B, \text{Eig}(\vec{F}, 1))$  the *axis set* of  $F$ , and call lines of the form

$$\{b + \lambda.x \mid \lambda \in \mathbb{K}\} \subseteq B$$

with  $b \in B$  the *axes* of  $F$ .

Our theorem thus states that any affine isometry can be represented as a linear map composed with a translation along the axis set of  $F$ .

## Chapter 9

# Revisiting Synthetic Geometry

### 9.1 Incidence Geometries

#### Definition 9.1: Incidence Geometries

An *incidence geometry* is a set  $X$  equipped with a set  $G \subseteq \mathcal{P}(X)$  of subsets of  $X$  known as *lines*, such that:

1. Each line contains at least two points,
2. For every pair of points, there exists exactly one line containing both.

Instead of " $x$  is contained in  $g \in G$ ", we often say " $x$  lies on  $g \in G$ ".

**Exercise 9.2.** Let  $R$  be a division ring. Then  $X := R^n$  can be turned into an incidence geometry by defining lines to be all sets of the form

$$g = \{p + \lambda v : \lambda \in R\},$$

where  $p, v \in R^n$ .

Moving forward, this will be the canonical choice of incidence geometry on  $R^n$ .

**Definition 9.3.** A set of points lying on the same line is called *colinear*.

**Definition 9.4.** A set of lines containing a common point is called *confluent*.

**Definition 9.5.** A set of three non-colinear points is known as a *triangle*.

**Definition 9.6: Betweenness**

A **betweenness relation** on an incidence geometry  $(X, G)$  is a subset  $Z \subseteq X^3$  such that the following properties hold:

1. **Induced Orderings on Lines:** On every line  $g \in G$ , there exists an ordering  $\leq$  such that  $(x, y, z) \in Z$  if and only if  $x \leq y \leq z$  or  $x \geq y \geq z$ .
2. **Pasch's Axiom:** Let  $p, q, r \in X$ . Let

$$[p, q] := \{x \in X : (p, x, q) \in Z\}$$

denote the **line segment between  $p$  and  $q$** . Then every line  $g \in G$  either intersects none of the line segments  $[p, q], [q, r], [r, p]$ , or at least two.

An incidence geometry equipped with a betweenness relation is known as an **ordered incidence geometry**.

**Exercise 9.7.** Let  $K$  be an ordered field. Then:

1. The set of all tuples  $(x, y, z)$ , where  $x, y, z \in K^n$  fulfill  $y = \lambda x + \mu z$  with  $\lambda, \mu \in [0, 1] \subset K$  and  $\lambda + \mu = 1$ , gives a betweenness relation on  $K^n$ .
2. There exists an affine isomorphism  $\varphi : K \cong (g \in G)$  between any line in  $G$  and  $K$ .
3. The ordering on  $K$  can be recovered from the betweenness relation on  $K^n$  by taking the unique ordering  $\leq$  on  $g \in G$  that fulfills  $\varphi(0) \leq \varphi(1)$ .

Moving forward, this will be the canonical choice of betweenness relation on  $K^n$ .

**Definition 9.8.** Given an ordered incidence geometry  $(X, G, Z)$ , a **ray** in  $X$  is a set  $A \subseteq X$  of the form

$$A = \{x \in g \mid x \leq p\},$$

where  $g \in G$  is a line and  $p \in g$  is a point on that line.

**Definition 9.9: Congruence Groups**

Given an ordered incidence geometry  $(X, G, Z)$ , a **congruence group** of  $(X, G, Z)$  is a group  $K$  of automorphism of  $(X, G, Z)$  such that every pair of rays in  $A, B \subseteq X$  has exactly two elements  $h, k \in K$  such that  $h(A) = k(A) = B$ .

Intuitively, one of these two elements will be a rotation and the other a rotation composed with reflection along an axis.

**Exercise 9.10.** Let  $K$  be an ordered field. Let  $s$  be the standard scalar product in  $K^2$ . Then the set of  $s$ -invariant affine automorphisms of  $K^2$  forms a congruence group of  $K^2$ .

Moving forward, this will be the canonical choice of congruence group on  $K^2$ .

**Definition 9.11.** We say that an ordered incidence geometry has the *least upper bound property* if every nonempty subset  $A \subset G$  that is bounded from above has a least upper bound.

**Definition 9.12.** We call an ordered incidence geometry  $(X, G, Z)$  with congruence group  $K$  *near-euclidean* if  $(X, G, Z)$  fulfills the supremum property and  $G$  is nonempty.

**Definition 9.13.** We say that an incidence geometry  $(X, G)$  fulfills the *parallel postulate* iff. for every line  $g \in G$  and every point  $p \notin g$ , there exists at most one line  $h \in G$  such that  $p \in h$  and  $h \cap g = \emptyset$ .

**Exercise 9.14.** Let  $K \subseteq \mathbb{R}$  be a proper subfield of  $\mathbb{R}$  containing square roots of every positive element. Then  $K^2$  fulfills the parallel postulate.

The primary goal of this chapter will be proving the following result:

**Theorem 9.15: Classification of near-euclidean geometries**

Let  $(X, G, Z)$  be a near-euclidean incidence geometry. Then:

1. If  $(X, G, Z)$  fulfills the parallel postulate, it is isomorphic to  $\mathbb{R}^2$ .
2. If  $(X, G, Z)$  doesn't fulfill the parallel postulate, it is isomorphic to the *hyperbolic plane*  $H$ .

## 9.2 Projection Maps

**Definition 9.16.** Let  $V$  be a vector space. Then a *linear hyperplane* is a vector subspace of  $V$  of codimension 1, i.e.

$$\text{codim}(W \subseteq V) := \dim(V/W) = 1$$

Equivalently,  $W \neq V$  such that  $V$  is generated by  $W$  and any vector  $v \in V \setminus W$ . If  $V$  is finite-dimensional, we have  $\dim(V/W) = \dim V - \dim W$ , and thus a linear hyperplane is a subspace  $W$  whose dimension is one lower than the dimension of  $V$ . An *affine Hyperplane* is defined entirely analogously as an affine subspace of codimension one.

**Exercise 9.17.** 1. In any affine space, the intersection of an affine line with another affine subspace is either empty, a single point, or the entire line.

2. For any affine hyperplane  $H$  and any line  $g$  with directional vector  $\vec{v} \notin \vec{H}$ , the intersection of  $g$  and  $H$  consists of a single point.

**Definition 9.18.** Given an affine space  $E$ , an affine hyperplane  $H \subset E$ , and a vector  $\vec{v} \in \vec{E} \setminus \vec{H}$ , the *parallel projection of  $E$  onto  $H$  in direction  $\vec{v}$*  is the map

$$\pi = \pi_{\vec{v}} = \pi_{\vec{v}, H} : E \rightarrow H$$

assigning to each point  $e$  the intersection of the affine line  $(e + \langle \vec{v} \rangle)$  with  $H$ .

**Definition 9.19.** Given an affine space  $E$ , an affine hyperplane  $H \subset E$ , and a point  $q \in E \setminus H$ , the **central projection of  $E$  onto  $H$  centered at  $q$**  is the map

$$\zeta = \zeta_q = \zeta_{q,H} : E \setminus (q + \vec{H}) \rightarrow H$$

assigning to each point  $e \in E$  the intersection of the line through  $e$  and  $q$  with  $H$ .

**Exercise 9.20.** The image of a line  $g$  under central projection is either:

1. A line,
2. a pointed line, i.e. a line with a single missing point (the **vanishing point**),
3. a single point,
4. or the empty set.

## 9.3 Projectivization

### Definition 9.21: Projectivization

Let  $\mathbb{k}$  be a field. Let  $V$  be a  $\mathbb{k}$ -vector space. Then the set of all lines through the origin in  $V$  is called the **projectivization of  $V$**  and denoted by  $\mathbb{P}V$ .

In the finite-dimensional case, we have  $\dim \mathbb{P}V = \dim V - 1$ . It is therefore common to write

$$\mathbb{P}^n \mathbb{k} := \mathbb{P}(\mathbb{k}^{n+1})$$

Beware, this means  $\mathbb{P}\mathbb{k} \neq \mathbb{P}^1 \mathbb{k}$ ! The set  $\mathbb{P}^1 \mathbb{k}$  is called the **projective line over  $\mathbb{k}$** , the set  $\mathbb{P}^2 \mathbb{k}$  is called the **projective plane over  $\mathbb{k}$** .

**Definition 9.22.** Given a vector space  $V$ , we call a subset  $g \subseteq \mathbb{P}V$  a **projective line** or  **$\mathbb{P}$ -line** if there exists a vector subspace  $g' \subseteq V$  such that  $\dim g' = 2$  and  $g = \mathcal{P}g'$ .

**Lemma 9.23.** For any vector space  $V$ , the set  $\mathbb{P}V$  forms an incidence geometry with  $\mathbb{P}$ -lines as the lines.

**Lemma 9.24.** In a projective plane over any field, any two different  $\mathbb{P}$ -lines intersect in exactly one point.

**Definition 9.25.** An **embedding of incidence geometries** is an injective map  $X \hookrightarrow Y$  such that points in  $X$  are colinear if and only if their images are colinear. A bijective embedding of incidence geometries is known as a **colineation**.

**Lemma 9.26.** Any injective vector space homomorphism  $W \hookrightarrow V$  induces an embedding  $\mathbb{P}W \hookrightarrow \mathbb{P}V$ .

**Definition 9.27**

Let  $E$  be an affine space. Then we can equip the disjoint union

$$\mathbb{V}E := E \sqcup \vec{\mathbb{P}}E$$

with an incidence structure by designating lines to be:

1. Sets  $\mathbb{V}g := g \sqcup \{\vec{g}\}$  for any affine line  $g \in E$ ,
2. and  $\mathbb{P}$ -lines  $l = \mathbb{P}l'$ .

We call this structure the *projective completion of  $E$* . We call points in  $\vec{\mathbb{P}}E$  *points at infinity*.

**Exercise 9.28.** If  $E$  is a two-dimensional affine space, then its projective completion  $\mathbb{V}E$  contains exactly one line containing only infinitely distant points. We call this line the **line at infinity**.

**Exercise 9.29.** Given an affine space  $E$ , the trivial embeddings  $E \hookrightarrow \mathbb{V}E$  and  $\vec{\mathbb{P}}E \hookrightarrow \mathbb{V}E$  are embeddings of incidence geometries. Any injective affine map  $F \hookrightarrow E$  of affine spaces is an embedding of incidence geometries and induces an embedding  $\mathbb{V}E \hookrightarrow \mathbb{V}F$  of their projective completions.

**Theorem 9.30**

Let  $V$  be a vector space. Let  $E$  be an affine hyperplane in  $V$  not containing the origin. Then

$$\mathbb{V}E \cong \mathbb{P}V$$

**Corollary 9.31: Projective lines are projective completions**

Let  $F$  be a field. Then we have  $\mathbb{P}^1 F \cong \mathbb{V}F = F \sqcup \{\infty\}$ .

**Exercise 9.32.** 1. For any field  $\mathbb{F}$ , we have

$$\mathbb{P}^n \mathbb{F} \cong \bigsqcup_{i=0}^n \mathbb{F}^i$$

2. For a finite Field  $\mathbb{F}_q$ , we have

$$|\mathbb{P}^n \mathbb{F}_q| = \sum_{i=0}^n q^i$$

## Chapter 10

# Hyperbolic Geometry

### 10.1 Circle Inversions

**Definition 10.1.** Let  $E$  be a Euclidean plane. Then the corresponding *extended Euclidean plane* is the set

$$\hat{E} = E \cup \{\infty\},$$

adding a "point at infinity" to  $E$ .

**Definition 10.2.** We call a subset  $L \subseteq \hat{E}$  a *generalized circle* if it is either a circle in  $E$  or the union of a line in  $E$  with  $\{\infty\}$ . We call the former *real circles* and the latter *extended lines*.

**Definition 10.3.** Let  $E$  be a euclidean plane. Then we assign to each generalized circle  $L \subseteq \hat{E}$  the *circle inversion map*

$$s_L : \hat{E} \rightarrow \hat{E},$$

which is given by the ordinary reflection at  $L$  with the additional rule  $s_L(\infty) = \infty$  if  $L$  is an extended line, and by

$$c + \vec{v} \mapsto \begin{cases} c + \left(\frac{r}{\|\vec{v}\|}\right)^2 \vec{v} & \vec{v} \neq \vec{0} \\ \infty & \vec{v} = \vec{0} \end{cases}$$

with  $\infty \mapsto c$  if  $L$  is a circle with center  $c$  and radius  $r$ .

**Exercise 10.4.**  $s_L$  is an involution, i.e. we have  $s_L^2 = \text{id}$ .

**Exercise 10.5.** For any two points  $p, q \in \hat{E}$ , there exists a circle inversion  $s_L$  with  $s_L(p) = q$  and  $s_L(q) = p$

**Exercise 10.6.** Every circle inversion maps generalized circles to generalized circles.

## 10.2 Möbius Transformations

**Definition 10.7.** A composition of circle inversions is known as a *Möbius transformation*.

**Exercise 10.8.** The set  $\text{Möb}(\hat{E})$  of Möbius transformations of a given extended Euclidean plane  $\hat{E}$  forms a group under composition.

**Exercise 10.9.** Any two generalized circles in a generalized Euclidean plane can be mapped onto each other by a Möbius transformation.

**Theorem 10.10.** Given an extended Euclidean plane  $E$ , its Möbius transformations are precisely the bijections  $\varphi : \hat{E} \xrightarrow{\sim} \hat{E}$  mapping generalized circles to generalized circles.

**Theorem 10.11.** Any Möbius transformation that leaves a given generalized circle  $L$  pointwise unchanged is either the circle inversion  $s_L$  at that circle or the identity map.

**Exercise 10.12.** Given a generalized Euclidean plane  $\hat{E}$ , a Möbius transformation  $\varphi : \hat{E} \rightarrow \hat{E}$  and a generalized circle  $L \subseteq \hat{E}$ , we have  $\varphi \circ s_L \circ \varphi^{-1} = s_{\varphi(L)}$ .

**Exercise 10.13.** Given a generalized Euclidean plane  $\hat{E}$ , a Möbius transformation  $\varphi : \hat{E} \rightarrow \hat{E}$ , and a generalized circle  $L \subseteq \hat{E}$ , we have  $\varphi(L) = L$  if and only if  $\varphi \circ s_L = s_L \circ \varphi$ .

**Definition 10.14.** Let  $\hat{E}$  be a generalized Euclidean plane. Given a pair of generalized circles  $L, M \subset \hat{E}$  we call  $L$  *orthogonal to*  $M$ , which we write  $L \perp M$ , if  $L \neq M$  and  $s_L(M) = M$ .

**Theorem 10.15.** Let  $L, M \subset \hat{E}$  be generalized circles in a generalized Euclidean plane  $\hat{E}$  such that  $L \perp M$ . Let  $\varphi : \hat{E} \rightarrow \hat{E}$  be a Möbius transformation. Then  $\varphi(L) \perp \varphi(M)$ .

**Proposition 10.16.** Let  $E$  be a Euclidean plane,  $g \subset E$  a line,  $H \subset E \setminus g$  an open Half-plane with  $g$  as its boundary, and

$$K_H = \text{Möb}_H(\hat{E}) := \{\varphi \in \text{Möb}(\hat{E}) \mid \varphi(H) = H\}$$

the set of Möbius transformations leaving  $H$  invariant. Then:

1. Any Möbius transformation that leaves  $H$  invariant also leaves the extended line  $\hat{g}$  invariant.
2.  $\text{Möb}_H(\hat{E})$  operates transitively on  $H$ , i.e. any point of  $H$  can be mapped onto any other by a Möbius transformation  $\varphi \in \text{Möb}_H(\hat{E})$ .
3.  $\text{Möb}_H(\hat{E})$  is generated by the set of circle inversions  $s_L$  with  $L \perp \hat{g}$ .
4. Different elements of  $K_H$  are still different when restricted to  $H$ .

## 10.3 Complex Möbius Geometry

**Definition 10.17.** We denote by  $\gamma$  the “double complex conjugation map”

$$\begin{aligned}\gamma : \mathbb{C}^2 &\rightarrow \mathbb{C}^2 \\ (w, z) &\mapsto (\overline{w}, \overline{z})\end{aligned}$$

**Theorem 10.18.** The set of all antilinear automorphisms of  $\mathbb{C}^2$  is given by

$$\mathrm{GL}(\mathbb{C}^2)_\gamma := \{F \circ \gamma \mid F \in \mathrm{GL}(\mathbb{C}^2)\}$$

**Definition 10.19.** We denote by

$$\mathrm{GL}(\mathbb{C}^2) \langle \gamma \rangle := \mathrm{GL}(\mathbb{C}^2) \sqcup \mathrm{GL}(\mathbb{C}^2)_\gamma$$

the set of all automorphisms of  $\mathbb{C}^2$  that are either linear or antilinear.

Since  $\mathbb{P}^1\mathbb{C}$  consists of subsets of  $\mathbb{C}^2$ , every map  $\mathbb{C}^2 \rightarrow \mathbb{C}^2$  mapping lines to lines acts on  $\mathbb{P}^1\mathbb{C}$  in a trivial way. By theorem 9.31  $\mathbb{P}^1\mathbb{C} \cong \hat{\mathbb{C}}$ , the map

$$\begin{aligned}\iota_{\mathbb{C}} : \hat{\mathbb{C}} &\xrightarrow{\sim} \mathbb{P}^1\mathbb{C} \hookrightarrow \mathbb{C}^2 \\ z &\mapsto (z, 1) \\ \infty &\mapsto (1, 0)\end{aligned}$$

is a bijection, which means we also get an action on  $\hat{\mathbb{C}}$ .

### Theorem 10.20: Action of $\mathrm{GL}(\mathbb{C}^2) \langle \gamma \rangle$ on $\hat{\mathbb{C}}$

The canonical action gives us a group homomorphism

$$\begin{aligned}\varphi : \mathrm{GL}(2; \mathbb{C}) \langle \gamma \rangle &\rightarrow \mathrm{Ens}^\times(\hat{\mathbb{C}}) \\ \gamma &\mapsto (z \mapsto \overline{z}) \\ M := \begin{pmatrix} a & b \\ c & d \end{pmatrix} &\mapsto \left( z \mapsto \begin{cases} \frac{az+b}{cz+d} & z \neq \infty \wedge cz+d \neq 0 \\ \frac{a}{c} & z = \infty \wedge c \neq 0 \\ \infty & z \neq \infty \wedge cz+d = 0 \\ \infty & z = \infty \wedge c = 0 \end{cases} \right)\end{aligned}$$

*Proof.* 1. Let  $z \in \mathbb{C}$ . Translating directly from  $\mathbb{C}$  to gives  $\mathbb{C}^2$   $\iota_{\mathbb{C}}(z) = \begin{pmatrix} z \\ 1 \end{pmatrix}$ , which means  $M(\iota_{\mathbb{C}}(z)) = \begin{pmatrix} az+b \\ cz+d \end{pmatrix}$ .

- (a) If  $cz+d \neq 0$ , then this point lies on the same line through the origin as  $\begin{pmatrix} \frac{az+b}{cz+d} \\ 1 \end{pmatrix}$ , which is then translated back from  $\mathbb{P}^1\mathbb{C}$  to  $\hat{\mathbb{C}}$  as the point  $\frac{az+b}{cz+d}$ .
- (b) If  $cz+d = 0$ , then this point lies on the same line through the origin as  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ , which is then translated back from  $\mathbb{P}^1\mathbb{C}$  to  $\hat{\mathbb{C}}$  as the point  $\infty$ .

- 2. If  $z = \infty$ , then  $\iota_{\mathbb{C}}(\infty) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ , which gets mapped to  $\begin{pmatrix} a \\ c \end{pmatrix}$ , from which we can proceed as before.

□

**Theorem 10.21.** Our canonical group homomorphism  $\varphi : \text{GL}(\mathbb{C}^2) \langle \gamma \rangle \rightarrow \text{Ens}^\times(\hat{\mathbb{C}})$  is a surjective group homomorphism

$$\varphi : \text{GL}(\mathbb{C}^2) \langle \gamma \rangle \rightarrow \text{Möb}(\hat{\mathbb{C}})$$

whose kernel is the group consisting of all scalar multiples of the identity.

*Proof.* 1. Since  $\text{GL}(\mathbb{C}^2)$  is generated by the diagonal matrices together with the matrices  $S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  and  $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ ,  $\text{GL}(2, \mathbb{C}) \langle \gamma \rangle$  is generated by these matrices along with  $\gamma$ . It therefore suffices to show that the images of these matrices are Möbius transformations. Now, according to our previous analysis of the maps generated by  $\varphi$ , these images are as follows:

- (a) A diagonal matrix with entries  $\lambda$  and  $\mu$  gets mapped to  $\frac{\lambda}{\mu} \cdot z$ , which corresponds to a scaled rotation.
- (b)  $S$  gets mapped to  $z \mapsto \frac{1}{z}$ , which is the circle inversion at the unit circle composed with a reflection along the real axis.
- (c)  $T$  gets mapped to the translation  $z \mapsto z + 1$ .
- (d)  $\gamma$  gets mapped to the conjugation  $z \mapsto \bar{z}$ , which is just the reflection along the real axis.

All of these maps are known Möbius transformations.

- 2. We now know that the image of  $\varphi$  contains inversion at the unit circle, all translations, and all scalings. Therefore, it contains all inversions along proper circles. It also contains reflection along the extended real axes and all rotations at the origin, so it also contains reflections along extended lines. Thus, the image contains the generators of  $\text{Möb}(\hat{\mathbb{C}})$ , and therefore the entire group.
- 3. An antilinear automorphism cannot stabilize every one-dimensional subspace. Thus, antilinear automorphisms cannot be mapped the identity, i.e. they cannot be contained in the kernel.

Meanwhile, the only linear automorphisms stabilizing every one-dimensional subspace are scalings, i.e. multiples of the identity.

□

### Definition 10.22: Projective Linear Groups

Given a vector space  $V$ , the *projective linear group*  $\text{PGL}(V)$  is defined as the quotient group of  $\text{GL}(V)$  by the group  $Z(V)$  of all scalar multiples of the identity. The *projective special linear group*  $\text{PSL}(V)$  is analogously defined as the quotient group of  $\text{SL}(V)$  by the group  $SZ(V)$  of all multiples of the identity by a unit scalar.

**Theorem 10.23: Möbius group as a projective linear group**

$$\text{Möb}^+(\hat{\mathbb{C}}) \cong \text{PGL}(\mathbb{C}^2)$$

**Definition 10.24.** We denote by  $\tau$  reflection along the axis  $\langle e_1 \rangle$ , i.e.

$$\begin{aligned} \tau : \mathbb{R}^2 &\rightarrow \mathbb{R}^2 \\ \begin{pmatrix} a \\ b \end{pmatrix} &\mapsto \begin{pmatrix} a \\ -b \end{pmatrix} \end{aligned}$$

**Theorem 10.25: Action of  $\text{SL}(\mathbb{C}^2)$  on  $\hat{\mathbb{C}}$**

Our surjective group homomorphism  $\varphi : \text{GL}(\mathbb{C}^2) \langle \gamma \rangle \twoheadrightarrow \text{Möb}(\mathbb{C}^2)$  induces a surjective group homomorphism

$$\text{SL}(\mathbb{R}^2) \langle \tau \gamma \rangle \twoheadrightarrow \text{Möb}_{\mathbb{H}}(\hat{\mathbb{C}})$$

whose kernel is  $\pm \text{Id}_{\mathbb{R}^2}$ .

**Theorem 10.26: Conjugations of  $\mathbb{H}$  as a projective special linear group**

$$\text{Möb}_{\mathbb{H}}^+(\hat{\mathbb{C}}) \cong \text{PSL}(\mathbb{R}^2)$$

**Part VI**

**Group Theory**

# Chapter 1

## Discrete Symmetry Groups

### 1.1 Finite subgroups of the orthogonal groups

#### Theorem 1.1

The finite subgroups of  $SO(3)$ , up to isomorphism, are:

1. The cyclic groups  $C_n$ ,
2. The dihedral groups  $D_n$ ,
3. The tetrahedral group  $T \cong A_4$ ,
4. The cube group  $W \cong S_4$ ,
5. The icosahedral group  $I \cong A_5$ .

#### Corollary 1.2

The finite subgroups of  $O(3)$ , up to isomorphism, are:

1. The finite subgroups of  $SO(3)$ :
  - (a) The cyclic groups  $C_n$ ,
  - (b) The dihedral groups  $D_n$ ,
  - (c) The tetrahedral group  $T \cong A_4$ ,
  - (d) The cube group  $W \cong S_4$ ,
  - (e) The icosahedral group  $I \cong A_5$ .
2. Products of these groups with  $C_2$ .

## 1.2 Lattice Groups

**Definition 1.3.** An  $n$ -dimensional real *lattice* is a subset of  $\mathbb{R}^n$  of the form

$$\Lambda = \left\{ \sum_{i=1}^n k_i \cdot v_i \mid k_i \in \mathbb{Z} \right\},$$

where  $v_1, \dots, v_n$  is a basis of  $\mathbb{R}^n$ . We say that the lattice  $\Lambda$  is *generated by the matrix*  $M = (v_1, \dots, v_n)$ .

### Theorem 1.4: Crystallographic restriction theorem

Every rotational symmetry of a two- or three-dimensional lattice has order 1, 2, 3, 4, or 6.

## 1.3 The Two-Dimensional Point Groups

**Definition 1.5.** An  $n$ -dimensional *crystallographic point group* is a subgroup of  $O(n)$  mapping an  $n$ -dimensional lattice to itself while leaving a given point unchanged.

### Theorem 1.6

The two-dimensional crystallographic point groups are:

1. The cyclic groups  $C_1, C_2, C_3, C_4$  and  $C_6$ ,
2. The dihedral groups  $D_1, D_2, D_3, D_4$  and  $D_6$ .

We have  $D_1 \cong C_2$ , so there are ten point groups up to equality and nine point groups up to equality.

*Proof Sketch.* Follows immediately from the classification of finite subgroups of  $O(2)$  and the crystallographic restriction theorem. □

## 1.4 The Three-Dimensional Point Groups

### Theorem 1.7

There are eighteen three-dimensional crystallographic point groups up to isomorphism, and they are:

1. Five cyclic groups:

- $C_1$
- $C_2 \cong D_1$
- $C_3$
- $C_4$
- $C_6 \cong C_3 \times C_2$

2. Four non-cyclic dihedral groups:

- $D_2 \cong C_2 \times C_2$
- $D_3$
- $D_4$
- $D_6 \cong D_3 \times C_2$

3. Two remaining products of cyclic groups with  $C_2$ :

- $C_4 \times C_2$
- $C_6 \times C_2$

4. Three remaining products of dihedral groups with  $C_2$ :

- $D_2 \times C_2$
- $D_4 \times C_2$
- $D_6 \times C_2$

5. Four remaining groups consisting of permutation groups and their products with  $C_2$ :

- $A_4$
- $A_4 \times C_2$
- $S_4$
- $S_4 \times C_2$

### Theorem 1.8

The following table gives a full classification of three-dimensional crystallographic point groups up to identity:

| Crystal System | Crystal Class             | Example Figure | HM Notation | Isomorphism      |
|----------------|---------------------------|----------------|-------------|------------------|
| triclinic      | triclinic-pedial          |                | 1           | $C_1$            |
|                | triclinic-pinacoidal      |                | $\bar{1}$   |                  |
| monoclinic     | monoclinic-sphenoidal     |                | 2           | $C_2$            |
|                | monoclinic-domatic        |                | $m$         |                  |
|                | monoclinic-prismatic      |                | $2/m$       |                  |
| orthorhombic   | orthorhombic-disphenoidal |                | 222         | $D_2$            |
|                | orthorhombic-pyramidal    |                | $mm2$       |                  |
|                | orthorhombic-dipyramidal  |                | $mmm$       | $D_2 \times C_2$ |
| tetragonal     | tetragonal-pyramidal      |                | 4           | $C_4$            |
|                | tetragonal-disphenoidal   |                | $\bar{4}$   |                  |
|                | tetragonal-dipyramidal    |                | $4/m$       | $C_4 \times C_2$ |
|                | tetragonal-trapezoidal    |                | 422         |                  |
|                | ditetragonal-pyramidal    |                | $4mm$       | $D_4$            |
|                | tetragonal-scalenoidal    |                | $\bar{4}2m$ |                  |
|                | ditetragonal-dipyramidal  |                | $4/mmm$     | $D_4 \times C_2$ |
| trigonal       | trigonal-pyramidal        |                | 3           | $C_3$            |
|                | rhombohedral              |                | $\bar{3}$   | $C_6$            |
|                | trigonal-trapezoidal      |                | 32          | $D_3$            |
|                | ditrigonal-pyramidal      |                | $3m$        |                  |
|                | ditrigonal-scalenohedral  |                | $\bar{3}m$  | $D_6$            |
| hexagonal      | hexagonal-pyramidal       |                | 6           | $C_6$            |
|                | trigonal-dipyramidal      |                | $\bar{6}$   |                  |
|                | hexagonal-dipyramidal     |                | $6/m$       | $C_6 \times C_2$ |
|                | hexagonal-trapezohedral   |                | 622         |                  |
|                | dihexagonal-pyramidal     |                | $6mm$       | $D_6$            |
|                | ditrigonal-dipyramidal    |                | $\bar{6}m2$ |                  |
|                | dihexagonal-dipyramidal   |                | $6/mmm$     | $D_6 \times C_2$ |
| cubic          | tetartoidal               |                | 23          | $A_4$            |
|                | diploidal                 |                | $m\bar{3}$  | $A_4 \times C_2$ |
|                | gyroidal                  |                | 432         | $S_4$            |
|                | hextetrahedral            |                | $\bar{4}3m$ |                  |
|                | hexoctahedral             |                | $m\bar{3}m$ | $S_4 \times C_2$ |

## 1.5 Classification of Lattices

## 1.6 Wallpaper Groups

## 1.7 Three-Dimensional Space Groups

## **Part VII**

# **Multivariable Calculus**

# Chapter 1

## Differentiation in $\mathbb{R}^n$

### 1.1 Differentials and Partial Derivatives

#### Definition 1.1: Fréchet Derivative

Let  $V, W$  be normed vector spaces. Let  $U$  be an open subset of  $V$ . Then  $f : U \rightarrow W$  is called *Fréchet differentiable* or just *differentiable* at  $p \in U$  if there exists a continuous linear map  $Df(p) : V \rightarrow W$  such that

$$\lim_{h \rightarrow 0} \frac{\|f(p+h) - f(p) - Df(p)(h)\|_W}{\|h\|_V} = 0.$$

We call  $Df(p)$  the *differential of  $f$  at  $p$* . We call  $f$  *differentiable* if it is differentiable at every point  $p \in U$ . We call  $f$  *continuously differentiable* if the map  $Df : p \mapsto Df(p)$  is continuous.

Recall that linear maps are continuous if and only if they are bounded, which is always the case if  $V$  is finite-dimensional.

**Exercise 1.2.** Verify that if we identify scalars  $c \in \mathbb{R}$  with linear maps  $x \mapsto c \cdot x$ , this definition agrees with the usual definition for  $f : \mathbb{R} \rightarrow \mathbb{R}$ .

**Theorem 1.3.** If  $f : U \rightarrow W$  is differentiable, it is continuous.

*Proof.* If  $f$  is differentiable, we have

$$\lim_{\|h\|_V \rightarrow 0} \frac{\|f(p+h) - f(p) - Df(p)(h)\|_W}{\|h\|_V} = 0.$$

This can only be the case if

$$\lim_{\|h\|_V \rightarrow 0} \|f(p+h) - f(p) - Df(p)(h)\|_W = 0,$$

i.e. if

$$\lim_{\|h\|_V \rightarrow 0} f(p+h) = \lim_{\|h\|_V \rightarrow 0} f(p) + Df(p)(h).$$

Since  $Df(p)$  is continuous with  $Df(0) = 0$ , this simplifies to

$$\lim_{\|h\|_V \rightarrow 0} f(p+h) = f(p),$$

which is the definition of sequential continuity.  $\square$

**Definition 1.4: Directional and Partial Derivates**

Let  $V$  and  $W$  be normed vector spaces. Let  $v \in V$ . Let  $U \subseteq V$  be open and  $f : U \rightarrow W$ . Then we call the limit

$$\frac{\partial}{\partial v} f(p) = \left. \frac{d}{dh} f(p + hv) \right|_{h=0} = \lim_{h \rightarrow 0} \frac{f(p + hv) - f(p)}{h}$$

the **directional derivative** of  $f$  at  $p$  in direction  $v$ . If  $(x_0, \dots, x_n)$  is a basis of  $V$ , we call the directional derivatives

$$\partial_i f(p) := \frac{\partial}{\partial x_i} f(p)$$

the **partial derivatives of  $f$  at  $p$** . If all partial derivatives exist, we call  $f$  **partially differentiable**. If all maps  $\partial_i f : p \mapsto \partial_i f(p)$  are continuous, we call  $f$  **continuously partially differentiable**.

Partial derivatives aren't actually any more complicated than standard one-dimensional derivatives - you can calculate partial derivatives by simply fixing all coordinates except one and then taking a regular derivative in the remaining coordinate. For example, if we have a function

$$\begin{aligned} f : \mathbb{R}^3 &\rightarrow \mathbb{R} \\ (x, y, z) &\mapsto f(x, y, z), \end{aligned}$$

we can define a function  $f|_1 : \mathbb{R} \rightarrow \mathbb{R}$  by restriction to the first coordinate:

$$f|_1(x) := f(x, y, z).$$

Using this function, we can simplify the partial derivative:

$$\begin{aligned} \partial_1 f(x, y, z) &= \lim_{h \rightarrow 0} \frac{f((x, y, z) + he_1) - f(x, y, z)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x + h, y, z) - f(x, y, z)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f|_1(x + h) - f|_1(x)}{h} \\ &= \frac{d}{dx} f|_1(x) \end{aligned}$$

This tells us that, for example, the function  $f(x, y, z) = x \cdot y \cdot z$  has partial derivatives

$$\begin{aligned} \partial_1 f(x, y, z) &= \frac{d}{dx} xyz = yz \\ \partial_2 f(x, y, z) &= \frac{d}{dy} xyz = xz \\ \partial_3 f(x, y, z) &= \frac{d}{dz} xyz = xy. \end{aligned}$$

**Exercise 1.5.** 1. Show that if  $f$  is differentiable, then all directional derivatives exist and fulfill  $\frac{\partial}{\partial v} f(p) = Df(p)(v)$ .

2. Conclude that  $Df(p)$  is represented in basis  $(x_0, \dots, x_n)$  by the **Jacobian matrix**

$$Jf(p) = \left( \frac{\partial}{\partial x_1} f(p), \dots, \frac{\partial}{\partial x_n} f(p) \right) = \begin{pmatrix} \frac{\partial}{\partial x_1} f_1(p) & \dots & \frac{\partial}{\partial x_n} f_1(p) \\ \vdots & & \vdots \\ \frac{\partial}{\partial x_1} f_n(p) & \dots & \frac{\partial}{\partial x_n} f_n(p) \end{pmatrix}$$

If the image of  $f$  is one-dimensional, the Jacobian matrix is simply a row vector. We call the transpose of this vector the **gradient of  $f$  at  $p$**  and denote it

$$\nabla f(p) := Jf(p)^\top = \begin{pmatrix} \frac{\partial}{\partial x_1} f(p) \\ \vdots \\ \frac{\partial}{\partial x_n} f(p) \end{pmatrix}$$

The transposition comes from the fact that we would like to be able to get the directional derivatives of  $f$  at  $p$  by taking a scalar product with the gradient, which gives us

$$\begin{aligned} \langle \nabla f(p), v \rangle &= \nabla f(p)^\top v \\ &= Jf(p) \cdot v \\ &= Df(p)(v) \\ &= \frac{\partial}{\partial v} f(p) \end{aligned}$$

There is one very important pitfall to watch out for when it comes to differentiability: We have just shown that if  $f$  is differentiable, we can get the differential by construction the Jacobian matrix from the partial derivatives. This makes it look like every partially differentiable function is differentiable, which turns out to not be true in general.

For example, there exist functions which are partially differentiable, but not continuous - we showed earlier that all differentiable functions are continuous, so these functions are quick counterexamples.

One example is

$$\begin{aligned} f : \mathbb{R}^2 &\rightarrow \mathbb{R} \\ (x, y) &\mapsto \begin{cases} \frac{xy}{(x^2+y^2)^2} & (x, y) \neq 0 \\ 0 & (x, y) = 0 \end{cases} \end{aligned}$$

This function has  $\partial_1 f(0) = \partial_2 f(0) = 0$ , since  $f(h, 0) = f(0, h) = 0$  for all  $h \in \mathbb{R}$ , but isn't continuous, since  $\lim_{n \rightarrow \infty} f\left(\frac{1}{n}, \frac{1}{n}\right) = \lim_{n \rightarrow \infty} \frac{n^2}{4} = \infty \neq f(0)$ .

For a more involved counterexample, consider the function

$$\begin{aligned} f : \mathbb{R}^2 &\rightarrow \mathbb{R} \\ (x, y) &\mapsto \begin{cases} \frac{y^3}{x^2+y^2} & (x, y) \neq 0 \\ 0 & (x, y) = 0 \end{cases} \end{aligned}$$

This function is continuous - for  $(x, y) \neq 0$ , it is a composition of continuous

functions, and thus continuous, and since  $x^2 + y^2 \geq y^2$ , we additionally have

$$\begin{aligned} |f(x, y)| &= \frac{|y^3|}{|x^2 + y^2|} \\ &= \frac{|y^3|}{x^2 + y^2} \\ &\leq \frac{|y^3|}{y^2} \\ &= |y|, \end{aligned}$$

so  $f(x, y)$  converges to  $f(0, 0) = 0$  as  $(x, y)$  converges to 0, implying that  $f$  is also continuous at the origin.

$f$  is also directionally differentiable in all directions:

$$\begin{aligned} \partial_v f(0) &= \lim_{h \rightarrow 0} \frac{f(h \cdot v) - f(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(h \cdot v)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\left( \frac{h^3 v_2^3}{h^2 v_1^2 + h^2 v_2^2} \right)}{h} \\ &= \lim_{h \rightarrow 0} \frac{h^3 v_2^3}{h^3 v_1^2 + h^3 v_2^2} \\ &= \lim_{h \rightarrow 0} \frac{v_2^3}{v_1^2 + v_2^2} \\ &= \frac{v_2^3}{v_1^2 + v_2^2} \\ &= f(v) \end{aligned}$$

In particular,  $\partial_1 f(0) = 0$  and  $\partial_2 f(0) = 1$ . However,  $f$  is not differentiable at 0, since otherwise our representation  $Df(0) = Jf(0)$  would give us:

$$\begin{aligned} \lim_{\vec{h} \rightarrow 0} \frac{|f(h) - f(0) - Df(0)(\vec{h})|}{\|\vec{h}\|_2} &= \lim_{\vec{h} \rightarrow 0} \frac{|f(h) - Df(0)(h)|}{\|\vec{h}\|_2} \\ &= \lim_{\vec{h} \rightarrow 0} \left| \frac{f(h) - \partial_1 f(0) \cdot h_1 + \partial_2 f(0) \cdot h_2}{\sqrt{h_1^2 + h_2^2}} \right| = \lim_{\vec{h} \rightarrow 0} \left| \frac{f(h) - h_2}{\sqrt{h_1^2 + h_2^2}} \right| \\ &= \lim_{\vec{h} \rightarrow 0} \left| \frac{1}{\sqrt{h_1^2 + h_2^2}} \cdot \left( \frac{h_2^3}{h_1^2 + h_2^2} - h_2 \right) \right| = \lim_{\vec{h} \rightarrow 0} \left| \frac{1}{\sqrt{h_1^2 + h_2^2}} \cdot \left( \frac{h_2^3 - h_2(h_1^2 + h_2^2)}{h_1^2 + h_2^2} \right) \right| \\ &= \lim_{\vec{h} \rightarrow 0} \left| \frac{1}{\sqrt{h_1^2 + h_2^2}} \cdot \left( \frac{h_2^3 - h_2 h_1^2 - h_2^3}{h_1^2 + h_2^2} \right) \right| = \lim_{\vec{h} \rightarrow 0} \left| \frac{1}{\sqrt{h_1^2 + h_2^2}} \cdot \left( \frac{-h_2 h_1^2}{h_1^2 + h_2^2} \right) \right| = \lim_{\vec{h} \rightarrow 0} \left| \frac{h_2 h_1^2}{\sqrt{h_1^2 + h_2^2}^3} \right| \end{aligned}$$

Since we are only interested in whether this expression converges to 0, we can

square it to simplify slightly:

$$\left| \frac{h_2 h_1^2}{\sqrt{h_1^2 + h_2^2}^3} \right|^2 = \left| \frac{h_2^2 h_1^4}{(h_1^2 + h_2^2)^3} \right|$$

If we now set  $h := (\frac{1}{n}, \frac{1}{n})$ , we get:

$$\left| \frac{h_2^2 h_1^4}{(h_1^2 + h_2^2)^3} \right| = \left| \frac{\frac{1}{n^6}}{\left(\frac{2}{n^2}\right)^3} \right| = \left| \frac{\frac{1}{n^6}}{\frac{8}{n^6}} \right| = \frac{1}{8} \neq 0$$

The important takeaway here is that directional differentiability only guarantees “differentiability along straight paths”, while full differentiability also requires “differentiability along curved paths”.

**Theorem 1.6**

$f : V \rightarrow W$  is continuously differentiable if and only if it is continuously partially differentiable.

*Proof.*  $\Rightarrow$ : We have  $\text{Hom}(\mathbb{R}^n, \mathbb{R}^m) \cong \mathbb{R}^{mn}$ . If the differential  $Jf : \mathbb{R}^n \mapsto \mathbb{R}^{mn}$  exists, it is continuous if and only if every component  $(Jf)_i : \mathbb{R}^n \mapsto \mathbb{R}$  is continuous. By the definition of  $Jf$ , these are the partial derivatives.

$\Leftarrow$ : Since every normed complex vector space is isomorphic to a normed real vector space, we can assume without loss of generality that  $W$  is real. Since all norms are equivalent, and since a sequence in  $W$  converges iff. all components converge, we can additionally assume that  $W$  is one-dimensional, i.e.  $W \cong \mathbb{R}$ .

We need to show that

$$\lim_{h \rightarrow \vec{0}} \frac{f(p+h) - f(p) - Jf(p)h}{\|h\|_2} = \vec{0}$$

Let  $h := \sum_{i=1}^n h_i e_i$ . We decompose the path from  $p$  to  $p+h$  into its partial steps, taking one coordinate direction at a time:

$\forall k \in [0..n] :$

$$p_k = p + \sum_{i=1}^k h_i e_i$$

Now, by the mean value theorem, there exists an  $s_k \in [0, 1]$  such that

$$\begin{aligned} f(p_k) - f(p_{k-1}) &= f(p_{k-1} + h_k e_k) - f(p_{k-1}) \\ &= \partial_k f(p_{k-1} - s_k h_n e_n) h_k \end{aligned}$$

Now, since

$$Jf(p)(h) = \sum_{i=1}^n \partial_i f(p) h_i,$$

we get:

$$\begin{aligned}
 & \frac{|f(p+h) - f(p) - Jf(p)(h)|}{\|h\|_2} \\
 &= \frac{1}{\|h\|_2} \left| f(p_n) - f(p_0) - \sum_{i=1}^n \partial_i f(p) h_i \right| \\
 &= \frac{1}{\|h\|_2} \left| \sum_{i=1}^n f(p_i) - f(p_{i-1}) - \partial_i f(p) h_i \right| \\
 &= \frac{1}{\|h\|_2} \left| \sum_{i=1}^n \partial_i f(p_{i-1} + s_i h_i e_i) h_i - \partial_i f(p) h_i \right| \\
 &= \frac{1}{\|h\|_2} \left| \sum_{i=1}^n (\partial_i f(p_{i-1} + s_i h_i e_i) - \partial_i f(p)) h_i \right| \\
 &= \frac{1}{\|h\|_2} \left| \sum_{i=1}^n \left( \partial_i f \left( p + \sum_{j=1}^{i-1} h_j e_j + s_i h_i e_i \right) - \partial_i f(p) \right) h_i \right| \\
 &\leq \frac{1}{\|h\|_2} \sum_{i=1}^n \left| \left( \partial_i f \left( p + \sum_{j=1}^{i-1} h_j e_j + s_i h_i e_i \right) - \partial_i f(p) \right) h_i \right| \\
 &= \sum_{i=1}^n \left| \partial_i f \left( p + \sum_{j=1}^{i-1} h_j e_j + s_i h_i e_i \right) - \partial_i f(p) \right| \cdot \frac{|h_i|}{\|h\|_2} \\
 &\leq \sum_{i=1}^n \left| \partial_i f \left( p + \sum_{j=1}^{i-1} h_j e_j + s_i h_i e_i \right) - \partial_i f(p) \right|
 \end{aligned}$$

Since  $s_i \in [0, 1]$ , we have

$$\lim_{h \rightarrow 0} p + \sum_{j=1}^{i-1} h_j e_j + s_i h_i e_i = p$$

And thus, since the partial derivatives  $\partial_i f$  are continuous, the right side converges to zero, so  $f$  is differentiable.

□

**Exercise 1.7.** Let  $V$  be an inner product space over a field  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{F} = \mathbb{C}$ . Let  $v \in V$ . Then the function

$$\begin{aligned}
 \langle \cdot, v \rangle : V &\rightarrow \mathbb{F} \\
 w &\mapsto \langle w, v \rangle
 \end{aligned}$$

is differentiable, with  $D \langle \cdot, v \rangle (w) = v^\top$  for all  $w \in V$ .

**Exercise 1.8.** Let  $b$  be a symmetric bilinear form. Let

$$q(v) := b(v, v)$$

be the corresponding quadratic form. Then  $q$  is differentiable, with

$$Dq(v)(w) = 2b(v, w)$$

## 1.2 Rules of Differentiation

**Exercise 1.9.** Show that the differential operator  $D$  is linear, i.e. that we have

$$D(\alpha f + \beta g)(p) = \alpha Df(p) + \beta Dg(p)$$

for  $f, g$  differentiable at  $p$ .

### Theorem 1.10: Chain Rule in $\mathbb{R}^n$

Let  $f : U \rightarrow \mathbb{R}^m$  and  $g : V \rightarrow \mathbb{R}^r$  with  $U \subset \mathbb{R}^n, V \subset \mathbb{R}^m$  open and  $f(U) \subseteq V$ . Let  $f$  be differentiable at a point  $p \in U$  and let  $g$  be differentiable at  $f(p)$ . Then

$$D(g \circ f)(p) = Dg(f(p)) \circ Df(p)$$

**Exercise 1.11.** Verify that this version of the chain rule agrees with the familiar chain rule for  $f, g : \mathbb{R} \rightarrow \mathbb{R}$ .

*Proof.* By definition of the differential, we have:

$$\begin{aligned} f(x) &= f(p) \\ &\quad + Df(p)(x - p) \\ &\quad + \varepsilon_f(x)\|x - p\| \\ g(f(x)) &= g(f(p)) \\ &\quad + Dg(f(p))(f(x) - f(p)) \\ &\quad + \varepsilon_g(f(x))\|f(x) - f(p)\| \end{aligned}$$

For some functions  $\varepsilon_f : U \rightarrow \mathbb{R}^m, \varepsilon_g : U \rightarrow \mathbb{R}^r$  which are continuous in  $p$  and  $f(p)$  and fulfill  $\varepsilon_f(p) = 0$  and  $\varepsilon_g(f(p)) = 0$ .

The second equation tells us:

$$\begin{aligned} g(f(x)) &= g(f(p)) \\ &\quad + Dg(f(p))(f(x) - f(p)) \\ &\quad + \varepsilon_g(f(x))\|f(x) - f(p)\| \end{aligned}$$

Plugging in our equation for  $f(x)$  now lets us regroup our terms into an

equation of the same form as the definition of the differential:

$$\begin{aligned}
 g(f(x)) &= g(f(p)) \\
 &\quad + Dg(f(p)) \left[ \overbrace{f(p) + Df(p)(x-p) + \varepsilon_f(x)\|x-p\| - f(p)}^{f(x)} \right] \\
 &\quad + \varepsilon_g(f(x)) \left\| \overbrace{f(p) + Df(p)(x-p) + \varepsilon_f(x)\|x-p\| - f(p)}^{f(x)} \right\| \\
 &= g(f(p)) \\
 &\quad + Dg(f(p)) \left[ Df(p)(x-p) + \varepsilon_f(x)\|x-p\| \right] \\
 &\quad + \varepsilon_g(f(x)) \left\| Df(p)(x-p) + \varepsilon_f(x)\|x-p\| \right\| \\
 &= g(f(p)) \\
 &\quad + Dg(f(p))Df(p)(x-p) \\
 &\quad + Dg(f(p))\varepsilon_f(x)\|x-p\| \\
 &\quad + \varepsilon_g(f(x)) \left\| Df(p)(x-p) + \varepsilon_f(x)\|x-p\| \right\| \\
 &= g(f(p)) \\
 &\quad + Dg(f(p))Df(p)(x-p) \\
 &\quad + \underbrace{\left( \varepsilon_f(x)Dg(f(p)) + \varepsilon_g(f(x)) \left\| Df(p) \left( \frac{x-p}{\|x-p\|} \right) + \varepsilon_f(x) \right\| \right)}_{:=\varepsilon_{g \circ f}(x)} \|x-p\| \\
 &= g(f(p)) \\
 &\quad + (Dg(f(p)) \circ Df(p))(x-p) \\
 &\quad + \varepsilon_{g \circ f}(x)\|x-p\|
 \end{aligned}$$

All that remains is to show that  $\varepsilon_{g \circ f}(x)$  is continuous if we set

$$\varepsilon_{g \circ f}(p) = 0.$$

Since we have  $\varepsilon_f(p) = 0$  and  $\varepsilon_g(f(p)) = 0$ , we have our desired result if we can show that

$$\left\| Df(p) \left( \frac{x-p}{\|x-p\|} \right) \right\| = \frac{\|Df(p)(x-p)\|}{\|x-p\|}.$$

is bounded. Since  $\mathbb{R}^n$  is finite-dimensional, we can find a constant  $c$  such that

$$\|Df(p)(x-p)\| \leq c \cdot \|x-p\|,$$

which means that our quotient is indeed bounded, completing the proof.  $\square$

**Theorem 1.12: Product Rule in  $\mathbb{R}^n$** 

Let  $U \subseteq \mathbb{R}^n$ . Let  $f, g : U \rightarrow \mathbb{R}$  at a point  $p \in U$ . Then

$$D(f \cdot g)(p) = Df(p) \cdot g(p) + f(p) \cdot Dg(p).$$

**Exercise 1.13.** Verify that this version of the product rule agrees with the familiar product rule for  $f, g : \mathbb{R} \rightarrow \mathbb{R}$ .

*Proof.* We define the multiplication function  $m(x, y) := x \cdot y$ . The partial derivatives are  $\partial_1 m(x, y) = y$  and  $\partial_2 m(x, y) = x$ , i.e.

$$Dm(x, y) = \begin{pmatrix} y & x \end{pmatrix}$$

Applying the chain rule now gives

$$\begin{aligned} D(f \cdot g)(p) &= D(m \circ (f, g))(p) \\ &= Dm((f, g)(p)) \circ D(f, g)(p) \\ &= Dm(f(p), g(p)) \circ (Df(p), Dg(p)) \\ &= Dm(f(p), g(p)) \cdot (Df(p), Dg(p)) \\ &= \begin{pmatrix} g(p) & f(p) \end{pmatrix} \cdot \begin{pmatrix} Df(p) \\ Dg(p) \end{pmatrix} \\ &= Df(p) \cdot g(p) + f(p) \cdot Dg(p). \end{aligned}$$

□

**Exercise 1.14.** Let  $f : V \rightarrow W$  be differentiable. Let  $\gamma : [0, 1] \rightarrow V$  be differentiable. Then  $f \circ \gamma$  is differentiable with

$$(f \circ \gamma)'(t) = Df(\gamma'(t)) \cdot \gamma'(t)$$

**Theorem 1.15.** Let  $U \subseteq V$  be open. Let  $f : U \rightarrow W$  be differentiable with  $Df = 0$ . Then  $f$  is locally constant.

**Corollary 1.16.** Let  $U \subseteq V$  be open and connected. Let  $f : U \rightarrow W$  be differentiable with  $Df = 0$ . Then  $f$  is constant.

## 1.3 Higher Partial Derivatives

### Theorem 1.17: Schwartz's Theorem

Let  $f : U \subseteq V \rightarrow W$  be twice continuously partially differentiable. Then the order of partial derivation does not matter, i.e.

$$\partial_i \partial_j f = \partial_j \partial_i f$$

for all  $i, j \in [1..n]$ .

### Theorem 1.18: First-Order Taylor Expansion in multiple Variables

Let  $f : U \subseteq V \rightarrow W$ . Let  $f$  be continuously differentiable. Let  $x \in U$ ,  $h \in V$  fulfill  $[x, x + h] \subseteq U$ . Then

$$f(x + h) = f(x) + Df(x)(h) + r(h),$$

with  $r \in o(\|h\|)$ .

## 1.4 Differential Operators

## Chapter 2

# The Two Queens of Multivariable Calculus

### 2.1 The Inverse Function Theorem

**Theorem 2.1: Inverse Function Theorem**

Let  $U \subseteq \mathbb{R}^n$  be open. Let  $f : U \rightarrow \mathbb{R}^n$  be continuously differentiable. Let  $x \in U$  be a point such that  $Df(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is invertible. Then there exists an open neighbourhood  $V$  of  $x$  such that:

1.  $f(V)$  is an open neighbourhood of  $f(x)$ .
2.  $f|_V$  is a  $C^1$ -diffeomorphism, i.e. invertible with continuously differentiable inverse.

In short: if a continuously differentiable function has an invertible differential at a given point, there exists an open neighbourhood around that point in which the function is invertible:

**Corollary 2.2.** Let  $U \subseteq \mathbb{R}^n$  be open. Let  $f : U \rightarrow \mathbb{R}^n$  be continuously differentiable. Let  $Df(x)$  be invertible for all  $x \in \mathbb{R}^n$ . Then  $f(U)$  is open.

## 2.2 The Implicit Function Theorem

### Theorem 2.3: Implicit Function Theorem

Let  $U \subseteq \mathbb{R}^m \times \mathbb{R}^k$  be open. Let  $f : U \rightarrow \mathbb{R}^k$  be continuously differentiable. Let  $(x, y) \in U$  with  $f(x, y) = z$ . Let  $Df|_{\mathbb{R}^k}(x, y) : \mathbb{R}^k \rightarrow \mathbb{R}^k$  be invertible. Then there exist

1. open neighbourhoods  $V$  of  $x$  and  $W$  of  $y$ ,
2. and a continuously differentiable function  $g : V \rightarrow W$ ,

such that the set of values in  $V \times W$  for which  $f$  takes the value  $z$  is the graph of  $g$ , i.e.

$$\{(v, w) \in V \times W \mid f(v, w) = z\} = \{(v, g(v)) \mid v \in V\}$$

In short: under mild assumptions, the level sets of a continuously differentiable function are graphs of continuously differentiable functions:

## Chapter 3

# Multiple Integrals

**Lemma 3.1.** Let  $K \subseteq \mathbb{R}^n$  be compact. Let  $f : K \times [a, b] \rightarrow \mathbb{R}$  be continuous. Then the function

$$x \mapsto \int_a^b f(x, t) dt$$

is uniformly continuous.

*Proof.*  $K \times [a, b]$  is a finite product of compact sets, i.e. compact by Theorem 2.33. Thus,  $f$  is uniformly continuous by Theorem 2.35, which means that for every  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that for any  $x, y \in \mathbb{R}, t \in \mathbb{R}^n$

$$d((x, t), (y, t)) < \delta \implies d(f(x, t), f(y, t)) < \varepsilon.$$

This tells us that for any  $x, y \in \mathbb{R}$ ,

$$\begin{aligned} d(x, y) < \delta &\implies d((x, t), (y, t)) < \varepsilon \\ &\implies |f(x, t) - f(y, t)| = d(f(x, t), f(y, t)) < \varepsilon \\ &\implies d\left(\int_a^b f(x, t) dt, \int_a^b f(y, t) dt\right) \\ &= \left| \int_a^b f(x, t) dt - \int_a^b f(y, t) dt \right| \\ &\leq \int_a^b |f(x, t) - f(y, t)| dt \\ &\leq (b - a)\varepsilon, \end{aligned}$$

since  $\varepsilon$  was arbitrary, our function is uniformly continuous as desired.  $\square$

**Definition 3.2.** Given a compact cuboid  $Q = \prod_{i=1}^n [a_i, b_i] \subseteq \mathbb{R}^n$  and a continuous function  $f : Q \rightarrow \mathbb{R}$ , we declare the *integral of  $f$  over  $Q$*  to be the nested integral

$$\int_Q f(x) d^n x := \int_{a_n}^{b_n} \dots \int_{a_1}^{b_1} f(x_1, \dots, x_n) dx_1 \dots dx_n$$

By Lemma 3.1, all involved functions are continuous, and thus their integrals are indeed well-defined. This definition will become a fully justified theorem

once we begin our study of the Lebesgue integral later on.

**Exercise 3.3.** Verify that linearity and monotonicity carry over from the single-variable case, i.e show that for any  $f, g : Q \rightarrow \mathbb{R}$  and  $\lambda \in \mathbb{R}$ , we have:

1.  $\int_Q f(x) + g(x) d^n x = \int_Q f(x) d^n x + \int_Q g(x) d^n x$
2.  $\int_Q \lambda f(x) d^n x = \lambda \int_Q f(x) d^n x$
3.  $f \leq g \implies \int_Q f(x) d^n x \leq \int_Q g(x) d^n x$
4.  $\left| \int_Q f(x) d^n x \right| \leq \int_Q |f(x)| d^n x$

**Definition 3.4.** We define

$$\text{vol } Q = \int_Q 1 = \prod_{i=1}^n |b_i - a_i|$$

**Definition 3.5.** By partitioning the intervals  $[a, b] \times [c, d] := Q \subseteq \mathbb{R}^2$  into  $r$  equidistant chunks with boundaries

$$\begin{aligned} a &= a_1 \leq a_2 \leq \dots \leq a_r = b \\ c &= c_1 \leq c_2 \leq \dots \leq c_r = d, \end{aligned}$$

we partition  $Q$  into  $r^2$  equally sized smaller rectangles

$$Q_{ij} := [a_i, c_{i+1}] \times [a_j, c_{j+1}]$$

which must each have area  $\text{vol } Q / r^2$ . We can thus define *two-dimensional Riemann sums* analogously to how we did it in one dimension:

$$S_r^2(f) := \sum_{i=0}^{r-1} \sum_{j=0}^{r-1} f(a_i, c_j) \frac{\text{vol } Q}{r^2}$$

#### Lemma 3.6: Multivariable Integration via Riemann Sums

Given a compact cuboid  $Q \subseteq \mathbb{R}^2$  and a continuous function  $f : Q \rightarrow \mathbb{R}$ , we have

$$\int_Q f(x) d^2 x = \lim_{r \rightarrow \infty} S_r^2(f)$$

*Proof.* Applying the single-variable version of this theorem twice, we get:

$$\begin{aligned} \int_Q f(x) d^2 x &= \int_c^d \int_a^b f(x_1, x_2) dx_1 dx_2 \\ &= \lim_{r \rightarrow \infty} S_r \left( \int_a^b f(x_1, x_2) dx_1 \right) \\ &= \lim_{r \rightarrow \infty} S_r^2(f(x_1, x_2)) \end{aligned}$$

□

## **Chapter 4**

# **Curves**

## **Chapter 5**

# **Surfaces**

## **Part VIII**

# **Point-Set Topology**

# Chapter 1

## Topological Spaces

### 1.1 Topologies

#### Definition 1.1: Accumulation points and isolated points

Let  $X$  be a topological space. Let  $A \subseteq X$ . Then we call  $x \in X$  an *accumulation point* of  $A$  if every neighbourhood of  $x$  contains a point in  $A$  other than  $x$  itself.

If a point  $x \in A$  is not an accumulation point of  $A$ , we call  $x$  an *isolated point*.

**Definition 1.2.** Let  $X$  be a topological space. Let  $A \subseteq X$ . Then the set of all accumulation points of  $A$  is denoted  $A'$ , and is known as the *derived set of  $A$* .

**Exercise 1.3.** Show that  $A' \subseteq \overline{A}$ .

## 1.2 Basic Separation Properties

*A very common axiom is the Hausdorff separation axiom, which says that any two distinct points can be housed orf from each other by disjoint open sets.*

“Topology via Logic”,  
Steven Vickers

**Definition 1.4.** Let  $(X, \mathcal{O})$  be a topological space. Let  $x, y \in X$ . Then we call  $x$  and  $y$  *topologically distinguishable* iff.  $\mathcal{U}(x) \neq \mathcal{U}(y)$ , i.e. one of the points has a neighborhood not containing the other.

### Definition 1.5: $T_0$ Space

Let  $(X, \mathcal{O})$  be a topological space. Then we call  $(X, \mathcal{O})$   *$T_0$*  if every pair of points is topologically distinguishable.

### Definition 1.6: $T_1$ Space

Let  $(X, \mathcal{O})$  be a topological space. Then we call  $(X, \mathcal{O})$   *$T_1$*  if for every pair of points, both of them have neighbourhoods not containing the other:

$$\begin{aligned} \forall x, y \in X : \\ \exists U_x \in \mathcal{U}(x), U_y \in \mathcal{U}(y) : \\ x \notin U_y, y \notin U_x. \end{aligned}$$

**Exercise 1.7.** 1. Find a topological space that isn't  $T_0$ .

2. Show that every  $T_1$  space is  $T_0$ .

3. Find a space that is  $T_0$ , but not  $T_1$ .

**Exercise 1.8.** Show that  $(X, \mathcal{O})$  is  $T_1$  if and only if every singleton set is closed.

**Exercise 1.9.** Show that  $(X, \mathcal{O})$  is  $T_1$  if and only if every subset of  $X$  is the intersection of all of its neighborhoods:

$$\forall Y \subseteq X : Y = \bigcap \{U \mid U \in \mathcal{U}(Y)\}$$

### Definition 1.10: Hausdorff Space

Let  $(X, \mathcal{O})$  be a topological space. Then we call  $(X, \mathcal{O})$   *$T_2$  or Hausdorff* if every pair of points can be housed orf from each other, i.e. they have disjoint neighbourhoods:

$$\begin{aligned} \forall x \neq y \in X : \\ \exists U_x \in \mathcal{U}(x), U_y \in \mathcal{U}(y) : \\ U_x \cap U_y = \emptyset \end{aligned}$$

**Exercise 1.11.** Show that every Hausdorff space is  $T_1$ .

**Exercise 1.12.** Let  $(X, \mathcal{O})$  be Hausdorff. Then every convergent sequence in  $X$  has a unique limit.

**Exercise 1.13.** Let  $X$  be a Hausdorff space. Let  $x$  be an accumulation point of  $A \subseteq X$ . Then every neighbourhood of  $x$  contains infinitely many points of  $A$ .

**Lemma 1.14.** Let  $X$  be a Hausdorff space. Then the set of accumulation points of any subset is closed.

*Proof.* Let  $A \subseteq X$ . Let  $A'$  be the set of accumulation points of  $A$ . Let  $x$  be an accumulation point of  $A'$ . Since  $X$  is Hausdorff, every neighbourhood of  $x$  must contain infinitely many points  $x_i \in A'$ . But since these neighbourhoods are also neighbourhoods of the  $x_i$ , they must also all contain infinitely many points in  $A$ .

Therefore,  $x$  is already an accumulation point of  $A$ , i.e.  $x \in A'$ . Therefore,  $A'$  contains all of its accumulation points, and is thus closed.  $\square$

**Exercise 1.15.** Find a space  $X$  and a subset  $A$  such that the set of accumulation points of  $A$  isn't closed.

**Theorem 1.16.** Let  $(X, \mathcal{O})$  be a topological space. Then  $(X, \mathcal{O})$  is Hausdorff if and only if every point is the intersection of the closures of all of its neighbourhoods:

$$\forall x \in X : \{x\} = \bigcap \left\{ \overline{U} \mid U \in \mathcal{U}(x) \right\}$$

*Proof.*  $\Rightarrow$  Let  $(X, \mathcal{O})$  be Hausdorff. Let  $x, y \in X$  and  $y \notin \{x\}$ , i.e.  $y \neq x$ . Then, by Hausdorff, there exist disjoint neighborhoods  $U_x$  and  $U_y$  of  $x$  and  $y$ . Since  $U_x$  and  $U_y$  are disjoint, we have  $U_x \subseteq X \setminus U_y$ . But then, by monotonicity of the closure operator, we get

$$\overline{U_x} \subseteq \overline{X \setminus U_y} = X \setminus U_y,$$

i.e.  $y \notin \overline{U_x}$ , i.e.

$$y \notin \bigcap \left\{ \overline{U} \mid U \in \mathcal{U}(x) \right\}.$$

$\Leftarrow$  Let  $x \neq y$ . Let  $\{x\} = \bigcap \left\{ \overline{U} \mid U \in \mathcal{U}(x) \right\}$ . Then there must be an open neighborhood  $U_x$  of  $x$  whose closure doesn't contain  $y$ . But then the complement  $U_y := X \setminus \overline{U_x}$  of that closure is an open neighborhood of  $y$  that is disjoint from  $U_x$ .

$\square$

## 1.3 Bases and Countability

### Definition 1.17

We call a topological space *first countable* if it has a countable neighbourhood basis around every point  $x$ , i.e. a set of neighbourhoods  $B_i$  of  $x$  such that every neighbourhood of  $x$  contains some  $B_i$ .

**Exercise 1.18.** Every metric space is first countable.

It turns out that first countable spaces are exactly the spaces where sequences are sufficient to maintain our intuitions about how sequence convergence interacts with topological properties:

### Theorem 1.19: Detecting openness and closedness using sequences

Let  $X$  be first countable. Let  $A \subseteq X$ . Let  $x \in X$ . Then:

1.  $x \in \text{int } A$  if and only if every sequence in  $X$  converging to  $x$  is eventually in  $A$ .
2.  $x \in \overline{A}$  if and only if there exists a sequence in  $A$  converging to  $x$ .

*Proof.* Copy the proof of the metric case, replacing open balls with our countable neighbourhood basis.  $\square$

**Corollary 1.20.** Let  $X$  be first countable. Let  $A \subseteq X$ . Let  $x \in X$ . Then:

1.  $A$  is closed in  $X$  if and only if it contains every limit of every convergent sequence of points in  $A$ .
2.  $A$  is open in  $X$  if and only if every sequence in  $X$  converging to a point of  $A$  is eventually in  $A$ .

### Definition 1.21

We call a topological space *separable* if it contains a countable dense subset.

**Exercise 1.22.**

1. Every countable space is separable.
2.  $\mathbb{R}^n$  is separable for every  $n \in \mathbb{N}$ .

### Definition 1.23

We call a topological space *second countable* if it has a countable basis.

**Exercise 1.24.**  $\mathbb{R}^n$  is second countable for every  $n \in \mathbb{N}$ .

**Exercise 1.25.** Let  $X$  be second countable. Then:

1.  $X$  is first countable.
2.  $X$  is separable.

**Theorem 1.26.** A metric space is second countable if and only if it is separable.

## Chapter 2

# New Spaces from Old

### 2.1 The Subspace Topology

#### Theorem 2.1

Let  $(X, \mathcal{O})$  be a topological space, and let  $S \subseteq X$  be any subset. Then the collection

$$\mathcal{O}_S := \{U \subseteq S \mid \exists V \in \mathcal{O} : U = S \cap V\},$$

consisting of all intersections of open sets of  $X$  with  $S$ , forms a topology, which we call the *subspace topology*.

**Exercise 2.2.** Let  $S \subseteq X$ . Show that  $B \subseteq S$  is closed in  $S$  if and only if it is equal to the intersection of  $S$  with some closed subset of  $X$ .

**Exercise 2.3.** Let  $X$  be a topological space. Let  $U \subseteq S \subseteq X$ .

1. If  $U$  is open in  $X$ , then  $U$  is open in  $S$ .
2. If  $U$  is closed in  $X$ , then  $U$  is closed in  $S$ .

**Exercise 2.4.** Let  $X$  be a topological space. Let  $U \subseteq S \subseteq X$ .

1. If  $U$  is open in  $S$ , and  $S$  is open in  $X$ , then  $U$  is open in  $X$ .
2. If  $U$  is closed in  $S$ , and  $S$  is closed in  $X$ , then  $U$  is closed in  $X$ .

If  $U \subseteq S \subseteq X$ , the notations  $\overline{U}$ ,  $\text{int } U$ , and  $\text{ext } U$  will always refer to the closure, interior, and exterior of  $U$  in  $X$ , not in  $S$ .

**Exercise 2.5.** Let  $X$  be a topological space. Let  $U \subseteq S \subseteq X$ .

1. The closure of  $U$  in  $S$  is equal to  $\overline{U} \cap S$ .
2. Find an example where the interior of  $U$  in  $S$  is not equal to  $\text{int } U \cap S$ .
3. Show that  $\text{int } U \cap S$  is a subset of the interior of  $U$  in  $S$ .

#### Theorem 2.6: Universal Property of the Subspace Topology

Let  $X$  and  $Y$  be topological spaces. Let  $S \subseteq Y$ . Then  $f : X \rightarrow S$  is continuous if and only if the embedded map  $\iota_S \circ f : X \rightarrow Y$  is continuous.

**Corollary 2.7.** Let  $S \subseteq X$ . Then the inclusion map  $\iota_S : S \hookrightarrow X$  is continuous.

**Corollary 2.8.** Let  $X$  and  $Y$  be topological spaces. Let  $f : X \rightarrow Y$  be continuous. Then:

1. If  $S \subseteq X$ , the restriction

$$f|_S : S \rightarrow Y$$

is continuous.

2. If  $f(X) \subseteq T \subseteq Y$ , the map

$$f : X \rightarrow T$$

is well-defined and continuous.

3. If  $Y \subseteq Z$ , then the embedded function

$$\iota_Y \circ f : X \rightarrow Z$$

is continuous.

**Exercise 2.9.** Let  $X$  be a topological space. Let  $R \subseteq S \subseteq X$ . Then the subspace topologies of  $R$  in  $S$  and  $X$  agree.

**Exercise 2.10.** Let  $(X, \mathcal{O})$  be a topological space. Let  $S \subseteq X$ . Then if  $\mathcal{B}$  is a basis of  $\mathcal{O}$ , the collection

$$\mathcal{B}_S = \{B \cap S \mid B \in \mathcal{B}\}$$

is a basis of the subspace topology  $\mathcal{O}_S$ .

**Exercise 2.11.**

1. Every subspace of a  $T_0$  space is  $T_0$ .
2. Every subspace of a  $T_1$  space is  $T_1$ .
3. Every subspace of a Hausdorff space is Hausdorff.
4. Every subspace of a first countable space is first countable.
5. Every subspace of a second countable space is second countable.

**Definition 2.12**

A map that is a homeomorphism onto its image is called a **topological embedding**, or just **embedding**.

**Exercise 2.13.** A homeomorphism is a surjective topological embedding.

**Exercise 2.14.** Let  $X$  be a topological space. Let  $S \subseteq X$ . Then the inclusion map  $\iota_S : S \hookrightarrow X$  is an embedding.

**Exercise 2.15.** Every embedding is continuous and injective.

**Exercise 2.16.** Every injective continuous map that is open or closed is an embedding.

**Exercise 2.17.** Find an embedding that is neither open nor closed.

**Theorem 2.18**

Let  $U \subseteq \mathbb{R}^n$  be open. Let  $f : U \rightarrow \mathbb{R}^k$  be continuous. Then the graph  $\Gamma(f)$  of  $f$  is homeomorphic to  $U$ .

*Proof.* Let  $\Phi_f : U \rightarrow \mathbb{R}^{n+k}$  be the canonical map

$$\Phi_f(x) = (x, f(x))$$

taking  $U$  to its graph.

By definition of the graph  $\Gamma(f)$ ,  $\Phi_f(x) : U \rightarrow \Gamma(f)$  is surjective.

Since  $\Phi_f$  is a composition of continuous maps, it is continuous.

Since the first  $n$  components of  $\Phi_f$  are the identity,  $\Phi_f$  is injective.

Finally, the projection  $p : \mathbb{R}^{n+k} \rightarrow \mathbb{R}^n$  is continuous, so the restriction  $p|_{\Gamma(f)}$  is a continuous inverse of  $\Phi_f$ .  $\square$

## 2.2 Product Spaces

**Lemma 2.19.** Let  $(X, \mathcal{O})$  be a topological space. Then  $(X, \mathcal{O})$  is Hausdorff if and only if the diagonal

$$\Delta := \{(x, x) \mid x \in X\}$$

is closed in the product topology.

*Proof.*  $\Rightarrow$  Assume  $(X, \mathcal{O})$  is Hausdorff. Let  $(x, y) \notin \Delta$ . Then  $x \neq y$  by definition of  $\Delta$ . Therefore, there exist disjoint neighbourhoods  $U_x, U_y$  of  $x$  and  $y$ . Since these neighborhoods are disjoint, we have

$$\Delta \cap (U_x \times U_y) = \emptyset$$

. Therefore, every point  $(x, y) \notin \Delta$  has an open neighborhood  $(U_x \times U_y)$  that doesn't intersect  $\Delta$ , i.e.  $\Delta$  is closed.

$\Leftarrow$  Let  $x \neq y$ . Assume that  $\Delta$  is closed. Then  $(x, y) \notin \Delta$  has a basic open neighborhood  $U_x \times U_y$  that doesn't intersect  $\Delta$ . Since  $U_x \times U_y$  doesn't intersect  $\Delta$ ,  $U_x$  and  $U_y$  must be disjoint.

□

## 2.3 Disjoint Union Spaces

## 2.4 Quotient Spaces

## 2.5 Adjunction Spaces

## Chapter 3

# Topological Manifolds

### Definition 3.1

We call a topological space  $X$  *locally Euclidean of dimension  $n$*  if every point  $x \in X$  has a neighbourhood that is homeomorphic to an open subset of  $\mathbb{R}^n$ .

**Exercise 3.2.** If a topological space  $X$  is a locally euclidean of dimension  $n$ , then:

1. Every point  $x \in X$  has a neighbourhood homeomorphic to an open ball in  $\mathbb{R}^n$ , or
2. Every point  $x \in X$  has a neighbourhood homeomorphic to  $\mathbb{R}^n$ , or

**Exercise 3.3.** Show that a topological space is locally Euclidean of dimension 0 if and only if it is discrete.

### Definition 3.4

A topological space is called an  *$n$ -dimensional topological manifold*, or just *topological  $n$ -manifold*, if it is:

1. locally euclidean of dimension  $n$ ,
2. second countable,
3. and Hausdorff.

The latter two requirements exclude some pathological spaces - the long line would be a topological manifold if it was second countable, and the bug-eyed line would be a topological manifold if it was Hausdorff. Most interesting results about topological manifolds require both of these properties, so little would be gained by omitting them.

Intuitively, it sounds very sensible that a nonempty topological space cannot be both a topological  $m$ -manifold and a topological  $n$ -manifold. This statement turns out to be true - however, we will save this proof for later, since it requires a substantial amount of machinery borrowed from algebraic topology.

**Lemma 3.5.** Every open subset of a topological  $n$ -manifold is a topological  $n$ -manifold.

**Exercise 3.6.** A topological space is a topological 0-manifold if and only if it is countable and discrete.

**Theorem 3.7.** Any separable metric space that is locally Euclidean of dimension  $n$  is a topological  $n$ -manifold.

**Definition 3.8**

We call a topological space an  *$n$ -dimensional manifold with boundary* if it is second countable, Hausdorff, and every open point has a neighbourhood homeomorphic to an open subset of either  $\mathbb{R}^n$ , or to the  $n$ -dimensional upper half-space

$$\mathbb{H}^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n \geq 0\}$$

**Definition 3.9**

Let  $M$  be a topological  $n$ -manifold with boundary. Then a *coordinate chart for  $M$*  is a pair  $(U, \varphi)$ , consisting of an open subset  $U \subseteq M$ , called a *coordinate domain*, and a homeomorphism  $\varphi$  from  $U$  to an open subset of  $\mathbb{R}^n$  or  $\mathbb{H}^n$ , called a *coordinate map*. If  $U$  is homeomorphic to an open ball, we call  $U$  a *coordinate ball*.

Note that if we can assign coordinate maps for some covering of a topological space  $X$ , then that space must by definition be locally euclidean.

**Definition 3.10: Manifold interior and manifold boundary**

If  $\varphi(U)$  is an open subset of  $\mathbb{R}^n$ , we call  $\varphi$  an *interior chart* and points  $p \in U$  *interior points*. If  $\varphi(U)$  is a subset of  $\mathbb{H}^n$  with  $\varphi(U) \cap \partial\mathbb{H}^n \neq \emptyset$ , we call  $\varphi$  a *boundary chart*. If there exists a boundary chart taking  $p$  to  $\partial\mathbb{H}^n$ , we call  $p$  a *boundary point of  $M$* .

Every manifold is a manifold with boundary, but the opposite might not hold. We sometimes use the redundant term "manifold without boundary" to refer to a manifold with boundary which has the empty set as its boundary. The word "manifold", without further qualifiers, always means a manifold without boundary.

The definition of a manifold's boundary only agrees with the topological definition once we find the right ambient space to embed  $M$  into, since  $M$  always has empty topological boundary when viewed as a topological space of its own.

For example, the manifold boundary of the closed unit disk  $\overline{\mathbb{B}}^2$  is the unit circle, which agrees with the topological boundary if the disk is viewed as a subset of  $\mathbb{R}^2$ , but not with the topological boundary of the disk when viewed as its own topological space (which is trivially empty, as stated), or with the topological boundary of the disk viewed as a subset of  $\mathbb{R}^3$  (which is the entire disk).

**Theorem 3.11.** If  $M$  is a topological  $n$ -manifold with boundary, then  $\text{int } M$  is both an open subset of  $M$  and a topological  $n$ -manifold without boundary.

*Proof.* Since  $\text{int } M$  is a union of coordinate domains, it is open. Second-countability and Hausdorffness are trivially inherited by subspaces. By definition,  $\text{int } M$  consists only of points homeomorphic to an open subset of  $\mathbb{R}^n$ , so  $\text{int } M$  is additionally locally euclidean of dimension  $n$ .  $\square$

**Proposition 3.12.** Let  $M$  be a topological manifold with boundary. Then a point  $x \in M$  cannot be both a boundary point and an interior point.

The general proof once again requires some tricky algebraic topology, so we save it for later.

**Corollary 3.13.** If  $M$  is a nonempty topological  $n$ -manifold with boundary, then  $\partial M$  is a closed subset of  $M$ , and  $M$  is a topological  $n$ -manifold if and only if  $\partial M = \emptyset$ .

*Proof.* Since  $\text{int } M$  is an open subset of  $M$ ,  $\partial M = M \setminus \text{int } M$  is closed. If  $M$  is a manifold, then every point is an interior point, so  $M = \text{int } M$ , so  $\partial M = \emptyset$ , and vice versa.  $\square$

## 3.1 Manifolds from Embeddings

### Theorem 3.14

Let  $U \subseteq \mathbb{R}^n$  be open. Let  $f : U \rightarrow \mathbb{R}^k$  be continuous. Then the graph  $\Gamma(f)$  of  $f$  is a topological  $n$ -manifold.

*Proof.* By 2.18,  $\Gamma(f)$  is homeomorphic to  $U \subseteq \mathbb{R}^n$ , making  $\Gamma(f)$  a trivial topological  $n$ -manifold coverable by a single coordinate domain.  $\square$

### Theorem 3.15

The unit  $n$ -spheres  $\mathbb{S}^n$  are topological  $n$ -manifolds.

*Proof.* For each  $i \in [1 \dots n + 1]$ , let  $U_i^+$  and  $U_i^-$  denote the sets

$$U_i^+ := \{x \in \mathbb{R}^{n+1} \mid x_i > 0\}$$

$$U_i^- := \{x \in \mathbb{R}^{n+1} \mid x_i < 0\},$$

which are open in  $\mathbb{R}^{n+1}$ .

Since  $\mathbb{S}^n$  doesn't contain the origin, every point on  $\mathbb{S}^n$  must have at least one non-zero coordinate, so these sets cover  $\mathbb{S}^{n+1}$ .

On these sets,  $\|x\| = 1$  has a unique solution, so we can find the intersection  $x \in \mathbb{S}^n \cap U_i^\pm$  by solving for  $x_i$ , which gives us

$$\begin{aligned} \sqrt{\sum_{j=1}^{n+1} x_j^2} &= 1 \\ \Leftrightarrow \sum_{j=1}^{n+1} x_j^2 &= 1 \\ \Leftrightarrow x_i^2 &= 1 - \sum_{j \neq i} x_j^2 \\ \Leftrightarrow x_i &= \begin{cases} \sqrt{1 - \sum_{j \neq i} x_j^2} & x \in U_i^+ \\ -\sqrt{1 - \sum_{j \neq i} x_j^2} & x \in U_i^- \end{cases} \end{aligned}$$

We can thus describe the intersections of  $\mathbb{S}^n$  with our sets  $U_i^\pm$  by taking every component except the  $i$ -th to be arbitrary, and setting the  $i$ -th component to the value taken by these continuous square root functions. Swapping the  $i$ -th and the  $n + 1$ -st components is continuous, so our intersections are the graphs of these functions. Since these intersections cover  $\mathbb{S}^n$ , and our functions are  $\mathbb{R}^n \rightarrow \mathbb{R}$ ,  $\mathbb{S}^n$  is a topological  $n$ -manifold.  $\square$

**Theorem 3.16.** Let  $p \in \mathbb{S}^n$ . Let  $H := \langle p \rangle^\perp$  be the hyperplane given as the orthog-

onal complement of the line through  $p$  and the origin. Then the *stereographic projection map*  $\sigma_p$  taking a point  $x \in \mathbb{S}^n \setminus \{p\}$  to the point where the line  $\vec{p}x$  intersects  $H$ , is a homeomorphism  $\mathbb{S}^n \setminus \{p\} \cong H(\cong \mathbb{R}^n)$

*Proof.* Let  $N := e_{n+1} = (0, \dots, n+1)^\top$  be the "north pole" of the sphere. Take a rotation  $R$  in the plane spanned by  $p$  and  $N$  mapping  $p$  to  $N$ . Since rotations in the origin preserve  $\mathbb{S}^n$ , we get

$$\sigma_p = R^{-1} \circ \sigma_N \circ R.$$

Since rotations are orthogonal maps, and thus homeomorphisms, it suffices to show that  $\sigma_N$  is a homeomorphism.

In this case, the hyperplane  $H$  orthogonal to  $\langle N \rangle$  is simply given by  $x_{n+1} = 0$ . Therefore, the intersection  $u := \sigma_N(x)$  of the line  $\vec{N}x = \{\lambda.(x - N) + N \mid \lambda \in \mathbb{R}\}$  can be found by finding the value of  $\lambda$  that makes the  $n+1$ st component zero. We get:

$$\begin{aligned} \lambda.(x_{n+1} - 1) + 1 &= 0 \\ \Leftrightarrow \lambda.(x_{n+1} - 1) &= -1 \\ \Leftrightarrow \lambda &= \frac{1}{1 - x_{n+1}} \end{aligned}$$

For  $i \neq n+1$ , we have  $N_i = 0$ , so  $u_i = \lambda.x_i$ . Substituting our  $\lambda$  into this equation gives

$$\sigma_N(x) = \frac{x_1, \dots, x_n, 0}{1 - x_{n+1}}$$

This map is continuous for  $x_{n+1} \neq 1$ , i.e.  $x \in \mathbb{S}^n \setminus \{N\}$ .

Solving our equations for  $x$  instead gives

$$\begin{aligned} x_i &= \frac{u_i}{\lambda}, \\ x_{n+1} &= \frac{\lambda - 1}{\lambda}, \end{aligned}$$

And since our preimage needs to lie on  $\mathbb{S}^n$ , we substitute into  $\|x\|^2 = 1$  and get

$$\begin{aligned} 1 &= \sum_{i=1}^{n+1} x_i^2 \\ &= \left( \sum_{i=1}^n \left( \frac{u_i}{\lambda} \right)^2 \right) + \left( \frac{\lambda - 1}{\lambda} \right)^2 \\ &= \frac{\sum_{i=1}^n u_i^2}{\lambda^2} + \frac{\lambda^2 - 2\lambda + 1}{\lambda^2} \\ &= \frac{\|u\|^2 + \lambda^2 - 2\lambda + 1}{\lambda^2} \\ \Leftrightarrow \lambda^2 &= \|u\|^2 + \lambda^2 - 2\lambda + 1 \\ \Leftrightarrow 2\lambda &= \|u\|^2 + 1 \\ \Leftrightarrow \lambda &= \frac{\|u\|^2 + 1}{2} \end{aligned}$$

Substituting back into our two equations gives

$$\sigma_N^{-1}(u_1, \dots, u_n) = \frac{(2u_1, \dots, 2u_n, \|u\|^2 - 1)^\top}{\|u\|^2 + 1},$$

which is continuous. Therefore,  $\sigma_N : \mathbb{S}^n \setminus \{N\} \cong H \cong \mathbb{R}^n$  is indeed a homeomorphism, implying that every  $\sigma_p$  is a homeomorphism.  $\square$

$\sigma_N$  and  $\sigma_S$  (where  $S = -N$  is the "south pole") give an alternate coordinate atlas on  $\mathbb{S}^n$ , which is sometimes easier to work with than our original atlas given by square root functions.

## Chapter 4

# Topologies on Algebraic Structures

### 4.1 Topological Groups

#### Definition 4.1

Let  $(G, \circ)$  be a group. Let  $\mathcal{O}$  be a topology on  $G$ . Then we call  $(G, \circ, \mathcal{O})$  a **topological group** iff. the maps  $^{-1} : g \rightarrow g^{-1}$  and  $\circ : G \times G \rightarrow G$  are continuous.

**Exercise 4.2.** Show that  $(G, \circ, \mathcal{O})$  is a topological group if and only if the map

$$\varphi : (g, h) \mapsto g \circ h^{-1}$$

is continuous.

**Exercise 4.3.** Let  $(G, \circ, \mathcal{O})$  be a topological group. Then for any  $g \in G$ , the left and right group operation maps  $l_g : h \mapsto g \circ h$  and  $r_g : h \mapsto h \circ g$  are homeomorphisms.

**Exercise 4.4.** Let  $(G, \circ, \mathcal{O}_G)$  be a topological group. Let  $(H, \circ, \mathcal{O}_H)$  be a normal subgroup equipped with the subspace topology, and let  $(G/H, \circ, \mathcal{O}_q)$  be the quotient group, equipped with the quotient topology. Then the quotient map  $p : G \rightarrow G/H$  is continuous and open.

**Corollary 4.5.** If  $H$  is closed in  $G$  and  $G$  is Hausdorff, then  $G/H$  is Hausdorff. If  $G/H$  is  $T_1$ , then  $H$  is closed.

## Chapter 5

# Connectedness

**Theorem 5.1.** Let  $\{C_i\}_{i \in I}$  be a collection of connected sets in a topological space  $X$  such that no two of the  $C_i$  are disjoint. Then  $\bigcup_{i \in I} C_i$  is connected.

**Theorem 5.2.** Let  $C$  be connected. Then every set  $D$  with  $C \subseteq D \subseteq \overline{C}$  is connected. In particular,  $\overline{C}$  is connected.

*Proof.* Assume that  $D$  isn't connected. Then we can write  $D = U \cup V$ , with  $U$  and  $V$  disjoint and open. Since  $C \subseteq D$ , we then have  $C = U' \cup V'$  for  $U' = U \cap C$  and  $V' = V \cap C$ . Both sets are open in  $C$  by definition of the subspace topology, and since every neighbourhood of a point in  $D \subseteq \overline{C}$  contains a point in  $C$ , neither of the two are empty. Therefore,  $C$  cannot be connected.  $\square$

**Corollary 5.3.** Connected components of a topological space are closed.

# Chapter 6

## Compactness

The familiar notions of open covers and compactness generalize directly from metric spaces to topological spaces. The most major difference is that sequential compactness is generally weaker than “covering compactness”, and has to be replaced with a notion of compactness based on ultrafilter convergence.

**Definition 6.1.** Let  $(X, \mathcal{O})$  be a topological space. Then we call a set  $\mathcal{U} \subseteq \mathcal{O}$  of open subsets an *open covering* of  $X$  if every point of  $X$  is contained in the union of all elements of  $\mathcal{U}$ , i.e. if

$$\bigcup_{U_i \in \mathcal{U}} U_i = X$$

### Definition 6.2: Compactness

Let  $(X, \mathcal{O})$  be a topological space. Then we call  $(X, \mathcal{O})$  *compact* if every open covering of  $X$  contains a finite open covering as a subset. We call a subset  $K \subseteq X$  compact if it is compact as a topological space equipped with the subspace topology.

**Exercise 6.3.** Every closed subset of a compact space is compact.

**Exercise 6.4.** Let  $(X, \mathcal{O})$  be compact and  $f : X \rightarrow Y$  be continuous. Then  $f[X]$  is compact.

**Exercise 6.5.** Let  $(X, \mathcal{O})$  be a nonempty topological space. Then the following are equivalent:

1.  $(X, \mathcal{O})$  is compact, i.e. every open covering has a finite subcovering.
2. Every set  $\mathcal{A}$  of closed sets whose intersection is the empty set has a finite subset whose intersection is the empty set.
3. If  $\mathcal{A}$  is a set of closed sets such that the intersection over any finite subset of  $\mathcal{A}$  is non-empty, the intersection of  $\mathcal{A}$  is non-empty.
4. Every filter on  $(X, \mathcal{O})$  has a limit point.
5. Every ultrafilter on  $(X, \mathcal{O})$  is convergent.

**Theorem 6.6. (Alexander Subbase Theorem)** Let  $(X, \mathcal{O})$  be a topological space. Let  $\mathcal{S}$  be a subbasis. Then  $X$  is compact if every open covering of  $X$  by sets of  $\mathcal{S}$  has a finite subcovering.

*Proof.* Assume that  $X$  is not compact. Then there exists a non-convergent ultrafilter  $\mathcal{F}$ , i.e. we have

$$\begin{aligned} \forall x \in X : \\ \exists U_x \in \mathcal{U}(x) : \\ (X \setminus U_x) \in \mathcal{F} \end{aligned}$$

By definition of subbases, every  $U_x \in \mathcal{U}(x)$  contains an open subset  $O_x$  of the form

$$O_x = \bigcap_{k=0}^n S_{k,x}$$

for some  $S_{k,x} \in \mathcal{S}$  and  $n \in \mathbb{N}$ . We therefore have

$$\begin{aligned} X \setminus U_x &\subseteq X \setminus O_x \\ &= X \setminus \bigcap_{k=0}^n S_{k,x} \end{aligned}$$

which implies  $X \setminus \bigcap_{k=0}^n S_{k,x} \in \mathcal{F}$  by the filter axioms. Since the intersection of this set with  $\bigcap_{k=0}^n S_{k,x} \in \mathcal{F}$  is empty, and filters are closed under finite intersection,  $\mathcal{F}$  cannot contain every  $S_{k,x}$ . Therefore, since  $\mathcal{F}$  is an ultrafilter, it must contain at least one set  $X \setminus S_{k,x}$  for every  $x$ . We will refer to these  $S_{k,x}$  with  $X \setminus S_{k,x} \in \mathcal{F}$  as  $S_x$ .

Since every  $S_x$  is an open neighborhood of  $x$ , and thus contains  $x$ , the set of all  $S_x$  is an open covering of  $X$ . By compactness, there must be a finite subcovering

$$\bigcup_{x \in X'} S_x = X.$$

Now, since  $\mathcal{F}$  contains every  $X \setminus S_x$ , it also contains

$$\bigcap_{x \in X'} (X \setminus S_x) = X \setminus \bigcup_{x \in X'} S_x = X \setminus X = \emptyset.$$

which contradicts  $\mathcal{F}$  being a filter.  $\square$

**Lemma 6.7.** Let  $(X, \mathcal{O})$  be a Hausdorff space. Let  $K \subseteq X$  be compact. Then  $K$  can be separated from any point  $x \in X \setminus K$  by open neighborhoods, i.e:

$$\begin{aligned} \forall x \in X : \\ \exists U_x \in \mathcal{U}(x), U_K \in \mathcal{U}(K) : \\ U_x \cap U_K = \emptyset \end{aligned}$$

**Exercise 6.8.** Let  $(X, \mathcal{O})$  be compact and Hausdorff. Let  $A, B \subseteq X$  be closed. Then  $A$  and  $B$  can be separated from each other by disjoint open neighborhoods.

**Exercise 6.9.** Every compact subset of a Hausdorff space is closed.

**Exercise 6.10.** Every continuous function from a compact space into a Hausdorff space is closed.

**Exercise 6.11.** *Every continuous injective function from a compact space into a Hausdorff space is an embedding.*

**Definition 6.12.** We call a topological space  $X$  *Lindelöf* if every open cover of  $X$  has a countable subcover.

**Theorem 6.13.** Every second-countable space is Lindelöf.

**Definition 6.14.** We call a topological space  $X$  *countably compact* if every countable open cover of  $X$  has a finite subcover.

**Exercise 6.15.** *A countably compact space is compact if and only if it is Lindelöf.*

**Theorem 6.16.** A metric space is Lindelöf if and only if it is second countable.

**Theorem 6.17.** Every sequentially compact space is countably compact.

**Theorem 6.18.** A first-countable space is countably compact if and only if it is sequentially compact.

**Theorem 6.19.** In a second countable Hausdorff space, compactness, countable compactness and sequential compactness are all equivalent to each other.

# Chapter 7

## Filters

### 7.1 Filter Convergence

It would be nice to find a generalization of sequence convergence to non-first-countable spaces that shares many of its useful properties.... It turns out that we can! The correct objects for the task are *filters*.

**Exercise 7.1.** Let  $(X, \mathcal{O})$  be a topological space. Then  $(X, \mathcal{O})$  is Hausdorff if and only if every convergent filter on  $X$  has a unique limit.

**Theorem 7.2.** Let  $X$  be a topological space. Let  $U \subseteq X$ . Then  $U$  is open iff.  $U$  is contained in every ultrafilter converging to some point in  $U$ .

*Proof.*  $\Rightarrow$ : By definition, if  $U$  is an open neighborhood of  $x$ , any filter converging to  $x$  must contain  $U$ .

$\Leftarrow$ : Let  $U \subseteq X$  and assume  $U \in \mathcal{U}$  for every ultrafilter converging to a point  $u \in U$ . Assume  $U$  is not open. Then there exists a point  $u \in U$  such that every open neighborhood of  $u$  contains at least one point in  $X \setminus U$ . But then any two members of the family

$$\{X \setminus U\} \cup \{U(u)\}$$

have nonempty intersection. Therefore, this family can be extended to an Ultrafilter  $\mathcal{U}_u$ . Since  $\mathcal{U}_u$  contains all open neighborhoods of  $u$  by construction, it converges to  $u$ , and must thus contain  $U$ . Since it also contains  $X \setminus U$ , we arrive at a contradiction.

□

### 7.2 Tychonoff's Theorem

**Definition 7.3.** Let  $\mathcal{F}$  be a filter on  $X$ . Let  $f : X \rightarrow Y$ . Then the *pushforward Filter of  $f$  on  $\mathcal{F}$*  is the set

$$f_*(\mathcal{F}) := \{A \subseteq Y \mid p_i^{-1}[A] \in \mathcal{F}\},$$

**Exercise 7.4.** Verify that the pushforward of a filter is a filter, and that the pushforward of an ultrafilter is an ultrafilter.

Intuitively, the pushforward of a filter under a function generalizes the sequence  $f(x_n)$  for some sequence  $x_n$  and function  $f$ . By extension, the following theorem generalizes the fact that in metric spaces, continuity and sequential continuity are equivalent:

**Theorem 7.5: Characterization of continuity in terms of ultrafilters**

Let  $f : X \rightarrow Y$ . Then  $f$  is continuous if and only if  $f_*(\mathcal{U})$  converges to  $f(x)$  for every ultrafilter  $\mathcal{U}$  that converges to  $x \in X$ .

*Proof.* 1. Assume  $f$  is continuous. Let  $\mathcal{U}$  be an ultrafilter on  $X$  converging to  $x \in X$ .

Let  $U \subseteq Y$  be an open neighbourhood of  $f(x)$ . Then, since  $f$  is continuous,  $f^{-1}(U)$  is an open neighbourhood of  $x$ . We thus have  $f^{-1}(U) \in \mathcal{U}$ , which gives us  $U \in f_*(\mathcal{U})$  by the definition of the pushforward filter.

Thus, every open neighbourhood of  $f(x)$  is contained in  $f_*(\mathcal{U})$ , so  $f_*(\mathcal{U})$  converges to  $f(x)$ .

2. Assume that the pushforward of any ultrafilter converging to  $x$  converges to  $f(x)$ . Let  $U \subseteq Y$  be open.

Let  $\mathcal{U}$  be an ultrafilter on  $X$  converging to some point  $x \in f^{-1}(U)$ . Then, by our hypothesis,  $f_*(\mathcal{U})$  converges to  $f(x)$ . Since  $U$  is an open neighborhood of  $f(x)$ , we have  $U \in f_*(\mathcal{U})$ , and thus  $f^{-1}(U) \in \mathcal{U}$ . Thus, by theorem 7.2,  $f^{-1}(U)$  is open, so  $f$  is continuous.  $\square$

This theorem generalizes the fact that a sequence in a metric product space converges if and only if every individual component converges.

**Theorem 7.6.** Let  $(X_i, \mathcal{O}_i)_{i \in I}$  be a family of topological spaces. Let  $\mathcal{F}$  be a filter on  $\prod_{i \in I} X_i$ . Then  $\mathcal{F}$  converges to a point  $x \in \prod_{i \in I} X_i$  if and only if each pushforward filter  $(p_i)_*(\mathcal{F})$  converges to  $p_i(x) = x_i$ .

*Proof.*  $\Rightarrow$ : Assume that  $\mathcal{F}$  converges to  $x \in \prod_{i \in I} X_i$ . Since all projections  $p_i$  are continuous, each  $(p_i)_*(\mathcal{F})$  converges to  $p_i(x) = x_i$  by theorem 7.5.

$\Leftarrow$ : Assume that each  $(p_i)_*(\mathcal{F})$  converges to  $p_i(x) = x_i$ . By definition of the product topology, the sets  $S_i := p_i^{-1}(U_i)$  for open  $U_i \in \mathcal{O}_i$  form a subbase of  $\prod_{i \in I} X_i$ . Therefore, any open neighborhood  $U \in \mathcal{U}(x)$  contains an open neighborhood  $B = \bigcap_{i \in J} p_i^{-1}(U_i)$  for some open neighborhoods  $U_i \in \mathcal{O}_i$ .

Since each  $(p_i)_*(\mathcal{F})$  converges to  $x_i$ , we have  $U_i \in (p_i)_*(\mathcal{F})$  for each  $i$ . Therefore, by definition of the pushforward filter, we have  $p_i^{-1}(U_i) \in \mathcal{F}$  for all  $i$ . Since filters are closed under intersection, we get  $B \in \mathcal{F}$ ,

and since  $B \subseteq U$  we also get  $U \in \mathcal{F}$  as desired.

□

### Theorem 7.7: Tychonoff's Theorem

Let  $(X_i, \mathcal{O}_i)_{i \in I}$  be a family of topological spaces. Then if all  $X_i$  are compact,  $\prod_{i \in I} X_i$  is compact.

*Proof.* 1. If there exists an  $i \in I$  such that  $X_i = \emptyset$ , then  $\prod X_i = \emptyset$  is compact.

2. Now, assume  $\prod_{i \in I} X_i \neq \emptyset$ . Then there exists an ultrafilter  $\mathcal{U}$  on  $\prod_{i \in I} X_i$ .

Since each  $X_i$  is compact, each pushforward  $(p_i)_*(\mathcal{U})$  converges to an  $x_i \in X_i$ . Thus, by theorem 7.6,  $\mathcal{U}$  converges to  $(x_i)_{i \in I}$  and  $\prod_{i \in I} X_i$  is compact.

□

## Chapter 8

# Higher Separation Properties

### 8.1 Regular Spaces

#### Definition 8.1: Regular Space

Let  $(X, \mathcal{O})$  be a topological space. Then we call  $(X, \mathcal{O})$  **regular** if every point can be housed orf from any closed set not containing it:

$$\forall (X \setminus C) \in \mathcal{O} :$$

$$\forall x \notin C :$$

$$\exists U_x \in \mathcal{U}(x), U_C \in \mathcal{U}(C) :$$

$$U_x \cap U_C = \emptyset$$

We call  $(X, \mathcal{O})$   $T_3$  or **regular Hausdorff**, if it is regular and Hausdorff.

**Exercise 8.2.** Find a topological space that is regular, but not  $T_0$ .

**Exercise 8.3.** Show that a topological space is regular Hausdorff ( $T_3$ ) if and only if it is regular and  $T_1$ .

#### Definition 8.4: Completely Regular Space

Let  $(X, \mathcal{O})$  be a topological space. Then we call  $(X, \mathcal{O})$  **completely regular** if every point can be housed orf from any closed set not containing it by a continuous function:

$$\forall (X \setminus C) \in \mathcal{O} :$$

$$\forall x \notin C :$$

$$\exists f \in C(X, [0, 1]_{\mathbb{R}}) :$$

$$f(x) = 1, f[A] \subseteq \{0\}$$

We call  $(X, \mathcal{O})$   $T_{3a}$ ,  $T_{3\frac{1}{2}}$ , or **completely regular Hausdorff**, if it is completely regular and Hausdorff.

**Exercise 8.5.** Show that every completely regular space is regular.

**Exercise 8.6.** Find a space that is completely regular, but not  $T_0$ .

**Exercise 8.7.** Show that a space is completely regular Hausdorff if and only if it is completely regular and  $T_1$ .

**Theorem 8.8.** Every zero-dimensional space  $(X, \mathcal{O})$  is completely regular.

*Proof.* Since  $X$  is zero-dimensional, it has a basis consisting of clopen sets. Given any clopen set  $U$ , the function

$$f_U : x \mapsto \begin{cases} 1 & x \in U \\ 0 & x \in X \setminus U \end{cases}$$

is continuous. Now, let  $C$  be a closed set not containing  $x$ . Then  $X \setminus C$  is an open neighbourhood of  $x$ . We can now choose a basic (i.e. clopen) neighbourhood  $U \subseteq X \setminus C$  of  $x$ , at which point  $f_U$  is our desired function separating  $x$  and  $C$ .  $\square$

**Theorem 8.9.** Let  $(X, \mathcal{O})$  be completely regular. Then the set

$$\{f^{-1}[U] \mid f \in C(X, [0, 1]), U \in \mathcal{O}_{\mathbb{R}}\}$$

is a basis of  $X$ .

*Proof.* 1. Since the  $f$  are continuous, every basis set is open.  
2. Let  $x \in X$ . Let  $U$  be an open neighbourhood of  $x$ . Since  $X$  is completely regular, there exists a continuous function  $f$  separating  $\{x\}$  and  $X \setminus U$ . Then the basic set  $f^{-1}[(0, 1]]$  must be a subset of  $U$ , since every element outside of  $U$  gets mapped to 0. Thus, every open neighbourhood contains a basic neighbourhood.  $\square$

**Theorem 8.10.** Every completely regular Hausdorff space  $(X, \mathcal{O})$  can be embedded into the cube

$$\prod_{f \in C(X, [0, 1])} [0, 1]$$

*Proof.* We choose the trivial embedding  $\iota : x \mapsto (f(x))_{f \in C(X, [0, 1])}$ .

1.  $\iota$  is continuous, since the projections of  $\iota$  are exactly the elements of  $C(X, [0, 1])$ , i.e. continuous.
2. Let  $x \neq y$ . By  $T_1$ , all singletons are closed. Therefore, by complete regularity, there exists a function  $f$  with  $f(x) = 1$  and  $f(y) = 0$ . Therefore, we have  $\iota(x)_f \neq \iota(y)_f$ , i.e.  $\iota(x) \neq \iota(y)$ .
3. By 8.9, we only need to show openness for the basis sets

$$\{f^{-1}[U] \mid f \in C((X, \mathcal{O}), [0, 1]_{\mathbb{R}}), U \in \mathcal{O}_{\mathbb{R}}\}.$$

For these sets, we get:

$$\begin{aligned}
 \iota[f^{-1}[U]] &= \iota[(p_f \circ \iota)^{-1}[U]] \\
 &= \iota[\iota^{-1}(p_f^{-1}[U])] \\
 &= \iota \left[ \iota^{-1} \left[ U \times \prod_{g \neq f} [0, 1] \right] \right] \\
 &= \left( U \times \prod_{g \neq f} [0, 1] \right) \cap \iota[X]
 \end{aligned}$$

Where  $\left( U \times \prod_{g \neq f} [0, 1] \right)$  is open, since every component besides the  $f$ -component is open.

□

## 8.2 Normal Spaces

### Definition 8.11: Normal Space

Let  $(X, \mathcal{O})$  be a topological space. Then we call  $(X, \mathcal{O})$  **normal** if every pair of disjoint closed sets can be housed orf from each other by disjoint neighbourhoods:

$$\begin{aligned} \forall X \setminus C_1, X \setminus C_2 \in \mathcal{O} : \\ C_1 \cap C_2 = \emptyset \implies \\ \exists U_1 \in \mathcal{U}(C_1), U_2 \in \mathcal{U}(C_2) : \\ U_1 \cap U_2 = \emptyset \end{aligned}$$

We call  $(X, \mathcal{O})$   **$T_4$** , or **normal Hausdorff**, if it is normal and Hausdorff.

**Exercise 8.12.** Find a topological space that is normal, but not  $T_0$ .

### Lemma 8.13: Urysohn's Lemma

A topological space is normal if and only if any two disjoint closed subsets can be separated by a continuous function.

**Corollary 8.14.** Every normal Hausdorff space is completely regular Hausdorff.

**Theorem 8.15.** A second-countable regular topological space is normal.

*Proof.* Let  $A, B$  be disjoint closed sets in a second-countable regular topological space  $X$ . Let  $\{U_i\}_{i \in I}$  be the collection of open sets such that  $U \cap A \neq \emptyset$  and  $\overline{U} \cap B = \emptyset$ . Since  $X$  is regular, this collection forms an open covering of  $A$ , and  $X$  is second-countable, we can take a countable subcovering  $\{U_n\}_{n \in \mathbb{N}}$ .

Analogously, we can construct a countable open covering  $\{V_n\}_{n \in \mathbb{N}}$  of  $B$  fulfilling  $\overline{V_n} \cap A = \emptyset$ .

Now, in order to make these two coverings disjoint from each other as well, we take

$$\begin{aligned} Y_n &:= U_n \cap \bigcap_{k=1}^n X \setminus \overline{V_k} \\ Z_n &:= V_n \cap \bigcap_{k=1}^n X \setminus \overline{U_k} \end{aligned}$$

By construction of our  $U_n$  and  $V_n$ , our  $Y_n$  form an open cover of  $A$  and our  $Z_n$  form an open cover of  $B$ . Furthermore, we have  $Y_m \cap Z_n = \emptyset$  for all  $m, n \in \mathbb{N}$ , since  $m \geq n$  implies

$$\begin{aligned} Y_m \cap Z_n &\supseteq Y_m \cap V_n \\ &\supseteq Y_m \cap \overline{V_n} \\ &\supseteq U_m \cap (X \setminus \overline{V_n}) \cap \overline{V_n} \\ &= \emptyset, \end{aligned}$$

and  $n \geq m$  implies the same, but with  $Y, Z$  and  $U, V$  swapped.  
Therefore,  $\bigcup_{m \in \mathbb{N}} Y_m$  and  $\bigcup_{n \in \mathbb{N}} Z_n$  are disjoint open sets containing  $A$  and  $B$ , completing the proof that  $X$  is normal.  $\square$

## 8.3 Urysohn's Metrization Theorem

### Corollary 8.16: Urysohn's Metrization Theorem

Every second-countable completely regular space Hausdorff space  $X$  can be embedded into the Hilbert Cube

$$\prod_{n \in \mathbb{N}} [0, 1]$$

*Proof.* Since  $X$  is second-countable completely regular Hausdorff,  $X$  is normal Hausdorff by Theorem 8.15. Now, we define a set

$$S = \{(U, V) \in \mathcal{B}^2 \mid \overline{U} \subseteq V\}.$$

Urysohn's Lemma 8.13 tells us that for each  $s = (U, V) \in S$ , there exists a continuous function  $g_s : X \rightarrow [0, 1]$  such that  $g_s[\overline{U}] = 1$  and  $g_s[X \setminus V] = 0$ . This lets us find a canonical continuous map  $X \rightarrow \prod_{s \in S} [0, 1]$ :

$$\begin{aligned} g : X &\rightarrow \prod_{s \in S} [0, 1] \\ x &\mapsto (s \mapsto g_s(x)) \end{aligned}$$

This map is injective, since for  $x \neq y$ , we can use complete regularity to find a pair  $s = (U, V) \in S$  with  $x \in U$  and  $y \notin V$  such that  $g_s(x) = 1$  and  $g_s(y) \in g_s[X \setminus V] = 0$ , i.e.  $g(x) \neq g(y)$ .

It remains to be shown that  $g$  is open. If  $W$  is an open neighbourhood of  $x \in X$ , we want the set  $g(W)$  to be open in  $\prod_{s \in S} [0, 1]$ , which follows if we can find for each  $g(y) \in g(W)$  a subbasis neighbourhood  $\pi^{-1}(S)$  of the product topology such that

$$g(y) \in g(X) \cap \pi^{-1}(S) \subseteq g(W)$$

Now, as earlier, we find basic neighbourhoods  $V \subseteq W \in \mathcal{B}$  and  $s = (U, V) \in S$  such that  $g_s(x) \in g_s[\overline{U}] = 1$  and  $g_s[X \setminus V] = 0$ , i.e.

$$g_s(y) \neq 0 \implies y \in V,$$

i.e.  $g_s^{-1}[(0, 1]] \subseteq V$ . Since we have  $g_s = \pi_s \circ g$ , we get  $g_s^{-1} = g^{-1} \circ \pi_s^{-1}$ . Fiddling around with images and preimages for a bit now gives us:

$$\begin{aligned} g_s^{-1}[(0, 1]] \cap X &= g^{-1}(\pi_s^{-1}[(0, 1]]) \cap X \subseteq V \\ &\implies g(g^{-1}(\pi_s^{-1}[(0, 1]]) \cap X) \subseteq g[V] \\ &\stackrel{g \text{ inj.}}{\implies} g(g^{-1}(\pi_s^{-1}[(0, 1]])) \cap g[X] \subseteq g[V] \\ &\implies \pi_s^{-1}[(0, 1]] \cap g[X] \subseteq g[V] \end{aligned}$$

□

## Chapter 9

# Lattice Theory

### 9.1 Lattices

**Definition 9.1.** A partially ordered set  $(L, \leq)$  is called a *lattice* if every pair of elements  $x, y \in L$  has a least upper bound  $x \vee y$ , also known as a *join*, and a greatest lower bound  $x \wedge y$ , also known as a *meet*. A *complete lattice* is a lattice in which every subset  $A \subseteq L$  has a join  $\sup A$  and a meet  $\inf A$ .

**Definition 9.2.** Given a complete lattice  $L$ , the *bottom element* is the element  $0 := \inf L$ , and the *top element* is the element  $\infty := \sup L$ .

**Definition 9.3.** Let  $L$  be a complete lattice. Let  $x, y \in L$ . Then we say that  $x$  is *well above*  $y$ , denoted  $x > y$ , iff for any subset  $A \subseteq L$ ,

$$y \geq \inf A \implies \exists a \in A : x \geq a$$

**Lemma 9.4.** Let  $L$  be a complete lattice. Let  $x, y, z \in L$ . Then:

1.  $x > y$  implies  $x \geq y$ ,
2.  $x > y$  and  $y \geq z$  implies  $x \geq z$ ,
3.  $x \geq y$  and  $y > z$  implies  $x > z$ .

*Proof.* 1. We have  $y \geq \inf \{y\}$ , so  $x > y$  implies  $x \geq y$ .  
2. Follows from applying transitivity to the first part of the lemma.  
3.  $z \geq \inf A \implies \exists a \in A : x \geq y \geq a$ .

□

**Lemma 9.5.** Let  $L$  be a complete lattice. Let  $x \in L$  and  $A \subseteq L$ . Then

$$x > \inf A \iff \exists a \in A : x > a$$

## **9.2 Value Quantaes**

## **9.3 Generalized metrics metrize every topological space**

## **Part IX**

# **Algebraic Topology**

# Chapter 1

## Homotopy

### Definition 1.1

Let  $X, Y$  be topological spaces. Let  $f, g : X \rightarrow Y$  be continuous. Then  $f$  is called **homotopic** to  $g$  via a map  $h$  iff.  $h$  is a "continuous interpolation between  $f$  and  $g$ , i.e:

1.  $h$  is a continuous map  $X \times [0, 1] \rightarrow Y$ ,
2. For all  $x \in X$ , we have  $h(x, 0) = f(x)$ ,
3. For all  $x \in X$ , we have  $h(x, 1) = g(x)$ .

We call  $h$  a **free homotopy**, or simply a **homotopy**, between  $f$  and  $g$ . We write  $f \sim_h g$  if  $f$  is a homotopy between  $f$  and  $g$ . We write  $f \sim g$  if there exists a homotopy between  $f$  and  $g$ .

**Exercise 1.2.** Show that the property of being homotopic is an equivalence relation.

**Definition 1.3.** We say that  $f$  and  $g$  are **homotopic relative to a subset  $A \subseteq X$**  and write  $f \sim_{\text{rel } A} g$  or  $f \sim_{h, \text{rel } A} g$  iff:

1.  $f \sim_h g$ ,
2. For all  $t \in [0, 1]$ , we have  $h(t, \cdot)|_A = f|_A = g|_A$

I personally find the designation "homotopic relative to  $A$ " quite unintuitive, given that the property of being homotopic relative to a given subset is stronger than the property of being homotopic.

**Exercise 1.4.** Show that  $f \sim_h g$  is equivalent to  $f \sim_{h, \text{rel } \emptyset} g$ .

**Exercise 1.5.** Let  $f_0, f_1 : X \rightarrow Y$  and  $g_0, g_1 : Y \rightarrow Z$  be continuous. Let  $h_0$  be a homotopy between  $f_0$  and  $f_1$  and let  $h_1$  be a homotopy between  $f_1$  and  $g_1$ . Then  $H : (x, t) \mapsto h_1(h_0(x, t), t)$  is a homotopy between  $g_0 \circ f_0$  and  $g_1 \circ f_1$ .

### Definition 1.6

Let  $f : X \rightarrow Y$  be continuous. Then  $f$  is called a **homotopy equivalence** between  $X$  and  $Y$  iff. there exists a map  $g : Y \rightarrow X$  such that  $g \circ f$  is homotopic to  $\text{id}_X$  and  $f \circ g$  is homotopic to  $\text{id}_Y$ . We write  $X \simeq_{f, g} Y$ ,  $X \simeq_f Y$ , or simply  $X \simeq Y$ .

**Exercise 1.7.** Show that the property of being homotopy equivalent is an equivalence relation.

**Exercise 1.8.** Show that homeomorphism implies homotopy equivalence.

**Definition 1.9.** A topological space  $X$  is called *contractible* iff.  $X$  is homotopy equivalent to a one-point space.

**Exercise 1.10.** 1. Show that

$$h : (x, t) \mapsto x - tx$$

is a homotopy between  $\text{id}_{\mathbb{R}^n}$  and  $f : \mathbb{R}^n \mapsto \{0\}$ .

2. Conclude that  $\mathbb{R}^n$  is contractible.

3. Conclude that homotopy equivalence does not imply homeomorphism.

**Exercise 1.11.** A topological space  $X$  is contractible if and only if there exists a point  $x_0 \in X$  such that  $\text{id}_X$  is homotopic to the constant map  $X \rightarrow x_0$ .

**Definition 1.12.** Let  $X$  be a subset of a topological vector space. Then we call  $X$  *star-shaped* iff. there exists a point  $x_0 \in X$  such that  $X$  contains the connecting line between  $x_0$  and any other point  $x \in X$ , i.e.

$$\begin{aligned} \forall x \in X : \\ \forall t \in [0, 1] : \\ (1 - t)x_0 + tx \in X \end{aligned}$$

**Exercise 1.13.** Show that any star-shaped set is contractible. Conclude that any convex set is contractible.

**Lemma 1.14.** Let  $X$  be a topological space. Let  $A \subseteq X$ . Then  $A$  and  $X$  are homotopy equivalent if there exists a homotopy between  $\text{id}_X$  and  $\text{id}_X|_A$ , i.e. a continuous map  $h : X \times [0, 1] \rightarrow X$  such that for all  $x \in X$ :

1.  $h(x, 0) = x$ ,
2.  $h(x, 1) \in A$ ,
3.  $x \in A \implies h(x, 1) = x$

## 1.1 The Fundamental Group

**Definition 1.15.** Let  $X$  be a topological space. Then a continuous map  $\gamma : [0, 1] \rightarrow X$  with  $\gamma(0) = \gamma(1)$  is known as a *loop*. We denote the set of all loops in  $X$  with  $\gamma(0) = \gamma(1) = x_0$  as  $\Omega(X, x_0)$ .

**Lemma 1.16.** Let  $\gamma : [0, 1] \rightarrow X$  be a loop. Let  $\varphi : [0, 1] \rightarrow [0, 1]$  be a continuous map such that  $\varphi(0) = 0$  and  $\varphi(1) = 1$ . Then  $\gamma \circ \varphi$  and  $\gamma$  are homotopic relative to  $\{0, 1\}$ .

*Proof.* The map

$$h : (x, t) \mapsto \gamma((1-t)x + t\varphi(x))$$

is a homotopy between  $\gamma$  and  $\gamma \circ \varphi$ .  $\square$

**Definition 1.17.** Let  $f, g \in \Omega(X, x_0)$ . Then we define the *concatenation of  $f$  and  $g$*  as the map

$$(f * g)(t) = \begin{cases} f(2t) & 0 \leq t \leq 1/2, \\ g(2t - 1) & 1/2 \leq t \leq 1. \end{cases}$$

**Definition 1.18.** Let  $\gamma \in \Omega(X, x_0)$ . We define the *inverse loop* of  $\gamma$  to be the loop that “moves through gamma backwards”, i.e.

$$\bar{\gamma}(t) := \gamma(1 - t)$$

**Exercise 1.19.** Verify that  $\bar{\bar{\gamma}} = \gamma$ .

**Exercise 1.20.** Let  $\gamma \in \Omega(X, x_0)$ . Then  $\gamma * \bar{\gamma}$  is homotopic to  $e_{x_0} : X \rightarrow \{x_0\}$  relative to  $\{0, 1\}$ .

### Theorem 1.21: Existence of the fundamental group

Let  $X$  be a topological space. Let  $x_0 \in X$ . Then the quotient space

$$\pi_1(X, x_0) = \Omega(X, x_0) / \sim_{\text{rel } (0,1)}$$

of loops modulo homotopy becomes a group, known as the *fundamental group of  $(X, x_0)$* , when equipped path concatenation

$$[f] * [g] = [f * g],$$

as the group operation. The neutral element of this group is  $e_{x_0} := (x \mapsto x_0)$ .

**Theorem 1.22.** If we additionally define

$$\begin{aligned} \pi_1(f) : \pi_1(X, x) &\mapsto \pi_1(Y, f(x)) \\ [\gamma] &\mapsto [f \circ \gamma] \end{aligned}$$

for all continuous  $f : X \rightarrow Y$ , the map  $\pi_1$  becomes a covariant functor from the category of topological spaces into the category of groups.

**Definition 1.23.** Whenever we need to make the base point clear, we write  $\pi_1(f, x) := \pi_1(f) : \pi_1(X, x) \mapsto \pi_1(Y, f(x))$

**Theorem 1.24: Fundamental groups at different base points**

Let  $X$  be a topological space. Let  $x, y \in X$ . Then every continuous path  $\delta : [0, 1] \rightarrow X$  induces a group isomorphism

$$J_\delta : \pi_1(X, \delta(0)) \cong \pi_1(X, \delta(1))$$

$$[\gamma] \mapsto [\bar{\delta} * \gamma * \delta]$$

*Proof.* 1. Since loop concatenation preserves homotopy,  $J_\delta$  sends different representatives of the same equivalence class to the same equivalence class.

2.  $J_\delta$  goes from  $\pi_1(X, x)$  to  $\pi_1(X, y)$ , since for any  $\gamma : [0, 1] \rightarrow X$  with  $\gamma(0) = \gamma(1) = x$ ,  $\bar{\delta}$  traces out a path from  $\delta(1)$  to  $\delta(0)$ , then  $\gamma$  traces out a loop from  $\delta(0)$  to  $\delta(0)$ , and then  $\delta$  traces out a path from  $\delta(0)$  back to  $\delta(1)$ , i.e.  $\bar{\delta} * \gamma * \delta$  is a loop from  $\delta(1)$  to  $\delta(1)$ .

3.  $J_\delta$  is a group homomorphism, since

$$\begin{aligned} J_\delta([\gamma_1 * \gamma_2]) &= [\bar{\delta} * \gamma_1 * \gamma_2 * \delta] \\ &= [\bar{\delta} * \gamma_1 * e_{x_0} * \gamma_2 * \delta] \\ &= [\bar{\delta} * \gamma_1 * \delta * \bar{\delta} * \gamma_2 * \delta] \\ &= [\bar{\delta} * \gamma_1 * \delta] * [\bar{\delta} * \gamma_2 * \delta] \\ &= J_\delta([\gamma_1]) * J_\delta([\gamma_2]) \end{aligned}$$

(Recall that any map that respects the group operation also respects the neutral element and inverses, and is thus a homomorphism.)

4.  $J_\delta$  is bijective with  $J_\delta^{-1} = J_{\bar{\delta}}$ :

$$\begin{aligned} J_\delta(J_{\bar{\delta}}(\gamma)) &= [\bar{\delta} * (\delta * \gamma * \bar{\delta}) * \delta] = [\gamma] \\ J_{\bar{\delta}}(J_\delta(\gamma)) &= [\delta * (\bar{\delta} * \gamma * \delta) * \bar{\delta}] = [\gamma] \end{aligned}$$

□

**Corollary 1.25.** If  $X$  is path connected,  $\pi_1(X, x) \cong \pi_1(X, y)$  for any  $x, y \in X$ .

**Definition 1.26.** Let  $\gamma_0 \in \Omega(X, x)$  and  $\gamma_1 \in \Omega(X, y)$ . Then we call  $h : [0, 1] \times [0, 1] \rightarrow X$  a **free homotopy of loops**, or simply **loop homotopy**, between  $\gamma_0$  and  $\gamma_1$  iff.  $h$  is a homotopy between  $\gamma_0$  and  $\gamma_1$  such that  $h_t(x) := h(x, t)$  is a loop for any fixed  $t \in [0, 1]$ .

**Exercise 1.27.** 1. Verify that  $h$  is a loop homotopy iff.  $h$  is a homotopy and  $h_t(0) = h_t(1)$  for any  $t \in [0, 1]$ .

2. Show that any  $\{0, 1\}$ -homotopy is a loop homotopy.

The additional constraint that  $h(\cdot, t)$  is a loop may seem superfluous, but there are in fact homotopies between loops that aren't loop homotopies:

- Exercise 1.28.**
1. Let  $\gamma_0 = \gamma_1 : [0, 1] \rightarrow \{0\}$  be constant loops. Let  $h(x, t) := xt(1-t)$ . Then  $h$  is a homotopy between  $\gamma_0$  and  $\gamma_1$ , but fails to be a loop homotopy.
  2. Let  $\gamma_0(x) = e^{2\pi ix}$ . Let  $\gamma_1(x) = e^{4\pi ix}$ . Then  $h(x, t) := e^{2\pi i(1+t)x}$  is a homotopy between  $\gamma_0$  and  $\gamma_1$ , but fails to be a loop homotopy.
  3. Assume that any loop  $\gamma : [0, 1] \rightarrow S^1$  with 1 as its base point is homotopic a loop of the form  $\gamma(x) := e^{2\pi i n x}$  for some  $n \in \mathbb{Z}$  (we will prove this later). Does 2. imply that the fundamental group  $\pi_1(S^1, 1)$  is trivial?

**Definition 1.29.** A loop  $\gamma \in \Omega(X, x)$  is called **contractible in  $X$**  iff. there exists an  $x \in X$  and a loop homotopy  $h$  such that  $\gamma \sim_h e_x$ .

**Lemma 1.30.**  $\gamma \in \Omega(X, x)$  is contractible in  $X$  if  $\gamma \sim_{\text{rel } 0,1} e_{\gamma(0)}$ .

**Theorem 1.31.** Let  $\gamma \in \Omega(X, x)$  be a loop. Let  $S^1 \cong I/\{0, 1\}$  be equipped with the quotient topology. Then  $\gamma$  is contractible if and only if  $\gamma \circ p : S^1 \rightarrow X$  has a continuous extension to the closed unit ball  $\bar{B}^2$ .

**Theorem 1.32.** Let  $h$  be a homotopy between  $f, g : X \rightarrow Y$ . Let  $x_1 \in X$ . Then

$$\begin{aligned} h(x_1, \cdot) : [0, 1] &\rightarrow Y \\ t &\mapsto h(x_1, t) \end{aligned}$$

is a path from  $f(x_1)$  to  $g(x_1)$  which fulfills

$$\pi_1(g, x_1) = J_{h(x_1, \cdot)} \circ \pi_1(f, x_1)$$

Recall that  $\pi_1(g, x_1)$  is the map  $[\gamma] \mapsto [g \circ \gamma]$ , and that  $J_\delta$  is the map  $[\gamma] \mapsto [\bar{\delta} * \gamma * \delta]$ , so this theorem states that

$$[g \circ \gamma] = \left[ \bar{h}(x_1, \cdot) * (f \circ \gamma) * h(x_1, \cdot) \right]$$

for any  $\gamma \in \pi_1(X, x_1)$ , or, equivalently, that the diagram

$$\begin{array}{ccc} & \pi_1(X, x_1) & \\ \swarrow \pi_1(f, x_1) & & \searrow \pi_1(g, x_1) \\ \pi_1(Y, f(x_1)) & \xrightarrow{J_{h(x_1, \cdot)}} & \pi_1(Y, g(x_1)) \end{array}$$

commutes.

*Proof.* Define

$$H_t(x) := h(x_1, t + (1-t)x).$$

We have  $(f \circ \gamma) \in \Omega(X, f(x_1))$  and  $(g \circ \gamma) \in \Omega(X, g(x_1))$ , since

$$f(\gamma(0)) = f(x_1) = f(\gamma(1))$$

$$g(\gamma(0)) = g(x_1) = g(\gamma(1))$$

Then a homotopy between  $g \circ \gamma$  and  $\bar{h}(x_1, \cdot) * (f \circ \gamma) * h(x_1, \cdot)$  is given by:

$$k(x, t) := (\bar{H}_t * (h(\cdot, t) \circ \gamma) * H_t)(x),$$

since we get:

$$\begin{aligned} k(x, 0) &= (\bar{H}_0 * (h(\cdot, 0) \circ \gamma) * H_0)(x) \\ &= (\bar{H}_0 * (f \circ \gamma) * H_0)(x) \\ &= (\bar{h}(x_1, \cdot) * (f \circ \gamma) * h(x_1, \cdot))(x) \end{aligned}$$

and

$$\begin{aligned} k(x, 1) &= (\bar{H}_1 * (h(\cdot, 1) \circ \gamma) * H_1)(x) \\ &= (h(x_1, (1-1) + (1-(1-1))(\cdot)) * (g \circ \gamma) * h(x_1, 1 + (1-1)(\cdot)))(x) \\ &= (h(x_1, \cdot) * (g \circ \gamma) * h(x_1, 1))(x) \\ &= (e_{g(x_1)} * (g \circ \gamma) * e_{g(x_1)})(x) \\ &= (g \circ \gamma)(x), \end{aligned}$$

where the second to last step follows since the loop  $(g \circ \gamma)$  ends on the base point  $g(x_1)$ .  $\square$

**Lemma 1.33.** Let  $\gamma : [0, 1] \rightarrow X$  be continuous. Let  $f : X \rightarrow Y$ . Then

$$\pi_1(f, \gamma(1)) \circ J_\gamma = J_{f \circ \gamma} \circ \pi_1(f, \gamma(0)),$$

i.e. the diagram

$$\begin{array}{ccc} \pi_1(X, \gamma(0)) & \xrightarrow{\pi_1(f, \gamma(0))} & \pi_1(Y, f(\gamma(0))) \\ \downarrow J_\gamma & & \downarrow J_{f \circ \gamma} \\ \pi_1(X, \gamma(1)) & \xrightarrow{\pi_1(f, \gamma(1))} & \pi_1(Y, f(\gamma(1))) \end{array}$$

commutes.

*Proof.* Let  $\sigma \in \Omega(X, \gamma(0))$ . Then we have

$$\begin{aligned} \pi_1(f, \gamma(1)) \circ J_\gamma([\sigma]) &= \pi_1(f, \gamma(1))([\bar{\gamma} * \sigma * \gamma]) \\ &= [f \circ (\bar{\gamma} * \sigma * \gamma)] \\ &= [(f \circ \bar{\gamma}) * (f \circ \sigma) * (f \circ \gamma)] \\ &= [\overline{(f \circ \gamma)} * (f \circ \sigma) * (f \circ \gamma)] \\ &= J_{f \circ \gamma}([f \circ \sigma]) \\ &= J_{f \circ \gamma} \circ \pi_1(f, \gamma(0))([\sigma]) \end{aligned}$$

$\square$

**Theorem 1.34.** Let  $f : X \rightarrow Y$  be a homotopy equivalence. Let  $x_0 \in X$ . Then the map  $\pi_1(f, x_0)$  is a group isomorphism  $\pi_1(X, x_0) \cong \pi_1(Y, f(x_0))$ .

## Chapter 2

# Fibrations and Coverings

### Definition 2.1: Locally Trivial Fibration

Let  $\pi : Y \rightarrow X$  be continuous. Then  $\pi$  is called a *locally trivial fibration* iff. for every  $x \in X$ , there exists an open neighborhood  $U$  such that there exists a homeomorphism  $h_U : \pi^{-1}[U] \cong U \times \pi^{-1}[x]$  fulfilling  $p_1 \circ h_U = \pi|_{\pi^{-1}[U]}$ .

$\pi^{-1}[x]$  is called the **Fiber of  $\pi$  at  $x$** , and sometimes written  $F_x$ .

$$\begin{array}{ccc} Y \supseteq \pi^{-1}[U] & \xrightarrow{h_U} & U \times \pi^{-1}[x] \\ \downarrow \pi & & \swarrow p_1 \\ X \supseteq U & & \end{array}$$

**Definition 2.2.** Let  $\pi : Y \rightarrow X$  be a locally trivial fibration. Then:

1. If all  $\pi^{-1}[x]$  are homeomorphic, we call  $\pi$  a **fiber bundle**.
2. If every  $\pi^{-1}[x]$  is discrete, we call  $\pi$  a **covering map**.
3. If we can pick  $U = X$ ,  $\pi$  is called a **trivial fibration**.

**Exercise 2.3.** If we replace  $\pi^{-1}[x]$  in the definition by an arbitrary topological space  $F_U$ , we get  $F_U \cong \pi^{-1}[x]$ . Thus, both definitions are equivalent.

**Lemma 2.4.** Let  $\pi : Y \rightarrow X$  be a locally trivial fibration. Let  $X$  be connected. Then  $\pi$  is a trivial fibration.

*Proof.* Let  $x_0 \in X$ . Let  $V = \{x \in X \mid F_x \cong F_{x_0}\}$  be the set of all points whose fibers are homeomorphic to the fiber of  $x_0$ . Let  $U$  be a neighborhood as in the definition. Then the fibers of all  $x \in U$  must be homeomorphic by definition. Thus, every point in  $V$  has a neighborhood contained in  $V$ , so  $V$  is open. We have

$$X \setminus V = \bigcup_{y \notin V} \{x \in X \mid F_x \cong F_y\} := \bigcup_{y \notin V} V_y$$

Since every  $V_y$  is open by the same argument,  $X \setminus V$  must be open. But since  $X$  is connected with  $(X \setminus V) \cup V = X$  and  $V$  nonempty,  $X \setminus V$  must

be empty. □

**Theorem 2.5.** A continuous function  $\pi : Y \rightarrow X$  is a covering map if and only if every  $x \in X$  has an open neighborhood  $U$  such that  $\pi^{-1}[U]$  can be decomposed into disjoint open sets  $V_i \subseteq Y$  such that  $\pi|_{V_i}$  is a homeomorphism  $V_i \cong U$ .

*Proof.*  $\Rightarrow$ : Let  $\pi$  be a covering map. Let  $I = F_U$ . Since  $h_U$  is bijective, the sets  $V_i := h_U^{-1}[U \times \{i\}]$  are disjoint. Since  $h_U$  is a homeomorphism and  $I = F_U$  is discrete, the  $V_i$  are also open. Since  $p_1|_{U \times \{i\}}$  is a homeomorphism  $U \times \{i\} \cong U$ , the composition

$$\pi|_{V_i} = (p_1|_{U \times \{i\}}) \circ h|_{V_i}$$

must be a homeomorphism  $V_i \mapsto U$ .

$\Leftarrow$ : Define

$$\begin{aligned} h_U : \pi^{-1}[U] &\cong U \times I \\ y \in V_i &\mapsto (\pi(y), i). \end{aligned}$$

Then the restriction of  $h_U$  to each  $V_i$  is by definition a homeomorphism  $V_i \cong U \cong U \times \{i\}$ , and so the whole of  $h_U$  is a homeomorphism  $\pi^{-1}[U] = \bigsqcup_{i \in I} V_i \cong \bigsqcup_{i \in I} U \times \{i\} = U \times I$ . □

### Definition 2.6: Lifting

Let  $\pi : Y \rightarrow X$  be a locally trivial fibration. Let  $f : Z \rightarrow X$  be continuous. Then a continuous map  $\hat{f} : Z \rightarrow Y$  is called a **lift from  $f$  to  $Y$**  if  $\pi \circ \hat{f} = f$ .

Any path can be lifted:

**Theorem 2.7.** Let  $\pi : Y \rightarrow X$  be a locally trivial fibration. Let  $\gamma : [0, 1] \rightarrow X$  be continuous. Let  $\pi(y_0) = \gamma(0)$ . Then there exists a lift  $\hat{\gamma} : [0, 1] \rightarrow Y$  with  $\hat{\gamma}(0) = y_0$  such that  $\pi \circ \hat{\gamma} = \gamma$ .

And lifts respect fundamental groups:

**Theorem 2.8.** Let  $\pi : Y \rightarrow X$  be a locally trivial fibration. Let  $f : Z \rightarrow X$ . Let  $\hat{f} : Z \rightarrow Y$  be a lift of  $f$ . Then we have:

$$\begin{aligned} \forall z_0 \in Z : \\ \exists y_0 \in Y : \\ \hat{f}(z_0) = y_0, \\ \pi_1(f)[\pi_1(Z, z_0)] \subseteq \pi_1(\pi)[\pi_1(Y, y_0)] \end{aligned}$$

*Proof.*

$$\begin{aligned}\pi_1(f)[\pi_1(Z, z_0)] &= (\pi_1(\pi \circ \hat{f}))[\pi_1(Z, z_0)] \\ &= (\pi_1(\pi) \circ \pi_1(\hat{f}))[\pi_1(Z, z_0)] \\ &= \end{aligned}$$

□

## **Part X**

# **Measure Theory**

# Contents

|          |                                                                           |            |
|----------|---------------------------------------------------------------------------|------------|
| <b>1</b> | <b>Measure Theory</b>                                                     | <b>177</b> |
| 1.1      | The Measure Problem .....                                                 | 177        |
| 1.1.1    | Vitali Sets .....                                                         | 178        |
| 1.1.2    | The Banach-Tarski Paradox .....                                           | 179        |
| 1.2      | Lattices and Boolean Algebras .....                                       | 180        |
| 1.2.1    | Boolean Algebras.....                                                     | 180        |
| 1.2.2    | Lattices.....                                                             | 182        |
| 1.3      | Set Algebras .....                                                        | 183        |
| 1.3.1    | Set Rings .....                                                           | 183        |
| 1.3.2    | Set Semirings .....                                                       | 184        |
| 1.4      | $\sigma$ -Algebras.....                                                   | 185        |
| 1.4.1    | The Borel $\sigma$ -Algebra .....                                         | 187        |
| 1.5      | Dynkin Systems.....                                                       | 188        |
| 1.6      | Monotone Classes .....                                                    | 189        |
| 1.7      | Measures .....                                                            | 191        |
| 1.8      | Carathéodory Extension .....                                              | 196        |
| 1.9      | Measures on $\mathbb{R}^n$ .....                                          | 197        |
| 1.9.1    | Jordan Content.....                                                       | 197        |
| 1.9.2    | Lebesgue Measure .....                                                    | 199        |
| <b>2</b> | <b>Measurable Functions</b>                                               | <b>202</b> |
| <b>3</b> | <b>Lebesgue Integration</b>                                               | <b>206</b> |
| 3.1      | Step Functions .....                                                      | 206        |
| 3.2      | Defining the Lebesgue Integral.....                                       | 208        |
| 3.3      | Comparing Riemann Integration and Lebesgue Integration .....              | 212        |
| 3.4      | Convergence Theorems .....                                                | 213        |
| 3.5      | $L^p$ -Spaces.....                                                        | 215        |
| 3.6      | Density Functions .....                                                   | 221        |
| <b>4</b> | <b>Integration in <math>\mathbb{R}^n</math></b>                           | <b>223</b> |
| 4.1      | Products of Measure Spaces.....                                           | 223        |
| 4.2      | Product Measures and Fubini's Theorem .....                               | 230        |
| 4.3      | Change of Variables .....                                                 | 238        |
| 4.4      | Alternative coordinate systems on $\mathbb{R}^2$ and $\mathbb{R}^3$ ..... | 241        |
| 4.4.1    | Polar Coordinates.....                                                    | 241        |
| 4.4.2    | Spherical Coordinates .....                                               | 245        |
| 4.4.3    | Cylindrical Coordinates .....                                             | 248        |
| 4.5      | The Surface Area Measure.....                                             | 250        |
| 4.6      | The Divergence Theorem.....                                               | 256        |
| <b>5</b> | <b>Convolutions</b>                                                       | <b>259</b> |

# Chapter 1

## Measure Theory

### 1.1 The Measure Problem

The most basic goal of measure theory is to establish a generalized notion of a “measure function”, which assigns a “volume” to a given set. In particular, we would like to establish a function that assigns volume functions to subsets of  $\mathbb{R}^n$  and has the following properties:

1. When given a subset with an easily intuitively definable volume, the volume function should agree with that volume. In particular, the volume of a cuboid should be the product of the lengths of its sides, and the length of a real interval  $(a, b)$  should be  $b - a$ .
2. The volume of a countable disjoint union of sets should be the sum of the individual volumes. This property is generally referred to as  $\sigma$ -additivity.
3. The volume should be invariant under isometries, i.e. functions like rotations, translations, and reflections should not change the volume of a set.
4. We want our volume to be a positive real number.

We call a  $\sigma$ -additive function a *measure* - we want to eventually find a measure on  $\mathbb{R}^n$  that also fulfills the other properties, but we want measures to be far more generally applicable, and as such we want to be able to define measures much more general spaces that may not have a defined notion of an interval or an isometry.

It will turn out that there exists exactly one function on  $\mathbb{R}^n$ , called the *Lebesgue measure*  $\lambda^n$ , that fulfills these conditions for a very large family of sets - enough to include every “somewhat reasonable” subset of  $\mathbb{R}^n$ . However, there are still counterexamples.

**Proposition 1.1.** Every subset of  $\mathbb{R}^n$  being Lebesgue-measurable is consistent with ZF (without the axiom of choice).

**Theorem 1.2.** Assuming the axiom of choice, there exist subsets of  $\mathbb{R}^n$  that cannot be assigned a volume without arriving at a contradiction.

It turns out that this result crucially relies on the full axiom of choice, and in particular is not implied by commonly used weaker forms of the axiom of choice such as the axiom of dependent choice.

The following two subsections deal with two different ways of proving this theorem by construction *non-measurable sets*: The *Vitali Sets*, and the decomposition of a sphere given in the *Banach-Tarski-Paradox*.

### 1.1.1 Vitali Sets

**Proposition 1.3.** The relation  $x \sim y \Leftrightarrow x - y \in \mathbb{Q}$  is an equivalence relation on the real numbers.

**Theorem 1.4.** There exist sets  $V \subset [0, 1]$  such that for each  $r \in \mathbb{R}$ , there exists exactly one number  $v \in V$  such that  $v - r$  is rational. We call such a set a **Vitali Set**.

*Proof.* Consider the aforementioned equivalence relation  $x \sim y$  on  $\mathbb{R}$ . Each equivalence class must contain at least one representative also contained in  $[0, 1]$ , since if  $x - y = q \in \mathbb{Q}$ , we have  $x - (y + q) \in \mathbb{Q}$  for all  $q \in \mathbb{Q}$ , letting us pick a  $q$  such that  $x - (y + q) \in [0, 1]$ . Being equivalence classes, they must also be disjoint.

This means we can use the axiom of choice to pick exactly one element of each equivalence class of  $\sim$ , giving us our set  $V$ .  $\square$

**Lemma 1.5.** Let  $q_1, q_2, \dots$  be an enumeration of  $\mathbb{Q} \cap [-1, 1]$ . Then for  $j \neq k$ , we have

$$(q_j + V) \cap (q_k + V) = \emptyset$$

*Proof.* Assume the intersection is non-empty. Then there exist  $v_1, v_2 \in V$  such that  $q_j + v_1 = q_k + v_2$ , meaning  $v_1 - v_2 \in \mathbb{Q}$ . Therefore,  $v_1$  must be in the same equivalence class as  $v_2$ . Since  $V$  contains exactly one element of each equivalence class, we have  $v_1 = v_2$ , and therefore we also have  $q_j = q_k$ , i.e.  $j = k$ .  $\square$

**Lemma 1.6.** We have

$$[0, 1] \subset \bigcup_{k \in \mathbb{N}} (q_k + V) \subset [-1, 2]$$

*Proof.* 1.  $\bigcup_{k \in \mathbb{N}} (q_k + V) \subset [-1, 2]$  follows trivially from  $q_k \in [-1, 1]$  and  $V \subset [0, 1]$ .

2.  $[0, 1] \subset \bigcup_{k \in \mathbb{N}} (q_k + V)$  follows from the definition of  $V$ , since for every  $y \in [0, 1]$  we have a unique  $v \in V$  such that  $y - v := q \in \mathbb{Q}$ , and since  $y \in [0, 1]$  and  $v \in [0, 1]$ , we have  $q \in [-1, 1]$ , i.e.  $q$  is contained in our enumeration.  $\square$

**Corollary 1.7.** Vitali sets are not measurable by a translation-invariant normed measure.

*Proof.* Assume that  $\lambda$  is translation-invariant, countably additive and that  $\lambda([a, b]) = b - a$ . Then we have:

$$1 = \lambda([0, 1]) \leq \lambda\left(\bigcup_{k \in \mathbb{N}} (q_k + V)\right) \leq \lambda([-1, 2]) = 3$$

Since our measure is countably additive and invariant under isometries, we can translate each individual set and preserve the measure:

$$\begin{aligned} \lambda\left(\bigcup_{k \in \mathbb{N}} (q_k + V)\right) &= \sum_{k \in \mathbb{N}} \lambda(q_k + V) \\ &= \sum_{k \in \mathbb{N}} \lambda(V) \end{aligned}$$

Now, if  $\lambda(V) \leq 0$ , we wouldn't have  $1 \leq \sum_{k \in \mathbb{N}} \lambda(V)$ , and if  $\lambda(V) > 0$ , we wouldn't have  $\sum_{k \in \mathbb{N}} \lambda(V) \leq 3$ . Therefore every possible measure we could assign to  $V$  leads to a contradiction.  $\square$

### 1.1.2 The Banach-Tarski Paradox

**Theorem 1.8.** Given any two sets  $A, B \subset \mathbb{R}^n$ , with  $n \geq 3$ , such that both  $A$  and  $B$  have a nonempty interior, there exist disjoint decompositions  $A_1 \sqcup \dots \sqcup A_k = A$  and  $B_1 \sqcup \dots \sqcup B_k = B$  such that for each  $i$ ,  $A_i$  and  $B_i$  can be transformed into each other by an isometry.

**Corollary 1.9.** The unit sphere can be transformed into two copies of the unit sphere by a finite disjoint decomposition followed by an isometry. The subsets of the decomposition therefore violate the countable union condition we expect from measure functions, and can therefore not be assigned a meaningful volume.

## 1.2 Lattices and Boolean Algebras

We have just seen that we cannot define our desired measure function on the full power set of  $\mathbb{R}^n$ . This means we will have to work on smaller systems of subsets of a given set. Naturally, we want to find the largest such systems that are still well-behaved enough to allow us to define a sensible notion of measure.

We will arrive at different algebraic structures on subsets of power sets, which will serve as the domains of our measure functions.

In order to gain a full birds-eye view of these definitions, we will first introduce the more general notion of *boolean algebras*:

### 1.2.1 Boolean Algebras

**Definition 1.10.** A **boolean algebra** is a set  $X$ , equipped with two binary operations  $\wedge$  and  $\vee$ , a unary operation  $\neg$ , and two elements  $0$  and  $1$ , such that:

1.  $\wedge$  and  $\vee$  are commutative,
2.  $1$  is a neutral element of  $\wedge$ , and  $0$  is a neutral element of  $\vee$ ,
3.  $\wedge$  distributes over  $\vee$  and  $\vee$  distributes over  $\wedge$ ,
4.  $x \wedge \neg x = 0$ , and  $x \vee \neg x = 1$ .

**Corollary 1.11.** Any boolean algebra also has the following properties:

1.  $\wedge$  and  $\vee$  are associative,
2.  $\wedge$  and  $\vee$  have the following **absorption property**:

$$a \wedge (a \vee b) = a$$

$$a \vee (a \wedge b) = a,$$

3.  $a = b \wedge a$  if and only if  $a \vee b = b$ .

There are three "central" boolean algebras, from which most of the terminology describing them is descended:

**Theorem 1.12.** The set of **propositional formulas** forms a boolean algebra, where  $0$  is the logical falsum ( $\perp$ , an unfulfillable formula),  $1$  is the logical verum ( $\top$ , a tautological formula), and  $\wedge$ ,  $\vee$  and  $\neg$  are logical "and", "or" and "not".

In computer science and circuit engineering, one often considers the subalgebra of this boolean algebra where every formula is directly evaluated to "0" or "1".

**Theorem 1.13. (Power set algebra):** The **power set**  $\mathcal{P}(X)$  of any set  $X$  forms a boolean algebra, where  $0 = \emptyset$ ,  $1 = X$ ,  $\wedge$  is the set intersection operation  $\cap$ ,  $\vee$  is the set union operation  $\cup$ , and  $\neg$  is the set complement operation  $M \rightarrow X \setminus M$ .

**Theorem 1.14. (Restrictions of Boolean algebras):** Let  $\mathcal{B}$  be a Boolean algebra on a set  $X$ , and let  $Y$  be a subset of  $X$ . Then the **restriction of  $\mathcal{B}$  to  $Y$** , defined as

$$\mathcal{B}|_Y := \{E \cap Y \mid E \in \mathcal{B}\},$$

is a boolean Algebra on  $Y$ .

**Theorem 1.15.** If  $Y \in \mathcal{B}$ , then

$$\mathcal{B}|_Y = \mathcal{B} \cap \mathcal{P}(Y) = \{E \subset Y \mid E \in \mathcal{B}\}$$

**Theorem 1.16. (Atomic algebra):** Let  $X$  be partitioned into a union

$$X = \bigcup_{\alpha \in I} A_\alpha$$

of disjoint sets  $A_\alpha$ , which we refer to as *atoms*. Then this partition forms a Boolean algebra

$$\mathcal{A}((A_\alpha)_{\alpha \in I}) := \left\{ E \mid E = \bigcup_{\alpha \in J} A_\alpha, J \subset I \right\}$$

of all the sets that can be represented as a union of atoms.

The power set Algebra on  $X$  is exactly the atomic algebra where  $X$  is partitioned into singleton atoms.

**Theorem 1.17.** Atomic algebras are uniquely determined by their atoms, up to relabeling. More precisely: Let  $(A_\alpha)_{\alpha \in I}$  and  $(B_\beta)_{\beta \in J}$  be two partitions of a set  $X$ . Then

$$\mathcal{A}((A_\alpha)_{\alpha \in I}) = \mathcal{A}((B_\beta)_{\beta \in J})$$

if and only if there exists a bijection  $\varphi : I \rightarrow J$  such that  $B_{\varphi(\alpha)} = A_\alpha$  for all  $\alpha \in I$ .

**Theorem 1.18.** Every finite Boolean algebra is an atomic algebra.

**Corollary 1.19.** Every finite Boolean algebra has cardinality  $2^n$ , where  $n \in \mathbb{N}$ .

**Corollary 1.20.** There is a one-to-one correspondence, up to relabeling, between finite Boolean algebras on a set  $X$  and finite partitions of  $X$  into non-empty sets.

**Theorem 1.21. (Dyadic algebras):** Let  $n, i_1, \dots, i_d \in \mathbb{Z}$ . The **dyadic algebra**  $\mathcal{D}_n(\mathbb{R}^d)$  at scale  $2^{-n}$  in  $\mathbb{R}^d$  is the atomic algebra generated by the products of the half-open dyadic intervals

$$I_j := \left[ \frac{i_j}{2^n}, \frac{i_j + 1}{2^n} \right)$$

of length  $2^{-n}$ .

This algebra consists exactly of the "grid figures" made up of a finite number of "pixels" of length  $2^{-n}$ .

**Theorem 1.22. (Intersection of Boolean Algebras):** The intersection of a family  $(\mathcal{B}_\alpha)_{\alpha \in I}$  of Boolean algebras on a set  $X$  is again a Boolean algebra, assuming the convention that, if  $I$  is empty, the intersection is the full power set. Furthermore, this Intersection is the finest Boolean algebra that is coarser than every  $\mathcal{B}_\alpha$ .

**Definition 1.23.** Let  $\mathcal{F}$  be any family of subsets of a set  $X$ . Then we define  $\langle \mathcal{F} \rangle_{\mathcal{B}}$  to be the intersection of all Boolean algebras that contain  $\mathcal{F}$ . We call this the **Boolean algebra generated by  $\mathcal{F}$** .

Equivalently,  $\langle \mathcal{F} \rangle_{\mathcal{B}}$  is the smallest Boolean algebra containing  $\mathcal{F}$ .

**Theorem 1.24.**  $\mathcal{F}$  is a Boolean algebra if and only if  $\langle \mathcal{F} \rangle_{\mathcal{B}} = \mathcal{F}$ .

### 1.2.2 Lattices

Boolean algebras themselves turn out to be specific instances of *lattices*, which play an important role in order theory and universal algebra.

**Definition 1.25.** A **lattice** is an algebraic structure  $(L, \vee, \wedge)$ , consisting of a set  $L$ , an operation  $\vee$ , called **join**, and an operation  $\wedge$ , called **meet**, such that the absorption laws  $a \vee (a \wedge b) = a$  and  $a \wedge (a \vee b) = a$ .

**Proposition 1.26.** Equivalently, a partially ordered set  $(L, \leq)$  is a lattice if every pair of elements has a least upper bound  $\sup(a, b) := a \vee b \in L$  and a greatest lower bound  $\inf(a, b) := a \wedge b \in L$ .

**Definition 1.27.** We call a lattice **bounded** if there exists a **least element**  $0$ , i.e.  $0$  fulfills  $a \vee 0 = a$ , and a **greatest element**  $1$ , which fulfills  $a \wedge 1 = a$ .

**Corollary 1.28.** A boolean algebra is a bounded lattice such that meet and join are distributive over each other and such that complements exist.

## 1.3 Set Algebras

**Definition 1.29.** Let  $X$  be a set and  $\mathcal{A} \subset \mathcal{P}(X)$ . Then we call  $\mathcal{A}$  a **set algebra** if it has the following properties:

1.  $\emptyset \in \mathcal{A}$
2. For any  $A \in \mathcal{A}$ , we have  $X \setminus A \in \mathcal{A}$  ( $\mathcal{A}$  is closed under the operation of taking complements).
3. For any  $F, G \in \mathcal{A}$ , we have  $F \cup G \in \mathcal{A}$  ( $\mathcal{A}$  is closed under binary unions).

**Corollary 1.30.** If  $\mathcal{A}$  is a set algebra on  $X$ , it also fulfills the following:

1.  $X \in \mathcal{A}$ ,
2. For any  $F, G \in \mathcal{A}$ , we have  $F \cap G \in \mathcal{A}$ ,
3. For any  $A_1, \dots, A_n \in \mathcal{A}$ , we have  $\bigcup_{i=1}^n A_i \in \mathcal{A}$ ,
4. For any  $A_1, \dots, A_n \in \mathcal{A}$ , we have  $\bigcap_{i=1}^n A_i \in \mathcal{A}$ .

Thus, we obtain the following more concise (but less readable) definition of set algebras:

**Corollary 1.31.** A set algebra is a subalgebra of the power set boolean algebra on  $X$ .

**Corollary 1.32.** A topology on  $X$  is “simply” a set algebra on  $X$  that is closed under arbitrary unions.

**Theorem 1.33. (Stone’s Representation Theorem for Boolean Algebras):** Every boolean algebra is isomorphic to a set algebra.

### 1.3.1 Set Rings

A very important weakening of the concept of set algebras is given by *set rings*, which contain the empty set and are closed under intersection and union, but don’t have to contain the full set  $X$  or be closed under complements.

**Theorem 1.34.** Let  $\mathcal{A}$  be a set ring. Then it is closed under finite symmetric difference, and forms a ring in the algebraic sense, with symmetric difference as addition and intersection as multiplication. If it contains the full set  $X$ , it forms a ring with identity.

**Definition 1.35.** Let  $I \subset \mathbb{R}$ . We call  $I$  an **interval** if there exist  $a, b \in \mathbb{R}$  such that  $(a, b) \subset I \subset [a, b]$ .

**Theorem 1.36.** The set of subsets of the real numbers which can be written as a finite union of intervals forms a set ring.

**Theorem 1.37.** Let  $\mathcal{R}$  and  $\mathcal{S}$  be set rings. Then the set of finite unions of cartesian products of elements of  $\mathcal{R}_i$  and  $\mathcal{S}_i$ , i.e. of elements of the form,

$$\bigcup_{i \in \mathbb{N}} R_i \times S_i,$$

is also a set ring, which we will denote by  $\mathcal{R} \boxtimes \mathcal{S}$ .

**Corollary 1.38.** The set of finite unions of cuboids in  $\mathbb{R}^n$ , where we define a cuboid to be a product of arbitrary intervals, i.e. we don't care if the boundary on any particular side is open or closed, forms a set ring. We call sets of this form **Elementary Sets**.

### 1.3.2 Set Semirings

**Theorem 1.39.** Let  $\mathcal{S} \subset \mathcal{P}(X)$ . We call  $\mathcal{S}$  a **set semiring**, or **semiring of sets**, if:

1.  $\emptyset \in \mathcal{S}$ ,
2.  $\mathcal{S}$  is closed under finite intersections,
3. For  $A, B \in \mathcal{S}$ , there exist disjoint sets  $S_1, \dots, S_n \in \mathcal{S}$  such that  $A \setminus B = \bigcup_{i=1}^n S_i$ .

This means that a set semiring is a weakened form of a set ring where complements are not necessarily contained in the semiring, but can still be "constructed" from elements of the ring. Any set ring is therefore immediately also a set semiring.

Sadly, unlike with rings of sets, there is absolutely no connection between set semirings and the algebraic notion of a semiring - a semiring of sets is exclusively a (semi)(ring of sets), and *not* a (semiring)(of sets). This makes it tempting for me to use an alternative name which makes this distinction more clear, but since I haven't encountered any good alternative names anywhere else (and because I already know I will forget to stick with this convention moving forwards) I will stick with the less than perfect established name.

Set semirings are of fundamental importance to measure theory because the set of cuboids in  $\mathbb{R}^n$  forms a set semiring, and we will end up defining our Lebesgue measure by approximating sets through coverings of the set with cuboids. Of course, we first have to establish a basic theory of set semirings and prove this claim.

**Theorem 1.40.** The set  $\mathcal{I}$  of real intervals forms a set semiring.

**Theorem 1.41.** The product of two set semirings is again a set semiring.

**Corollary 1.42.** The set  $\mathcal{Q}$  of cuboids in  $\mathbb{R}^n$  (once again with both open and closed sides allowed) forms a set semiring.

## 1.4 $\sigma$ -Algebras

The most important type of set algebra for the purposes of measure theory is the  $\sigma$ -Algebra, on which we will eventually define the notion of a “measure” in our desired final form.

### Definition 1.43: $\sigma$ -Algebra

Let  $X$  be an arbitrary set and  $\mathcal{A} \subset \mathcal{P}(X)$ . We call  $\mathcal{A}$  a  $\sigma$ -Algebra on  $X$  if:

1.  $X \in \mathcal{A}$
2. For all  $A \in \mathcal{A}$ , we have  $X \setminus A \in \mathcal{A}$  ( $\mathcal{A}$  is closed under the operation of taking a complement).
3. For all  $A_i \in \mathcal{A}$ , we have  $\bigcup_{i \in \mathbb{N}} A_i \in \mathcal{A}$  ( $\mathcal{A}$  is closed under the operation of taking countable unions).

If  $\mathcal{A}$  is a  $\sigma$ -algebra on  $X$ , we call  $(X, \mathcal{A})$  a **measure space**, and any set  $A \in \mathcal{A}$   **$\mathcal{A}$ -measurable**.

The “ $\sigma$ ” here once again stands for “countable sum”, as it also did for  $\sigma$ -additivity and  $F_\sigma$  sets.

**Corollary 1.44.** Any  $\sigma$ -algebra contains the empty set and is closed under countable intersection.

**Corollary 1.45.** A  $\sigma$ -algebra can be more concisely defined as a set algebra that is closed under *countable* union and intersection, not just finite ones.

This also trivially makes every  $\sigma$ -algebra a Boolean algebra.

**Theorem 1.46.** Every atomic algebra is a  $\sigma$ -algebra.

### Theorem 1.47: Induced $\sigma$ -algebra

Just like for Boolean algebras, the restriction  $\mathcal{A}|_Y$  of a  $\sigma$ -algebra  $\mathcal{A}$  on  $X$  to a subset  $Y \subset X$ , defined by

$$\mathcal{A}|_Y = \{A \cap Y \mid A \in \mathcal{A}\},$$

is again a  $\sigma$ -algebra on  $Y$ .

**Theorem 1.48.** We have  $\mathcal{A}|_Y \subset \mathcal{A}$  if and only if  $Y \in \mathcal{A}$ .

*Proof.*

1. Assume  $\mathcal{A}|_Y \subset \mathcal{A}$ . We have  $X \cap Y = Y \in \mathcal{A}|_Y$ , and thus we get  $Y \in \mathcal{A}$ .
2. Assume  $Y \in \mathcal{A}$ . Then since  $\sigma$ -algebras are closed under intersection, we have  $A \cap Y \in \mathcal{A}$  for all  $A \in \mathcal{A}$ , i.e. we have  $\mathcal{A}|_Y \subset \mathcal{A}$ .

□

**Theorem 1.49.** Let  $X \neq \emptyset$ , let  $(Y, \mathcal{B})$  be a measurable space and let  $f : X \rightarrow Y$  be an arbitrary map. Then

$$f^{-1}(\mathcal{B}) = \{f^{-1}(B) \mid B \in \mathcal{B}\}$$

is a  $\sigma$ -algebra.

Note however that we don't necessarily have  $f^{-1}(\mathcal{B}) = \mathcal{A}$ , or even  $f^{-1}(\mathcal{B}) \subset \mathcal{A}$ ! We call maps that fulfill the latter property *measurable*. Measurable maps play a key role in measure theory and especially in integration theory, entirely analogous to the role continuous functions play in topology. We will study them in more detail in a later chapter.

**Theorem 1.50.** The intersection of arbitrarily many  $\sigma$ -algebras is on the same set  $X$  is once again a  $\sigma$ -algebra.

**Corollary 1.51: Generated  $\sigma$ -algebra**

Let  $\mathcal{E} \subset \mathcal{P}(X)$ . Then  $\langle \mathcal{E} \rangle_\sigma$  denotes the intersection of all  $\sigma$ -algebras containing  $\mathcal{E}$ .

In particular, note that if  $\mathcal{A}$  is a  $\sigma$ -Algebra containing  $\mathcal{E}$ , we have

$$\langle \mathcal{E} \rangle_\sigma \subseteq \mathcal{A}.$$

**Theorem 1.52.** We have  $\langle \mathcal{F} \rangle_{\mathcal{B}} = \langle \mathcal{F} \rangle_\sigma$  if and only if  $\langle \mathcal{F} \rangle_{\mathcal{B}}$  is a  $\sigma$ -algebra.

**Theorem 1.53.** Let  $X$  be any set. Let  $\mathcal{E}_i \subset \mathcal{P}(X)$  be a family of sets of subsets of  $X$  indexed by  $i \in I$ . Then we have:

$$\left\langle \bigcup_{i \in I} \langle \mathcal{E}_i \rangle_\sigma \right\rangle_\sigma = \left\langle \bigcup_{i \in I} \mathcal{E}_i \right\rangle_\sigma$$

### 1.4.1 The Borel $\sigma$ -Algebra

**Definition 1.54.** Let  $X$  be a topological space. The **Borel  $\sigma$ -algebra**  $\mathcal{B}[X]$  of  $X$  is the  $\sigma$ -Algebra generated by the open subsets of  $X$ .

**Theorem 1.55.** The Borel  $\sigma$ -Algebra  $\mathcal{B}[\mathbb{R}^d]$  is equivalently generated by any of the following:

1. The closed subsets of  $\mathbb{R}^d$ ,
2. The compact subsets of  $\mathbb{R}^d$ ,
3. The open balls of  $\mathbb{R}^d$ ,
4. The boxes in  $\mathbb{R}^d$ ,
5. The elementary sets in  $\mathbb{R}^d$ .

## 1.5 Dynkin Systems

### Definition 1.56: Dynkin System

Let  $\mathcal{D} \subset \mathcal{P}(X)$ . Then we call  $\mathcal{D}$  a **Dynkin system** iff:

1.  $X \in \mathcal{D}$
2.  $\mathcal{D}$  is closed under complements  $B \setminus A$  as long as  $A \subset B$
3.  $\mathcal{D}$  is closed under countable pairwise disjoint union

### Theorem 1.57

A Dynkin system is a  $\sigma$ -algebra iff. it is closed under intersection.

**Theorem 1.58.** Let  $\mathcal{E} \subset \mathcal{P}(X)$  be closed under intersection. Then  $\langle \mathcal{E} \rangle_{\mathcal{D}} = \langle \mathcal{E} \rangle_{\sigma}$ .

## 1.6 Monotone Classes

### Definition 1.59: Monotone classes

Let  $\mathcal{M} \subset \mathcal{P}(X)$ . Then we call  $\mathcal{M}$  a **monotone class** if:

1. For any family  $(A_i)_{i \in \mathbb{N}} \subset \mathcal{M}$  such that  $A_i \subset A_{i+1}$  holds for all  $i \in \mathbb{N}$ , we have

$$\bigcup_{i=1}^{\infty} A_i \in \mathcal{M}$$

2. For any family  $(A_i)_{i \in \mathbb{N}} \subset \mathcal{M}$  such that  $A_i \supset A_{i+1}$  holds for all  $i \in \mathbb{N}$ , we have

$$\bigcap_{i=1}^{\infty} A_i \in \mathcal{M}$$

We denote the smallest monotone class containing a set  $\mathcal{S} \subset \mathcal{P}(X)$  as  $\langle \mathcal{S} \rangle_{\mathcal{M}}$ .

**Corollary 1.60.** Every  $\sigma$ -algebra is a monotone class.

**Corollary 1.61.** Given  $\mathcal{F} \subset \mathcal{P}(X)$ , we have  $\langle \mathcal{F} \rangle_{\mathcal{M}} \subseteq \langle \mathcal{F} \rangle_{\sigma}$

Recall that a *set algebra* is a set  $\mathcal{A} \subset \mathcal{P}(X)$  that contains the empty set and is closed under complements and binary unions.

**Theorem 1.62.** Let  $\mathcal{F} \subset \mathcal{P}(X)$  be a set algebra and a monotone class. Then  $\mathcal{F}$  is a  $\sigma$ -algebra.

*Proof.* We need to show that  $\mathcal{F}$  is closed under countable unions. Let  $(F_j)_{j \in \mathbb{N}} \subset \mathcal{F}$  be a family of elements of  $\mathcal{F}$ . Since  $\mathcal{F}$  is closed under finite union, the increasing sequence of sets

$$A_i := \bigcup_{j=1}^i F_j$$

must be contained in  $\mathcal{F}$ . Therefore, since  $\mathcal{F}$  is a monotone class, the union

$$\bigcup_{i \in \mathbb{N}} A_i = \bigcup_{i \in \mathbb{N}} \left( \bigcup_{j=1}^i F_j \right) = \bigcup_{j \in \mathbb{N}} F_j$$

must also be contained in  $\mathcal{F}$ . Therefore,  $\mathcal{F}$  is closed under countable union, making it a  $\sigma$ -algebra.  $\square$

The relevance of monotone classes is given by the following theorem, which gives us an additional way of verifying whether a given subset of a power set forms a  $\sigma$ -algebra:

**Theorem 1.63: Monotone class theorem**

Let  $\mathcal{A} \subset \mathcal{P}(X)$  be a set algebra. Then

$$\langle \mathcal{A} \rangle_{\mathcal{M}} = \langle \mathcal{A} \rangle_{\sigma}$$

*Proof.* We have already observed that, since any  $\sigma$ -algebra is a monotone class, we have  $\langle \mathcal{A} \rangle_{\mathcal{M}} \subset \langle \mathcal{A} \rangle_{\sigma}$ . Therefore, it suffices to show that we also have  $\langle \mathcal{A} \rangle_{\mathcal{M}} \supset \langle \mathcal{A} \rangle_{\sigma}$ . This holds automatically if  $\langle \mathcal{A} \rangle_{\mathcal{M}}$  is a  $\sigma$ -algebra, and since we already know that any monotone class is a  $\sigma$ -algebra if it is a set algebra, it suffices to show that  $\langle \mathcal{A} \rangle_{\mathcal{M}}$  is a set algebra.

For any  $E \in \langle \mathcal{A} \rangle_{\mathcal{M}}$ , let  $\mathcal{M}_E$  be the system of "good subsets" of  $X$  which can be combined with  $E$  as desired, i.e.

$$\mathcal{M}_E := \left\{ F \subset X \mid \begin{array}{l} F \setminus E \in \langle \mathcal{A} \rangle_{\mathcal{M}} \\ E \setminus F \in \langle \mathcal{A} \rangle_{\mathcal{M}} \\ E \cup F \in \langle \mathcal{A} \rangle_{\mathcal{M}} \end{array} \right\}$$

1. First, we want to show that  $\mathcal{M}_E$  is a monotone class.
2. Next, we want to show that for every  $E \in \mathcal{A}$ , we have  $\langle \mathcal{A} \rangle_{\mathcal{M}} \subset \mathcal{M}_E$ . Since  $\mathcal{A}$  is a set algebra, every element of  $\mathcal{A}$  combines with  $E$  as desired, giving us  $\mathcal{A} \subset \mathcal{M}_E$ . Therefore, since  $\mathcal{M}_E$  is a monotone class, we have

$$\langle \mathcal{A} \rangle_{\mathcal{M}} \subset \langle \mathcal{M}_E \rangle_{\mathcal{M}} = \mathcal{M}_E.$$

3. Next, we want to show that for every  $F \in \langle \mathcal{A} \rangle_{\mathcal{M}}$ , we have  $\langle \mathcal{A} \rangle_{\mathcal{M}} \subset \mathcal{M}_F$ . We just showed that  $\langle \mathcal{A} \rangle_{\mathcal{M}} \subset \mathcal{M}_F$ , which means we have  $F \in \mathcal{M}_E$  for any  $E \in \mathcal{A}$ . Since the definition of  $\mathcal{M}_E$  is entirely symmetric in  $E$  and  $F$ , we also have  $E \in \mathcal{M}_F$ . Therefore, since all  $E \in \mathcal{A}$  are contained in  $\mathcal{M}_F$ , we have  $\mathcal{A} \subset \mathcal{M}_F$ , implying

$$\langle \mathcal{A} \rangle_{\mathcal{M}} \subset \langle \mathcal{M}_F \rangle_{\mathcal{M}} = \mathcal{M}_F.$$

4. We have just shown that, for any  $E, F \in \langle \mathcal{A} \rangle_{\mathcal{M}}$ , we have  $E \in \mathcal{M}_F$ , implying that  $\langle \mathcal{A} \rangle_{\mathcal{M}}$  is closed under complementation and binary union. Since  $X \in \mathcal{A}$ , we also have  $X \in \langle \mathcal{A} \rangle_{\mathcal{M}}$ . Therefore,  $\langle \mathcal{A} \rangle_{\mathcal{M}}$  is a set algebra.

Therefore,  $\langle \mathcal{A} \rangle_{\mathcal{M}}$  is indeed both a set algebra and a monotone class, making it a  $\sigma$  algebra.  $\square$

## 1.7 Measures

With our different subset systems in place, we can finally give a general formal definition of a measure. Along the way, we will encounter outer measures, contents, and premeasures, which are “weakened” measures that we can use to generate proper ones.

### Definition 1.64: Outer measure

Let  $X$  be a set and  $\mu$  be a function  $\mathcal{P}(X) \rightarrow [0, \infty]$ . We call  $\mu$  an **outer measure on  $X$**  if it is  $\sigma$ -subadditive.

**Theorem 1.65.** Let  $X$  be any set. Then the constant function  $\mu = 0$  is an outer measure on  $X$ , which is known as the **trivial measure**.

**Theorem 1.66.** Let  $X$  be any set. Let  $A \subset X$  and  $x \in X$ . Then the function

$$\delta_x(A) := \begin{cases} 1 & x \in A \\ 0 & x \in X \setminus A \end{cases}$$

is an outer measure on  $X$ , which is known as the **Dirac measure**.

### Theorem 1.67: Counting measure

Let  $X$  be any set. Let  $A \subset X$ . Then the function

$$\text{card}(A) := \begin{cases} |A| & |A| \text{ is finite} \\ \infty & |A| \text{ is infinite} \end{cases}$$

is an outer measure on  $X$ , which is known as the **counting measure**.

### Definition 1.68: Measurable set

Let  $\mu$  be an outer measure on a set  $X$ . Then we call a subset  $A \subset X$   **$\mu$ -measurable**, or just **measurable**, if for all  $S \subseteq X$  we have

$$\mu(S) = \mu(S \cap A) + \mu(S \setminus A)$$

The system of all  $\mu$ -measurable sets is sometimes denoted  $\mathcal{M}(\mu)$ .

By this definition,  $\mu$ -measurable sets are exactly the sets for which  $\mu$  is  $\sigma$ -additive. This means that if we restrict  $\mu$  to its measurable sets, we get a proper measure.

Note that by the subadditivity of outer measures, we already get that the left side is at most as large as the right side, meaning that this condition can equivalently be weakened to

$$\mu(S) \geq \mu(S \cap A) + \mu(S \setminus A).$$

### Theorem 1.69: Importance of $\sigma$ -algebras

Let  $\mu$  be an outer measure. Then the set of  $\mu$ -measurable sets forms a  $\sigma$ -Algebra.

Since the restriction of an outer measure to its measurable sets forms a proper measure, this theorem tells us that  $\sigma$ -algebras give us the most sensible definition for which properties the domain of a measure should fulfill. Therefore, we finally get a proper definition of a measure:

**Definition 1.70: Measure**

Let  $\mathcal{A}$  be a  $\sigma$ -Algebra. Then we call a  $\sigma$ -additive function  $\mu : \mathcal{A} \rightarrow [0, \infty]$  a **measure**.

**Definition 1.71.** If  $\mathcal{A}$  is a  $\sigma$ -algebra on  $X$ , and  $\mu$  is a measure on  $\mathcal{A}$ , we call the triple  $(X, \mathcal{A}, \mu)$  a **measure space**.

**Definition 1.72.** Let  $(X, \mathcal{A}, \mu)$  be a measure space. Then we call this space a **probability space**, and  $\mu$  a **probability measure**, iff  $\mu(X) = 1$ .

**Definition 1.73: Content, Premeasure**

Let  $\mathcal{S}$  be a set semiring. Then we call a finitely additive function  $\mathcal{S} \rightarrow [0, \infty]$  a **content**, and a  $\sigma$ -additive function  $\mathcal{S} \rightarrow [0, \infty]$  a **premeasure**.

In effect, a premeasure is a measure whose domain might not be as big as it could be. Every measure is trivially also a content and a premeasure, and every measure defined on  $\mathcal{P}(X)$  is an outer measure.

**Theorem 1.74: Restriction of a measure**

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $Y \in \mathcal{A}$ . Then if we define

$$\mu|_Y(A \cap M) := \mu(A \cap M),$$

$\mu|_Y$  defines a measure on  $\mathcal{A}|_Y$ .

**Theorem 1.75: Image measure**

Let  $X, Y$  be sets. Let  $f : X \rightarrow Y$  be a function. Let  $\mu$  be an outer measure on  $X$ . then the function

$$\begin{aligned} f(\mu) : \mathcal{P}(Y) &\rightarrow [0, \infty] \\ f(\mu)(B) &= \mu(f^{-1}(B)) \end{aligned}$$

is a measure on  $Y$ , which is known as the **image measure of  $\mu$  under  $f$** .

*Proof.* 1. First, note that we have:

$$\begin{aligned} f(\mu)(\emptyset) &= \mu(f^{-1}(\emptyset)) \\ &= \mu(\emptyset) \\ &= 0 \end{aligned}$$

2. Furthermore, for  $B \subset \bigcup_{i \in \mathbb{N}} B_i$ , we have  $f^{-1}(B) \subset \bigcup_{i \in \mathbb{N}} B_i$ . By the

subadditivity of  $\mu$ , we get:

$$\begin{aligned} f(\mu)(B) &= \mu(f^{-1}(B)) \\ &\leq \sum_{i \in \mathbb{N}} \mu(f^{-1}(B_i)) \\ &= \sum_{i \in \mathbb{N}} f(\mu)(B_i). \end{aligned}$$

Therefore,  $f(\mu)$  is also subadditive and thus an outer measure.  $\square$

**Lemma 1.76.** Let  $X, Y$  be sets. Let  $f : X \rightarrow Y$  be a function. Let  $\mu$  be an outer measure on  $X$  and let  $f(\mu)$  be the corresponding image measure. Then  $B \subset Y$  is  $f(\mu)$  measurable if  $f^{-1}(B)$  is  $\mu$ -measurable.

*Proof.* By the definition of measurability:

1.  $B \subset Y$  is  $f(\mu)$ -measurable if and only if for all  $T \subset Y$ , we have:

$$f(\mu)(T) \geq f(\mu)(T \cap B) + f(\mu)(T \setminus B)$$

i.e.

$$\mu(f^{-1}(T)) \geq \mu(f^{-1}(T \cap B)) + \mu(f^{-1}(T \setminus B))$$

Since  $f^{-1}(T \cap B) = f^{-1}(T) \cap f^{-1}(B)$  and  $f^{-1}(T \setminus B) = f^{-1}(T) \setminus f^{-1}(B)$ , this is equivalent to:

$$\mu(f^{-1}(T)) \geq \mu(f^{-1}(T) \cap f^{-1}(B)) + \mu(f^{-1}(T) \setminus f^{-1}(B))$$

2.  $f^{-1}(B)$  is  $\mu$ -measurable if and only if for all  $S \subset X$ , we have:

$$\mu(S) \geq \mu(S \cap f^{-1}(B)) + \mu(S \setminus f^{-1}(B)).$$

In particular, if  $f^{-1}(B)$  is  $\mu$ -measurable, this equation must hold for  $S = f^{-1}(T)$ , implying that  $B$  is  $f(\mu)$ -measurable.  $\square$

### Definition 1.77: ( $\sigma$ )-finite measure

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Then:

1. We call  $\mu$  **finite** if  $\mu(X) < \infty$ .
2. We call  $\mu$   **$\sigma$ -finite** if  $X$  can be written as a countable union of sets of finite measure.

**Theorem 1.78: Measures are  $\sigma$ -subadditive**

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $(A_i)_{i \in \mathbb{N}} \subset \mathcal{A}$  be a family of (not necessarily disjoint!) measurable subsets of  $X$ . Then we have:

$$\mu\left(\bigcup_{i \in \mathbb{N}} A_i\right) \leq \sum_{i \in \mathbb{N}} \mu(A_i)$$

**Theorem 1.79: Continuity from above and below**

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $(A_i)_{i \in \mathbb{N}} \subset \mathcal{A}$ . Then we have:

1. **Continuity from below:** If  $A_i \subset A_{i+1}$  for every  $i \in \mathbb{N}$ , we have

$$\lim_{i \rightarrow \infty} \mu(A_i) = \mu\left(\bigcup_{i \in \mathbb{N}} A_i\right)$$

2. **Continuity from above:** If  $\mu(A_k) < \infty$  for any  $k \in \mathbb{N}$  and  $A_i \supset A_{i+1}$  for every  $i \in \mathbb{N}$ , we have

$$\lim_{i \rightarrow \infty} \mu(A_i) = \mu\left(\bigcap_{i \in \mathbb{N}} A_i\right)$$

**Definition 1.80.** Let  $(X, \mathcal{A}, \mu)$ . We call a set  $A \in \mathcal{A}$  a  **$\mu$ -zero set**, or simply zero set, iff  $\mu(A) = 0$ . We denote the set of all  $\mu$ -zero sets by  $\mathcal{N}(\mu)$ .

**Definition 1.81:  $\mu$ -almost everywhere**

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $P$  be some property. We say that  $P$  holds  **$\mu$ -almost everywhere on  $M$** , or that  $P$  holds for  **$\mu$ -almost all  $x \in M$** , iff the set of  $x \in M$  such that  $P(x)$  is false is a zero set.

**Definition 1.82: Complete measure**

We call a measure  $\mu$  **complete** if any subset of a  $\mu$ -zero set is also a  $\mu$ -zero set.

Note that, in particular, any subset of a  $\mu$ -zero set needs to be  $\mu$ -measurable in the first place.

**Lemma 1.83.** Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $\mathcal{Z}_\mu$  be the system of all subsets of  $X$  that are also subsets of a  $\mu$ -zero set. Then  $\mu$  is complete if and only if  $\mathcal{Z}_\mu \subset \mathcal{A}$ .

**Corollary 1.84.** Let  $\mu$  be an outer measure. Then  $\mu|_{\mathcal{M}(\mu)}$  is complete.

**Theorem 1.85: Completion of a Measure**

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let

$$\overline{\mathcal{A}} = \{A \cup N \mid A \in \mathcal{A}, N \in \mathcal{Z}_\mu\}.$$

and

$$\overline{\mu}(A \cup N) := \mu(A).$$

Then  $\overline{\mathcal{A}}_\mu$  is a  $\sigma$ -algebra and  $\overline{\mu}$  is a complete measure on  $\overline{\mathcal{A}}_\mu$ .

**Theorem 1.86. (Uniqueness of Completion):** Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $(X, \mathcal{B}, \nu)$  be a complete measure space such that  $\mathcal{A} \subset \mathcal{B}$  and  $\mu(A) = \nu(A)$  for all  $A \in \mathcal{A}$ . Then we have  $\overline{\mathcal{A}}_\mu \subset \mathcal{B}$  and  $\overline{\mu} = \nu$  on  $\overline{\mathcal{A}}_\mu$ .

## 1.8 Carathéodory Extension

We can construct outer measures from a very large class of functions by the following construction:

### Theorem 1.87: Carathéodory Extension

Let  $\mathcal{S}$  be a system of subsets of a set  $X$  containing the empty set. Let  $\lambda : \mathcal{S} \rightarrow [0, \infty]$  be a function such that  $\lambda(\emptyset) = 0$ . Then the function

$$\mu(E) := \inf \left\{ \sum_{i=1}^{\infty} \lambda(P_i) \mid P_i \in \mathcal{S}, E \subset \bigcup_{i=1}^{\infty} P_i \right\}$$

is an outer measure on  $X$ .

### Definition 1.88: Regular Measure

Let  $\mu$  be an outer measure on  $X$ . Then we call  $\mu$  **regular** iff. for every  $M \subset X$ , we can find a  $\mu$ -measurable set  $D \supset M$  such that  $\mu(M) = \mu(D)$ .

**Lemma 1.89.** The Carathéodory extension of a premeasure is regular.

### Theorem 1.90: Characterization of measurable sets

Let  $\lambda : \mathcal{R} \rightarrow [0, \infty]$  be a  $\sigma$ -finite premeasure on a ring  $\mathcal{R} \subset \mathcal{P}(X)$ . Let  $\bar{\lambda}$  be its Carathéodory extension. Then a set  $D \subset X$  is  $\mu$  measurable if and only if either of the following hold:

1. There exists an  $E \supset D$  such that  $E \in \langle \mathcal{R} \rangle_{\sigma}$  and  $\bar{\lambda}(E \setminus D) = 0$ .
2. There exists a  $C \subset D$  such that  $C \in \langle \mathcal{R} \rangle_{\sigma}$  and  $\bar{\lambda}(D \setminus C) = 0$ .

## 1.9 Measures on $\mathbb{R}^n$

### 1.9.1 Jordan Content

**Definition 1.91.** Let  $E \subset \mathbb{R}^n$  be an elementary set. Then we can assign to it the **elementary volume**  $\text{vol}(E)$ , where the volume of a cuboid is the product of its side lengths and the volume of a finite union of cuboids is the sum of the volumes of the individual cuboids making up  $E$ .

**Definition 1.92. (Inner and outer Jordan content):** Let  $E \subset \mathbb{R}^n$ .

1. The **inner Jordan measure**  $J_*(E)$  is

$$J_*(E) := \sup_{\substack{Q_i \in \mathcal{Q}, \\ \bigcup_{i=1}^n Q_i \subset E}} \text{vol}(Q)$$

2. The **outer Jordan measure**  $J^*(E)$  is

$$J^*(E) := \inf_{\substack{Q_i \in \mathcal{Q}, \\ \bigcup_{i=1}^n Q_i \supset E}} \text{vol}(Q)$$

**Definition 1.93. (Jordan measurable set, Jordan content):** We call  $E$  **Jordan-measurable** if  $J_*(E) = J^*(E)$ . Then we call  $J(E) = J_*(E) = J^*(E)$  the **Jordan content**.

**Theorem 1.94.** The Jordan content is  $\sigma$ -additive.

**Theorem 1.95.** The following are equivalent:

1.  $E$  is Jordan measurable,
2. For every  $\varepsilon > 0$ , there exist elementary sets  $A \subset E \subset B$  such that  $\text{vol}(B \setminus A) \leq \varepsilon$ ,
3. For every  $\varepsilon > 0$ , there exists an elementary set  $A$  such that  $J^*(A \Delta E) \leq \varepsilon$ .

**Theorem 1.96.** The collection of subsets of  $\mathbb{R}^n$  that are either Jordan measurable or have a Jordan-measurable complement form a Boolean algebra, known as the Jordan algebra.

**Theorem 1.97.** The Jordan algebra is non-atomic.

**Theorem 1.98. (Regions under continuous Graphs are Jordan measurable):**

Let  $B$  be a closed box in  $\mathbb{R}^n$ , and let  $f : B \rightarrow \mathbb{R}$  be a continuous function. Then the set

$$\{(x, t) \mid x \in B, 0 \leq t \leq f(x)\} \subset \mathbb{R}^{n+1}$$

is Jordan measurable.

**Theorem 1.99.** Triangles are Jordan measurable.

**Theorem 1.100.** Convex polytopes in  $\mathbb{R}^n$  are Jordan measurable.

**Theorem 1.101.** Open and closed Euclidean balls are Jordan measurable.

**Theorem 1.102.** Every subset of a Jordan null set is Jordan measurable and also a Jordan null set.

**Theorem 1.103.** The sets  $[0, 1]^2 \setminus \mathbb{Q}^2$  and  $[0, 1]^2 \cap \mathbb{Q}^2$  are both not Jordan measurable.

Informally, sets with a lot of “holes” or with very messy, fractal-like boundaries are generally not Jordan-measurable.

**Theorem 1.104.** There exist countable unions, and countable intersections, of Jordan measurable sets which are not Jordan measurable.

### 1.9.2 Lebesgue Measure

One can extend the Jordan measure to a significantly larger number of subsets of  $\mathbb{R}^n$  by simply allowing countable unions of cuboids (instead of finite unions of cuboids). The Lebesgue measure is simply this generalization of the Jordan measure. Formally:

**Definition 1.105.** Let  $E \subset \mathbb{R}^n$ . The **Lebesgue outer measure** of  $E$  is given by

$$\lambda^*(E) := \inf_{\substack{E \subset \bigcup_{n=1}^{\infty} Q_n, \\ Q_n \in \mathcal{Q}}} \left\{ \sum_{n=1}^{\infty} \text{vol}^n Q_n \right\},$$

i.e. the Lebesgue outer measure of  $E$  is the greatest lower bound of the measures of all coverings of  $E$  by cuboids.

**Theorem 1.106.** We have  $\lambda^0 = \text{card}$  - the zero-dimensional Lebesgue measure is the counting measure.

*Proof.*  $\mathbb{R}^0$  is by definition cartesian product of  $\mathbb{R}$  with a set of cardinality 0 i.e. the empty set, as its index set:

$$\mathbb{R}^0 = \prod_{\emptyset} \mathbb{R}$$

By definition of cartesian products, its members must be functions  $\emptyset \rightarrow \mathbb{R}$ . There is exactly one such "empty function", which we will denote  $f_{\emptyset}$ . Thus, the only subsets of  $\mathbb{R}^0$  are the empty set and  $\{f_{\emptyset}\}$ .

1. Since  $\lambda^0$  and  $\text{card}$  are measures, we have

$$\lambda^0(\emptyset) = 0 = \text{card}(\emptyset)$$

2. By definition, a 0-dimensional cuboid must be a function  $\emptyset \rightarrow I$  to an interval  $I \subset \mathbb{R}$ . Since all functions from the empty set to any set are identical,  $f_{\emptyset}$  qualifies as a cuboid. We thus have:

$$\begin{aligned} \lambda^0(\{f_{\emptyset}\}) &= \text{vol}^0(\{f_{\emptyset}\}) \\ &= \prod_{\emptyset} \text{vol}(I) \\ &= 1 \\ &= \text{card}(\{f_{\emptyset}\}) \end{aligned}$$

Where the last step follows since empty products evaluate to the multiplicative identity, in this case the number 1.

□

**Theorem 1.107.** Let  $U \subset \mathbb{R}^n$  be open. Then  $U$  is a countable union of cuboids.

Therefore, we can define Lebesgue measurability similarly to Jordan measurability - a set is Lebesgue measurable if it is "almost" an open set.

**Definition 1.108.** A set  $E \subset \mathbb{R}^n$  is **Lebesgue measurable** if for every  $\varepsilon > 0$ , there exists an open set  $U \subset \mathbb{R}^n$  such that  $U \subset E$  and  $\lambda^*(U \setminus E) \leq \varepsilon$ . If  $E$  is Lebesgue measurable, we refer to  $\lambda(E) := \lambda^*(E)$  as the **Lebesgue measure** of  $E$ . If the dimension  $n$  should be emphasized, we sometimes write  $\lambda(E)$  as  $\lambda^n(E)$ .

**Theorem 1.109.** The Lebesgue measure fulfills:

$$E \subset \bigcup_{i \in \mathbb{N}} E_i \implies \lambda(E) \leq \sum_{i \in \mathbb{N}} \lambda(E_i)$$

This is a weakened form of  $\sigma$ -additivity, which is known as  $\sigma$ -subadditivity.

**Theorem 1.110.** The Lebesgue measure defines a  $\sigma$ -additive function on the Lebesgue measurable sets.

**Theorem 1.111.** The collection of subsets of  $\mathbb{R}^n$  that are either Lebesgue measurable or have a Lebesgue-measurable complement form a Boolean algebra.

**Theorem 1.112.** There exist Lebesgue measurable sets which are not Borel.

You may have heard in linear algebra that the determinant of a matrix  $S$  tells us how the matrix scales the volume of the unit cube. Since the Lebesgue measure is defined using volumes of unit cubes, this intuition also holds for the image of any set  $E$  under  $S$ :

**Theorem 1.113: Linear Transformation Equation**

Let  $S \in \mathbb{R}^{n \times n}$ . Then for all  $E \subset \mathbb{R}^n$ , we have:

$$\lambda^n(S(E)) = |\det(S)| \cdot \lambda^n(E)$$

**Definition 1.114: Smith-Volterra-Cantor Set**

The Smith-Volterra-Cantor set is the set obtained by removing the middle  $1/8$  of the interval  $[0, 1]$ , and then iteratively removing the middle  $1/8$  of any remaining intervals.

**Theorem 1.115: Compact does not imply Jordan measurable**

The Smith-Volterra-Cantor set is compact, but not Jordan measurable.

## Chapter 2

# Measurable Functions

### Definition 2.1: Measurable Functions

Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be measurable spaces. Then we call a map  $f : X \rightarrow Y$   **$\mathcal{A}$ - $\mathcal{B}$ -measurable**, or just **measurable**, if the preimage of every measurable set is again measurable, i.e.

$$B \in \mathcal{B} \implies f^{-1}(B) \in \mathcal{A}$$

Once again, note the similarities between this definition and the topological definition of a continuous function.

If the  $\sigma$ -algebra on one of the sets is supposed to be clear from context, many authors only specify one of the two sigma algebras. For example, for a function  $f : X \rightarrow \overline{\mathbb{R}}$ , many authors talk about  $\mathcal{A}$ -measurability when they implicitly mean  $\mathcal{A} - \mathcal{B}(\overline{\mathbb{R}})$ -measurability.

### Corollary 2.2

The composition of two measurable functions is again measurable.

**Lemma 2.3.** Let  $(X, \mathcal{A}), (Y, \mathcal{B})$  be measurable spaces and  $f : X \rightarrow Y$ . Let  $\mathcal{E} \subset \mathcal{B}$ . Then

$$f^{-1}(\langle \mathcal{E} \rangle_\sigma) = \langle f^{-1}(\mathcal{E}) \rangle_\sigma$$

### Corollary 2.4

Let  $\mathcal{E}$  be a base of  $\mathcal{B}$ , i.e.  $\langle \mathcal{E} \rangle_\sigma = \mathcal{B}$ . Then  $f : X \rightarrow Y$  is  $\mathcal{A}$ - $\mathcal{B}$  measurable if and only if

$$E \in \mathcal{E} \implies f^{-1}(E) \in \mathcal{A},$$

This means we don't need to check the preimage of every single set in  $\mathcal{B}$  to show that  $f$  is measurable - it suffices to check a base.

**Corollary 2.5.** Every continuous function between topological spaces is measurable in the corresponding Borel- $\sigma$ -algebras.

*Proof.* Let  $f : X \rightarrow Y$  be continuous. Then the preimage of every open set of  $Y$  is open in  $X$ , i.e. contained in the Borel  $\sigma$ -algebra on  $X$ , and the open sets of  $Y$  form a base of the Borel  $\sigma$ -algebra on  $Y$ .  $\square$

### Theorem 2.6: Simple criteria for measurability

Let  $(X, \mathcal{A})$  be a measurable space. Let  $f : X \rightarrow \overline{\mathbb{R}}$ . Then the following are equivalent:

1.  $f$  is  $\mathcal{A}$ - $\mathcal{B}(\overline{\mathbb{R}})$  measurable,
2.  $\forall c \in \mathbb{R} : \{f > c\} \in \mathcal{A}$ ,
3.  $\forall c \in \mathbb{R} : \{f \geq c\} \in \mathcal{A}$ ,
4.  $\forall c \in \mathbb{R} : \{f < c\} \in \mathcal{A}$ ,
5.  $\forall c \in \mathbb{R} : \{f \leq c\} \in \mathcal{A}$ ,

*Proof.* 1. The latter four conditions are equivalent to each other, since:

- (a)  $\{f \geq c\} = \bigcap_{k \in \mathbb{N}} \{f > c - \frac{1}{k}\}$ ,
- (b)  $\{f > c\} = \bigcup_{k \in \mathbb{N}} \{f \geq c + \frac{1}{k}\}$ ,
- (c)  $\{f < c\} = X \setminus \{f \geq c\}$ ,
- (d)  $\{f \leq c\} = X \setminus \{f > c\}$ .

2. The intervals  $[c, \infty]$  form a base of  $\mathcal{B}(\overline{\mathbb{R}})$ , since:

- (a)  $\{\infty\} = \bigcap_{k \in \mathbb{N}} [k, \infty]$
- (b)  $\{-\infty\} = \bigcap_{k \in \mathbb{N}} [-\infty, -k]$
- (c)  $(a, b) = [a, \infty] \setminus ([b, \infty] \cap [-\infty, a])$

$\square$

### Theorem 2.7

Let  $f, g : X \rightarrow \overline{\mathbb{R}}$  be  $\mathcal{A}$ - $\mathcal{B}(\overline{\mathbb{R}})$  measurable. Then the following sets are contained in  $\mathcal{A}$ :

1.  $\{f > g\}$ ,
2.  $\{f \geq g\}$ ,
3.  $\{f = g\}$ ,
4.  $\{f \neq g\}$ .

**Theorem 2.8.** Let  $f, g : X \rightarrow \overline{\mathbb{R}}$  be  $\mathcal{A}$ - $\mathcal{B}[\overline{\mathbb{R}}]$  measurable. Then the following functions are also  $\mathcal{A}$ -measurable:

1.  $cf$ , for all  $c \in \mathbb{R}$ ,
2.  $|f|^p$ , for all  $p \in \mathbb{R}_{>0}$ ,
3.  $f + g$ , assuming the sum is defined everywhere on  $X$ , i.e. there exists no  $x \in X$  such that  $f(x) = \infty$  and  $g(x) = -\infty$  or vice versa,
4.  $f \cdot g$ .

**Theorem 2.9.** Let  $(X, \mathcal{A})$  be a measurable space and  $\mathbb{1}_E : X \rightarrow \mathbb{R}$  be the indicator function of a set  $E \subset X$ . Then  $\mathbb{1}_E$  is  $\mathcal{A}$ - $\mathcal{B}[\mathbb{R}]$  measurable if and only if  $E \in \mathcal{A}$ .

*Proof.* We have  $\{1\} = (-\infty, 1) \cup (1, \infty) \in \mathcal{B}[\mathbb{R}]$  and  $\mathbb{1}_E^{-1}(\{1\}) = E$ , so  $E$  has to be in  $\mathcal{A}$  for  $\mathbb{1}_E$  to be measurable.

$E \in \mathcal{A}$  is also a sufficient condition for  $\mathbb{1}_E$  to be measurable, since the only other possible preimages are  $\emptyset \in \mathcal{A}$ ,  $X \in \mathcal{A}$ ,  $X \setminus E \in \mathcal{A}$  □

**Theorem 2.10.** Let  $(X, \mathcal{A})$  be a measurable space, let  $D \in \mathcal{A}$ , and let  $f_k : D \rightarrow \mathbb{R}$  be  $\mathcal{A}$  measurable. Then the following functions are  $\mathcal{A}$ -measurable:

1.  $\inf_{n \in \mathbb{N}} f_n$
2.  $\sup_{n \in \mathbb{N}} f_n$
3.  $\liminf_{n \rightarrow \infty} f_n$
4.  $\limsup_{n \rightarrow \infty} f_n$

*Proof.* For  $s \in \mathbb{R}$ , we have:

1.  $\left\{ \left( \inf_{n \in \mathbb{N}} f_n \right) \geq s \right\} = \bigcap_{k=1}^{\infty} \{f_k \geq s\} \in \mathcal{A}$
2.  $\left\{ \left( \sup_{n \in \mathbb{N}} f_n \right) \leq s \right\} = \bigcap_{k=1}^{\infty} \{f_k \leq s\} \in \mathcal{A}$

Therefore  $\inf_{n \in \mathbb{N}} f_n$  and  $\sup_{n \in \mathbb{N}} f_n$  are measurable. The same argument holds for sup and inf over subsets of  $\mathbb{N}$ . Therefore, the compositions

$$\liminf_{n \rightarrow \infty} f_n = \sup_{n \in \mathbb{N}} \left( \inf_{k \geq n} f_k \right)$$

and

$$\limsup_{n \rightarrow \infty} f_n = \inf_{n \in \mathbb{N}} \left( \sup_{k \geq n} f_k \right)$$

are also measurable. □

**Corollary 2.11**

Let  $f_n$  be a sequence of  $\mathcal{A}$ -measurable functions with a pointwise limit  $f$ . Then  $f$  is  $\mathcal{A}$ -measurable.

**Definition 2.12:  $\mu$ -measurable**

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $D \in \mathcal{A}$ . Then we call a function  $f : D \rightarrow \overline{\mathbb{R}}$   $\mu$ -measurable iff:

1.  $X \setminus D$  is a  $\mu$ -zero set, and
2.  $f$  is  $\mathcal{A}|_D$ -measurable.

**Lemma 2.13.** Let  $f : D \rightarrow \overline{\mathbb{R}}$  be  $\mu$ -measurable. Then the function

$$g(x) = \begin{cases} f(x) & x \in D \\ 0 & x \notin D \end{cases}$$

is  $\mathcal{A}$ -measurable.

**Lemma 2.14**

Let  $(X, \mathcal{A}, \mu)$  be a **complete** measure space. Let  $f$  be  $\mu$ -measurable. Then every function  $g$  such that  $g = f$   $\mu$ -almost everywhere is  $\mu$ -measurable.

**Theorem 2.15:  $\mu$ -measurable pointwise limits**

Let  $(X, \mathcal{A}, \mu)$  be a **complete** measure space. Let  $(f_k)_{k \in \mathbb{N}}$  be a family of  $\mu$ -measurable functions. Then if  $f_k$  converges pointwise to a function  $f$   $\mu$ -almost everywhere,  $f$  is  $\mu$ -measurable.

# Chapter 3

## Lebesgue Integration

### 3.1 Step Functions

**Definition 3.1.** Let  $(X, \mathcal{A})$  be a measurable space. Then we call a function  $f : Y \rightarrow \mathbb{R}$  a **step function** if it can be represented as a finite linear combination of indicator functions of sets in  $\mathcal{A}$ , i.e there exist  $\alpha_i \in \mathbb{R}$  and  $A_i \in \mathcal{A}$  such that:

$$f = \sum_{i \leq k} \alpha_i \cdot \mathbb{1}_{A_i}$$

**Proposition 3.2.** Step functions form a vector space over  $\mathbb{R}$ .

**Proposition 3.3.** Step functions can only take finitely different values.

**Proposition 3.4.** Every step function is (extensionally) equal to a step function over pairwise disjoint sets.

*Proof.* Let

$$s = \sum_{i \leq k} \alpha_i \cdot \mathbb{1}_{A_i}$$

Then if  $\beta_1, \dots, \beta_m$  are all of the finitely many different values that  $s$  can take,  $s$  is by definition equal to

$$t := \sum_{i \leq m} \beta_i \cdot \mathbb{1}_{\{s=\beta_i\}}$$

And since a function cannot take two values at once, the sets  $\{s = \beta_i\}$  are disjoint.  $\square$

**Lemma 3.5. (Sum of Step Functions):** Let  $s_1$  and  $s_2$  be step functions

$$s_1 = \sum_{i=0}^m \alpha_i \mathbb{1}_{A_i}, \quad s_2 = \sum_{j=0}^n \beta_j \mathbb{1}_{B_j}$$

defined on pairwise disjoint sets. Then we have:

$$s_1 + s_2 = \sum_{i=0}^m \sum_{j=0}^n (\alpha_i + \beta_j) \mathbb{1}_{A_i \cap B_j}$$

*Proof.* 1. Assume  $s_1$  and  $s_2$  are nonnegative step functions

$$s_1 = \sum_{i=0}^m \alpha_i \cdot \mathbb{1}_{A_i}, \quad s_2 = \sum_{j=0}^n \beta_j \cdot \mathbb{1}_{B_j}$$

defined on pairwise disjoint sets.

Then:

$$s_1(x) + s_2(x) = \sum_{i=0}^m \alpha_i \cdot \mathbb{1}_{A_i}(x) + \sum_{j=0}^n \beta_j \cdot \mathbb{1}_{B_j}(x)$$

Since the  $A_i$  and  $B_j$  are disjoint coverings of  $X$ , for every  $x \in X$ , there exists exactly one  $i$  such that  $x \in A_i$  and exactly one  $j$  such that  $x \in B_j$ . Therefore, every  $x$  contributes exactly one  $\alpha_i$  for  $x \in A_i$  and one  $\beta_j$  for  $x \in B_j$ , i.e.  $x \in A_i \cap B_j$  and  $s_1(x) + s_2(x) = \alpha_i + \beta_j$ . Since this is the only intersection which contains  $x$ , we can put everything together to get our desired formula:

$$s_1(x) + s_2(x) = \sum_{i=0}^m \sum_{j=0}^n (\alpha_i + \beta_j) \cdot \mathbb{1}_{A_i \cap B_j}(x)$$

□

The integral of an indicator function is already clear from our intuition: The points contained in the area under an indicator function  $\mathbb{1}_A$  are exactly the cartesian product of  $A$  with the interval  $[0, 1]$ , forming a "rectangle with gaps" whose side lengths are 1 and  $\mu(A)$ . Therefore the integral should be:

**Definition 3.6. (Lebesgue integral of an indicator function):**

$$\int_X \mathbb{1}_A d\mu = \mu(A)$$

The definition for the integral of a step function should follow naturally - a step function is just a finite sum of scaled characteristic functions, therefore the integral of a step function is the sum of the scaled integrals of the step functions.

**Definition 3.7. (Lebesgue integral of a step function):** Let  $D \subset X$ ,  $A_i \subset X$  pairwise disjoint, and  $\alpha_i \geq 0$ . Then:

$$\begin{aligned} \int_D s d\mu &= \int_D \left( \sum_{i \leq k} \alpha_i \cdot \mathbb{1}_{A_i} \right) d\mu \\ &:= \sum_{i \leq k} \left( \alpha_i \cdot \int_D \mathbb{1}_{A_i} d\mu \right) \\ &= \sum_{i \leq k} (\alpha_i \cdot \mu(D \cap A_i)) \end{aligned}$$

**Theorem 3.8. (Linearity of the step function integral):**

**Theorem 3.9. (Monotonicity of the step function integral):**

## 3.2 Defining the Lebesgue Integral

We can now use the step function integral to define the integral of more general functions:

### Theorem 3.10: Approximation via step functions

Let  $(X, \mathcal{A})$  be a measurable space. Let  $f : X \rightarrow \overline{\mathbb{R}}$  be a nonnegative  $\mathcal{A}$ -measurable function. Then there exists a monotonically increasing sequence  $s_n$  of nonnegative step functions whose pointwise limit is  $f$ .

*Construction 1.* For  $n \in \mathbb{N}$  and  $k \in \{0, \dots, n \cdot 2^n\}$ , we set:

$$F_{n,k} := \left\{ x \in X \mid \frac{k}{2^n} \leq f(x) < \frac{k+1}{2^n} \right\}.$$

Then

$$s_n(x) = \begin{cases} \frac{k}{2^n} & x \in F_{n,k} \\ 0 & \text{otherwise} \end{cases}$$

are step functions converging to  $f$ . □

*Construction 2.* Define  $s_0 = 0$ . Then we can inductively define

$$E_n := \left\{ s_{n-1} + \frac{1}{n} \leq f \right\}$$

and

$$s_n := s_{n-1} + \frac{1}{n} \cdot \mathbb{1}_{E_n},$$

meaning we have

$$s_n = \sum_{k=1}^n \frac{1}{k} \cdot \mathbb{1}_{E_k}.$$

We now need to show that this series converges pointwise to  $f$ . Let  $n \in \mathbb{N}$ .

1. Assume  $x \in E_n$ . Then, by definition,

$$f_n(x) = s_{n-1}(x) + \frac{1}{n} \leq f(x)$$

2. Assume  $x \notin E_n$ . Then, by induction, we have

$$f_n(x) = s_0(x) = 0 \leq f(x).$$

Therefore, we have  $f_0 \leq f_1 \leq \dots$  and  $f_n \leq f$  for all  $n$ , meaning that

$$\lim_{n \rightarrow \infty} f_n(x) \leq f(x).$$

If  $f(x) = \infty$ , then  $x \in E_n$  for every  $n$ , and we have

$$\lim_{n \rightarrow \infty} f_n(x) = \sum_{k=1}^{\infty} \frac{1}{k} = \infty$$

If  $f(x) < \infty$ , then there must be an infinite number of  $n \in \mathbb{N}$  such that  $f_{n-1}(x) > f_n(x) - \frac{1}{n}$ , implying

$$\lim_{n \rightarrow \infty} f_n(x) \geq f(x)$$

□

**Definition 3.11: Lebesgue integral of a positive function**

Let  $f : D \rightarrow [0, \infty]$  be  $\mathcal{A}$ -measurable. Then:

$$\int_D f \, d\mu = \sup_{\substack{s \text{ is a step function,} \\ 0 \leq s \leq f}} \left\{ \int_D s \, d\mu \right\}$$

**Definition 3.12. (Lebesgue integral of an arbitrary measurable function):** Let  $f$  be  $\mathcal{A}$ -measurable and

$$f^+ = f \cdot \mathbb{1}_{f \geq 0}, \quad f^- = -f \cdot \mathbb{1}_{f < 0}$$

Note that  $f^+ \geq 0$  and  $f^- > 0$ . Then, as long as  $f^+$  or  $f^-$  have a finite integral, we can define:

$$\int_D f \, d\mu = \int_D f^+ \, d\mu - \int_D f^- \, d\mu$$

**Corollary 3.13. (Integrating over subsets):**

$$\int_M f \, d\mu = \int_X f \cdot \mathbb{1}_M \, d\mu = \int_M f \, d\mu|_M$$

It is common to derive from this corollary a slight abuse of notation: Assume that  $f$  is not  $\mathcal{A}$ -measurable, but it is  $\mu$ -measurable, i.e.  $f : D \rightarrow \overline{\mathbb{R}}$  such that  $\mu(X \setminus D) = 0$  and  $f$  is  $\mathcal{A}|_D$ -measurable. Then it is common to implicitly expand the domain of  $D$  to  $X$  by setting  $f(x) = 0$  on  $X \setminus D$ , and to therefore write:

$$\int_X f \, d\mu := \int_D f \, d\mu$$

**Corollary 3.14. (Integrating over zero sets):** Let  $N$  be a set such that  $\mu(N) = 0$ . Then

$$\int_N f \, d\mu = 0.$$

**Definition 3.15: Integrable function**

We call a function  $f : X \rightarrow \overline{\mathbb{R}}$  **integrable** with regards to a measure  $\mu$  if it is  $\mu$ -measurable and

$$\int_X f \, d\mu \in \mathbb{R}$$

**Proposition 3.16. (Integrating with the counting measure):** Let  $X$  be an arbitrary set. Let  $\text{card}$  be the counting measure on  $\mathcal{P}(X)$ . Let  $f : X \rightarrow \overline{\mathbb{R}}$ . Then  $f$  is integrable with respect to  $\text{card}$  if and only if  $\sum_{x \in X} f(x)$  is absolutely convergent, and we have

$$\int_X f \, d\text{card} = \sum_{x \in X} f(x)$$

**Theorem 3.17: Monotonicity of the Lebesgue integral**

Let  $\mu$  be an outer measure on  $X$  such that  $f, g : X \rightarrow \overline{\mathbb{R}}$  are  $\mu$ -measurable. Then if  $f \leq g$   $\mu$ -almost everywhere and  $\int_X f \, d\mu > -\infty$ , then the integral of  $g$  exists and we have

$$\int_X f \, d\mu \leq \int_X g \, d\mu$$

*Proof.* 1. Assume  $f, g$  are nonnegative. Then if  $s$  is a step function such that  $s \leq f$ . Then if we define  $t := \mathbb{1}_{f \leq g} \cdot s$ , we have  $t \leq g$ . Furthermore, for all  $c \geq 0$ , we have:

$$\mu(\{s = c\}) = \mu(\{s = c\} \cap \{f \leq g\})$$

2.

□

**Theorem 3.18**

Let  $f, g : X \rightarrow \overline{\mathbb{R}}$  and let  $f$  be  $\mu$ -measurable. Then if  $g = f$   $\mu$ -almost everywhere, then  $g$  is  $\mu$ -measurable, and

$$\int_X g \, d\mu = \int_X f \, d\mu$$

as long as the right integral exists.

**Lemma 3.19. (Chebyshev Inequality):** Let  $f : X \rightarrow [0, \infty]$  be  $\mu$ -measurable with  $\int_X f \, d\mu < \infty$ . Let  $s \in (0, \infty]$ . Then:

$$\mu(\{x : f(x) \geq s\}) \leq \begin{cases} \frac{1}{s} \int_X f \, d\mu & s \in (0, \infty) \\ 0 & s = \infty \end{cases}$$

**Corollary 3.20.** Let  $f : X \rightarrow \overline{\mathbb{R}}$  be  $\mu$ -measurable. Then:

1. If  $\int_X f \, d\mu < \infty$ ,  $\{x : f(x) = \infty\}$  is a  $\mu$ -zero set.
2. If  $f \geq 0$  and  $\int_X f \, d\mu = 0$ ,  $\{x : f(x) > 0\}$  is a  $\mu$ -zero set.

**Theorem 3.21: Monotone Convergence Theorem**

Let  $f_n : X \rightarrow [0, \infty]$  be  $\mu$ -measurable such that  $f_i \leq f_{i+1}$ . Let  $\lim_{n \rightarrow \infty} f_n = f$ . Then

$$\int_X \lim_{n \rightarrow \infty} f_n \, d\mu = \lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \sup_{n \in \mathbb{N}} \int_X f_n \, d\mu$$

**Theorem 3.22: Linearity of the Lebesgue integral**

Let  $\mu$  be a measure on  $X$ . Let  $f, g : X \rightarrow \overline{\mathbb{R}}$  and  $\alpha, \beta \in \mathbb{R}$ . Then we have:

$$\int_X (\alpha f + \beta g) \, d\mu = \alpha \int_X f \, d\mu + \beta \int_X g \, d\mu$$

**Corollary 3.23.** Let  $f : X \rightarrow \overline{\mathbb{R}}$  be  $\mu$ -measurable. Then  $f$  is integrable if and only if  $|f|$  is integrable.

**Corollary 3.24.** Let  $f : X \rightarrow \overline{\mathbb{R}}$  be  $\mu$ -measurable. Then, assuming the integral of  $f$  exists, we have

$$\left| \int_X f \, d\mu \right| \leq \int_X |f| \, d\mu$$

**Corollary 3.25: Dominated Convergence Theorem**

Let  $g : X \rightarrow [0, \infty]$  be  $\mu$ -measurable such that  $|f| \leq g$   $\mu$ -almost everywhere. Then if

$$\int_X g \, d\mu < \infty,$$

$f$  is integrable.

### 3.3 Comparing Riemann Integration and Lebesgue Integration

**Theorem 3.26.** The Dirichlet function  $\mathbb{1}_{\mathbb{Q}}$  is Lebesgue integrable, but not Riemann integrable.

#### Theorem 3.27

A Lebesgue-integrable function is not necessarily “almost Riemann-integrable”. More formally, there exists a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  with the following properties:

1.  $f$  is Lebesgue-integrable.
2. There exists no Riemann-integrable function  $g$  such that  $f = g$  almost everywhere.

*Proof.* Consider the indicator function  $\mathbb{1}_{\text{SVC}}$  of the Smith-Volterra-Cantor set. □

#### Theorem 3.28

Let  $f : [a, b] \rightarrow \mathbb{R}$ . Then  $f$  is Riemann integrable if and only if it is the limit of a uniformly convergent sequence of step functions over Jordan measurable sets.

## 3.4 Convergence Theorems

### Theorem 3.29: Levi's Stronger Monotone Convergence Theorem

The monotone convergence theorem continues to hold if  $f_n \leq f_{n+1}$  and  $f_n \nearrow f$  only hold  $\mu$ -almost everywhere.

### Corollary 3.30: Exchange of Integrals and Sums

Let  $f_n$  be nonnegative and  $\mu$ -measurable. Then

$$\int_X \left( \sum_{n=1}^{\infty} f_n \right) d\mu = \sum_{n=1}^{\infty} \left( \int_X f_n d\mu \right)$$

### Theorem 3.31: Fatou's Lemma

Let  $(f_n)$  be a sequence of  $\mu$ -measurable functions  $X \rightarrow \overline{\mathbb{R}}$ . Let  $g \in \mathcal{L}^1$ . Then if we have  $f_n \geq g$   $\mu$ -almost everywhere for all  $n \in \mathbb{N}$ , we have

$$\int_X \left( \liminf_{n \rightarrow \infty} f_n \right) d\mu \leq \liminf_{n \rightarrow \infty} \int_X f_n d\mu$$

### Theorem 3.32: Lebesgue's Dominated Convergence Theorem

Let  $f, f_k$  be  $\mu$ -measurable functions such that  $f_k \rightarrow f$   $\mu$ -almost everywhere. Assume there exists a nonnegative function  $g \in \mathcal{L}^1$  such that  $\sup_k |f_k| \leq g$   $\mu$ -almost everywhere. Then we have  $f \in \mathcal{L}^1$  and

$$\int_X f d\mu = \lim_{k \rightarrow \infty} \int_X f_k d\mu$$

**Lemma 3.33. (Continuity of Parametrized Integrals):** Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $U \subset \mathbb{R}^n$  be open. Let  $f : U \times X \rightarrow \mathbb{R}$ . Then

$$\begin{aligned} \varphi : U &\rightarrow \mathbb{R} \\ \varphi(y) &= \int_X f(y, x) d\mu(x) \end{aligned}$$

is continuous if the following hold:

1.  $f(\cdot, x)$  is continuous in  $U$  for  $\mu$ -almost all  $x \in X$ ,
2.  $f(y, \cdot)$  is  $\mu$ -measurable for all  $y \in U$ ,
3. there exists a function  $g \in \mathcal{L}^1(\mu)$  such that for all  $y \in U$  and  $\mu$ -almost all  $x \in X$ , we have

$$\|f(y, x)\| \leq g(x)$$

**Theorem 3.34: Differentiation under the Integral Sign**

Let  $(X, \mathcal{A}, \mu)$  be a measure space,  $U \subset \mathbb{R}^n$  open, and  $f : U \times X \rightarrow \mathbb{R}$ . Then

$$\varphi(y) := \int_X f(y, x) d\mu(x)$$

is continuously differentiable with

$$\frac{\partial}{\partial y_i} \int_X f(y, x) d\mu(x) = \int_X \frac{\partial}{\partial y_i} f(y, x) d\mu(x)$$

if the following hold:

1.  $f(\cdot, x)$  is continuously differentiable in  $U$  for almost all  $x$ ,
2.  $f(y, \cdot)$  is  $\mu$ -measurable for almost all  $y \in U$ ,
3. There exists a function  $g \in \mathcal{L}^1(\mu)$  such that

$$\left| \frac{\partial}{\partial y_i} f(y, x) \right| \leq g(x)$$

for all  $y \in U$ ,  $\mu$ -almost all  $x \in X$  and  $i \in \{1, \dots, n\}$ .

**Theorem 3.35. (Absolute continuity of the Lebesgue integral):** Let  $f \in \mathcal{L}^1(\mu)$ . Let  $\varepsilon > 0$ . Then there exists a  $\delta > 0$  such that, for all  $A \in \mathcal{A}$ :

$$\mu(A) < \delta \implies \int_A |f| d\mu < \varepsilon$$

## 3.5 $L^p$ -Spaces

### Definition 3.36: $\mathcal{L}^p$

Let  $1 \leq p < \infty$ . We denote by  $\mathcal{L}^p = \mathcal{L}^p(X, \mathcal{A}, \mu)$  the set of all  $\mu$ -measurable functions  $f$  such that

$$\int_X |f|^p d\mu < \infty$$

### Definition 3.37: $L^p$ Norm

Let  $1 \leq p < \infty$  and  $f \in \mathcal{L}^p(X, \mathcal{A}, \mu)$ . We call

$$\|f\|_p = \sqrt[p]{\int_X |f|^p d\mu}$$

The  $L^p$ -Norm, or just  $p$ -Norm of  $f$ .

Note that, if we view integration as the natural "continuous analogue of summation"<sup>1</sup> and  $f$  as a vector with uncountably many components, this is a direct generalization of the familiar  $p$ -norm of vectors, such as the euclidean norm

$$\|\vec{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

### Theorem 3.38: $L^\infty$ -Norm

We have

$$\|f\|_\infty := \lim_{p \rightarrow \infty} \|f\|_p = \inf_{s > 0} \mu\{|f| > s\}$$

**Lemma 3.39. (Young Inequality):** Let  $p, q \in (1, \infty)$  such that  $\frac{1}{p} + \frac{1}{q} = 1$ . Then, for nonnegative  $a, b \geq 0$ , we have:

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}$$

*Proof.* From the concavity of the logarithm, we get:

$$\ln\left(\frac{a^p}{p} + \frac{b^q}{q}\right) \geq \frac{1}{p} \ln(a^p) + \frac{1}{q} \ln(b^q)$$

Which we can immediately simplify to get:

$$\frac{1}{p} \ln(a^p) + \frac{1}{q} \ln(b^q) = \ln(a) + \ln(b) = \ln(ab)$$

Exponentiating both sides gives us our desired result.  $\square$

<sup>1</sup>We can actually formalize this by going the other way - summation is just integration over a discrete space, using the counting measure as the canonical measure.

**Comment 3.40.** Keep in mind that:

$$\begin{aligned}\frac{1}{p} + \frac{1}{q} &= 1 \\ \iff \frac{1}{q} &= 1 - \frac{1}{p} \\ &= \frac{p-1}{p} \\ \iff q &= \frac{p}{p-1}\end{aligned}$$

Now, letting  $\frac{1}{\infty} = 0$ , we get:

**Corollary 3.41: Hölder Inequality**

Let  $p, q \in [1, \infty]$  such that  $\frac{1}{p} + \frac{1}{q} = 1$ . Let  $f \in \mathcal{L}^p$  and  $g \in \mathcal{L}^q$ . Then we have  $fg \in \mathcal{L}^1$  and, in particular,

$$\begin{aligned}\left| \int_X f(x) \cdot g(x) \, d\mu(x) \right| \\ \leq \|fg\|_1 \leq \|f\|_p \cdot \|g\|_q\end{aligned}$$

*Proof.*

$p = 1$ : In this case, we have  $q = \infty$ . We have

$$g(x) < \inf_{s>0} \{\mu(|g| > s) = 0\} = \|g\|_\infty$$

$\mu$ -almost everywhere, and thus

$$\begin{aligned}\int_X |f(x) \cdot g(x)| \, d\mu(x) &\leq \int_X |f(x)| \cdot \|g\|_\infty \, d\mu(x) \\ &= \int_X |f(x)| \, d\mu(x) \cdot \|g\|_\infty \\ &= \|f\|_1 \cdot \|g\|_\infty\end{aligned}$$

$p > 1$ : Assume that  $\|f\|_p \cdot \|g\|_q > 0$ . Then, applying the Young inequality

with  $a = \frac{|f(x)|}{\|f\|_p}$  and  $b = \frac{|g(x)|}{\|g\|_q}$ , we get

$$\begin{aligned}\frac{f(x)g(x)}{\|f\|_p \|g\|_q} &\leq \frac{|f(x)||g(x)|}{\|f\|_p \|g\|_q} \\ &\leq \frac{|f(x)|^p}{p\|f\|_p^p} + \frac{|g(x)|^q}{q\|g\|_q^q}\end{aligned}$$

Integrating both sides gives:

$$\begin{aligned}
 \frac{\|fg\|_1}{\|f\|_p\|g\|_q} &= \frac{1}{\|f\|_p\|g\|_q} \cdot \int_X f(x)g(x) \, d\mu(x) \\
 &\leq \frac{1}{\|f\|_p\|g\|_q} \cdot \int_X |f(x)||g(x)| \, d\mu(x) \\
 &\leq \frac{1}{p\|f\|_p^p} \cdot \int_X |f(x)|^p \, d\mu(x) + \frac{1}{q\|g\|_q^q} \cdot \int_X |g(x)|^q \, d\mu(x) \\
 &= \frac{1}{p\|f\|_p^p} \cdot \|f\|_p^p + \frac{1}{q\|g\|_q^q} \cdot \|g\|_q^q \\
 &= \frac{1}{p} + \frac{1}{q} \\
 &= 1
 \end{aligned}$$

Which gives us our desired inequality.  $\square$

**Lemma 3.42.**  $\mathcal{L}^p$  is closed under addition.

*Proof.* 1.  $p = 1$ : Follows from the triangle inequality on  $\mathbb{R}$ :

$$\begin{aligned}
 \|f + g\|_1 &= \int_X |f(x) + g(x)| \, d\mu(x) \\
 &\leq \int_X |f(x)| + |g(x)| \, d\mu(x) \\
 &= \int_X |f(x)| \, d\mu(x) + \int_X |g(x)| \, d\mu(x) \\
 &= \|f\|_1 + \|g\|_1
 \end{aligned}$$

2.  $p \in (1, \infty)$ : Since the function  $t \mapsto t^p$  is convex on  $[0, \infty)$ , we have:

$$\begin{aligned}
 |f + g|^p &= 2^p \left| \frac{f + g}{2} \right|^p \\
 &\leq 2^{p-1}(|f|^p + |g|^p)
 \end{aligned}$$

3.  $p = \infty$ : We have  $|f(x)| \leq \|f\|_\infty$  and  $|g(x)| \leq \|g\|_\infty$  for  $\mu$ -almost all  $x$ , which immediately gives us

$$|f(x) + g(x)| \leq |f(x)| + |g(x)| \leq \|f\|_\infty + \|g\|_\infty$$

$\mu$ -almost everywhere and thus

$$\|f + g\|_\infty = \inf_{s>0} \{ |f + g| \leq s \text{ } \mu\text{-almost everywhere} \} \leq \|f\|_\infty + \|g\|_\infty$$

$\square$

We now want to show that the  $L^p$  norm defines a seminorm, which we will then easily be able to turn into a proper norm.

**Theorem 3.43: Minkowski Inequality**

Let  $f, g \in \mathcal{L}^p$ . Then we have

$$\|f + g\|_p \leq \|f\|_p + \|g\|_p$$

*Proof.* We already showed the cases  $p = 1$  and  $p = \infty$  while proving  $\mathcal{L}^p$  is closed under addition. Thus, let  $p \in (1, \infty)$ . Then the triangle inequality on  $\mathbb{R}$  gives:

$$\begin{aligned} |f(x) + g(x)|^p &= |f(x) + g(x)| \cdot |f(x) + g(x)|^{p-1} \\ &\leq (|f(x)| + |g(x)|) \cdot |f(x) + g(x)|^{p-1} \\ &= |f(x)| |f(x) + g(x)|^{p-1} + |g(x)| |f(x) + g(x)|^{p-1} \end{aligned}$$

Applying the Hölder inequality with  $q = \frac{p}{p-1}$  gives:

$$\begin{aligned} &\int_X |f(x)| |f(x) + g(x)|^{p-1} d\mu(x) \\ &\leq \|f\|_p \cdot \| |f(x) + g(x)|^{p-1} \|_q \\ &= \|f\|_p \cdot \left( \int_X |f(x) + g(x)|^{(p-1)q} d\mu(x) \right)^{\frac{1}{q}} \\ &= \|f\|_p \cdot \left( \int_X |f(x) + g(x)|^p d\mu(x) \right)^{\frac{p-1}{p}} \\ &= \|f\|_p \cdot \|f + g\|_p^{p-1} \end{aligned}$$

Switching the roles of  $f$  and  $g$  similarly gives:

$$\int_X |g(x)| |f(x) + g(x)|^{p-1} d\mu(x) \leq \|g\|_p \cdot \|f + g\|_p^{p-1}$$

And putting everything together, we get:

$$\begin{aligned} \|f + g\|_p^p &= \int_X |f(x) + g(x)|^p d\mu(x) \\ &\leq \int_X |f(x)| |f(x) + g(x)|^{p-1} d\mu(x) + \int_X |g(x)| |f(x) + g(x)|^{p-1} d\mu(x) \\ &\leq \|f\|_p \cdot \|f + g\|_p^{p-1} + \|g\|_p \cdot \|f + g\|_p^{p-1} \\ &= \|f + g\|_p \cdot \|f + g\|_p^{p-1} \end{aligned}$$

From which we get our desired result by dividing out the factor of  $\|f + g\|_p^{p-1}$ .  $\square$

**Corollary 3.44**

The  $\mathcal{L}^p$ -norm is a seminorm on  $\mathcal{L}^p$ . In particular, we have:

1.  $\|f\|_p \geq 0$ , with  $\|f\|_p = 0$  iff.  $f = 0$   $\mu$ -almost everywhere
2.  $\|\lambda f\|_p = |\lambda| \cdot \|f\|_p$
3.  $\|f + g\|_p \leq \|f\|_p + \|g\|_p$

**Corollary 3.45:  $L^p$ -space**

Define  $L^p$ -space to be the quotient space of  $\mathcal{L}^p$  under  $\mu$ -almost-everywhere equality. Then  $\|\cdot\|_p$  forms a proper norm on  $L^p$ .

**Lemma 3.46.** Let  $1 \leq p < \infty$ . Let  $u_j \in L^p$  and

$$f_k = \sum_{j=1}^k u_j$$

Then, if  $\sum_{j=1}^{\infty} \|u_j\|_p < \infty$ , we have:

1. The pointwise limit  $f(x) = \lim_{k \rightarrow \infty} f_k(x)$  exists for  $\mu$ -almost all  $x \in X$ ,
2.  $f \in L^p$ ,
3.  $\|f - f_k\|_p \rightarrow 0$

**Lemma 3.47: Convergence in  $L^p$**

If we have  $f_k \rightarrow f$  in  $L^p$  Norm for  $1 \leq p \leq \infty$ , then there exists a subsequence  $f_{k_i}$  such that  $f_{k_i} \rightarrow f$  pointwise  $\mu$ -almost everywhere.

**Theorem 3.48: Riesz-Fischer**

For  $1 \leq p \leq \infty$ ,  $(L^p, \|\cdot\|_p)$  is a Banach space.

**Definition 3.49.** Let  $1 \leq p < \infty$ . Let the sequence  $(f_n) \subset L^p$  converge pointwise  $\mu$ -almost everywhere against the  $\mu$ -measurable function  $f$ . Let

$$v(A) := \limsup_{n \rightarrow \infty} \int_A |f_n|^p d\mu$$

Then we call  $f_k$  **uniformly integrable** iff, for any  $A \in \mathcal{A}$ , we have:

1.  $\forall \varepsilon > 0 : \exists \delta > 0 : \forall A \in \mathcal{A} : \mu(A) < \delta \implies v(A) < \varepsilon$
2.  $\forall \varepsilon > 0 : \exists E \in \mathcal{A} : \mu(E) < \infty, v(X \setminus E) < \varepsilon$

**Theorem 3.50: Vitali Convergence Theorem**

Let  $1 \leq p < \infty$ . Let the sequence  $(f_n) \subset L^p$  converge pointwise  $\mu$ -almost everywhere against the  $\mu$ -measurable function  $f$ . Then the following are equivalent:

1.  $f \in L^p$  and  $\|f_n - f\|_p \rightarrow 0$ ,
2.  $f_k$  is uniformly integrable.

**Definition 3.51.** Let  $\Omega \subset \mathbb{R}^n$  be open. Then the **support of a function**  $f : \Omega \rightarrow \mathbb{R}$  is the set

$$\text{spt} f := \overline{\{x \in \Omega : f(x) \neq 0\}}$$

The space of  $k$  times continuously differentiable functions with compact support in  $\Omega$  is denoted  $C_c^k(\Omega)$ .

**Theorem 3.52:  $C_c^0(\Omega)$  is dense in  $L^p(\Omega)$**

Let  $\Omega \subset \mathbb{R}^n$  be open. Let  $1 \leq p < \infty$ . Then, for every  $f \in L^p(\Omega)$ , there exists a sequence  $f_k$  in  $C_c^0(\Omega)$  such that  $\|f - f_k\|_p \rightarrow 0$ .

## 3.6 Density Functions

### Theorem 3.53

Let  $(X, \mathcal{A}, \mu)$  be a measure space. Let  $\theta : X \rightarrow \overline{\mathbb{R}}$  be nonnegative and  $\mu$ -measurable. Then the map

$$\begin{aligned} \nu : \mathcal{A} &\rightarrow \overline{\mathbb{R}} \\ A &\mapsto \int_A \theta \, d\mu \end{aligned}$$

is a measure, which we denote  $\mu_\theta$ . We call  $\theta$  the **density of  $\nu$  with respect to  $\mu$** .

**Corollary 3.54.** The following hold for  $\mu_\theta$ :

1.  $\mu(A) = 0$  implies  $\mu_\theta(A) = 0$ .
2. For every nonnegative  $\mu$ -measurable function  $f$ , we have

$$\int_X f \, d\mu_\theta = \int_X f \cdot \theta \, d\mu.$$

3.  $\theta$  is unique up to equality  $\mu$ -almost everywhere.

### Definition 3.55

Let  $\mu$  and  $\nu$  be measures on  $(X, \mathcal{A})$ . Then we call  $\nu$  **absolutely continuous with respect to  $\mu$** , which we denote  $\nu \ll \mu$ , if  $\mu(A) = 0$  implies  $\nu(A) = 0$ .

**Lemma 3.56.** Let  $\sigma$  and  $\nu$  be finite measures on  $(X, \mathcal{A})$  such that  $\nu(A) \leq \sigma(A)$  for every  $A \in \mathcal{A}$ . Then there exists a density function  $\theta$  such that  $\nu = \sigma_\theta$ .

This lemma is actually a special case of the significantly more general Riesz representation theorem for Hilbert spaces, which I sadly do not have the time to go into at this point (but look it up, it's really neat!).

**Lemma 3.57.** Let  $\mu, \nu$  be measures on  $(X, \mathcal{A})$ . Let  $\sigma := \mu + \nu$ . Then if  $f \in \mathcal{L}^*(\sigma)$ , then  $f \in \mathcal{L}^*(\mu)$  and  $f \in \mathcal{L}^*(\nu)$  and we have

$$\int_X f \, d\sigma = \int_X f \, d\mu + \int_X f \, d\nu$$

**Theorem 3.58. (Mini-Radon-Nikodym):** Let  $\mu, \nu$  be finite measures on  $(X, \mathcal{A})$  such that  $\nu \ll \mu$ . Then there exists a density function  $\theta \in L^1(\mu)$  such that  $\nu = \mu_\theta$ .

In this case,  $\theta$  is sometimes also called the **Radon-Nikodym derivative of  $\nu$  with respect to  $\mu$**  and denoted  $\frac{d\nu}{d\mu}$ .

**Theorem 3.59.** Let  $\mu$  and  $\nu$  be finite measures on  $(X, \mathcal{A})$ . Then the following are equivalent:

1.  $\nu \ll \mu$ ,
2. There exists a density function  $\theta \in L^1(\mu)$  such that  $\nu = \mu_\theta$ ,
3. For all  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that  $\mu(A) < \delta$  implies  $\nu(A) < \varepsilon$ .

*Proof.* (i)  $\implies$  (ii) is our mini-Radon-Nikodym theorem. (ii)  $\implies$  (iii) follows from absolute continuity of the Lebesgue integral. (iii)  $\implies$  (i) follows immediately from the definition of  $\ll$ .  $\square$

#### Theorem 3.60: Radon-Nikodym

Let  $\mu, \nu$  be  $\sigma$ -finite measures on  $(X, \mathcal{A})$  such that  $\nu \ll \mu$ . Then there exists a density function  $\theta \in L^1(\mu)$  such that  $\nu = \mu_\theta$ , i.e.

$$\int_X f \, d\nu = \int_X f \cdot \theta \, d\mu$$

#### Definition 3.61

Let  $\mu$  and  $\nu$  be measures on  $(X, \mathcal{A})$ . Then we call  $\mu$  and  $\nu$  **singular with respect to each other**, which we denote  $\mu \perp \nu$ , if there exists a set  $M \in \mathcal{A}$  such that

$$\mu(M) = \nu(X \setminus M) = 0.$$

#### Theorem 3.62: Lebesgue's decomposition theorem

Let  $\mu$  and  $\nu$  be measures on  $(X, \mathcal{A})$ , and let  $\nu$  be  $\sigma$ -finite. Then there exists a unique decomposition  $\nu = \nu_a + \nu_s$  such that  $\nu_a \ll \mu$  and  $\nu_s \perp \mu$ .

# Chapter 4

## Integration in $\mathbb{R}^n$

### 4.1 Products of Measure Spaces

In this section, we will be interested in finding a canonical  $\sigma$ -Algebra on cartesian products of measurable spaces. Category theory tells us the correct solution directly: The product  $\sigma$ -algebra of an indexed family of  $\sigma$ -algebras should be the sigma algebra fulfilling the universal property of product objects.

We are working in the category of measurable spaces, where morphisms are measurable functions. Therefore, our definition simply amounts to defining product sigma algebras as the  $\sigma$ -algebras induced by the projection functions, which is in turn the smallest  $\sigma$ -algebra such that all projections are measurable.

#### Definition 4.1: Product $\sigma$ -Algebra

Let  $(X_i, \mathcal{A}_i)_{i \in I}$  be a family of measurable spaces. Then the product  $\sigma$ -algebra on the cartesian product  $\prod_{i \in I} X_i$  is the smallest  $\sigma$ -Algebra  $\bigotimes_{i \in I} \mathcal{A}_i$  such that every projection  $\pi_j$  is  $\bigotimes_{i \in I} \mathcal{A}_i - \mathcal{A}_j$ -measurable.

Explicitly, this means we have:

$$\bigotimes_{i \in I} \mathcal{A}_i = \left\langle \bigcup_{i \in I} \pi_i^{-1}(\mathcal{A}_i) \right\rangle_{\sigma}$$

Note that this is also entirely analogous to the definition of the product topology, which is the coarsest (i.e. "smallest") topology such that projections are continuous.

If the cardinality of  $I$  is a small finite number, we may sometimes write out the product  $\sigma$ -algebra as:

$$\bigotimes_{i \in I} \mathcal{A}_i = \mathcal{A}_1 \otimes \mathcal{A}_2 \otimes \dots$$

#### Theorem 4.2

Let  $(X, \mathcal{A})$  and  $(Y_i, \mathcal{B}_i)$  be measurable spaces. Let  $g : X \rightarrow \prod_{i \in I} Y_i$ . Then  $g$  is  $\mathcal{A} - \bigotimes_{i \in I} \mathcal{B}_i$ -measurable if and only if all maps  $\pi_i \circ g$  are  $\mathcal{A} - \mathcal{B}_i$ -measurable.

*Proof.* 1. Assume  $g$  is measurable. We know that  $\pi_i$  are measurable by definition, and we know that compositions of measurable functions are measurable. Therefore,  $\pi_i \circ g$  are measurable.

2. Assume  $\pi_i \circ g$  are measurable. Then we have:

$$\begin{aligned} g^{-1}\left(\bigcup_{i \in I} \pi_i^{-1}(\mathcal{A}_i)\right) &= \bigcup_{i \in I} g^{-1}(\pi_i^{-1}(\mathcal{A}_i)) \\ &= \bigcup_{i \in I} (\pi_i \circ g)^{-1}(\mathcal{A}_i) \subset \mathcal{A} \end{aligned}$$

Where the last step follows from the measurability of  $\pi_i \circ g$ . Therefore, since  $\bigcup_{i \in I} \pi_i^{-1}(\mathcal{A}_i)$  forms a basis of  $\bigotimes_{i \in I} \mathcal{A}_i$ , we have that  $g$  is measurable.

□

As a corollary, we get that our product space actually fulfills the universal property of product objects:

**Corollary 4.3.** Let  $(X, \mathcal{A})$  and  $(Y_i, \mathcal{B}_i)_{i \in I}$  be measurable spaces. Let  $g_i : X \rightarrow Y_i$  be a family of  $\mathcal{A} - \mathcal{B}_i$ -measurable functions. Then there exists a  $\mathcal{A} - \mathcal{B}_I$ -measurable function  $g : X \rightarrow \prod_{i \in I} Y_i$  such that, for each  $i \in I$ ,  $g_i = \pi_i \circ g$ .

*Proof.* Define  $g$  as the cartesian product of all  $g_i$ :

$$\begin{aligned} g : I &\rightarrow \left( X \rightarrow \prod_{i \in I} Y_i \right) \\ i &\mapsto g_i \end{aligned}$$

Then our previous lemma tells us that, since all  $g_i = \pi_i \circ g$  are measurable,  $g$  is measurable as well.

□

**Corollary 4.4.** Let  $(X_i, \mathcal{A}_i)_{i \in I}$  be measurable spaces, and let  $J \subset I$  be nonempty. Then the projection  $\pi_J$  is  $\bigotimes_{i \in I} \mathcal{A}_i - \bigotimes_{j \in J} \mathcal{A}_j$ -measurable.

**Definition 4.5. (Embedding into a larger product space):** Let  $p_j \in X_j$ . Then we define the **embedding of  $\prod_{i \in I \setminus j} X_i$  into  $\prod_{i \in I} X_i$  by  $x_j$**  to be the map  $e_{x_j}$  which takes a product  $(x_i)_{i \in I \setminus j}$  and "adds in  $x_j$  as a new component", i.e. it maps  $(x_i)_{i \in I \setminus j}$  to the product  $(x'_i)_{i \in I}$  with  $x'_j = x_j$  and  $(x'_i)_{i \in I \setminus j} = (x_i)_{i \in I \setminus j}$ .

**Definition 4.6: Cuts of subsets**

Let  $M \subseteq \prod_{i \in I} X_i$ , and let  $x_j \in X_j$ . Then we define the **cut of  $M$  through  $x_j$**  to be the set  $M^{x_j} = e_{x_j}^{-1}(M)$ .

More explicitly,  $M^{x_j}$  is the set such that for every  $y \in M$ , we have  $y|_{I \setminus \{j\}} \in M^{x_j}$  if and only if  $y_j = x_j$ .

**Corollary 4.7.** Let  $(X_i, \mathcal{A}_i)_{i \in I}$  be measurable spaces, let  $M \in \bigotimes_{i \in I} \mathcal{A}_i$ , and let

$x_j \in X_j$  for  $j \in I$ . Then we have

$$M^{x_j} \in \bigotimes_{i \in I \setminus j} \mathcal{A}_i$$

*Proof.* Let  $x_j \in X_j, j \in I$ . We want to show that  $e_{x_j}$  is measurable, which would follow immediately if we could show that all projections  $\pi_i^I \circ e_{x_j}$  are measurable.

1. Let  $i \in I \setminus j$ . Then  $\pi_i^I \circ e_{x_j} = \pi_i^{I \setminus j}$ , which is measurable.
2. Let  $i = j$ . Then  $\pi_i^I \circ e_{x_j} = \pi_j^I \circ e_{x_j}$ , which is just the constant map onto  $x_j$ , which is measurable.

Since all projections are measurable,  $e_{x_j}$  itself must also be measurable. Therefore, since  $M$  is measurable, the preimage  $M^{x_j} = e_{x_j}^{-1}(M)$  must also be measurable.  $\square$

Note that, in particular, this tells us that if  $(X_1, \mathcal{A}_1)$  and  $(X_2, \mathcal{A}_2)$  are measurable spaces, then for any  $M \in \mathcal{A}_1 \otimes \mathcal{A}_2$ ,  $x_1 \in X_1$ , and  $x_2 \in X_2$ , we have

$$M^{x_2} = \{x \in X_1 \mid (x, x_2) \in M\} \in \mathcal{A}_1$$

and

$$M^{x_1} = \{x \in X_2 \mid (x_1, x) \in M\} \in \mathcal{A}_2$$

#### Theorem 4.8: A basis of the product $\sigma$ -Algebra

Let  $(X_i, \mathcal{A}_i)_{i \in I}$  be a family of measurable spaces. For each  $i \in I$ , let  $\mathcal{E}_i$  be a basis of  $\mathcal{A}_i$ . Then we have

$$\bigotimes_{i \in I} \mathcal{A}_i = \left\langle \bigcup_{i \in I} \pi_i^{-1}(\mathcal{E}_i) \right\rangle_{\sigma}.$$

*Proof.* Recall that, for any map  $f$  and any system of subsets  $\mathcal{S} \subset \mathcal{A}$  of a  $\sigma$ -algebra  $\mathcal{A}$ , we have

$$f^{-1}(\langle \mathcal{S} \rangle_{\sigma}) = \langle f^{-1}(\mathcal{S}) \rangle_{\sigma}$$

and

$$\left\langle \bigcup_{i \in I} \mathcal{S}_i \right\rangle_{\sigma} = \left\langle \bigcup_{i \in I} \langle \mathcal{S}_i \rangle_{\sigma} \right\rangle_{\sigma},$$

which gives us:

$$\begin{aligned}\bigotimes_{i \in I} \mathcal{A}_i &= \left\langle \bigcup_{i \in I} \pi_i^{-1}(\mathcal{A}_i) \right\rangle_{\sigma} \\ &= \left\langle \bigcup_{i \in I} \pi_i^{-1}(\langle \mathcal{E}_i \rangle_{\sigma}) \right\rangle_{\sigma} \\ &= \left\langle \bigcup_{i \in I} \langle \pi_i^{-1}(\mathcal{E}_i) \rangle_{\sigma} \right\rangle_{\sigma} \\ &= \left\langle \bigcup_{i \in I} \pi_i^{-1}(\mathcal{E}_i) \right\rangle_{\sigma}\end{aligned}$$

□

If  $I$  is finite, we get the following representation of the product  $\sigma$ -algebra:

**Corollary 4.9.** For each  $i \in \{1, \dots, n\}$ , let  $(X_i, \mathcal{A}_i)$  be a measurable space such that  $\mathcal{E}_i$  is a basis of  $\mathcal{A}_i$  and such that  $X_i$  can be represented as a countable union of basis elements  $E_i^k \in \mathcal{E}_i$ . Then

$$\mathcal{Q}_0 := \left\{ \prod_{i=1}^n E_i \mid E_i \in \mathcal{E}_i \right\}$$

is a basis of  $\bigotimes_{i=1}^n \mathcal{A}_i$ .

*Proof.* We need to show  $\langle \mathcal{Q}_0 \rangle_{\sigma} = \bigotimes_{i=1}^n \mathcal{A}_i$ .

1. Let  $E_i \in \mathcal{E}_i$ . We know that

$$\pi_i^{-1}(E_i) = E_i \times \prod_{j \neq i} X_j,$$

which gives us the following representation for the basis  $\mathcal{Q}$  we just established in the last theory:

$$\mathcal{Q} = \bigcup_{i=1}^n \pi_i^{-1}(\mathcal{E}_i) = \bigcup_{i=1}^n \left\{ E_i \times \prod_{j \neq i} X_j \mid E_i \in \mathcal{E}_i \right\}$$

Furthermore,  $\cap$ -stability of  $\sigma$ -algebras gives us that any element

$$\prod_{i=1}^n E_i = \bigcap_{i=1}^n \pi_i^{-1}(E_i)$$

of  $\mathcal{Q}_0$  must be contained in  $\langle \mathcal{Q} \rangle_{\sigma}$ .

Therefore, we have

$$\text{t } \langle \mathcal{Q}_0 \rangle_{\sigma} \subset \langle \mathcal{Q} \rangle_{\sigma} = \bigotimes_{i=1}^n \mathcal{A}_i$$

2. Now, let  $E_i \in \mathcal{E}_i$  and  $k \in \mathbb{N}$ . By definition of  $\mathcal{Q}_0$ , we have

$$E_i \times \prod_{j \neq i} E_j^k \in \mathcal{Q}_0,$$

which gives us:

$$\pi_i^{-1}(E_i) = \bigcup_{k=1}^{\infty} \left( E_i \times \prod_{j \neq i} E_j^k \right) \in \langle \mathcal{Q}_0 \rangle_{\sigma},$$

and by extension:

$$\mathcal{Q} = \bigcup_{i=1}^n \pi_i^{-1}(E_i) \subset \langle \mathcal{Q}_0 \rangle_{\sigma}$$

which means we also have

$$\bigotimes_{i=1}^n \mathcal{A}_i = \langle \mathcal{Q} \rangle_{\sigma} \subset \sigma(\mathcal{Q}_0).$$

□

**Corollary 4.10.** The Borel  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R}^n)$  is the product of the Borel  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R})$ , i.e. we have:

$$\mathcal{B}(\mathbb{R}^n) = \bigotimes_{i=1}^n \mathcal{B}(\mathbb{R})$$

*Proof.* We know that the set  $\mathcal{I}$  of real intervals forms a basis of  $\mathcal{B}(\mathbb{R})$ . Given this basis, the set  $\mathcal{Q}_0$  is exactly the set of  $n$ -dimensional real cuboids, which forms a basis of  $\mathcal{B}(\mathbb{R}^n)$ . Therefore, we have:

$$\mathcal{B}(\mathbb{R}^n) = \langle \mathcal{Q}_0 \rangle_{\sigma} = \bigotimes_{i=1}^n \mathcal{B}(\mathbb{R})$$

□

For the final part of this chapter, we establish some notation for commonly encountered subsets of product  $\sigma$ -algebras.

**Definition 4.11.** Let  $I$  be a set. Then we denote by  $\mathcal{P}_0(I)$  the finite, nonempty subsets of  $I$ .

**Definition 4.12: Measurable Cuboids**

Let  $(X_i, \mathcal{A}_i)_{i \in I}$  be a family of measurable spaces. Then we define the set of **measurable cuboids** in  $\prod_{i \in I} X_i$  as:

$$\mathcal{Q} := \bigcup_{J \in \mathcal{P}_0(I)} \left\{ \prod_{j \in J} A_j \times \prod_{i \in I \setminus J} X_i \mid A_j \in \mathcal{A}_j \right\},$$

i.e. a finite number of the sides of our cuboid can be arbitrary measurable sets, and the remaining sides must correspond to the remaining domain sets  $(X_i)_{i \in I \setminus J}$ .

**Definition 4.13: Cylindrical Sets**

Let  $(X_i, \mathcal{A}_i)_{i \in I}$  be a family of measurable spaces. Then we define the set of **cylinder sets** in  $\prod_{i \in I} X_i$  as:

$$\begin{aligned} \mathcal{Z} &:= \bigcup_{J \in \mathcal{P}_0(I)} \left\{ A_J \times \prod_{i \in I \setminus J} X_i \mid A_J \in \bigotimes_{j \in J} \mathcal{A}_j \right\} \\ &= \bigcup_{J \in \mathcal{P}_0(I)} \pi_J^{-1} \left( \bigotimes_{j \in J} \mathcal{A}_j \right), \end{aligned}$$

i.e. the base of the cylinder is a measurable set  $A_J \subset \prod_{i \in I} X_i$  and all other sides are once again the remaining domain sets  $(X_i)_{i \in I \setminus J}$ .

Note that if we view an everyday geometric cylinder  $Z \subset \mathbb{R}^3$  as the product of the circle at its base with a finite real interval, it doesn't actually qualify as a cylindrical set in  $\mathbb{R}^3$ , since it would have to be a product of the base circle with all of  $\mathbb{R}$ , i.e. only "infinite" cylinders immediately qualify.

By technicality, normal cylinders are still cylindrical sets though, since we can set  $J = I$  and have view the entire cylinder as the base, without adding any additional "sides".

Therefore, this definition is not very useful if  $I$  is finite, since every measurable set is a cylindrical with itself as its base and every cylindrical set is a product of measurable sets, i.e. measurable.

**Theorem 4.14**

Let  $(X_i, \mathcal{A}_i)_{i \in I}$  be a family of measurable spaces. Then the measurable cuboids in  $\prod_{i \in I} X_i$  and the cylindrical sets in  $\prod_{i \in I} X_i$  form bases of  $\bigotimes_{i \in I} \mathcal{A}_i$ .

*Proof.* We will show  $\mathcal{Q} \subset \mathcal{Z} \subset \bigotimes_{i \in I} \mathcal{A}_i \subset \langle \mathcal{Q} \rangle_\sigma$ , which implies  $\langle \mathcal{Q} \rangle_\sigma \subset \langle \mathcal{Z} \rangle_\sigma \subset \bigotimes_{i \in I} \mathcal{A}_i \subset \langle \mathcal{Q} \rangle_\sigma$ , proving the theorem.

1.  $\mathcal{Q} \subset \mathcal{Z}$ : Let  $A \in \mathcal{Q}$ . Then  $A$  is given as

$$\begin{aligned} A &= \prod_{j \in J}^n A_j \times \prod_{i \notin J} X_i \\ &= \pi_J^{-1} \left( \prod_{j \in J} A_j \right). \end{aligned}$$

Therefore, since  $\prod_{j \in J} A_j \in \bigotimes_{j \in J} \mathcal{A}_j$ ,  $A$  is a cylinder set.

2.  $\mathcal{Z} \subset \bigotimes_{i \in I} \mathcal{A}_i$ : Recall that  $\pi_J$  is  $\bigotimes_{i \in I} \mathcal{A}_i - \mathcal{A}_J$ -measurable. Therefore, for any  $J \in \mathcal{P}_0(I)$ , we have that  $\pi_J^{-1}(\mathcal{A}_J) \subset \bigotimes_{i \in I} \mathcal{A}_i$ , which means any cylinder set is contained in  $\bigotimes_{i \in I} \mathcal{A}_i$ .

3.  $\bigotimes_{i \in I} \mathcal{A}_i \subset \langle \mathcal{Q} \rangle_\sigma$ : For any  $i \in I$  and  $A_i \in \mathcal{A}_i$ , we have:

$$\pi_i^{-1}(A_i) = A_i \times \prod_{j \neq i} X_j \in \mathcal{Q},$$

i.e.  $\pi_i^{-1}(\mathcal{A}_i) \subset \mathcal{Q}$  for all  $i \in I$ , which means we also have:

$$\bigcup_{i \in I} \pi_i^{-1}(\mathcal{A}_i) \subset \mathcal{Q}.$$

By definition of  $\bigotimes_{i \in I} \mathcal{A}_i$ , we get:

$$\bigotimes_{i \in I} \mathcal{A}_i = \left\langle \bigcup_{i \in I} \pi_i^{-1}(\mathcal{A}_i) \right\rangle_\sigma \subset \langle \mathcal{Q} \rangle_\sigma.$$

□

## 4.2 Product Measures and Fubini's Theorem

In this section, given two measure spaces  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$ , we want to derive a canonical measure  $\mu \otimes \nu$  on  $\mathcal{A} \otimes \mathcal{B}$ . Our intuition already tells us what the measure of a cuboid should be: Given a cuboid

$$Q = A \times B \subset X \times Y,$$

where  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$ , we want the measure of the cuboid to be the product of the measures of its sides:

$$(\mu \otimes \nu)(Q) = \mu(A) \cdot \nu(B)$$

Recall that, for any  $M \in \mathcal{A} \otimes \mathcal{B}$ , the cuts through  $M$  by  $x \in X$  and  $y \in Y$  are simply defined as

$$\begin{aligned} M^x &= \{y \in Y \mid (x, y) \in M\}, \\ M^y &= \{x \in X \mid (x, y) \in M\}. \end{aligned}$$

### Theorem 4.15

Let  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$  be  $\sigma$ -finite measure spaces. Let  $M \in \mathcal{A} \otimes \mathcal{B}$ . Then:

1. The function  $x \mapsto \nu(M^x)$  is  $\mathcal{A}$ -measurable,
2. The function  $y \mapsto \mu(M^y)$  is  $\mathcal{B}$ -measurable,
3. We have

$$\int_X \nu(M^x) d\mu(x) = \int_Y \mu(M^y) d\nu(y)$$

*Proof.* Let  $\mathcal{S}$  be the system of all sets  $M \in \mathcal{A} \otimes \mathcal{B}$  for which this theorem holds.

Let  $\mathcal{Q}$  be the system of all measurable cuboids in  $X \times Y$ , which we have just established are simply given by  $Q = A \times B$  such that  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$ .

Let  $\mathcal{F}$  be the system of finite unions of cuboids (which by our definition of cuboids inherently includes complements of cuboids). Since we count  $X \times Y$  as a cuboid,  $\mathcal{F}$  forms a set algebra.

We will show that  $\mathcal{S}$  is a monotone class containing  $\mathcal{F}$ , which will give us

$$\mathcal{A} \otimes \mathcal{B} = \langle \mathcal{Q} \rangle_\sigma = \langle \mathcal{F} \rangle_\sigma = \langle \mathcal{F} \rangle_{\mathcal{M}} \subseteq \mathcal{S}$$

since  $\mathcal{S} \subseteq \mathcal{A} \otimes \mathcal{B}$  by definition, this implies  $\mathcal{A} \otimes \mathcal{B} = \mathcal{S}$ .

The proof the  $\mathcal{S}$  is a monotone class is however left as an exercise to the reader :) □

**Theorem 4.16: Product measure**

Let  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$  be  $\sigma$ -finite measure spaces. Then there exists exactly one measure  $\mu \otimes \nu$  on  $\mathcal{A} \otimes \mathcal{B}$  such that for any  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$ , we have

$$(\mu \otimes \nu)(A \times B) = \mu(A)\nu(B)$$

This measure is given by

$$(\mu \otimes \nu)(M) = \int_X \nu(M^x) d\mu(x) = \int_Y \mu(M^y) d\nu(y)$$

**Corollary 4.17: Cavalieri's Principle**

Let  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$  be  $\sigma$ -finite measure spaces. Then, for all  $M \in \mathcal{A} \otimes \mathcal{B}$ , we have

$$\begin{aligned} (\mu \otimes \nu)(M) &= \int_X \nu(M^x) d\mu(x) \\ &= \int_X \int_Y \chi_M(x, y) d\nu(y) d\mu(x) \\ &= \int_Y \mu(M^y) d\nu(y) \\ &= \int_Y \int_X \chi_M(x, y) d\mu(x) d\nu(y) \end{aligned}$$

**Corollary 4.18.** Let  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$  be  $\sigma$ -finite measure spaces. Then, for any  $M \in \mathcal{A} \otimes \mathcal{B}$ , the following are equivalent:

1.  $(\mu \otimes \nu)(M) = 0$ ,
2.  $\mu(M^y) = 0$   $\nu$ -almost everywhere,
3.  $\nu(M^x) = 0$   $\mu$ -almost everywhere.

**Lemma 4.19.** Let  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$  be  $\sigma$ -finite measure spaces. Let  $f : X \times Y \rightarrow \mathbb{R}$  be  $\mathcal{A} \otimes \mathcal{B}$  measurable. Then the function  $f(x, \cdot) : Y \rightarrow \mathbb{R}$  is  $\mathcal{B}$ -measurable for all  $x \in X$ .

**Theorem 4.20: Fubini's Theorem**

Let  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$  be  $\sigma$ -finite measure spaces. Let  $f \in \mathcal{L}^*(\mu \otimes \nu)$ . Then:

$$\begin{aligned} \int_{X \times Y} f d(\mu \otimes \nu) &= \int_X \int_Y f(x, y) d\nu(y) d\mu(x) \\ &= \int_Y \int_X f(x, y) d\mu(x) d\nu(y) \end{aligned}$$

**Corollary 4.21.** The volume of the open ball

$$B_r^n(0) = \{\vec{x} \in \mathbb{R}^n \mid \|\vec{x}\|_2 < r\}.$$

is given by:

$$\lambda^n(B_r^n(0)) = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} \cdot r^n = \begin{cases} \frac{\pi^k}{k!} \cdot r^n & n = 2k \\ \frac{\pi^k}{\prod_{i=1}^k \frac{1}{2} + k - i} \cdot r^n & n = 2k + 1 \end{cases}$$

In particular:

1. The 1-dimensional unit ball is simply the interval  $(-1, 1)$  and thus has length 2,
2. The 2-dimensional unit ball is a circle of radius 1 and has area  $\pi$
3. The 3-dimensional unit ball has volume  $\frac{4\pi}{3}$
4. The 4-dimensional unit ball has volume  $\frac{\pi^2}{2}$
5. The 5-dimensional unit ball has volume  $\frac{8\pi^2}{15}$
6. ...

*Proof.* By Cavalieri's principle and the linear transformation formula for the Lebesgue measure, we get:

$$\begin{aligned} \lambda^n(B_r^n(0)) &= \int_{\mathbb{R}} \lambda^{n-1}(B_r^n(0)^y) dy \\ &= \int_{(-r, r)} \lambda^{n-1}(B_r^n(0)^y) dy \\ &= \int_{-1}^1 = \det \left( \sqrt{r^2 - y^2} \cdot I_n \right) \cdot \lambda^{n-1} \{x \in \mathbb{R}^{n-1} \mid \|x\|_2 < 1\} dy \\ &= r^n \cdot \int_{-1}^1 \left( \sqrt{r^2 - y^2} \right)^{n-1} \cdot \lambda^{n-1} \{x \in \mathbb{R}^{n-1} \mid \|x\|_2 < 1\} dy \\ &= r^n \cdot \lambda^{n-1}(B_1^{n-1}(0)) \cdot \int_{-1}^1 \left( \sqrt{1 - y^2} \right)^{n-1} dy \end{aligned}$$

We can thus get our result by solving the integral

$$I_n := \int_{-1}^1 \left( \sqrt{1 - y^2} \right)^{n-1} dy$$

Substituting  $\cos(\theta) = y$  gives:

$$\begin{aligned} I_n &= \int_{-1}^1 \left( \sqrt{1 - y^2} \right)^{n-1} dy \\ &= \int_{-1}^1 \left( \sqrt{1 - \cos(\theta)^2} \right)^{n-1} \cdot \sin(\theta) d\theta \\ &= \int_0^\pi \sin^n(\theta) d\theta \end{aligned}$$

Now, partial integration gives:

$$\begin{aligned}
 I_n &= \int_0^\pi \sin^n(\theta) d\theta \\
 &= \int_0^\pi \sin(\theta) \cdot \sin^{n-1}(\theta) d\theta \\
 &= \left[ -\cos(\theta) \cdot \sin^{n-1}(\theta) \right]_0^\pi + (n-1) \cdot \int_0^\pi \cos^2(\theta) \cdot \sin^{n-2}(\theta) d\theta \\
 &= (n-1) \cdot \int_0^\pi \cos^2(\theta) \cdot \sin^{n-2}(\theta) d\theta \\
 &= (n-1) \cdot \int_0^\pi (1 - \sin^2) \cdot \sin^{n-2}(\theta) d\theta \\
 &= (n-1) \cdot (I_{n-2} - I_n),
 \end{aligned}$$

which means:

$$\begin{aligned}
 I_n &= (n-1) \cdot I_{n-2} - (n-1) \cdot I_n \\
 \implies nI_n &= (n-1) \cdot I_{n-2} \\
 \implies I_n &= \frac{n-1}{n} \cdot I_{n-2}
 \end{aligned}$$

Unrolling this recursive formula, we get:

$$\begin{aligned}
 I_{2k} &= I_0 \cdot \prod_{i=1}^k \frac{2i-1}{2i} \\
 I_{2k+1} &= I_1 \cdot \prod_{i=1}^k \frac{2i}{2i+1}
 \end{aligned}$$

We can easily calculate  $I_0$  and  $I_1$  explicitly:

$$\begin{aligned}
 I_0 &= \int_0^\pi \sin^0(\theta) d\theta \\
 &= \int_0^\pi 1 d\theta \\
 &= \pi \\
 I_1 &= \int_0^\pi \sin(\theta) d\theta \\
 &= (-\cos(\pi)) - (-\cos(0)) \\
 &= 2
 \end{aligned}$$

Thus, we have:

$$\begin{aligned}
 I_{2k} &= \pi \cdot \prod_{i=1}^k \frac{2i-1}{2i} \\
 I_{2k+1} &= 2 \cdot \prod_{i=1}^k \frac{2i}{2i+1}
 \end{aligned}$$

This gives us:

$$\begin{aligned} I_{2k+1}I_{2k} &= 2\pi \cdot \prod_{i=1}^k \frac{2i-1}{2i} \cdot \prod_{i=1}^k \frac{2i}{2i+1} \\ &= 2\pi \cdot \prod_{i=1}^k \frac{2i-1}{2i+1} \\ &= \frac{2\pi}{2k+1} \\ &= \frac{\pi}{k + \frac{1}{2}} \end{aligned}$$

$$\begin{aligned} I_{2k}I_{2k-1} &= 2\pi \cdot \prod_{i=1}^k \frac{2i-1}{2i} \cdot \prod_{i=1}^{k-1} \frac{2i}{2i+1} \\ &= 2\pi \cdot \frac{2k-1}{2k} \cdot \prod_{i=1}^{k-1} \frac{2i-1}{2i} \cdot \prod_{i=1}^{k-1} \frac{2i}{2i+1} \\ &= 2\pi \cdot \frac{2k-1}{2k} \cdot \frac{1}{2k-1} \\ &= \frac{\pi}{k} \end{aligned}$$

We know that

$$\lambda(B_1^1(0)) = \lambda((-1, 1)) = 2$$

And, recalling the fact that  $\lambda^0$  is the counting measure, we also know that

$$\lambda^0(B_1^0(0)) = \lambda^0(\{0\}) = 1$$

Thus, at long last, we can calculate the volumes of our unit balls:

$$\begin{aligned} \lambda^{2k}(B_1^{2k}(0)) &= (I_{2k}I_{2k-1}) \cdot (I_{2k-2}I_{2k-3}) \cdot \dots \cdot (I_2I_1) \cdot \lambda^0(B^0(0)) \\ &= \left( \prod_{i=1}^k \frac{\pi}{i} \right) \cdot 1 \\ &= \frac{\pi^k}{k!} = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} \end{aligned}$$

$$\begin{aligned}
 \lambda^{2k+1}(B_1^{2k+1}(0)) &= (I_{2k+1}I_{2k}) \cdot (I_{2k-1}I_{2k-2}) \cdot \dots \cdot (I_3I_2) \cdot \lambda^1(B^1(0)) \\
 &= \left( \prod_{i=1}^k \frac{\pi}{i + \frac{1}{2}} \right) \cdot 2 \\
 &= \frac{\pi^k}{\prod_{i=1}^k i + \frac{1}{2}} \cdot 2 \\
 &= \frac{\pi^k}{\prod_{i=0}^k i + \frac{1}{2}} \\
 &= \frac{\pi^k \cdot \sqrt{\pi}}{\left( \prod_{i=0}^k i + \frac{1}{2} \right) \cdot \sqrt{\pi}} \\
 &= \frac{\pi^{\frac{n}{2}}}{\left( \prod_{i=0}^k i + \frac{1}{2} \right) \cdot \Gamma(\frac{1}{2})} \\
 &= \frac{\pi^{\frac{n}{2}}}{\left( \prod_{i=1}^{k+1} i - \frac{1}{2} \right) \cdot \Gamma(\frac{1}{2})} \\
 &= \frac{\pi^{\frac{n}{2}}}{\Gamma(k + \frac{1}{2} + 1)} \\
 &= \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)}
 \end{aligned}$$

Which gives us

$$\lambda^n(B_r^n(0)) = \frac{\pi^{n/2}}{\Gamma(\frac{n}{2} + 1)} \cdot r^n$$

as desired.  $\square$

**Proposition 4.22.** The 5-dimensional unit ball has the highest volume out of any unit ball. As the dimension increases, the volume of the unit ball converges to zero.

*Proof Sketch.* Plugging small natural numbers into our formula, we see that the formula is monotonically increasing for  $n \leq 5$  and monotonically decreasing for  $n \geq 5$ . Furthermore, we have  $\Gamma(\frac{n}{2} + 1) \geq \Gamma(k + 1) = k!$ , which grows asymptotically faster than  $\pi^{n/2} \leq \sqrt{\pi} \cdot \pi^k$ .  $\square$

**Lemma 4.23**

Let  $j \in \{1, \dots, n\}$ . Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be continuously differentiable with  $f, \partial_j f \in L^1(\mathbb{R}^n)$ . Then

$$\int_{\mathbb{R}^n} \partial_j f(x) d\lambda^n(x) = 0$$

*Proof.* 1. First, assume there is an  $R > 0$  such that  $f(x) = 0$  for all  $\|x\|_2 \geq R$ . Let  $x = (y, x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$ . Then by Fubini we get:

$$\begin{aligned} \int_{\mathbb{R}^n} \partial_j f(x) d\lambda^n(x) &= \int_{\mathbb{R}^n} \partial_j f(y, x_n) d\lambda^n(y, x_n) \\ &= \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} \partial_j f(y, x_n) d\lambda^1(x_n) d\lambda^{n-1}(y). \end{aligned}$$

For  $j = n$ , we apply the fundamental theorem of calculus and our assumption that  $f(x) = 0$  for all  $\|x\|_2 \geq R$  and get:

$$\begin{aligned} \int_{\mathbb{R}} \partial_j f(y, x_n) d\lambda^1(x_n) &= \lim_{x_n \rightarrow \infty} f(y, x_n) - f(y, -x_n) \\ &= 0 \end{aligned}$$

By induction, the same holds for all  $j$ .

2. Now, let  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  be a smooth function fulfilling:

$$\varphi(t) = \begin{cases} 1 & |t| \leq 1, \\ 0 & |t| \geq 2. \end{cases}$$

And let

$$\begin{aligned} \varphi_R : \mathbb{R}^n &\rightarrow \mathbb{R} \\ \varphi_R(x) &= \varphi\left(\frac{\|x\|_2}{R}\right) \end{aligned}$$

Then we have

$$(\varphi_R f)(x) = 0$$

for all  $\|x\|_2 \geq 2R$ , meaning that by the product rule of differentiation and step 1 we get:

$$\begin{aligned} \int_{\mathbb{R}^n} \varphi_R(x) \cdot \partial_j f(x) d\lambda^n(x) + \int_{\mathbb{R}^n} \partial_j \varphi_R(x) \cdot f(x) d\lambda^n(x) \\ = \int_{\mathbb{R}^n} \partial_j (\varphi_R(x) \cdot f(x)) d\lambda^n(x) \\ = 0, \end{aligned}$$

i.e.

$$\int_{\mathbb{R}^n} \varphi_R(x) \cdot \partial_j f(x) d\lambda^n(x) = - \int_{\mathbb{R}^n} \partial_j \varphi_R(x) \cdot f(x) d\lambda^n(x)$$

Now, notice we also have  $\varphi_R(x) = 1$  for  $\|x\| \leq R$  - in particular, we have  $\varphi_R \rightarrow 1$  and  $\partial_j \varphi_R \rightarrow 0$  pointwise on  $\mathbb{R}^n$  for  $R \rightarrow \infty$ , which means we have

$$\int_{\mathbb{R}^n} \partial_j f(x) d\lambda^n(x) = \int_{\mathbb{R}^n} \lim_{R \rightarrow \infty} \varphi_R(x) \cdot \partial_j f(x) d\lambda^n(x)$$

and

$$\int_{\mathbb{R}^n} \lim_{R \rightarrow \infty} \partial_j \varphi_R(x) \cdot f(x) d\lambda^n(x) = 0$$

Finally, we also have:

$$|\varphi_R \cdot (\partial_j f)| \leq |\partial_j f| \in L^1(\mathbb{R}^n)$$

and

$$\begin{aligned} |(\partial_j \varphi_R) \cdot f| &\leq \frac{\max |\varphi'|}{R} \cdot |f| \\ &\leq (\max |\varphi'|) \cdot |f| \in L^1(\mathbb{R}^n) \end{aligned}$$

Thus, we can apply Lebesgue's dominated convergence theorem to pull out the limits and connect the two equalities, giving us:

$$\begin{aligned} \int_{\mathbb{R}^n} \partial_j f(x) \, d\lambda^n(x) &= \int_{\mathbb{R}^n} \lim_{R \rightarrow \infty} \varphi_R(x) \cdot \partial_j f(x) \, d\lambda^n(x) \\ &= \lim_{R \rightarrow \infty} \int_{\mathbb{R}^n} \varphi_R(x) \cdot \partial_j f(x) \, d\lambda^n(x) \\ &= - \lim_{R \rightarrow \infty} \int_{\mathbb{R}^n} \partial_j \varphi_R(x) \cdot f(x) \, d\lambda^n(x) \\ &= - \int_{\mathbb{R}^n} \lim_{R \rightarrow \infty} \partial_j \varphi_R(x) \cdot f(x) \, d\lambda^n(x) \\ &= 0 \end{aligned}$$

□

#### Corollary 4.24: Partial Integration-ish Formula

Let  $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$  be continuously differentiable such that  $f, \partial_j f \in L^p(\mathbb{R}^n)$  and  $g, \partial_j g \in L^q(\mathbb{R}^n)$  with  $\frac{1}{p} + \frac{1}{q} = 1$ . Then we have

$$\int_{\mathbb{R}^n} (\partial_j f(x)) \cdot g(x) \, d\lambda^n(x) = 0 - \int_{\mathbb{R}^n} f(x) \cdot (\partial_j g(x)) \, d\lambda^n(x)$$

*Proof.* Applying the product rule of differentiation, and then the last lemma, we get:

$$\begin{aligned} \int_{\mathbb{R}^n} (\partial_j f(x)) \cdot g(x) \, d\lambda^n(x) + \int_{\mathbb{R}^n} f(x) \cdot (\partial_j g(x)) \, d\lambda^n(x) \\ = \int_{\mathbb{R}^n} \partial_j (f(x) \cdot g(x)) \, d\lambda^n(x) \\ = 0 \end{aligned}$$

□

## 4.3 Change of Variables

**Lemma 4.25.** Let  $U \subset \mathbb{R}^n$  and  $x_0 \in U$ . Let  $\varphi : U \rightarrow \mathbb{R}^n$  be a function such that  $D\varphi(x_0)$  is invertible.

Then for a sequence  $Q_j = Q(x_j, r_j) \subset U$  of cuboids of sidelength  $r_j$  with center  $x_j$  such that  $r_j \rightarrow 0$  and  $x_0 \in Q_j$ , we have:

$$\limsup_{j \rightarrow \infty} \frac{\lambda^n(\varphi(Q_j))}{\lambda^n(Q_j)} \leq |\det D\varphi(x_0)|$$

*Proof.* We can assume  $x_0 = 0$  and  $\varphi(0) = 0$ , since otherwise we can translate space as needed before doing any calculations without breaking any of our assumptions.

1. Assume  $D(\varphi(0)) = E_n$ . Then, by definition of differentiability and equivalence of norms on finite-dimensional vector spaces, we have

$$\begin{aligned} 0 &= \lim_{x \rightarrow 0} \frac{\|\varphi(x) - \varphi(0) - D\varphi(0)x\|_\infty}{\|x\|_\infty} \\ &= \lim_{x \rightarrow 0} \frac{\|\varphi(x) - x\|_\infty}{\|x\|_\infty}, \end{aligned}$$

Let  $\varepsilon > 0$ . Then, by the definition of convergence, for every  $x$  with a sufficiently small norm, we have:

$$\frac{\|\varphi(x) - x\|_\infty}{\|x\|_\infty} \leq \varepsilon,$$

which means

$$\|\varphi(x) - x\|_\infty \leq \varepsilon \|x\|_\infty.$$

Furthermore, for  $x \in Q_j$ , we have:

$$\begin{aligned} \|x\|_\infty &= \|x - \vec{0}\|_\infty \\ &= \|x - x_0\|_\infty \\ &\leq \|x - x_j\| + \|x_j - x_0\| \\ &\leq 2r_j \end{aligned}$$

For sufficiently large  $j$ , these imply:

$$\begin{aligned} \|\varphi(x) - x\|_\infty &\leq \varepsilon \|x\|_\infty \\ &\leq 2\varepsilon r_j. \end{aligned}$$

Further applying the triangle inequality, we get:

$$\begin{aligned} \|\varphi(x) - \varphi(x_j)\| &\leq \|\varphi(x) - x\|_\infty \\ &\quad + \|x - x_j\|_\infty \\ &\quad + \|x_j - \varphi(x_j)\|_\infty \\ &\leq 2\varepsilon r_j + r_j + 2\varepsilon r_j \\ &\leq (1 + 4\varepsilon)r_j, \end{aligned}$$

which means that  $\varphi$  increases the side length of our cube by a factor of at most  $(1 + 4\varepsilon)$ . Therefore, it increases the volume by a factor at most  $(1 + 4\varepsilon)^n$ , i.e:

$$\frac{\lambda^n(\varphi(Q_j))}{\lambda^n(Q_j)} \leq (1 + 4\varepsilon)^n$$

Letting  $j \rightarrow \infty$  and  $\varepsilon \searrow 0$ , we have

$$\begin{aligned} \limsup_{j \rightarrow \infty} \frac{\lambda^n(\varphi(Q_j))}{\lambda^n(Q_j)} &\leq \lim_{\varepsilon \rightarrow 0} (1 + 4|\varepsilon|)^n \\ &= 1 \\ &= |\det E_n| \end{aligned}$$

2. Now, let  $S := D\varphi(0)$  and  $\varphi_0 := S^{-1} \circ \varphi$ , i.e.  $\varphi = S \circ \varphi_0$ . Then  $D\varphi_0(0) = E_n$ . By the linear transformation equation  $\lambda^n(S(E)) = |\det(S)|\lambda^n(E)$  (1.9.2), we have:

$$\begin{aligned} \limsup_{j \rightarrow \infty} \frac{\lambda^n(\varphi(Q_j))}{\lambda^n(Q_j)} &= \limsup_{j \rightarrow \infty} \frac{\lambda^n(S(\varphi_0(Q_j)))}{\lambda^n(Q_j)} \\ &= |\det S| \limsup_{j \rightarrow \infty} \frac{\lambda^n(\varphi_0(Q_j))}{\lambda^n(Q_j)} \\ &\leq |\det S| \\ &= |\det D\varphi(0)| \end{aligned}$$

□

#### Theorem 4.26: Multivariable Substitution Formula

Let  $U \subset \mathbb{R}^n$  be open. Let  $\varphi : U \rightarrow \mathbb{R}^n$  be  $C^1$ . Then if  $f : V \rightarrow \mathbb{R}$  is  $\lambda^n$ -measurable, we have:

$$\int_V f(y) dy = \int_{\varphi^{-1}(V)} f(\varphi(x)) \cdot |\det D\varphi(x)| dx.$$

#### Corollary 4.27

Let  $U, V \subset \mathbb{R}^n$  be open. Let  $\varphi : U \rightarrow \mathbb{R}^n$  be  $C^1$ . Then if  $A \subset U$  is  $\lambda^n$ -measurable, so is  $\varphi(A)$ , and we have

$$\lambda^n(\varphi(A)) = \int_A |\det D\varphi(x)| dx.$$

*Proof.* Apply the previous equation to  $f = \mathbb{1}_{\varphi(A)}$ .

□

**Definition 4.28: Gram Matrix of a Diffeomorphism**

Let  $\varphi : U \rightarrow V$  be a  $C^k$ -diffeomorphism between open subsets of  $\mathbb{R}^n$ . Then the **Gram matrix**  $g(\varphi) \in C^{k-1}(U, \mathbb{R}^{n \times n})$  is given by:

$$g(\varphi)(x) = D\varphi(x)^\top D\varphi(x),$$

or equivalently:

$$g(\varphi)_{ij}(x) = \left\langle \frac{\partial}{\partial x_i} \varphi(x), \frac{\partial}{\partial x_j} \varphi(x) \right\rangle$$

Since  $g(\varphi)(x)$  is the product of a matrix with its own transpose, it is symmetric and strictly positive definite. This lets us alternatively write the multivariable substitution formula as:

$$\int_V f(y) dy = \int_{\varphi^{-1}(V)} f(\varphi(x)) \sqrt{\det g(\varphi)(x)} dx$$

## 4.4 Alternative coordinate systems on $\mathbb{R}^2$ and $\mathbb{R}^3$

### 4.4.1 Polar Coordinates

A commonly used alternative coordinate system on  $\mathbb{R}^2$  is given by **polar coordinates**, which describes a point  $x \in \mathbb{R}^2$  based on its distance  $r$  from the origin and the *polar angle*  $\omega$  between  $x$  and  $(0, r)^\top$ . These might be familiar from dealing with complex numbers, where any  $z \in \mathbb{C}$  can be represented as

$$z = r \cdot e^{\omega i}$$

#### Definition 4.29: Polar Coordinates on $\mathbb{R}^2$

Let  $U = (0, \infty) \times (0, 2\pi)$  and  $V = \mathbb{R}^2 \setminus \{(x, 0)^\top : x \geq 0\}$ . The **polar coordinate transformation** is the map:

$$\begin{aligned} \varphi : U &\rightarrow V \\ \begin{pmatrix} r \\ \omega \end{pmatrix} &\mapsto \begin{pmatrix} r \cos \omega \\ r \sin \omega \end{pmatrix} \end{aligned}$$

which maps a point given in polar coordinates to its corresponding point in the cartesian coordinate system.

The Jacobian of the polar coordinate transform is:

$$\begin{aligned} D\varphi(r, \omega) &= \begin{pmatrix} \frac{\partial}{\partial r} r \cos \omega & \frac{\partial}{\partial \omega} r \cos \omega \\ \frac{\partial}{\partial r} r \sin \omega & \frac{\partial}{\partial \omega} r \sin \omega \end{pmatrix} \\ &= \begin{pmatrix} \cos \omega & -r \sin \omega \\ \sin \omega & r \cos \omega \end{pmatrix} \end{aligned}$$

Which means the determinant is:

$$\begin{aligned} \det D\varphi(r, \omega) &= \det \begin{pmatrix} \cos \omega & -r \sin \omega \\ \sin \omega & r \cos \omega \end{pmatrix} \\ &= r \cos(\omega)^2 + r \sin(\omega)^2 \\ &= r \end{aligned}$$

#### Corollary 4.30: Polar Coordinate Integral Transform

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be integrable. Then we have:

$$\int_{\mathbb{R}^2} f(x, y) d\lambda(x, y) = \int_0^{2\pi} \int_0^\infty r \cdot f(r \cos(\omega), r \sin(\omega)) dr d\omega$$

**Proposition 4.31.** The inverse of the polar coordinate transform  $\varphi$  is:

$$\varphi^{-1}(x, y) = \begin{cases} (r, \arccos(\frac{x}{r}))^\top & y \geq 0 \\ (r, 2\pi - \arccos(\frac{x}{r}))^\top & y < 0 \end{cases}$$

where  $r = \sqrt{x^2 + y^2}$ .

**Example 4.32. (The Gaussian Integral):** We want to find the area under the Gaussian bell curve pre-normalization, i.e.

$$\int_{\mathbb{R}} e^{-x^2} dx$$

To do this, we add an additional dimension and exploit the resulting rotational symmetry. By Fubini's Theorem, we have:

$$\begin{aligned} \int_{\mathbb{R}^2} e^{-(x^2+y^2)} d\lambda^2(x, y) &= \int_{\mathbb{R}} \left( \int_{\mathbb{R}} e^{-(x^2+y^2)} d\lambda(y) \right) d\lambda(x) \\ &= \int_{\mathbb{R}} \left( \int_{\mathbb{R}} e^{-x^2} \cdot e^{-y^2} d\lambda(y) \right) d\lambda(x) \\ &= \int_{\mathbb{R}} e^{-x^2} \cdot \left( \int_{\mathbb{R}} e^{-y^2} d\lambda(y) \right) d\lambda(x) \\ &= \int_{\mathbb{R}} e^{-x^2} d\lambda(x) \cdot \int_{\mathbb{R}} e^{-y^2} d\lambda(y) \\ &= \left( \int_{\mathbb{R}} e^{-x^2} d\lambda(x) \right)^2 \end{aligned}$$

Therefore, we have

$$\int_{\mathbb{R}} e^{-x^2} d\lambda(x) = \sqrt{\int_{\mathbb{R}^2} e^{-(x^2+y^2)} d\lambda^2(x, y)}$$

We can now calculate the two-dimensional integral using multivariable substitution to transform to polar coordinates:

$$\begin{aligned} \int_{\mathbb{R}^2} e^{-(x^2+y^2)} d\lambda^2(x, y) &= \int_{(0, \infty) \times (0, 2\pi)} r e^{-r^2} d\lambda^2(r, \omega) \\ &= \int_0^\infty \left( \int_0^{2\pi} r e^{-r^2} d\omega \right) dr \\ &= \int_0^\infty 2\pi r e^{-r^2} dr \\ &= 2\pi \int_0^\infty r e^{-r^2} dr \\ &= 2\pi \int_0^\infty \frac{1}{2} e^{-r^2} 2r dr \\ &= 2\pi \int_{-\infty}^0 \frac{1}{2} e^s ds \\ &= \pi \int_{-\infty}^0 e^s ds \\ &= \pi \end{aligned}$$

Which means that the area under the bell curve is  $\sqrt{\pi}$ .

**Example 4.33. (Beta function):** For  $p, q \in \mathbb{R}_{>0}$ , the beta function  $B$  is defined as:

$$B(p, q) = \int_0^1 t^{p-1} \cdot (1-t)^{q-1} dt$$

The beta function is of interest primarily because of its close relationship to the gamma function:

**Theorem 4.34.** We have:

$$B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

*Proof.* Substituting  $t = g(\omega) := \sin(\omega)^2$  into the definition of the beta function gives us:

$$\begin{aligned} g'(\omega) &= \frac{d}{d\omega} \sin(\omega)^2 \\ &= \sin(\omega) \cos(\omega) + \cos(\omega) + \sin(\omega) \\ &= 2 \sin(\omega) \cos(\omega) \end{aligned}$$

and:

$$1 = \sin(\omega)^2 \Leftrightarrow \omega = \frac{\pi}{2},$$

which means we have:

$$\begin{aligned} B(p, q) &= \int_0^1 t^{p-1} \cdot (1-t)^{q-1} dt \\ &= \int_0^{\pi/2} \sin(\omega)^{2p-2} \cdot (1 - \sin(\omega)^2)^{q-1} \cdot 2 \sin(\omega) \cos(\omega) d\omega \\ &= \int_0^{\pi/2} \sin(\omega)^{2p-2} \cdot \cos(\omega)^{2q-2} \cdot 2 \sin(\omega) \cos(\omega) d\omega \\ &= 2 \cdot \int_0^{\pi/2} \sin(\omega)^{2p-1} \cdot \cos(\omega)^{2q-1} d\omega \end{aligned}$$

Meanwhile, performing a reverse substitution with  $t := g(r) = r^2$  in the definition of the gamma function gives:

$$\begin{aligned} \Gamma(m) &= \int_0^\infty t^{m-1} e^{-t} dt \\ &= \int_0^\infty (r^2)^{m-1} e^{-r^2} \cdot 2r dr \\ &= 2 \int_0^\infty r^{2m-1} e^{-r^2} dr \end{aligned}$$

Now, we get:

$$\begin{aligned} B(p, q)\Gamma(p+q) &= 2 \int_0^\infty B(p, q) \cdot r^{2p+2q-1} \cdot e^{-r^2} dr \\ &= 4 \int_0^\infty \int_0^{\pi/2} (\sin \omega)^{2p-1} \cdot r^{2p-1} \cdot (\cos \omega)^{2q-1} \cdot r^{2q-1} \cdot r \cdot e^{-r^2} d\omega dr \\ &= 4 \int_0^\infty \int_0^{\pi/2} (r \sin \omega)^{2p-1} \cdot (r \cos \omega)^{2q-1} \cdot r \cdot e^{-r^2} d\omega dr \end{aligned}$$

By applying the polar coordinate integral transform in reverse, we get:

$$\begin{aligned}
 B(p, q)\Gamma(p + q) &= 4 \int_0^\infty \int_0^{\pi/2} (r \sin \omega)^{2p-1} \cdot (r \cos \omega)^{2q-1} \cdot r \cdot e^{-r^2} d\omega dr \\
 &= 4 \int_0^\infty \int_0^\infty x^{2p-1} \cdot y^{2q-1} \cdot e^{-\sqrt{x^2+y^2}^2} dy dx \\
 &= 4 \int_0^\infty \int_0^\infty x^{2p-1} \cdot y^{2q-1} \cdot e^{-x^2-y^2} dy dx \\
 &= \left( 2 \int_0^\infty x^{2p-1} e^{-x^2} dx \right) \cdot \left( 2 \int_0^\infty y^{2q-1} e^{-y^2} dy \right) \\
 &= \Gamma(p)\Gamma(q)
 \end{aligned}$$

□

### 4.4.2 Spherical Coordinates

The most natural generalization of polar coordinates to  $\mathbb{R}^3$  are *spherical coordinates*, in which  $\theta \in (0, \pi)$  describes the *polar angle* between a point  $P$  and the  $z$ -Axis and  $\omega \in (0, 2\pi)$  describes the *azimuthal angle* between the  $x$ -axis and the projection  $P'$  of  $P$  onto the  $xy$ -plane. We can derive spherical coordinates from two dimensional coordinates as follows:

1. Take a point  $P = (x, y, z) \in \mathbb{R}^3$ . Notice that  $z = r \cos(\theta)$ .
2. Project  $P$  to a point  $P' = (x, y)$ . The distance of  $P'$  from  $(0, 0)$  is exactly  $\rho := r \sin \theta$ . Therefore,  $P'$  has polar coordinates  $(\rho, \omega)$ , which gives us:

$$P = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \rho \cos \omega \\ \rho \sin \omega \\ r \cos \theta \end{pmatrix} = \begin{pmatrix} r \sin \theta \cos \omega \\ r \sin \theta \sin \omega \\ r \cos \theta \end{pmatrix}$$

#### Definition 4.35: Spherical Coordinates on $\mathbb{R}^3$

Let  $U = (0, \infty) \times (0, \pi) \times (0, 2\pi)$  and  $V = \mathbb{R}^3 \setminus \{(x, 0, z)^\top : x \geq 0\}$ . The **spherical coordinate transformation** is the map:

$$\begin{aligned} \varphi : U &\rightarrow V \\ \begin{pmatrix} r \\ \theta \\ \omega \end{pmatrix} &\mapsto \begin{pmatrix} r \sin \theta \cos \omega \\ r \sin \theta \sin \omega \\ r \cos \theta \end{pmatrix} \end{aligned}$$

which maps a point given in spherical coordinates to its corresponding point in the cartesian coordinate system.

The Jacobian of the spherical coordinate transform is:

$$\begin{aligned} D\varphi(r, \theta, \omega) &= \begin{pmatrix} \frac{\partial}{\partial r} r \sin \theta \cos \omega & \frac{\partial}{\partial \theta} r \sin \theta \cos \omega & \frac{\partial}{\partial \omega} r \sin \theta \cos \omega \\ \frac{\partial}{\partial r} r \sin \theta \sin \omega & \frac{\partial}{\partial \theta} r \sin \theta \sin \omega & \frac{\partial}{\partial \omega} r \sin \theta \sin \omega \\ \frac{\partial}{\partial r} r \cos \theta & \frac{\partial}{\partial \theta} r \cos \theta & \frac{\partial}{\partial \omega} r \cos \theta \end{pmatrix} \\ &= \begin{pmatrix} \sin \theta \cos \omega & r \cos \theta \cos \omega & -r \sin \theta \sin \omega \\ \sin \theta \sin \omega & r \cos \theta \sin \omega & r \sin \theta \cos \omega \\ \cos \theta & -r \sin \theta & 0 \end{pmatrix} \end{aligned}$$

And a long, tedious and unenlightening calculation eventually gives the determinant as:

$$\det D\varphi(r, \theta, \omega) = r^2 \sin(\theta)$$

Which gives us:

#### Corollary 4.36: Spherical Coordinate Integral Transform

Let  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  be integrable. Then we have:

$$\begin{aligned} &\int_{\mathbb{R}^3} f(x, y, z) d\lambda^3(x, y, z) \\ &= \int_0^{2\pi} \int_0^\pi \int_0^\infty r^2 \sin \theta \cdot f(r \sin \theta \sin \omega, r \sin \theta \cos \omega, r \cos \theta) dr d\theta d\omega \end{aligned}$$

**Example 4.37. (Another derivation of the volume of  $B_R^3$ ):**

$$\begin{aligned}
 \lambda^3(B_R^3) &= \int_{\mathbb{R}^3} \mathbb{1}_{B_R^3}(x, y, z) \, d\lambda^3(x, y, z) \\
 &= \int_{\mathbb{R}^3} \mathbb{1}_{\{\sqrt{x^2+y^2+z^2} < R\}}(x, y, z) \, d\lambda^3(x, y, z) \\
 &= \int_0^{2\pi} \int_0^\pi \int_0^\infty \mathbb{1}_{\{r < R\}}(r) \cdot r^2 \sin \theta \, dr \, d\theta \, d\omega \\
 &= \int_0^{2\pi} \int_0^\pi \int_0^\infty \mathbb{1}_{(0,R)}(r) \cdot r^2 \sin \theta \, dr \, d\theta \, d\omega \\
 &= \left( \int_0^\infty \mathbb{1}_{(0,R)}(r) \cdot r^2 \, dr \right) \cdot \left( \int_0^\pi \sin \theta \, d\theta \right) \cdot \left( \int_0^{2\pi} 1 \, d\omega \right) \\
 &= \left( \int_0^R r^2 \, dr \right) \cdot \left( \int_0^\pi \sin \theta \, d\theta \right) \cdot \left( \int_0^{2\pi} 1 \, d\omega \right) \\
 &= \frac{1}{3} R^3 \cdot (-\cos(\pi) + \cos(0)) \cdot 2\pi \\
 &= \frac{4}{3} \pi R^3
 \end{aligned}$$

**Example 4.38. (Gravitational Potential of a Sphere:)** The gravitational potential that a body  $B \subset \mathbb{R}^3$  with mass distribution  $m : B \rightarrow \mathbb{R}$  exerts on a point  $x \in \mathbb{R}^3 \setminus B$  is given by:

$$V(x) = -G \cdot \int_B \frac{m(y)}{\|x - y\|_2} \, d\lambda^3(y)$$

We will derive a closed form expression for the case  $B = B_R^3$ . If we assume that the mass distribution is rotationally symmetric, we can replace  $m : \mathbb{R}^3 \rightarrow \mathbb{R}$  with a function  $\tilde{m} = m(\|y\|_2) : \mathbb{R} \rightarrow \mathbb{R}$ . We will commit a slight abuse of notation here and let  $m$  be this new function  $\tilde{m}$  from now on. Because of this rotational symmetry, we also have  $V(x) = V(0, 0, \|x\|)$ .

Now, if  $y$  has spherical coordinates  $(r, \theta, \omega)$ , then assuming  $y$  lies outside of our ball we can use the Pythagorean theorem to get:

$$\begin{aligned}
 \|x - y\|_2^2 &= (\|x\|_2 - r \cos \theta)^2 + r^2 \sin^2 \theta \\
 &= \|x\|_2^2 + r^2 - 2r\|x\|_2 \cos \theta
 \end{aligned}$$

Thus, we have:

$$\begin{aligned}
 V(x) &= -G \cdot \int_{B_R^3} \frac{m(r)}{\|x - y\|_2} \, d\lambda^3(y) \\
 &= -G \int_0^{2\pi} \int_0^\pi \int_0^R \left( \frac{m(r)}{\sqrt{\|x\|_2^2 + r^2 - 2r\|x\|_2 \cos \theta}} \cdot r^2 \sin \theta \right) dr \, d\theta \, d\omega \\
 &= -2\pi G \int_0^R r^2 \cdot m(r) \cdot \left( \int_0^\pi \frac{\sin \theta}{\sqrt{\|x\|_2^2 + r^2 - 2r\|x\|_2 \cos \theta}} \, d\theta \right) dr
 \end{aligned}$$

Substituting  $t = g(\theta) = -\cos \theta$  into the inner integral, we get:

$$\begin{aligned}
 & \int_0^\pi \frac{\sin \theta}{\sqrt{\|x\|_2^2 + r^2 - 2r\|x\|_2 \cos \theta}} d\theta \\
 &= \int_{-1}^1 \frac{1}{\sqrt{\|x\|_2^2 + r^2 + 2r\|x\|_2 t}} dt \\
 &= \left[ \frac{1}{r \cdot \|x\|_2} \cdot \sqrt{\|x\|_2^2 + r^2 + 2r\|x\|_2 t} \right]_{t=-1}^{t=1} \\
 &= \frac{1}{r \cdot \|x\|_2} \cdot \left( \sqrt{\|x\|_2^2 + r^2 + 2r\|x\|_2} - \sqrt{\|x\|_2^2 + r^2 - 2r\|x\|_2} \right) \\
 &= \frac{1}{r \cdot \|x\|_2} \cdot ((\|x\|_2 + r) - (\|x\|_2 - r)) \\
 &= \frac{2}{\|x\|_2}
 \end{aligned}$$

Therefore, we have:

$$V(x) = \frac{4\pi}{\|x\|_2} \cdot \int_0^R m(r)r^2 dr$$

Now, we can once again use the rotational symmetry of  $m$  to describe the total mass  $M$  of our body in terms of spherical coordinates:

$$\begin{aligned}
 M &= \int_{B_R^3(0)} m(y) d\lambda^3(y) \\
 &= \int_0^{2\pi} \int_0^\pi \int_0^R m(r)r^2 \sin \theta dr d\theta d\omega \\
 &= 4\pi \int_0^R m(r)r^2 dr
 \end{aligned}$$

Thus, the total mass distribution of our sphere is simply:

$$V(x) = -\frac{GM}{\|x\|}$$

which is exactly the same mass distribution we would get if we concentrated the entire mass of our sphere at its center. This relation is part of a theorem in classical mechanics known as the **shell theorem**.

### 4.4.3 Cylindrical Coordinates

Alternatively, one can trivially extend polar coordinates into three dimensions by simply leaving the  $z$  coordinate unchanged, leading to *cylindrical coordinates*:

**Definition 4.39: Cylindrical Coordinates on  $\mathbb{R}^3$**

Let  $U = (0, \infty) \times (0, 2\pi) \times \mathbb{R}$  and  $V = \mathbb{R}^3 \setminus \{(x, 0, z)^\top : x \geq 0\}$ . The **spherical coordinate transformation** is the map:

$$\begin{aligned} \varphi : U &\rightarrow V \\ \begin{pmatrix} r \\ \omega \\ z \end{pmatrix} &\mapsto \begin{pmatrix} r \cos \omega \\ r \sin \omega \\ z \end{pmatrix} \end{aligned}$$

which maps a point given in cylindrical coordinates to its corresponding point in the cartesian coordinate system.

Since the cylindrical coordinate transform leaves the  $z$  coordinate unchanged, its Jacobian and its determinant end up being almost identical:

$$\begin{aligned} D\varphi(r, \omega, z) &= \begin{pmatrix} \frac{\partial}{\partial r} r \cos \omega & \frac{\partial}{\partial \omega} r \cos \omega & \frac{\partial}{\partial z} r \cos \omega \\ \frac{\partial}{\partial r} r \sin \omega & \frac{\partial}{\partial \omega} r \sin \omega & \frac{\partial}{\partial z} r \sin \omega \\ \frac{\partial}{\partial r} z & \frac{\partial}{\partial \omega} z & \frac{\partial}{\partial z} z \end{pmatrix} \\ &= \begin{pmatrix} \cos \omega & -r \sin \omega & 0 \\ \sin \omega & r \cos \omega & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \det D\varphi(r, \omega, z) &= \det \begin{pmatrix} \cos \omega & -r \sin \omega & 0 \\ \sin \omega & r \cos \omega & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \det \begin{pmatrix} \cos \omega & -r \sin \omega \\ \sin \omega & r \cos \omega \end{pmatrix} \\ &= r \cos(\omega)^2 + r \sin(\omega)^2 \\ &= r \end{aligned}$$

**Corollary 4.40: Cylindrical Coordinate Integral Transform**

Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be integrable. Then we have:

$$\int_{\mathbb{R}^3} f(x, y, z) d\lambda(x, y, z) = \int_{\mathbb{R}} \int_0^{2\pi} \int_0^\infty r \cdot f(r \cos(\omega), r \sin(\omega), z) dr d\omega dz$$

**Proposition 4.41.** The inverse of the cylindrical coordinate transform  $\varphi$  is:

$$\varphi^{-1}(x, y, z) = \begin{cases} (r, \arccos(\frac{x}{r}), z)^\top & y \geq 0 \\ (r, 2\pi - \arccos(\frac{x}{r}), z)^\top & y < 0 \end{cases}$$

where  $r = \sqrt{x^2 + y^2}$ .

**Theorem 4.42: Volume of a Solid of Revolution**

Let  $S$  be a solid constructed by rotating a function  $\rho(z)$  around the  $z$  axis, i.e.

$$S = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 < \rho(z)^2\}$$

Then we have:

$$\lambda^3(S) = \pi \int_{\mathbb{R}} \rho(z)^2 d\lambda^1(z)$$

*Proof.*

$$\begin{aligned} \lambda^3(S) &= \int_{\mathbb{R}^3} \mathbb{1}_{\{x^2+y^2 < \rho(z)^2\}}(x, y, z) d\lambda^3(x, y, z) \\ &= \int_{\mathbb{R}} \int_0^{2\pi} \int_0^{\infty} r \cdot \mathbb{1}_{\{r^2 < \rho(z)^2\}}(r, \omega, z) dr d\omega dz \\ &= \int_{\mathbb{R}} \int_0^{2\pi} \int_0^{\infty} r \cdot \mathbb{1}_{\{r < \rho(z)\}}(r, \omega, z) dr d\omega dz \\ &= \int_{\mathbb{R}} \int_0^{2\pi} \int_0^{\rho(z)} r dr d\omega dz \\ &= \int_{\mathbb{R}} \int_0^{2\pi} \frac{1}{2} \rho(z)^2 d\omega dz \\ &= \int_{\mathbb{R}} \pi \rho(z)^2 dz \\ &= \pi \int_{\mathbb{R}} \rho(z)^2 dz \end{aligned}$$

□

## 4.5 The Surface Area Measure

**Definition 4.43.** Let  $U \subset \mathbb{R}^n$  be open. We call a map  $f : U \rightarrow \mathbb{R}^{n+k}$  an *immersion* iff  $f$  is continuously differentiable and  $Df(x)$  is invertible.

**Definition 4.44.** Let  $M \subset \mathbb{R}^{n+k}$  be an  $n$ -dimensional  $C^1$  submanifold. Then a *local parametrization* of  $M$  is an injective immersion  $f : \mathbb{R}^n \supset U \rightarrow M \subset \mathbb{R}^{n+k}$

### Definition 4.45: Gram Matrix

Let  $U \subset \mathbb{R}^n$  and  $f : U \rightarrow \mathbb{R}^{n+k}$ . Then the *Gram matrix*  $g_f(x)$  is the matrix

$$g_f(x) = Df(x)^\top Df(x)$$

Our goal now is to find a sensible definition of *surface area*, i.e. of an  $n$ -dimensional measure associated with an object embedded into  $n+k$ -dimensional space.

### Corollary 4.46: Area of an Embedded Diffeomorphism of a Set

Let  $f = S \circ \varphi$ , where  $\varphi : U \rightarrow V$  is a diffeomorphism with  $U, V \subset \mathbb{R}^n$  and  $S$  is a linear isometry. Then we have

$$\lambda^n(\varphi(E)) = \int_E \sqrt{\det g_f}$$

This theorem suggests that it is sensible to use the same definition for arbitrary immersions:

### Definition 4.47: Area of an Immersion of a Set

Let  $f \in C^1(U, \mathbb{R}^{n+k})$  be an  $n$ -dimensional immersion. Let  $E \subset U$  be  $\lambda^n$ -measurable. Then the  $n$ -dimensional area of  $f(E)$  is given by:

$$A(f, E) = \int_E \sqrt{\det g_f}$$

$\sqrt{\det g_f}$  is known as the *Jacobian* of  $f$ , and often denoted  $Jf$ , but I dislike notation that hides the actual calculation being done behind multiple layers of nested notation and will thus be sticking to  $\sqrt{\det g_f}$ .

### Corollary 4.48: Length of a Regular Curve

An Immersion  $f : I = (a, b) \rightarrow \mathbb{R}^n$  is known as a *regular curve*. The one-dimensional area of the curve, i.e. its length, is given by

$$L(f) := A(f, I) = \int_a^b \|Df(t)\|_2 dt$$

*Proof.* We have:

$$Df(t) = \begin{pmatrix} f'_1(t) \\ \vdots \\ f'_n(t) \end{pmatrix}$$

and thus

$$g_f(t) = Df(t)^\top Df(t) = \|Df(t)\|_2^2$$

□

Note that if some areas of the curve are covered multiple times, then this length formula counts them multiple times as well.

#### Corollary 4.49: Area of a Regular Surface

An immersion  $f : U \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$ , with  $U$  open, is known as a *regular surface*. Its area is given by

$$\begin{aligned} A(f) &:= A(f, U) = \int_U \sqrt{\left\| \frac{\partial}{\partial x} f \right\|_2^2 \cdot \left\| \frac{\partial}{\partial y} f \right\|_2^2 - \left\langle \frac{\partial}{\partial x} f, \frac{\partial}{\partial y} f \right\rangle^2} d\lambda^2(x, y) \\ &= \int_U \left\| \frac{\partial}{\partial x} f \times \frac{\partial}{\partial y} f \right\| d\lambda^2(x, y) \end{aligned}$$

*Proof.* We have

$$Df(x, y) = \begin{pmatrix} \frac{\partial}{\partial x} f(x, y) & \frac{\partial}{\partial y} f(x, y) \end{pmatrix},$$

therefore our Gram matrix is

$$g_f = \begin{pmatrix} \left\| \frac{\partial}{\partial x} f \right\|_2^2 & \left\langle \frac{\partial}{\partial x} f, \frac{\partial}{\partial y} f \right\rangle_2 \\ \left\langle \frac{\partial}{\partial x} f, \frac{\partial}{\partial y} f \right\rangle & \left\| \frac{\partial}{\partial y} f \right\|_2^2 \end{pmatrix},$$

giving our desired result for  $\sqrt{\det g_f}$ .

□

**Theorem 4.50: Area of a Graph**

Let  $U \subset \mathbb{R}^n$  be open,  $u : U \rightarrow \mathbb{R}^k$  and

$$\begin{aligned} f : U &\rightarrow \mathbb{R}^{n+k} \\ x &\mapsto (x, u(x)) \end{aligned}$$

Then we have:

$$A(f, U) = \int_U \sqrt{\det(E_n + Du(x)^\top Du(x))} d\lambda^n(x)$$

If  $k = 1$ , i.e.  $U \subset \mathbb{R}^n$ ,  $u : U \rightarrow \mathbb{R}$ , then we can simplify significantly and get:

$$A(f, U) = \int_U \sqrt{1 + \|Du(x)\|^2} d\lambda^n(x)$$

**Theorem 4.51: Surface Measure**

Let  $M$  be an  $n$ -dimension  $C^1$ -submanifold of  $\mathbb{R}^{n+k}$  locally parametrized by  $f : U \rightarrow M$ . Let  $E \subset M$ . Then there exists a measure  $\omega_M$  such that

$$\omega_M(E) = A(f, E)$$

**Theorem 4.52: Surface of Revolution**

Let  $S$  be a solid of revolution generated by  $\rho(z)$ . Then the surface area of  $S$  is:

$$\omega_S(S) = 2\pi \int_{\mathbb{R}} \rho(z) \sqrt{1 + \left(\frac{d}{dz} \rho(z)\right)^2} dz$$

*Proof.* This follows from our formula for the surface area of a regular surface. However, as additional practice, let us derive the formula from scratch. We can parametrize  $S$  up to a zero set via:

$$f : \begin{pmatrix} \theta \\ z \end{pmatrix} \mapsto \begin{pmatrix} \rho(z) \cos(\theta) \\ \rho(z) \sin(\theta) \\ z \end{pmatrix}$$

Which gives us:

$$\begin{aligned} Df(\theta, z) &= \begin{pmatrix} \frac{\partial}{\partial \theta} \rho(z) \cos(\theta) & \frac{\partial}{\partial z} \rho(z) \cos(\theta) \\ \frac{\partial}{\partial \theta} \rho(z) \sin(\theta) & \frac{\partial}{\partial z} \rho(z) \sin(\theta) \\ \frac{\partial}{\partial \theta} z & \frac{\partial}{\partial z} z \end{pmatrix} \\ &= \begin{pmatrix} -\rho(z) \sin(\theta) & \cos(\theta) \cdot \frac{\partial}{\partial z} \rho(z) \\ \rho(z) \cos(\theta) & \sin(\theta) \cdot \frac{\partial}{\partial z} \rho(z) \\ 0 & 1 \end{pmatrix} \end{aligned}$$

This gives us the Gram matrix:

$$\begin{aligned} g_f(\theta, z) &= \begin{pmatrix} -\rho(z) \sin(\theta) & \rho(z) \cos(\theta) & 0 \\ \cos(\theta) \cdot \frac{\partial}{\partial z} \rho(z) & \sin(\theta) \cdot \frac{\partial}{\partial z} \rho(z) & 1 \end{pmatrix} \cdot \begin{pmatrix} -\rho(z) \sin(\theta) & \cos(\theta) \cdot \frac{\partial}{\partial z} \rho(z) \\ \rho(z) \cos(\theta) & \sin(\theta) \cdot \frac{\partial}{\partial z} \rho(z) \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} \rho(z)^2 \cdot (\sin^2(\theta) + \cos^2(\theta)) & -\rho(z) \sin(\theta) \cos(\theta) \frac{d}{dz} \rho(z) \\ -\rho(z) \sin(\theta) \cos(\theta) \frac{d}{dz} \rho(z) & +\rho(z) \sin(\theta) \cos(\theta) \frac{d}{dz} \rho(z) \\ +\rho(z) \sin(\theta) \cos(\theta) \frac{d}{dz} \rho(z) & (\cos^2(\theta) + \sin^2(\theta)) \left( \frac{\partial}{\partial z} \rho(z) \right)^2 + 1 \end{pmatrix} \\ &= \begin{pmatrix} \rho(z)^2 & 0 \\ 0 & \left( \frac{\partial}{\partial z} \rho(z) \right)^2 + 1 \end{pmatrix} \end{aligned}$$

and thus:

$$\begin{aligned} \sqrt{|\det g_f(\theta, z)|} &= \sqrt{\rho(z)^2 \left( \frac{\partial}{\partial z} \rho(z) \right)^2 + \rho(z)^2} \\ &= \rho(z) \sqrt{1 + \left( \frac{\partial}{\partial z} \rho(z) \right)^2} \end{aligned}$$

Which leaves us with:

$$\begin{aligned} \omega_S(S) &= \int_0^{2\pi} \int_{\mathbb{R}} \rho(z) \sqrt{1 + \left( \frac{\partial}{\partial z} \rho(z) \right)^2} dz d\theta \\ &= 2\pi \int_{\mathbb{R}} \rho(z) \sqrt{1 + \left( \frac{\partial}{\partial z} \rho(z) \right)^2} dz \end{aligned}$$

□

**Example 4.53. (Gabriel's Horn):** We want to show that the solid of rotation  $G$  generated by  $z \in [1, \infty)$  and  $\rho(z) = \frac{1}{z}$  has finite volume, but infinite surface area.

1. We already showed in the section on cylindrical coordinates that the volume is simply:

$$\begin{aligned} \lambda^3(G) &= \pi \cdot \int_1^\infty \rho(z)^2 dz \\ &= \pi \cdot \int_1^\infty \frac{1}{z^2} dz \\ &= -\frac{\pi}{2} \cdot \left[ \frac{1}{z} \right]_{z=1}^{z=\infty} \\ &= -\frac{\pi}{2} \cdot (0 - 1) \\ &= \frac{\pi}{2} \end{aligned}$$

2. We have just shown that the surface area is given by:

$$\begin{aligned}\omega_S(S) &= 2\pi \int_{\mathbb{R}} \rho(z) \sqrt{1 + \left(\frac{\partial}{\partial z} \rho(z)\right)^2} dz \\ &= 2\pi \int_1^\infty \frac{1}{z} \sqrt{1 - \frac{1}{z^2}} dz \\ &> 2\pi \int_1^\infty \frac{1}{z} dz \\ &= 2\pi(\ln(\infty) - \ln(1)) \\ &= \infty\end{aligned}$$

**Theorem 4.54: Surface Integral**

Let  $M \subset \mathbb{R}^{n+k}$  be an  $n$ -dimensional  $C^1$  submanifold of  $\mathbb{R}^{n+k}$ , and let  $M = \bigsqcup_{i \in I} M_i$  be a decomposition of  $M$  into countably many pairwise disjoint measure sets such that  $M_i \subset V_i$  for a local parametrization  $f_i : U_i \rightarrow V_i$ . Then for any measurable function  $u : M \rightarrow \overline{\mathbb{R}}$ , we have:

$$\int_M u(x) d\omega_M(x) = \sum_{i \in I} \int_{f_i^{-1}(M_i)} u(f_i(x)) \cdot \left| \sqrt{\det g_{f_i}(x)} \right| d\lambda^n(x)$$

In particular, if we can find a local parametrization  $f : U \rightarrow V$  such that  $M \setminus V$  is a zero set, we have:

$$\int_M u(x) d\omega_M(x) = \int_{f^{-1}(M)} u(f(x)) \cdot |\det Df(x)| d\lambda^n(x)$$

**Theorem 4.55: Onion Formula**

Integrating a function  $f$  over  $\mathbb{R}^n$  is the same thing as integrating over all surface integrals of the function over all balls  $B_r^n$ , i.e. we have:

$$\int_{\mathbb{R}^n} f(x) d\lambda^n(x) = \int_0^\infty \left( \int_{\partial B_r^n} f(y) d\omega_{\partial B_r^n}(y) \right) dr$$

Note that via multivariable substitution we also get:

$$\int_0^\infty \left( \int_{\partial B_r^n} f(y) d\omega_{\partial B_r^n}(y) \right) dr = \int_0^\infty \left( \int_{\partial B_1^n} r^n f(ry) d\omega_{\partial B_1^n}(y) \right) dr$$

**Corollary 4.56.** We have:

$$\omega_{\partial B_1^n}(\partial B_1^n) = (n+1) \cdot \lambda^{n+1}(B_1^{n+1}),$$

i.e. the surface area of the  $n$ -dimensional unit ball is  $(n+1)$  times the volume of the  $n+1$ -dimensional unit ball.

*Proof.* Applying the onion formula and then Fubini twice, we get:

$$\begin{aligned}
 \lambda^{n+1}(B_1^{n+1}) &= \int_{\mathbb{R}^n} \mathbb{1}_{B_1^n}(x) d\lambda^n(x) \\
 &= \int_0^\infty \int_{\partial B_r^n} \mathbb{1}_{B_1^n}(y) d\omega_{\partial B_r^n}(y) dr \\
 &= \int_{\partial B_r^n} \int_0^\infty \mathbb{1}_{B_1^n}(y) dr d\omega_{\partial B_r^n}(y) \\
 &= \int_{\partial B_r^n} \int_0^1 1 dr d\omega_{\partial B_r^n}(y) \\
 &= \int_0^1 \int_{\partial B_r^n} 1 d\omega_{\partial B_r^n}(y) dr \\
 &= \int_0^1 \omega_{\partial B_r^n}(\partial B_r^n) dr \\
 &= \int_0^1 r^n \cdot \omega_{\partial B_1^n}(\partial B_1^n) dr \\
 &= \frac{\omega_{\partial B_1^n}(\partial B_1^n)}{n+1}
 \end{aligned}$$

□

## 4.6 The Divergence Theorem

**Definition 4.57.** Let  $\Omega \subset \mathbb{R}^n$ . We say  $\Omega$  has a  $C^1$  boundary if  $\partial\Omega$  is an  $n - 1$  dimensional  $C^1$ -manifold such that  $\Omega$  locally lies on only one side of  $\partial\Omega$ .

More precisely,  $\Omega$  has a  $C^1$  boundary iff for every point  $p \in \partial\Omega$ , there exists an open set  $U \subset \mathbb{R}^n$ , an open interval  $I \subset \mathbb{R}$ , an a  $C^1$ -function  $u : U \rightarrow I$  such  $p \in U \times I$  and

$$\Omega \cap (U \times I) = \{(x, y) \in U \times I : y < u(x)\}$$

**Lemma 4.58.** Let  $\Omega \subset \mathbb{R}^n$  be open with  $C^1$  boundary. Then we have:

1.  $\partial\Omega \cap (U \times I) = \{(x, y) \in U \times I : y = u(x)\}$
2.  $(\mathbb{R}^n \setminus \overline{\Omega}) \cap (U \times I) = \{(x, y) \in U \times I : y \geq u(x)\}$

In particular,  $\partial\Omega$  is an  $n - 1$  dimensional manifold given via the graph criterion as  $(x, u(x))$ .

### Theorem 4.59: Outer Normal Field

Let  $\Omega \subset \mathbb{R}^n$  be open with  $C^1$  boundary. Then there exists exactly one map  $\nu_\Omega : \partial\Omega \rightarrow \mathbb{R}^n$  such that:

1.  $\nu_\Omega(p) \perp T_p(\partial\Omega)$ ,
2.  $\|\nu_\Omega(p)\|_2 = 1$ ,
3.  $\exists \varepsilon > 0 : t \in (0, \varepsilon) \implies p + t\nu_\Omega(p) \notin \Omega$ .

This map is continuous and unique and is known as the *outer normal field* of  $\Omega$ . If  $u$  is given as in the definition of  $C^1$  boundary, and  $p = (x, u(x))$ , then we have:

$$\nu_\Omega(p) = \frac{(-Du(x), 1)^\top}{\sqrt{1 + \|Du(x)\|_2^2}}$$

### Lemma 4.60: Partition of Unity

Let  $W_i, i \in I$  be an open cover of a compact set  $K \subset \mathbb{R}^n$ . Then there exists a finite family of smooth functions  $\chi_j : \mathbb{R}^n \rightarrow \mathbb{R}$  with compact support such that

1. For all  $x \in K$ , we have  $\sum_{j \in J} \chi_j(x) = 1$
2. For all  $j \in J$ , there exists an  $i$  such that  $\text{spt}\chi_j \subset W_i$ .

**Theorem 4.61: Divergence Theorem / Gauss's Theorem**

Let  $\Omega \subset \mathbb{R}^n$  be a bounded open set with  $C^1$  boundary and outer normal  $\nu_\Omega : \partial\Omega \rightarrow \mathbb{R}^n$ . Let  $f : \overline{\Omega} \rightarrow \mathbb{R}$  be continuously differentiable. Then we have:

$$\int_{\Omega} \operatorname{div} f(x) \, d\lambda^n(x) = \int_{\partial\Omega} \langle f(x), \nu(x) \rangle \, d\omega(x).$$

**Theorem 4.62: Multidimensional Partial Integration**

Let  $\Omega \subset \mathbb{R}^n$  be a bounded open set with  $C^1$  boundary and outer normal  $\nu_\Omega : \partial\Omega \rightarrow \mathbb{R}^n$ . Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $G : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be continuously differentiable. Then we have:

$$\begin{aligned} & \int_{\Omega} f \cdot \operatorname{div}(G) \, d\lambda^n(x) \\ &= \int_{\partial\Omega} f \cdot \langle G, \nu_\Omega \rangle \, d\omega(x) - \int_{\Omega} \langle \vec{\nabla} f, G \rangle \, d\lambda^n(x) \end{aligned}$$

*Proof.* By the product rule for divergence, we have:

$$\begin{aligned} & f(x) \cdot \operatorname{div}(G(x)) + \langle \vec{\nabla} f(x), G(x) \rangle \\ &= \operatorname{div}(f(x) \cdot G(x)) \end{aligned}$$

Now, simply apply the divergence theorem, together with basic properties of scalar products, to get:

$$\begin{aligned} & \int_{\Omega} f(x) \cdot \operatorname{div}(G(x)) + \langle \vec{\nabla} f(x), G(x) \rangle \, d\lambda^n(x) \\ &= \int_{\Omega} \operatorname{div}(f(x) \cdot G(x)) \, d\lambda^n(x) \\ &= \int_{\partial\Omega} \langle f(x) \cdot G(x), \nu_\Omega(x) \rangle \, d\omega(x) \\ &= \int_{\partial\Omega} f(x) \cdot \langle G(x), \nu_\Omega(x) \rangle \, d\omega(x), \end{aligned}$$

which we can rearrange to get our desired identity.  $\square$

If we plug  $G(x) := \vec{\nabla} g(x)$  into this formula, we get:

**Corollary 4.63: Green's First Identity**

Let  $\Omega \subset \mathbb{R}^n$  be open and bounded with  $C^1$  boundary. Let  $f : \overline{\Omega} \rightarrow \mathbb{R}$  be continuously differentiable and let  $g : \overline{\Omega} \rightarrow \mathbb{R}$  be twice continuously differentiable. Then we have:

$$\begin{aligned} & \int_{\Omega} f \cdot \Delta g + \langle \vec{\nabla} f, \vec{\nabla} g \rangle \, d\lambda^n \\ &= \int_{\partial\Omega} f \cdot \langle \nu_{\Omega}, \vec{\nabla} g \rangle \, d\omega \\ & \left( = \int_{\partial\Omega} f \cdot \frac{\partial}{\partial \nu_{\Omega}} g \, d\omega(x) \right) \end{aligned}$$

**Theorem 4.64: Green's Second Identity**

Let  $\Omega \subset \mathbb{R}^n$  be open and bounded with  $C^1$  boundary. Let  $f : \overline{\Omega} \rightarrow \mathbb{R}$  and  $g : \overline{\Omega} \rightarrow \mathbb{R}$  be twice continuously differentiable. Then we have:

$$\begin{aligned} & \int_{\Omega} f \cdot \Delta g - g \cdot \Delta f \, d\lambda^n \\ &= \int_{\partial\Omega} f \cdot \langle \nu_{\Omega}, \vec{\nabla} g \rangle - g(x) \cdot \langle \nu_{\Omega}(x), \vec{\nabla} f(x) \rangle \, d\omega(x) \end{aligned}$$

*Proof.* This is just two instances of Green's first identity subtracted from each other:

$$\begin{aligned} & \int_{\Omega} f(x) \cdot \Delta g(x) - g(x) \cdot \Delta f(x) \, d\lambda^n \\ &= \int_{\Omega} f(x) \cdot \Delta g(x) + \langle \vec{\nabla} f(x), \vec{\nabla} g(x) \rangle \, d\lambda^n \\ & \quad - \int_{\Omega} g(x) \cdot \Delta f(x) + \langle \vec{\nabla} f(x), \vec{\nabla} g(x) \rangle \, d\lambda^n \\ &= \int_{\partial\Omega} f(x) \cdot \langle \nu_{\Omega}(x), \vec{\nabla} g(x) \rangle \, d\omega(x) \\ & \quad - \int_{\partial\Omega} g(x) \cdot \langle \nu_{\Omega}(x), \vec{\nabla} f(x) \rangle \, d\omega(x) \\ &= \int_{\partial\Omega} f(x) \cdot \langle \nu_{\Omega}(x), \vec{\nabla} g(x) \rangle - g(x) \cdot \langle \nu_{\Omega}(x), \vec{\nabla} f(x) \rangle \, d\omega(x) \end{aligned}$$

□

## Chapter 5

# Convolutions

Convolution is a way of creating a function by taking a weighted average of two functions. Among other things, it can be used to smooth a given function.

**Lemma 5.1.**  $\tau_h : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be the function that translates a vector by the vector  $h$ , i.e.  $\tau_h(x) = x + h$ . Let  $f \in L^p(\mathbb{R}^n)$  for  $1 \leq p < \infty$ . Then the following hold:

1.  $f \circ \tau_h \in L^p(\mathbb{R}^n)$
2.  $\|f \circ \tau_h\|_p = \|f\|_p$
3.  $\lim_{h \rightarrow 0} \|f \circ \tau_h - f\|_p = 0$

### Definition 5.2: Convolution

Let  $1 \leq p < \infty$ ,  $f \in L^p(\mathbb{R}^n)$  and  $g \in L^1(\mathbb{R}^n)$ . The *convolution of  $f$  and  $g$*  is defined to be:

$$\begin{aligned} f * g : \mathbb{R}^n &\rightarrow \overline{\mathbb{R}} \\ (f * g)(x) &= \int_{\mathbb{R}^n} f(x - y)g(y) d\lambda^n(y) \end{aligned}$$

**Theorem 5.3.** Let  $1 \leq p < \infty$ ,  $f \in L^p(\mathbb{R}^n)$  and  $g \in L^1(\mathbb{R}^n)$ . Then we have:

1.  $f * g \in L^p(\mathbb{R}^n)$
2.  $\|f * g\|_p \leq \|f\|_p \|g\|_1$
3.  $f * g = g * f$

**Theorem 5.4. (Approximation through convolution):** Let  $\eta \in L^1(\mathbb{R}^n)$  such that

$$\int_{\mathbb{R}^n} \eta(x) d\lambda^n(x).$$

Let  $\rho > 0$  and

$$\eta_\rho(x) := \rho^{-n} \cdot \eta\left(\frac{x}{\rho}\right).$$

Let  $1 \leq p < \infty$  and  $f \in L^p(\mathbb{R}^n)$ . Then we have:

1.  $f * \eta_\rho \in L^p(\mathbb{R}^n)$
2.  $\|f * \eta_\rho\|_p \leq \|f\|_p \cdot \|\eta\|_1$
3.  $\|(f * \eta_\rho) - f\|_p \rightarrow 0$

**Definition 5.5: Multi-index Notation**

Let  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ . Then we define:

1.  $|\alpha| = \sum_{i=1}^n \alpha_i$
2.  $\partial^\alpha f(x) = \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \dots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}} f(x)$

**Theorem 5.6: Smoothing**

Let  $\eta \in C^k(\mathbb{R}^n)$  such that for  $|\alpha| \leq k$ , we have

$$\|\partial^\alpha \eta\|_{C^0(\mathbb{R}^n)} \leq K.$$

Let  $f \in L^1(\mathbb{R}^n)$ . Then we have:

1.  $f * \eta \in C^k(\mathbb{R}^n)$ ,
2.  $\partial^\alpha (f * \eta) = f * (\partial^\alpha \eta)$ ,
3.  $\|\partial^\alpha (f * \eta)\|_{C^0(\mathbb{R}^n)} \leq K \|f\|_1$

Smoothing lets us strengthen our result on the density of continuous functions in  $L^p(\Omega)$ :

**Corollary 5.7: Density of  $C_0^\infty$  in  $L^p(\Omega)$**

Let  $\Omega \subset \mathbb{R}^n$  be open. Let  $1 \leq p < \infty$ . Then, for every  $f \in L^p(\Omega)$ , there exists a sequence  $f_k \in C_0^\infty(\Omega)$  such that  $\|f - f_k\|_{L^p(\Omega)} \rightarrow 0$ .

**Definition 5.8: Locally integrable Function**

Let  $\Omega \subset \mathbb{R}^n$  open and  $1 \leq p \leq \infty$ . Then we call a function  $f : \Omega \rightarrow \mathbb{R}$  *locally integrable* iff for any compact set  $K \subset \Omega$ , we have  $\mathbb{1}_K \cdot f \in L^p(\Omega)$ . We denote the set of all locally integrable functions  $f : \Omega \rightarrow \mathbb{R}$  by

$$L_{\text{loc}}^p(\Omega)$$

**Lemma 5.9: Fundamental Lemma of the Calculus of Variations**

Let  $\Omega \subset \mathbb{R}^n$  be open. Let  $f \in L^1_{\text{loc}}(\Omega)$  such that

$$\begin{aligned} &\forall \varphi \in C_0^\infty(\Omega) : \\ &\varphi \geq 0 \implies \int_{\Omega} f \varphi \, d\lambda^n(x) \geq 0 \end{aligned}$$

Then we have  $f(x) \geq 0$  for  $\lambda^n$ -almost all  $x \in \Omega$ .

**Corollary 5.10: There is no identity of convolution.**

There exists no function  $\eta \in L^1(\mathbb{R}^n)$  such that for all  $f \in C_0^\infty(\mathbb{R}^n)$ , we have  $f * \eta = f$ .