

Reflexology: A randomised controlled trial investigating the effects on beta-endorphin, cortisol and pregnancy related stress[☆]

Julie E.M. McCullough^{a, *}, Sarah Dianne Liddle^a, Ciara Close^b, Marlene Sinclair^a,
Ciara M. Hughes^a

^a Institute of Nursing and Health Research, Ulster University, Shore Road, Newtownabbey, Co Antrim, Northern Ireland BT37 0QB, United Kingdom

^b Centre for Public Health, Queens University, Belfast, Northern Ireland BT9 7BL, United Kingdom

ARTICLE INFO

Article history:

Received 13 August 2017

Received in revised form

31 January 2018

Accepted 31 January 2018

Keywords:

Reflexology

Beta-endorphin

Cortisol

Pregnancy

Low back pain

Pelvic girdle pain

ABSTRACT

Reflexology is used for various pregnancy related complaints. A three-armed, pilot randomised controlled trial was conducted to test changes in physiological and biochemical stress parameters. Ninety primiparous volunteers experiencing low back and/or pelvic girdle pain (LBPGP) were recruited to receive either six reflexology or footbath treatments or usual care. Primary outcome data included pain intensity and frequency measured on a visual analog scale (VAS), and salivary beta-endorphin and cortisol levels. 61 (68%) women completed the intervention. A clinically important reduction of 1.63 cm occurred for VAS pain frequency following reflexology. Beta-endorphin levels increased by 8.8% and 10.10% in the footbath and usual care groups respectively and decreased by 15.18% for the reflexology group. Cortisol increased by 31.78% for footbath participants, 31.42% in usual care and 18.82% in the reflexology group. Reflexology during pregnancy may help reduce LBPGP, and associated stress. However, antenatal reflexology is under researched and requires further investigation.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

The mechanism of action of reflexology is not yet understood, however, studies have found a positive effect on quality of life, stress levels and painful conditions [1–4] and it may be helpful for reducing low back pain (LBP) in the general population [5,6]. Reflexology has also been shown to reduce stress markers such as salivary amylase [7] and cortisol [3,8] and a recent systematic review and meta-analysis has concluded that it may have a blood pressure (BP) lowering effect [9]. A popular hypothesis states that reflexology causes a release of endorphins in the body [10], and in this way promotes the associated feelings of health and wellbeing. However, to date, no studies have been identified that have tested this.

Cortisol and beta-endorphin are two of a number of hormones,

involved in stress and the pain response. They are synthesised within the hypothalamus-pituitary-adrenal (HPA) cortex due to the release of corticotropin-releasing hormone (CRH) [11], which leads to the secretion of the pro-hormone pro-opiomelanocortin (POMC) by the pituitary gland. Cleavage of POMC leads to the production of ACTH in the pituitary, which stimulates the production of cortisol in the adrenal cortex and the release of beta-endorphin in the anterior pituitary gland.

Cortisol is released in response to chronic, prolonged physiological and psychological stress such as pain and apprehension [12]. Physiological effects of stress related increased levels of cortisol include decreased pain threshold [13] and elevated heart rate [14]. A correlation in cortisol levels between different body fluids (plasma, blood, urine and saliva) has been documented in both the general population and in pregnant women [15–17].

Beta-endorphin is released into the blood stream from the pituitary gland in response to episodes of acute pain and stress [18,19], and is involved in reducing pain [20], the perception of pain and increasing pain threshold [21]. Beta-endorphin concentrations are commonly evaluated in blood, plasma or cerebrospinal fluid (CSF). There is conflicting evidence regarding what constitutes 'normal' levels of beta-endorphin in saliva for the general population [22,23] and expected plasma beta-endorphin changes during

[☆] This research was part funded by the Department of Employment and Learning, Northern Ireland.

^{*} Corresponding author.

E-mail addresses: j.mccullough@ulster.ac.uk (J.E.M. McCullough), sd.liddle@ulster.ac.uk (S.D. Liddle), c.close@qub.ac.uk (C. Close), m.sinclair1@ulster.ac.uk (M. Sinclair), cm.hughes@ulster.ac.uk (C.M. Hughes).

Twitter@jem_mccullough (J.E.M. McCullough)

pregnancy [24–27] with no previous studies having been identified investigating salivary beta-endorphin in the pregnant population.

One of the most common pregnancy related symptoms is low back pain and/or pelvic girdle pain (LBPGP) which is reported to cause considerable distress and functional limitations for women [28]. The incidence of pregnancy related LBP ranges from 66% up to 71% [29,30] and of pelvic girdle pain from 20% to 65% [29,30] with a typical increase in symptoms as pregnancy progresses [31,32]. This leads to an increase in physiological and psychological stress which alter BP, heart rate (HR), cortisol and beta-endorphin levels.

In order to evaluate any changes related to reflexology treatments during pregnancy a group of low risk pregnant women who were experiencing pain as a result of LBPGP were recruited. This investigation was part of a large pilot study reported elsewhere [33] to evaluate strategies aimed at preventing or managing LBPGP during pregnancy as few guidelines are available, and with physiotherapy being the primary intervention in most cases. Literature reviews have highlighted that CAM may be helpful for managing this type of pain [30,34]. However, these reviews also indicated that reflexology had not yet been investigated as a management strategy. This lack of evidence is of concern to healthcare professionals given that reflexology is currently being used in maternity settings. As many women are unlikely to disclose their use of CAM to their midwife or doctor, the effectiveness and the safety of this treatment requires proper investigation [35,36]. This trial is the first of its kind to investigate the potential changes in salivary cortisol and beta-endorphin levels in pregnant women following the delivery of a complementary and alternative medicine (CAM) intervention.

2. Materials and methods

This trial was listed with the International Standard Randomised Controlled Trial Number Register (ISRCTN26607527), and ethical approval was granted by the Office for Research Ethics Committees Northern Ireland (12/NI/0052, 5 July 2012). The trial, known as the CAM (*Complementary and Alternative Medicine*) in Pregnancy study, ran from July 2012 until December 2013, with primiparous participants recruited from a large inner city maternity out-patients department at their routine 20 week anomaly scan. Using computer generated block randomisation and sealed opaque envelopes, all eligible participants (Table 1) were randomised into one of three arms of the study; reflexology (intervention), footbath (sham treatment), to control for the patient-therapist interaction, and usual antenatal care (control group). This study has been reported elsewhere providing additional study outcome data, including feasibility, recruitment and retention, and details of the reflexology and footbath procedures [37–39].

The reflexology and footbath group participants were asked to provide a 2 ml saliva sample via passive drool into cryovials approximately 10 min before and after their first and last treatment. Those in the usual care group were asked to provide 2 ml saliva samples, at the initial baseline interview and 6 weeks later. All saliva samples were kept on ice to avoid any microbial growth. Samples were held for a maximum of 5 h before being transported to a –80 °C freezer. Participants and therapists were not blinded to treatment allocation in this study, however, to minimise bias, an independent researcher otherwise uninvolved in the study recoded all of the completed questionnaires and saliva samples prior to analysis by the researchers.

2.1. Outcome measures

Patient reported outcome measures were completed by the participant alone and analysed by a researcher blinded to the intervention. A 0–10 cm Visual Analog Scale (VAS) for pain intensity and frequency was completed at baseline and weekly by those in the intervention group. A change on the VAS scale of 15 mm is regarded as clinically significant [40]. The State Trait Anxiety Inventory S-subscale Questionnaire (STAI-S) [41] was completed by all participants at baseline and end of intervention. A score of 39–40 on the S-anxiety scale may detect clinically significant symptoms [42,43] and is categorised as highly anxious [44].

Before and after each treatment, participants in the footbath and reflexology groups had their BP and HR checked and recorded (Omron M2 Basic Blood Pressure Monitor, Omron Corporation, Kyoto, Japan), with participants resting either seated (footbath group) or reclined (reflexology group). In order to reduce the possibility of white coat syndrome the therapists wore black tunics while carrying out all the treatments to negate any effect of increasing BP [45]. Usual care participants did not have their BP and HR measured weekly as it would not be routine or normal practice and, therefore, it would have created an unnecessary burden on participants.

Before each treatment a screening questionnaire was completed to ensure that no contraindications to treatment had developed and to collect data on any medications or interventions that had been used by the participant in the previous week to manage their pain, or any changes or benefits they had noticed. This information was also collected from women in the usual care group via completion of a weekly diary.

2.2. Saliva sampling

All participants had their non-smoking status confirmed

Table 1
Inclusion and exclusion criteria for the CAM in pregnancy study.

Inclusion Criteria	Exclusion Criteria
First time pregnant women	Women pregnant with more than one baby
≥18 years of age	Smokers
Presence of low back pain and/or pelvic girdle pain	Women with neurological diseases
26–29 weeks gestation	Deep Vein Thrombosis (DVT) sufferers
Able to understand written and verbal English	Fungal foot infections or verrucae
	Currently using CAM therapies
	Placenta Previa Grade 3 or 4
	Already participating in a research study
	Any serious spinal pathology e.g. cancer, cauda equina, infection in the spine
	Previous road traffic collision
	Previous surgery to the hip, back or pelvic region
	Inflammatory arthritis, e.g. rheumatoid arthritis
	Diabetes/Gestational diabetes
	Cardiac related problems
	Women whom the midwife deems unable to participate




Fig. 1. CONSORT flow and retention diagram of participants progress throughout the pilot RCT.

(Bedfont Pico Smokerlyzer, Bedfont Scientific Ltd., Maidstone, England) as cigarette smoking can interfere with salivary cortisol analysis [46]. Participants were asked to fast for one hour before sampling in order to limit any fluctuations in pH which may interfere with the assay procedure [47]. Participants were also asked to avoid brushing their teeth for three hours before sampling to avoid any likelihood of oral damage leading to blood contamination and to refrain from alcohol for 24 hours. Before sampling the mouth was rinsed with water in order to clean away any particulates or residue to avoid contamination and interference with chemical biomarkers. A saliva sample was produced into a cryovial (Salimetrics) by passive drool. Samples were stored in a -80°C freezer until analysis at which point they were slowly brought back to room temperature (RT), vortexed and centrifuged at 3000 g for 15 minutes. Enzyme Immunosorbant Assay (EIA) was used to determine the cortisol (Salimetrics, State College, Pennsylvania/Suffolk England) and beta-endorphin (Bachem, Peninsula Laboratories, LLC, San Carlos, California) concentrations in samples.

2.3. Interventions

The reflexology routine was based on the works of Enzer [48], Marquardt [49] and Tiran [50]. All reflexology treatments were carried out by one experienced maternity reflexologist and lasted

for approximately 30 min. Details of the treatment routine have been reported elsewhere [33].

The foot bath intervention participants placed their feet in a purpose-made plastic basin containing comfortably warm water and decorative stones to increase the aesthetic value of the treatment. The water was topped up as required to maintain the temperature and treatments also lasted approximately 30 min. All other conditions were standardised between the two intervention groups including the music played during the treatments, the attire of the therapists, room temperature, waiting area, and public facilities available. Usual care participants followed the usual antenatal care provided by their maternity health care providers.

2.4. Statistics

No power calculation was completed for this study as the main objective was to pilot the methods and procedures. Descriptive statistics, correlations, ANOVA and paired samples t-tests were completed. Data was analysed using Statistical Package for the Social Sciences (SPSS) Version 24. Data was analysed for completeness and whether missing values were missing completely at random (MCAR) using Little's MCAR test. To reduce bias in the analysis, where an outcome measure had less than 10% missing

Table 2

Mean baseline characteristics of the CAM in Pregnancy Study participants at 27 weeks gestation.

	Usual care (n = 23)	Reflexology (n = 23)	Foot bath (n = 15)	P-Value
Age (+/- SD)	30.4 (5.8)	30.4 (4.9)	31.3 (5.5)	0.854
BMI (+/- SD)	25.7 (4.4)	26.1 (3.9)	25.1 (4.1)	0.778
Gestational age (+/- SD)	27.1 (0.9)	27.9 (0.9)	28.3 (1.0)	0.994
Pain type (%)				
LBP ^a	52.2 (12/23)	52.2 (12/23)	53.3 (8/15)	0.989
PP	4.3 (1/23)	4.3 (1/23)	0	
LBPGP	43.5 (10/23)	43.5 (10/23)	46.7 (7/15)	
Gestation at pain onset (+/- SD)	17.2 (5.3)	18.4 (4.9)	18.7 (3.9)	0.845
% with previous LBPGP	47.8 (11/23)	60.2 (14/23)	33.3 (5/15)	0.304
State anxiety (+/- SD)	46.5 (4.3)	43.5 (5.3)	44.2 (3.8)	0.263
Mean heart rate (+/-SD) 95% CI		81.5 (8.8) [80.0–83.0]	81.3 (10.2) [79.0–83.5]	0.737
Mean systolic blood pressure (+/-SD) 95% CI		118.4 (8.9) [116.9–120.0]	113.4 (10.7) [111.0–115.8]	0.190
Mean diastolic blood pressure (+/-SD) 95% CI		70.4 (7.3) [69.1–71.7]	65.6 (6.7) [64.0–67.0]	0.833

data the missing values were input using the multiple imputation (MI) method [51]. Analysis using SPSS showed that the cortisol and beta-endorphin baseline data was not normally distributed. Therefore, non-parametric tests were carried out.

3. Results

An eligible sample of 428 pregnant women were informed of the study at their 20-week appointment and 262 information and consent forms were distributed. Of the 100 women who consented to take part one was excluded as she was a smoker and 90 attended for the baseline meeting and randomisation. Of these 61 (67.8%) completed at least four treatments, end of intervention questionnaires and the requested saliva samples. Final numbers were 23 participants each in the reflexology and usual care groups and 15 footbath participants (Fig. 1).

3.1. Baseline data

Table 2 shows the demographic data collected at the first saliva sample collection time point. Analysis of baseline data using paired-samples *t*-test and One Way ANOVA demonstrated no difference between the groups at baseline.

3.2. Physiological outcome measures

The results from this study found that there were no changes in HR for either treatment group or between groups ($p = 0.865$). Minimal changes in sBP were recorded, with an increase of 3 mmHg for the footbath group and no change for the reflexology group. Results for dBP showed an increase of 1.4 mmHg for both treatment groups over the course of a six-week intervention. There was no significant difference between groups for sBP ($p = 0.311$) or dBP ($p = 0.085$).

3.3. Pain and stress outcomes

There was a clinically significant reduction (1.63 cm) in VAS pain frequency for the reflexology group over the six-week intervention period but not for pain intensity (Table 3). No clinically important changes were observed in the footbath or usual care groups.

State anxiety was high at baseline (Table 2) indicated by a STAI result greater than or equal to 40 [52,53]. Over the course of the trial anxiety levels remained fairly static with no change for the usual care participants and a small increase for the footbath and reflexology participants.

3.4. Biochemical outcome measures

Individual participant samples were collected at the same time of day to reduce within participant variation. Over the six-week intervention period beta-endorphin levels increased by 8.8% and 10.10% in the footbath and usual care groups respectively and decreased by 15.18% for the reflexology group (Table 4/Fig. 2). Cortisol increased by 31.78% in the footbath group, 31.42% in usual care and 18.82% in the reflexology group from baseline to the end of the six-week intervention period (Table 5/Fig. 3). However, there were no significant differences between groups for cortisol ($p = 0.935$) or beta-endorphin ($p = 0.251$). Saliva samples collected before and after treatment one and treatment six showed a within treatment trend for a reduction in salivary beta-endorphin concentration for reflexology and an increase in response to a footbath treatment; this trend was not observed for cortisol response. However, the standard deviations for all the saliva analysis results were very high and indicate that further investigation is required.

There was no correlation between beta-endorphin or cortisol levels and any other outcome variable.

Table 3

Mean VAS frequency and intensity scores (+/-SD) at baseline and week six of the CAM in pregnancy study intervention period from mean gestation 27 weeks–33 weeks.

	VAS frequency baseline	VAS frequency week 6	Change over 6 week period	VAS intensity baseline	VAS intensity week 6	Change over 6 week period
Usual Care (n = 23)	5.86 (2.54)	5.50 (2.81)	-0.36 (2.94)	5.12 (2.19)	5.33 (2.69)	0.21 (3.07)
Reflexology (n = 23)	7.05 (2.36)	5.42 (2.80)	-1.64 (2.97) ^a	5.81 (2.02)	5.14 (2.65)	-0.67 (2.73)
Footbath (n = 15)	5.65 (2.28)	6.13 (2.86)	0.48 (3.28)	4.68 (2.31)	5.63 (2.27)	0.95 (2.86)

^a Denotes a clinically important change.

Table 4
Beta-endorphin concentrations (ng/ml) (+/-SD) at baseline and week six of the CAM in pregnancy study intervention period from mean gestation 27 weeks–33 weeks.

	Beta-endorphin baseline	Beta-endorphin post treatment 1	Beta-endorphin pre- treatment 6	Beta-endorphin post treatment 6/end of intervention
Usual Care (n = 23)	4.29 (3.24) 95% CI 2.88–5.69			4.72 (3.28) 95% CI 3.30–6.14
Reflexology (n = 23)	4.94 (2.62) 95% CI 4.50–5.38	4.60 (3.21) 95% CI 4.06–5.14	4.47 (3.10) 95% CI 3.95–4.99	4.19 (2.99) 95% CI 3.69–4.69
Footbath (n = 15)	3.79 (2.74) 95% CI 3.22–4.37	3.85 (2.97) 95% CI 3.22–4.47	3.52 (2.75) 95% CI 2.94–4.10	4.13 (3.30) 95% CI 3.44–4.83




Fig. 2. Average % change (SEM) in salivary beta-endorphin concentration from baseline to the end of the 6-week CAM in Pregnancy Study intervention period.

Table 5
Cortisol concentrations (μg/dL) (+/-SD) at baseline and week six of the CAM in pregnancy study intervention period from mean gestation 27 weeks–33 weeks.

	Cortisol baseline	Cortisol post treatment 1	Cortisol pre-treatment 6	Cortisol post treatment 6/end of intervention
Usual Care (n = 23)	0.226 (0.33) 95% CI 0.084–0.269			0.297 (0.38) 95% CI 0.133–0.460
Reflexology (n = 23)	0.170 (0.107) 95% CI 0.129–0.224	0.146 (0.060) 95% CI 0.120–0.173	0.187 (0.047) 95% CI 0.166–0.208	0.202 (0.081) 95% CI 0.166–0.238
Footbath (n = 15)	0.129 (0.063) 95% CI 0.094–0.164	0.145 (0.062) 95% CI 0.112–0.180	0.197 (0.065) 95% CI 0.161–0.232	0.170 (0.073) 95% CI 0.130–0.210




Fig. 3. Average % change (SEM) in cortisol concentration from baseline to the end of the 6-week CAM in Pregnancy Study intervention period.

3.5. Other findings

Additional data was collected from weekly screening

questionnaires from the intervention participants and diaries from the usual care participants. Forty percent of the participants used medications to manage their LBPGP and there was little difference

between groups. However, women with some form of PGP tended to use more over the counter (OTC) medications which included paracetamol, co-codamol, ibuprofen and Volterol (diclofenac).

Producing saliva was difficult for 100% of participants at some stage during the study and resulted in one woman in the reflexology group being lost to follow up as she could not produce her final sample.

Women in the footbath and particularly the reflexology groups reported that they had found the treatments they received and the time spent with the therapist to be very relaxing and enjoyable. Some of the participants reported mild tenderness on certain areas of the feet during the reflexology treatments. No other adverse effects occurred.

4. Discussion

This pilot RCT was designed to investigate the use of complementary therapies in a typical clinical setting. This complex intervention investigated quantitative physiological and biochemical outcomes. Close et al. [33,37,38] and McCullough et al. [39] have reported elsewhere on findings relating to the feasibility of the study, recruitment and retention, and additional data including patient reported pain and disability and labour outcomes. However, the small sample size was insufficient to demonstrate statistically significant differences between groups or the true effect of reflexology versus a sham treatment and, therefore, a larger study is warranted.

4.1. Physiological findings

While a reduction in BP due to increased levels of progesterone in the first 18–20 weeks of pregnancy is well documented, BP gradually increases to pre-pregnancy levels during the third trimester [54–56]. Reflexology has been reported to have a BP lowering effect [9], however, the six week course of reflexology and footbaths in the third trimester of pregnancy found only a small increase for sBP and dBp. HR increases throughout pregnancy [57,58]; however, there was no recorded increase, or a notable decrease in heart rate, which is a common effect of relaxation, for either of the treatment groups. It is possible that the cardiovascular effects of reflexology and footbath treatments may have been masked by participants advancing gestation.

4.2. Psychological findings

Elevated stress levels are a feature of pregnancy [59] and LBPGP has been shown to be positively correlated with increased stress and anxiety [60,61]. Therefore, the experience of LBPGP and the associated impact on daily life increases stress, exacerbating painful conditions [62,63] and increasing pain intensity [64]. A cycle of physiological pain and psychological stress ensues leading to biochemical responses. Firstly, due to initial low level pain and active stress, beta-endorphin is released [65–67], followed by cortisol release as the cycle continues into the chronic phase [68,69]. Meanwhile HR and BP become elevated due to the effects of stress [70].

Researchers investigating the use of CAM interventions during pregnancy have repeatedly reported that stress is reduced [71]. However, in the current study no correlation between STAI and stress hormones was found. As the STAI scores remained fairly static across the three groups this would indicate that the STAI was not a sensitive outcome measure for this population. Reductions in pain were observed as VAS for pain frequency increased in the footbath group and decreased in the reflexology and usual care groups, however, the reduction in the reflexology group was a

clinically important change of 1.63 cm. A change on the VAS pain scale of 1.5 cm or more is classified as a clinically important change [40,72], this is a change which can be felt by the patient and provides them with some relief [73]. However, what constitutes a clinically important change in LBPGP during pregnancy is unknown as this has not yet been investigated. Pain perception varies widely between individuals and women's perceptions of pain are altered during pregnancy due to biochemical changes [74,75].

4.3. Biochemical findings

The results from this study have been the first attempt to test the popular hypothesis that reflexology works to reduce pain by initiating the release of beta-endorphin [10]. Kaada and Torsteinbø [76] found an increase in beta-endorphin with connective tissue massage, however, this is a deeper, often uncomfortable manipulative therapy [77] which would not be of the type performed by reflexologists. While the current study found a reduction in pain, it did not support the endorphin release hypothesis as a mechanism of action for this effect, as the results demonstrated a reduction in beta-endorphin levels after a single treatment and also following a course of six weekly treatments. No definitive conclusion can be drawn due to the small numbers included in this study and the high within group variation in results; however, this beta-endorphin reducing effect corresponds to recent research involving healthy participants who received massage [78] which is an integral part of a reflexology treatment. Beta-endorphin was also found to be reduced as pain decreased after temporal mandibular joint (TMJ) surgery [79].

Massage has also been found to reduce cortisol [80,81]. Cortisol is known to increase as a normal part of pregnancy; the increase of 31.42% in cortisol levels for the usual care participants correlated to the findings of Suri et al. [16] who demonstrated a 30.56% increase in salivary cortisol from the second to third trimester. Cortisol is essential for foetal organ maturation [82] during normal healthy pregnancy and levels are well documented [26,83,84]. Therefore, while this study aimed to evaluate whether reflexology could reduce cortisol synthesis, an overall reduction from baseline was not expected. Instead, a reduced cortisol concentration in the treatment groups compared to the usual care group was found, suggesting that the nature of the footbath and to a greater extent reflexology led to a reduction in cortisol response. The reduced cortisol synthesis in the reflexology group is unlikely to have any detrimental effects on the foetus or mother as the detected levels were well within normal limits. The results suggest that reflexology induced positive feelings of wellbeing, which is linked to a lowered cortisol response [85], and the observed reduction in pain may have led to a deactivation of the HPA axis reducing the synthesis of beta-endorphin and cortisol. Therefore, the overall positive experience of reflexology may have been the mechanism affecting the HPA axis.

The patient-therapist interaction certainly had an impact on the outcomes of this trial. This would not only be expected but encouraged as rapport is critical to the overall delivery of reflexology and CAM in general. It is likely that associated reactions to the social support facet of reflexology, which have been found to increase pain threshold [86], may have altered women's experience of their pain during the trial. Biochemical and physiological changes associated with touch therapies, most likely led to the positive outcomes and high participant satisfaction of this study. This or hormonal changes as a result of normal pregnancy and elevated levels may have altered the process by which beta-endorphin is released in the body. Any changes in biochemical and physiological responses due to participant-therapist interaction or the placebo effect further demonstrate that they are integral

to the mode of action of CAM therapies in general and is an area requiring further investigation.

Reflexology has been found to significantly improve tiredness during pregnancy [87] and since many of the women reported better sleep after reflexology this may have also led to the reduction in stress hormones. Sleep was not formally measured as part of the current study; therefore, future investigations into this benefit are warranted given that sleep disruption negatively affects health and wellbeing.

4.4. Limitations

There are several limitations to this study notably the small sample size of self-selecting women who may not be representative of the general pregnant population. Pain frequency at baseline was higher for the reflexology participants compared to the other two groups. This was not statistically significant; however, it may have influenced results. Confounding factors such as spontaneous remission, patient or medical personnel biases, regression to the mean or the effect of unidentified co-interventions may have been responsible for the changes reported. The consumption of alcohol and caffeine [88,89] and waking time may have had an impact on cortisol and beta-endorphin levels. The influence that any of these factors may have had on outcomes was not investigated as part of this study. In addition, Suri, et al. [16] and DiPietro et al. [90] state that daily cortisol levels fluctuate but overall they increase so, given that the cortisol results only give levels for a single point in time, these specific time points may have been minor natural fluctuations in cortisol levels unrelated to the treatment received. Also, cortisol response to stress during pregnancy is related to the stage of pregnancy [91]. The wide variation in beta-endorphin concentrations within the groups prevents any firm conclusions being drawn at this time. In addition, it would have been preferable to also measure BP and HR in the usual care group to compare and contrast with the findings from the reflexology and foot bath groups.

Pregnant women were keen to experience reflexology. A lack of participant blinding to the trial interventions under investigation, which could not be avoided given the nature of this study, resulted in 97% of those women randomised to assert that they would have preferred to be allocated to the reflexology arm of the trial. Prior to randomisation there was significantly lower expectations for the helpfulness of footbath compared with reflexology. Blinding in trials of this nature is known to be problematic and is a factor contributing to bias in future trials where the control measure is not mutually acceptable to participants. Furthermore, the participant-therapist interaction, which is an integral part of CAM therapies, will also have affected physiological and biochemical results. The high rates of attrition and the reasons for this may also have influenced results and this has been discussed elsewhere [37].

Collection and analysis of saliva is known to be quick and simple to carry out and, therefore, was chosen as it is less invasive than blood collection, which may have had an impact on the stress hormones under investigation. However, all of the pregnant participants found the process of expectorating saliva into the collection tubes troublesome, frustrating and 'messy'. Many also had difficulty producing enough saliva, with one woman failing to do so, stating that it made them feel nauseous. This condition had not been previously reported during pregnancy or saliva collection from pregnant women.

5. Conclusion

These results suggest that the nature of reflexology may have acted to reduce pain, disrupting the cycle of pain and stress leading

to a reduction in the stress related release of cortisol and beta-endorphin over the six-week intervention period. It is also possible that other biochemicals released in response to reflexology lead to a down regulation of beta-endorphin.

This study was underpowered to provide definitive conclusions regarding findings, however, the trends in the reduction in stress hormones adds much needed quantitative evidence for the effects of reflexology and the use of reflexology during pregnancy and highlights the need for further investigation in the area. The extent of any benefits that this inexpensive, non-pharmacological, non-invasive, safe, patient preferred treatment can offer to pregnant women and the general population have clearly yet to be elucidated.

Competing interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

Acknowledgments

The authors wish to thank all of the women who took part in this research, and the midwives and staff at the XXXXXX XXXXXX. Thanks also to the late Professor George Lewith for his expert advice with this work and Dr Paul Slater and Professor Ian Bradbury for their statistical advice.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.ctcp.2018.01.018>.

References

- [1] N.L.N. Stephenson, S.P. Weinrich, Abbas S. Tavakoli, The Effects of Foot Reflexology on Anxiety and Pain in Patients With Breast and Lung Cancer, *Oncol. Nurs. Forum* 27 (1) (2000).
- [2] C.M. Hughes, S. Smyth, A.S. Lowe-Strong, Reflexology for the treatment of pain in people with multiple sclerosis: a double-blind randomised sham-controlled clinical trial, *Mult. Scler. J.* 15 (11) (2009) 1329–1338.
- [3] N.A. Hodgson, D. Lafferty, Reflexology versus Swedish massage to reduce physiologic stress and pain and improve mood in nursing home residents with cancer: a pilot trial, *Evid. base Compl. Alternative Med.* (2012), 456897, <https://doi.org/10.1155/2012/456897>.
- [4] G. Wyatt, A. Sikorskii, M.H. Rahbar, D. Victorson, M. You, Health related quality-of-life outcomes: a reflexology trial with patients with advanced-stage breast cancer, *Oncol. Nurs. Forum* 39 (6) (2012) 568–577.
- [5] F. Quinn, C.M. Hughes, G.D. Baxter, Reflexology in the management of low back pain: a pilot randomised controlled trial, *Compl. Ther. Med.* 16 (1) (2008) 3–8.
- [6] M. Eghbali, R. Safari, F. Nazari, S. Adboli, The effects of reflexology on chronic low back pain intensity in nurses employed in hospitals affiliated with Isfahan University of Medical Sciences, Iran, *J. Nurs. Midwifery Res.* 17 (3) (2012) 239–243.
- [7] N.A. Hodgson, S. Andersen, The clinical efficacy of reflexology in nursing home residents with dementia, *J. Alternative Compl. Med.* 14 (3) (2008) 269–275.
- [8] P.A. Mackereth, K. Booth, V.F. Hillier, A.L. Caress, Reflexology and progressive muscle relaxation training for people with multiple sclerosis: a crossover trial, *Compl. Ther. Clinical Practice* 15 (1) (2009) 14–21.
- [9] J.E.M. McCullough, S.D. Liddle, M. Sinclair, C. Close, C.M. Hughes, The physiological and biochemical outcomes associated with a reflexology treatment: a systematic review, *Evid. base Compl. Alternative Med.* (2014). Available online at: <http://www.hindawi.com/journals/ecam/2014/502123/cta/>.
- [10] D. Tiran, H. Chummun, Complementary therapies to reduce physiological stress in pregnancy, *Compl. Ther. Nurs. Midwifery* 10 (2004) 162–167.
- [11] H. Kunugi, H. Hori, N. Adachi, T. Numakawa, Interface between hypothalamic-pituitary-adrenal axis and brain-derived neurotrophic factor in depression, *Psychiatr. Clin. Neurosci.* 64 (5) (2010) 447–459.
- [12] E. Russell, G. Koren, M. Rieder, S. Van Uum, Hair cortisol as a biological marker of chronic stress: current status, future directions and unanswered questions, *Psychoneuroendocrinology* 37 (5) (2012) 589–601.
- [13] J.C. Choi, M.I. Chung, Y.D. Lee, Modulation of pain sensation by stress-related testosterone and cortisol, *Anaesthesia* 67 (10) (2012) 1146–1151.
- [14] R.R. Looser, P. Metzenthin, S. Helfricht, B.M. Kudielka, A. Loerbroks, J.F. Thayer,

- J.E. Fischer, Cortisol is significantly correlated with cardiovascular responses during high levels of stress in critical care personnel, *Psychosom. Med.* 72 (3) (2010) 281–289.
- [15] T. Umeda, R. Hiramatsu, T. Iwaoka, T. Shimada, F. Miura, T. Sato, Use of saliva for monitoring unbound free cortisol levels in serum, *Clin. Chim. Acta* 110 (1981) 245–253.
 - [16] D. Suri, J. Moran, J.U. Hibbard, K. Kasza, R.E. Weiss, Assessment of adrenal reserve in pregnancy: defining the normal response to the adrenocorticotropin stimulation test, *J. Clin. Endocrinol. Metabol.* 91 (2006) 3866–3872.
 - [17] S. Sakihara, K. Kageyama, Y. Oki, M. Doi, Y. Iwasaki, S. Takayasu, T. Moriyama, K. Terui, T. Nigawara, Y. Hirata, K. Hashimoto, T. Suda, Evaluation of plasma, salivary, and urinary cortisol levels for diagnosis of Cushing's syndrome, *Endocr. J.* 57 (4) (2010) 331–337.
 - [18] D.S. Sheps, M.N. Ballenger, G.E. De Gent, R. Krittaphong, E. Dittman, W. Maixner, W. McCartney, R.N. Golden, G. Koch, K.C. Light, Psychophysical responses to a speech stressor: correlation of plasma beta-endorphin levels at rest and after psychological stress with thermally measured pain threshold in patients with coronary artery disease, *J. Am. Coll. Cardiol.* 25 (7) (1995) 1499–1503.
 - [19] A.S. Sprouse-Blum, G. Smith, D. Sugai, D.F. Parsa, Understanding endorphins and their importance in pain management, *Hawaii Med. J.* 69 (3) (2010) 70–71.
 - [20] A.C. Hartwig, Peripheral beta-endorphin and pain modulation, *Anesth. Prog.* 38 (3) (1991) 75–78.
 - [21] N.F. Jarmukli, J. Ahn, A. Iranmanesh, D.C. Russell, Effect of raised plasma beta-endorphin on peripheral pain and angina thresholds in patients with stable angina, *Heart* 82 (1999) 204–209.
 - [22] N. Takai, M. Yamaguchi, T. Aragaki, K. Eto, K. Uchihashi, Y. Nishikawa, Gender-specific differences in salivary biomarker responses to acute psychological stress, *Ann. N. Y. Acad. Sci.* 1098 (2007) 510–515.
 - [23] B. Rai, J. Kaur, Salivary stress markers and psychological stress in simulated microgravity: 21 days in 6° head-down tilt, *J. Oral Sci.* 53 (1) (2011) 1343–4934.
 - [24] D.I. Hoffman, T.K. Abboud, H.R. Haase, T.T. Hung, U. Goebelsmann, Plasma beta-endorphin concentrations prior to and during pregnancy, in labor, and after delivery, *Am. J. Obstet. Gynecol.* 150 (5 Pt 1) (1984) 492–496.
 - [25] T.K. Abboud, Maternal and fetal beta endorphin: effects of pregnancy and labour, *Arch. Dis. Child.* 63 (1988) 707–709.
 - [26] E.C. Chan, R. Smith, T. Lewin, M.W. Brinsmead, H.P. Zhang, J. Cubis, K. Thornton, D. Hurt, Plasma corticotropin-releasing hormone, beta-endorphin and cortisol inter-relationships during human pregnancy, *Acta Endocrinol (Copenh)* 128 (4) (1993) 339–344.
 - [27] C. De Punzio, E. Neri, P. Metelli, M.S. Bianchi, M. Venticinque, M. Ferdeghini, P. Fioretti, The relationship between maternal relaxation and plasma beta-endorphin levels during parturition, *J. Psychosom. Obstet. Gynaecol.* 15 (4) (1994) 205–210.
 - [28] M. Sinclair, C. Close, J.E.M. McCullough, C.M. Hughes, S.D. Liddle, How do women manage pregnancy-related low back and/or pelvic pain? Descriptive findings from an online survey, *Evid. base Midwifery* 12 (3) (2014) 76–82.
 - [29] F.M. Kovacs, E. Garcia, A. Royuela, L. González, V. Abaira, Spanish Back Pain Research Network, Prevalence and factors associated with low back pain and pelvic girdle pain during pregnancy: a multicenter study conducted in the Spanish National Health Service, *Spine (Phila Pa 1976)* 37 (17) (2012) 1516–1533, 2012 Aug 1.
 - [30] S.D. Liddle, V. Pennick, Interventions for preventing and treating low-back and pelvic pain during pregnancy, *Cochrane Database Syst. Rev.* 9 (2015). CD001139.
 - [31] E. Vermani, R. Mittal, A. Weeks, Pelvic girdle pain and low back pain in pregnancy: a review, *Pain Pract.* 10 (1) (2010) 60–71.
 - [32] E. Nazik, G. Eryilmaz, Incidence of pregnancy-related discomforts and management approaches to relieve them among pregnant women, *J. Clin. Nurs.* 23 (2013) 1736–1750.
 - [33] C. Close, J.E.M. McCullough, M. Sinclair, S.D. Liddle, C.M. Hughes, A pilot randomised controlled trial investigating the effectiveness of reflexology for managing pregnancy low back and pelvic pain, *Compl. Ther. Clin. Pract.* (2015). <https://doi.org/10.1016/j.ctcp.2015.05.00>.
 - [34] C. Close, M. Sinclair, S.D. Liddle, E. Madden, J.E.M. McCullough, C.M. Hughes, A systematic review investigating the effectiveness of Complementary and Alternative Medicine (CAM) for the management of low back and/or pelvic pain (LBPP) in pregnancy, *J. Adv. Nurs.* 70 (8) (2014) 1702–1716.
 - [35] H.R. Hall, K. Jolly, Women's use of complementary and alternative medicines during pregnancy: a cross-sectional study, *Midwifery* 30 (5) (2014) 499–505.
 - [36] S. Warriner, K. Bryan, A.M. Brown, Women's attitude towards the use of complementary and alternative medicines (CAM) in pregnancy, *Midwifery* 30 (1) (2014) 138–143.
 - [37] C. Close, M. Sinclair, J.E.M. McCullough, D. Liddle, C. Hughes, Factors affecting recruitment and attrition in randomised controlled trials of complementary and alternative medicine for pregnancy-related issues, *Evid. base Compl. Alternative Med.* 2016 (2016). <https://doi.org/10.1155/2016/6495410>.
 - [38] C. Close, M. Sinclair, D. Liddle, J. McCullough, C. Hughes, Women's experience of low back and/or pelvic pain (LBPP) during pregnancy, *Midwifery* 37 (2016) 1–8.
 - [39] J.E.M. McCullough, C. Close, S.D. Liddle, M. Sinclair, C.M. Hughes, Results from a pilot randomised controlled trial exploring the effects of antenatal reflexology on labour outcomes, *Midwifery* 55 (2017) 137–144.
 - [40] E.J. Gallagher, M. Liebman, E. Polly, P.E. Bijur, Prospective validation of clinically important changes in pain severity measured on a visual analog scale, *Ann. Emerg. Med.* 38 (6) (2001) 633–638.
 - [41] C.D. Spielberger, R.L. Gorsuch, P.R. Lushene, P.R. Vagg, A.G. Jacobs, Manual for the State-trait Anxiety Inventory (Form Y), 1983. Available at: faculty.sjcn.edu/~State%20Trait%20Anxiety%20Inventory.doc. (Accessed 10 January 2012).
 - [42] R.G. Knight, H.J. Waal-Manning, G.F. Spears, Some norms and reliability data for the state-trait anxiety inventory and the zung self-rating depression scale, *Br. J. Clin. Psychol.* 22 (4) (1983) 245–249.
 - [43] G. Addolorato, C. Ancona, E. Capristo, R. Graziosetto, L. Di Rienzo, M. Maurizi, G. Gasbarrini, State and trait anxiety in women affected by allergic and vasomotor rhinitis, *J. Psychosom. Res.* 46 (3) (1999) 283–289.
 - [44] K.A. Grant, C. McMahon, M.P. Austin, Maternal anxiety during the transition to parenthood: a prospective study, *J. Affect. Disord.* 108 (1–2) (2008) 101–111.
 - [45] E.S. Muxfeldt, K.V. Bloch, A.R. Nogueira, G.F. Salles, Twenty-four hour ambulatory blood pressure monitoring pattern of resistant hypertension, *Blood Press. Monit.* 8 (5) (2003) 181–185.
 - [46] C. Kirschbaum, S. Wust, C.J. Strasburger, 'Normal' cigarette smoking increases free cortisol in habitual smokers, *Life Sci.* 50 (6) (1992) 435–442.
 - [47] E.B. Schwartz, D.A. Granger, E.J. Susman, M.R. Gunnar, B. Laird, Assessing salivary cortisol in studies of child development, *Child Dev.* 69 (6) (1998) 1503–1513.
 - [48] S. Enzer, *Maternity Reflexology Manual*, second ed., Soul to Sole Reflexology Limited, England, 2004.
 - [49] H. Marquardt, *Reflex Zone Therapy of the Feet: A Comprehensive Guide for Health Professionals*, Healing Arts Press, Vermont, 2007.
 - [50] D. Tiran, *Reflexology in Pregnancy and Childbirth*, Churchill Livingstone Elsevier, London, 2010.
 - [51] D.A. Bennett, How can I deal with missing data in my study? *Aust. N. Z. J. Publ. Health* 25 (5) (2001) 464–469.
 - [52] J.M.A. Teixeira, N.M. Fisk, V. Glover, Association between maternal anxiety in pregnancy and increased uterine artery resistance index: cohort based study, *BMJ* 318 (7177) (1999) 153–157.
 - [53] J.J. Newham, A. Wittkowski, J. Janine Hurley, J.D. Aplin, M. Westwood, Effects of antenatal yoga on maternal anxiety and depression: a randomized controlled trial, *Depress. Anxiety* 00 (2014) 1–10, 18/7/14, <http://onlinelibrary.wiley.com/doi/10.1002/da.22268/full>.
 - [54] T.M. Deak, J.B. Moskovitz, Hypertension and pregnancy, *Emerg. Med. Clin. N Am* 30 (4) (2012) 903–917.
 - [55] R. Mustafa, S. Ahmed, A. Gupta, R.C. Venuto, A comprehensive review of hypertension in pregnancy, *J. Pregnancy* 2012 (2012) 105918, <https://doi.org/10.1155/2012/105918>. Published online May 23, 2012. (Accessed 13 May 2014).
 - [56] A.G. Kattah, V.D. Garovic, The management of hypertension in pregnancy, *Adv. Chron. Kidney Dis.* 20 (3) (2013) 229–239.
 - [57] M.E. Hall, E.M. George, J.P. Granger, The heart during pregnancy, *Rev. Esp. Cardiol. (English Edition)* 64 (11) (2011) 1045–1050.
 - [58] J.G. Ouzounian, U. Elkayam, Physiologic changes during normal pregnancy and delivery, *Cardio. Clinics Cardiovasc. Dis. Pregnancy* 30 (3) (2012) 317–329.
 - [59] R.M. El-Bsat, Study regarding the benefits of applying hospital anxiety-depression scale (HAD) in athletes during pregnancy, 3rd international congress on physical education, sport and kineotherapy (ICPESK 2013), *Procedia - Soc. Behav. Sci.* 117 (19) (2014) 566–569.
 - [60] G.B.J. Andersson, Epidemiological features of chronic low-back pain, *Lancet* 354 (9178) (1999) 581–585.
 - [61] A. Bener, M. Verjee, E.E. Dafeeah, O. Falah, T. Al-Juhaishi, J. Schlogl, A. Sedeeq, S. Khan, Psychological factors: anxiety, depression, and somatization symptoms in low back pain patients, *J. Pain Res.* 6 (2013) 95–101.
 - [62] J.L. Rhudy, M.W. Meagher, Fear and anxiety: divergent effects on human pain thresholds, *Pain* 84 (1) (2000) 65–75.
 - [63] T. Reinhardt, N. Kleindienst, R.-D. Treede, Individual modulation of pain sensitivity under stress, *Pain Med. US* 14 (5) (2013) 676–685.
 - [64] J. Tang, S.J. Gibson, A Psychophysical evaluation of the relationship between trait anxiety, pain perception, and induced state anxiety, *J. Pain* 6 (9) (2005) 612–619.
 - [65] R.I.M. Dunbar, The social role of touch in humans and primates: behavioural function and neurobiological mechanisms, *Neurosci. Biobehav. Rev.* 34 (2) (2010) 260–268.
 - [66] R. Graignic-Philippe, J. Dayan, S. Chokron, A.-Y. Jacquet, S. Tordjman, Effects of prenatal stress on fetal and child development: a critical literature review, *Neurosci. Biobehav. Rev.* 43 (2014) 137–162.
 - [67] J.G. Veening, H.P. Barendregt, The effects of beta-endorphin: state change modification, *Fluids Barriers CNS* 12 (3) (2015).
 - [68] M. Boudarene, J.J. Legros, M. Timsit-Berthier, Study of the stress response: role of anxiety, cortisol and DHEAs, *L'Encephale* 28 (2) (2002) 139–146.
 - [69] N. Jacobs, I. Myin-Germeys, C. Derom, P. Delespaul, J. van Os, N.A. Nicolson, A momentary assessment study of the relationship between affective and adrenocortical stress responses in daily life, *Biol. Psychol.* 74 (1) (2007) 60–66.
 - [70] M. Esler, N. Eikelis, M. Schlaich, G. Lambert, M. Alvarenga, T. Dawood, D. Kaye, D. Barton, C. Pier, L. Guo, C. Brencley, G. Jennings, E. Lambert, Chronic mental stress is a cause of essential hypertension: presence of biological markers of stress, *Clin. Exp. Pharmacol. Physiol.* 35 (4) (2008) 498–502.

- [71] J.J. Newham, et al., State–trait anxiety inventory (STAI) scores during pregnancy following intervention with complementary therapies, *J. Affect. Disord.* (2012). <https://doi.org/10.1016/j.jad.2012.04.027>. (Accessed 26 March 2014).
- [72] R.W. Ostelo, R.A. Deyo, P. Stratford, G. Waddell, P. Croft, K.M. Von, L.M. Bouter, H.C. de Vet, Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change, *Spine* 33 (1) (2008) 90–94.
- [73] J.T. Farrar, R.K. Portenoy, J.A. Berlin, J.L. Kinman, B.L. Strom, Defining the clinically important difference in pain outcome measures, *Pain* 88 (3) (2000) 287–294.
- [74] A.R. Gintzler, N.H. Liu, The maternal spinal cord: biochemical and physiological correlates of steroid-activated antinociceptive processes, *Prog. Brain Res.* 133 (2001) 83–97.
- [75] G. Draisci, S. Catarci, C. Vollono, B.A. Zanfini, C. Pazzaglia, C. Cadeddu, D. Virdis, M. Valeriani, Pregnancy-induced analgesia: a combined psychophysical and neurophysiological study, *Eur. J. Pain* 16 (2012) 1389–1397.
- [76] B. Kaada, O. Torsteinbø, Increase of plasma beta-endorphins in connective tissue massage, *Gen. Pharmacol.* 20 (4) (1989) 487–489.
- [77] G.C. Goats, K.A. Keir, Connective tissue massage, *Br. J. Sports Med.* 25 (3) (1991) 131–133.
- [78] V. Morhenn, L.E. Beavin, P.J. Zak, Massage increases oxytocin and reduces adrenocorticotropin hormone in humans, *Alternative Ther. Health Med.* 18 (6) (2012) 11–18.
- [79] A. Feldreich, M. Ernberg, A. Rosén, Reduction in maximum pain after surgery in temporomandibular joint patients is associated with decreased beta-endorphin levels—a pilot study, *Int. J. Oral Maxillofac. Surg.* 46 (1) (2017) 97–103.
- [80] T. Field, M. Hernandez-Reif, M. Diego, S. Schanberg, C. Kuhn, Cortisol decreases and serotonin and dopamine increase following massage therapy, *Int J Neurosci.* 115 (10) (2005) 1397–1413.
- [81] L. Lindgren, S. Rundgren, O. Winsö, S. Lehtipalo, U. Wiklund, M. Karlsson, H. Stenlund, C. Jacobsson, C. Brulin, Physiological responses to touch massage in healthy volunteers, *Auton. Neurosci. Basic Clin.* 158 (1–2) (2010) 105–110.
- [82] J.R. Challis, D. Sloboda, S.G. Matthews, A. Holloway, N. Alfaidy, F.A. Patel, W. Whittle, M. Fraser, T.J. Moss, J. Newnham, The fetal placental hypothalamic–pituitary–adrenal (HPA) axis, parturition and postnatal health, *Mol. Cell. Endocrinol.* 185 (1–2) (2001) 135–144.
- [83] G.C. Liggins, The role of cortisol in preparing the fetus for birth, *Reprod. Fertil. Dev.* 6 (2) (1994) 141–150.
- [84] C. Buss, E.P. Davis, B. Shahbaba, J.C. Pruessner, K. Head, C.A. Sandman, Maternal cortisol over the course of pregnancy and subsequent child amygdala and hippocampus volumes and affective problems, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) E1312–E1319.
- [85] P. Lindfors, U. Lundberg, Is low cortisol release an indicator of positive health? *Stress Health* 18 (4) (2002) 153–160.
- [86] K. Uvnäs-Moberg, I. Arn, D. Magnusson, The psychobiology of emotion: the role of the oxytocinergic system, *Int. J. Behav. Med.* 12 (2) (2005) 59–65.
- [87] F. Ghaffari, T.P. Ghaznein, The reflexology of sole on tiredness intensity in pregnant women, *Caspian J. Intern. Med.* 1 (2) (2010) 58–62.
- [88] C. Gianoulakis, X. Dai, T. Brown, Effect of chronic alcohol consumption on the activity of the hypothalamic–pituitary–adrenal Axis and pituitary β -endorphin as a function of alcohol intake, age, and gender, *Alcohol Clin. Exp. Res.* 27 (3) (2003) 410–423.
- [89] W.R. Lovallo, T.L. Whitsett, M. al'Absi, B.H. Sung, A.S. Vincent, M.F. Wilson, Caffeine stimulation of cortisol secretion across the waking hours in relation to caffeine intake levels, *Psychosom. Med.* 67 (5) (2005) 734–739.
- [90] J.A. DiPietro, K.A. Costigan, K.T. Kivlighan, P. Chen, M.L. Laudenslager, Maternal salivary cortisol differs by fetal sex during the second half of pregnancy, *Psychoneuroendocrinology* 36 (4) (2011) 588–591.
- [91] C. Obel, M. Hedegaard, T.B. Henriksen, N.J. Secher, J. Olsen, S. Levine, Stress and salivary cortisol during pregnancy, *Psychoneuroendocrinology* 30 (7) (2005) 647–656.