

Seminar Computeralgebra 2018

Geck

- Introduction of
 - algebraic sets in affine space
 - affine algebras and algebraic groups
- Section 1.1
 - basic definitions
 - version of Hilbert's Nullstellensatz for hypersurfaces
- Section 1.2
 - Groebner bases and applications
 - define dimension of an algebraic set by Hilbert polynomial of an ideal

Kapitel 1.1

- k is a field and $k[X_1, \dots, X_n]$ a polynomial ring with X_i independent- commutative ring A with 1 is noetherian ring if every ideal of A is generated by finite number of elements
- Hilbert's Basis Theorem: If A is noetherian and X an indeterminate over A , then $A[X]$ is also noetherian. If k is a field, then $k[X_1, \dots, X_n]$ is noetherian.
 - $x \in k^n \Rightarrow \exists \varepsilon_x : k[X_1, \dots, X_n] \rightarrow k$ (Einsetzungshomomorphismus)
 - f is uniquely determined by \bar{f} if $|k| = \text{infinity}$
- $V(S)$ ist die Menge aller gemeinsamen Nullstellen aller Polynome aus S
 - genannt algebraisches Set
 - $V(S) = V(I)$, wenn $I =$ das von S generierte Ideal ist
 - $I(V)$ sind alle Polynome, für die alle $x \in V$ Nullstellen sind (genannt *vanishing ideal*)
 - S ist Teilmenge, algebraisches Set wird gefunden, daraus wird wieder eine Teilmenge des Polynomrings kreiert (diese ist größer/gleich S und ein Ideal)
 - $A[V] := k[X_1, \dots, X_n]/I(V)$ heißt affine Algebra von V
 - Schnitte von algebraischen Sets sind wieder ein algebraisches Set, unendliche Vereinigungen jedoch nicht
- Zariski topology
 - beliebige Schnittmengen und endliche Vereinigungen von algebraischen Sets sind algebraische Sets, genauso wie \emptyset and k^n
 - $X \subseteq k^n$ is open, if $k^n \setminus X$ is closed (algebraic)
 - $V \subseteq k^n$ (not necessarily algebraic) then $\bar{V} = V(I(V))$ is its closure in the Zariski topology $\Rightarrow V(I(V)) = V$ if V is algebraic set, hence $I : \{V \subseteq k^n | V \text{ algebraic}\} \rightarrow \{I \subseteq k[X_1, \dots, X_n] | I \text{ ideal}\}$ is injective
 - I is not surjective: f polynomial and non-constant, then $H_f := V(f) \subseteq k^n$ is called hypersurface ($n = 2 \Rightarrow$ plane curve), it may happen that

- $H_f = \emptyset$, but only if k is algebraically closed and $n \geq 2$
- Hilbert's Nullstellensatz for hypersurfaces: k algebraically closed field, $f \in k[X_1, \dots, X_n]$ non-constant and $\emptyset \neq H_f \subseteq k^n$ the corresponding hypersurface. $f = f_1^{n_1} \cdots f_r^{n_r}$ with f_1, \dots, f_r irreducible and pairwise coprime, then $H_f = H_{f_1} \cup \cdots \cup H_{f_r}$ and $I(H_f) = (f_1 \cdots f_r)$.
 - Topological space $Z \neq \emptyset$ is reducible, if $Z = Z_1 \cup Z_2$ where Z_1, Z_2 are non-empty closed subsets with $Z_i \neq Z$
 - $Y \subseteq Z$ is irreducible if Y is irreducible with induced topology
 - \emptyset is not considered irreducible
 - Z is noetherian topological space if every chain of closed sets $Z_1 \supseteq Z_2 \supseteq \cdots$ in Z becomes stationary
 - $Z \neq \emptyset$ is noetherian topological space, then there are only finitely many maximal closed irreducible subsets in Z ; they are called the irreducible components of Z . If these are Z_1, \dots, Z_r , we have $Z = Z_1 \cup \cdots \cup Z_r$.
 - $\emptyset \neq V \subseteq k^n$ is algebraic set
 - V is noetherian with respect to the Zariski topology. Thus $V = V_1 \cup \cdots \cup V_r$ where V_i are maximal closed irreducible subsets of V .
 - V is irreducible (in Zariski topology) $\Leftrightarrow I(V)$ is prime ideal.
 - Principal open subsets: V non-empty algebraic set, $f \in k[X_1, \dots, X_n]$ such that $f \notin I(V)$ and $V_f := \{v \in V \mid f(v) \neq 0\} \subseteq V$. Then V_f is non-empty open set in V and called *principal open set*. Any open set in V is a finite union of principal open sets.
 - Indeed let $U \subseteq V$ be open. Then V minus U is closed and so there exists an algebraic set W such that $V \setminus U = W \cap V$. Now, by Hilbert's Basis Theorem, $W \cap V$ is defined by a finite collection of polynomials, f_1, \dots, f_r . Then we have $U = V_{f_1} \cup \cdots \cup V_{f_r}$.
 - Principle open sets can be regarded as algebraic sets in their own right.

Kapitel 1.2

Thoughts

- Algebraische Geometrie: Menge von Nullstellen einer Menge von Polynomen
 - Algebraisches Set: K Körper, S Menge von Polynomen in n Variablen, $V(S)$ Menge von gemeinsamen Nullstellen und $V(S) \subset K^n$
 - Algebraische Gruppe: Gruppe bzgl. Polynomfunktionen und algebraisches Set
 - Historisch: Analytischer Kontext und Lie Gruppen
- Algebraisch abgeschlossener Körper K : Jedes Polynom mit Koeffizienten aus K hat eine Nullstelle in K , z.B. \mathbb{C}
- Es gibt Garben, Halme und Keime! :)
- Definition of "algebraic" (see e.g. Example 1.1.3(c))
- "maximal closed" (Proposition 1.1.11)

- Principal open subsets: Warum sind das Subsets? Das stimmt gar nicht...
- Lemma 1.1.15

CLOS

Kapitel 1

Kapitel 2

Kapitel 9