

Optimal Neural Network Feature Selection for Forecasting of Spatial-Temporal Series

Eurico Covas, others, and many,

Abstract—We show empirical evidence on how to construct the optimal feature selection or architecture for the input layer on a neural network used for forecasting of spatial-temporal series. The approach is based on results from dynamical systems, namely the non-linear embedding theorems. We demonstrate it for a two dimensional signal with one spatial dimension and one time dimension, and show that the optimal input layer seems to consist of a two dimensional grid, with spatial/temporal lags determined by the minimum of the mutual information of the spatial/temporal signal subsets and the number of points taken in space/time decided by the embedding dimensional of signal. We support our evidence by running a Monte Carlo simulation of several combinations of input layer architecture and showing that the one decided by the non-linear embedding theorems seems to be optimal or close of optimal.

Index Terms—IEEE, IEEEtran, journal, L^AT_EX, paper.

I. INTRODUCTION

IN this paper we show empirical evidence for an optimal architecture of the input layer of a neural network which tries to forecast or predict a spatial temporal signal. Here we show the empirical evidence for two particular cases of two dimensional data series s_m^n (one spatial, one temporal dimension). By two dimensional data series we mean a scalar field which can be defined by a $N \times M$ matrix with components $s_m^n \in \mathbb{R}$.

A. Neural networks for time-series forecasting

DO WE REALLY NEED THIS section....

B. Neural networks for spatial temporal forecasting

There has been several attempts to forecast spatial-temporal data series in the literature (add references). The references above confine themselves to articles that attempt to forecast the actual full scalar field s_m^n , as opposed to the ongoing research on pattern recognition in moving images (2D and 3D), which attempt to pick particular features in images (e.g. car, pedestrian, bicycle, person) and to forecast where those features will be in subsequent images within the particular moving sequence.

E. Covas is with the CITEUC, Geophysical and Astronomical Observatory, University of Coimbra, 3040-004, Coimbra, Portugal, and Queen Mary University London, 10 Godward Square, Mile End Rd, London E1 4FZ, e-mail: (eurico.covas@mail.com)

Manuscript received December 31, 2017; revised December 31, 2017.

C. Sunspot and solar data series forecasting using neural networks

Neural networks, among other forecasting methods (add references) has been used extensively to try to predict sunspots and related data series. Mostly this has been done in time only (add references). There are a few examples (add references) of actual spatial temporal forecasts using neural networks (see mine and the one on MDI/HDI data).

D. Input layer architecture for neural networks for spatial temporal forecasting

All of the references above on neural network forecasting of spatial temporal data series either use a simple delay based architecture for the input layer, or use a time delay based on the first minima of the *mutual information* as proposed in [1]–[3]) and/or use the number of points dictated by the embedding theorems by [4], [5], [6] and [7] using the method of *method of false nearest neighbours* detection suggested by [8] and reported in detail in [2], [9]–[11].

However, as far as we are aware, all the references to using the embedding theorems and the related mutual information method and the false nearest neighbours method seems to be not justified, i.e., the approach is explained but not proven either theoretically or empirically.

Here we show two cases, one based on the articles of one of us on sunspot forecasting ([12]–[14]) and one based on the Kuramoto-Sivashinsky (KS) equation (add references and add parlitz paper here)

Several authors [15]–[66] have already attempted to use neural networks to forecast aspects of the sunspot cycle, although none in both space and time, having restricted themselves to using these neural networks to forecast mostly either the sunspot number or the sunspot areas as a function of time.

Takens established the theoretical background [5] in the Takens embedding theorem. Further developments established a series of theorems by [4], [5], [6] and [7].

Some authors discuss the use of either mutual information and/or embedding dimension as a constraint on the input layer, i.e. the feature selection. [24], [27], [30], [55], [57], [63], [66]–[81] used the mutual information and/or the false nearest neighbours' methods to estimate the suitable embedding parameters.

[21], [22], [27], [57], [63], [66], [68] attempted to forecast the solar sunspot number using neural networks and they used the embedding dimension of the sunspot time series as way to define the architecture of the input layer.

[82] generalize the mutual information approach to higher dimensions but do not connect it to the problem of the neural network input layer architecture optimization.

There are also papers [83] that try to use neural networks to determine the optimal embedding and time delay for the purposes of local reconstruction of states with a view to forecast.

There are also papers [84] that use Support Vector Machines (SVMs) to forecast in space and time and use time delays and embedding approaches to define the states vectors.

In fact Parlitz and Merkwirth [85] in the ESANN 2000 meeting proceedings paper mentioned that local reconstruction of states "...may also serve as a starting point for deriving local mathematical models in terms of polynomials, radial basis functions or neural networks.". Here we attempt to show empirical evidence that this is not just a starting point, but the optimal neural network input architecture.

Eurico Covas
December 31, 2017

E. Neural Network architecture




Fig. 1. Overall architecture of the neural network.

1) Monte Carlo results: Subsubsection text here.

II. CONCLUSION

We have shown empirical evidence for the existence of an optimal architecture of the input layer of neural networks used for forecasting spatial-temporal series.

ACKNOWLEDGMENT

We would like to thank Reza Tavakol for very useful discussions on forecasting. We also thank Dr. David Hathaway for providing the data that this article is based upon. CITEUC is funded by National Funds through FCT - Foundation for Science and Technology (project: UID/Multi/00611/2013) and FEDER - European Regional Development Fund through COMPETE 2020 - Operational Programme Competitiveness and Internationalization (project: POCI-01-0145-FEDER-006922).

REFERENCES

- [1] A. M. Fraser and H. L. Swinney, "Independent coordinates for strange attractors from mutual information," *Phys. Rev. A*, vol. 33, pp. 1134–1140, Feb. 1986.
- [2] H. Abarbanel, *Analysis of Observed Chaotic Data*, ser. Institute for Nonlinear Science. Springer New York, 1997.
- [3] H. Kantz and T. Schreiber, *Nonlinear time series analysis*, ser. Cambridge nonlinear science series. Cambridge, New York: Cambridge University Press, 1997, originally published: 1997.
- [4] H. Whitney, "Differentiable manifolds," *Ann. Math. (2)*, vol. 37, no. 3, pp. 645–680, July 1936, mR:1503303. Zbl:0015.32001. JFM:62.1454.01.
- [5] F. Takens, "Detecting strange attractors in turbulence," *Lecture Notes in Mathematics, Berlin Springer Verlag*, vol. 898, p. 366, 1981.
- [6] R. Mañé, "On the dimension of the compact invariant sets of certain non-linear maps," *Lecture Notes in Mathematics, Berlin Springer Verlag*, vol. 898, p. 230, 1981.
- [7] T. Sauer, J. A. Yorke, and M. Casdagli, "Embedology," *Journal of Statistical Physics*, vol. 65, pp. 579–616, Nov. 1991.
- [8] M. B. Kennel, R. Brown, and H. D. I. Abarbanel, "Determining embedding dimension for phase-space reconstruction using a geometrical construction," *Phys. Rev. A*, vol. 45, pp. 3403–3411, Mar. 1992.
- [9] J. M. Martinerie, A. M. Albano, A. I. Mees, and P. E. Rapp, "Mutual information, strange attractors, and the optimal estimation of dimension," *Phys. Rev. A*, vol. 45, pp. 7058–7064, May 1992.
- [10] H. D. I. Abarbanel, R. Brown, J. J. Sidorowich, and L. S. Tsimring, "The analysis of observed chaotic data in physical systems," *Reviews of Modern Physics*, vol. 65, pp. 1331–1392, Oct. 1993.
- [11] H. D. I. Abarbanel and J. P. Gollub, "Analysis of Observed Chaotic Data," *Physics Today*, vol. 49, p. 86, Nov. 1996.
- [12] E. Covas and F. Mena, "Forecasting of yield curves using local state space reconstruction," in *Dynamics, Games and Science I*, ser. Springer Proceedings in Mathematics, M. M. Peixoto, A. A. Pinto, and D. A. Rand, Eds. Springer Berlin Heidelberg, 2011, vol. 1, pp. 243–251.
- [13] E. Covas, "Spatial-temporal forecasting the sunspot diagram," *A&A*, vol. 605, p. A44, Sep. 2017.
- [14] E. Covas, N. Peixinho, and J. Fernandes, "Neural network forecast of the sunspot diagram," 2017, submitted.
- [15] H. C. Koons and D. J. Gorney, "A sunspot maximum prediction using a neural network," *EOS Transactions*, vol. 71, p. 677, May 1990.
- [16] A. S. Weigend, "Connectionist Architectures for Time Series Prediction of Dynamical Systems," Ph.D. dissertation, STANFORD UNIVERSITY, 1991.
- [17] A. S. Weigend, B. A. Huberman, and D. E. Rumelhart, "Predicting Sunspots and Exchange Rates with Connectionist Networks," in *Non-linear modeling and forecasting*, M. Casdagli and S. Eubank, Eds. Addison-Wesley, 1992, pp. 395–432.
- [18] K. MacPherson, "Neural network computation techniques applied to solar activity prediction," *Advances in Space Research*, vol. 13, pp. 447–450, Sep. 1993.
- [19] K. P. Macpherson, A. J. Conway, and J. C. Brown, "Prediction of solar and geomagnetic activity data using neural networks," *J. Geophys. Res.*, vol. 100, pp. 21 735–21 744, Nov. 1995.
- [20] A. Conway, "Echoed time series predictions, neural networks and genetic algorithms," *Vistas in Astronomy*, vol. 38, pp. 351–356, 1994.
- [21] H. D. Navone and H. A. Ceccatto, "Forecasting chaos from small data sets: a comparison of different nonlinear algorithms," *Journal of Physics A: Mathematical and General*, vol. 28, no. 12, p. 3381, 1995.
- [22] R. A. Calvo, H. A. Ceccatto, and R. D. Piacentini, "Neural network prediction of solar activity," *ApJ*, vol. 444, pp. 916–921, May 1995.
- [23] T. Koskela, M. Lehtokangas, J. Saarinen, and K. Kaski, "Time series prediction with multilayer perceptron, fir and elman neural networks," in *In Proceedings of the World Congress on Neural Networks*. Press, 1996, pp. 491–496.
- [24] F. Fessant, C. Pierret, and P. Lantos, "Comparison of Neural Network and McNish and Lincoln Methods for the Prediction of the Smoothed Sunspot Index," *Sol. Phys.*, vol. 168, pp. 423–433, Oct. 1996.
- [25] F. Fessant, S. Bengio, and D. Collobert, "On the prediction of solar activity using different neural network models," *Annales Geophysicae*, vol. 14, pp. 20–26, Jan. 1996.
- [26] Y. R. Park, T. J. Murray, and C. Chen, "Predicting sunspots using a layered perceptron neural network," *IEEE Transactions on Neural Networks*, vol. 7, pp. 501–505, Mar. 1996.
- [27] D. R. Kulkarni, A. S. Pandya, and J. C. Parikh, "Modeling and predicting sunspot activity-state space reconstruction + artificial neural network methods," *Geophys. Res. Lett.*, vol. 25, pp. 457–460, 1998.




Fig. 2. Neural network architecture showing the feature selection grid.




Fig. 3. Monte Carlo simulation of different architectures of the input layer for the neural network forecast. It shows the average of the sunspot area over the next cycle against the structural similarity (SSIM). The perfect architecture would be the one with SSIM = 1 and the 24-month running average over the cycle being the same (red lines).




Fig. 4. Monte Carlo simulation of different architectures of the input layer for the neural network forecast. It shows the structural similarity (SSIM) against how far (in an Euclidean space metric) the particular parameters of a particular run were from the supposedly optimal architecture parameters (red dot). The green line (trendline) seems to show that as the parameters of a randomly chosen architecture get close to the supposedly optimal architecture ones, the SSIM converges to what seems to be the best possible forecast value given the limited (and noisy) dataset.

- [28] A. J. Conway, K. P. Macpherson, G. Blacklaw, and J. C. Brown, "A neural network prediction of solar cycle 23," *J. Geophys. Res.*, vol. 103, pp. 29 733–29 742, Dec. 1998.
- [29] A. J. Conway, "Time series, neural networks and the future of the Sun," *New A Rev.*, vol. 42, pp. 343–394, Oct. 1998.
- [30] P. Verdes, M. Parodi, P. Granitto, H. Navone, R. Piacentini, and H. Ceccatto, "Predictions of the maximum amplitude for solar cycle 23 and its subsequent behavior using nonlinear methods," *Solar Physics*, vol. 191, no. 2, pp. 419–425, 2000.
- [31] P. F. Verdes, P. M. Granitto, and H. A. Ceccatto, "Secular Behavior of Solar Magnetic Activity: Nonstationary Time-Series Analysis of the Sunspot Record," *Sol. Phys.*, vol. 221, pp. 167–177, May 2004.
- [32] H. Lundstedt, "Solar activity predicted with artificial intelligence," *Washington DC American Geophysical Union Geophysical Monograph Series*, vol. 125, pp. 201–204, 2001.
- [33] M. Small and C. K. Tse, "Minimum description length neural networks for time series prediction," *Phys. Rev. E*, vol. 66, no. 6, p. 066701, Dec. 2002.

- [34] A. V. Mordvinov, N. G. Makarenko, M. G. Ogurtsov, and H. Jungner, "Reconstruction of Magnetic Activity of the Sun and Changes in Its Irradiance on a Millennium Timescale Using Neurocomputing," *Sol. Phys.*, vol. 224, pp. 247–253, Oct. 2004.
- [35] A. Gholipour, C. Lucas, B. N. Araabi, and M. Shafiee, "Solar activity forecast: Spectral analysis and neurofuzzy prediction," *Journal of Atmospheric and Solar-Terrestrial Physics*, vol. 67, pp. 595–603, Apr. 2005.
- [36] A.-F. A. Attia, R. H. Abdel-Hamid, and M. Quassim, "A genetic-based neuro-fuzzy approach for prediction of solar activity," in *Modeling and Systems Engineering for Astronomy*, ser. Proc. SPIE, S. C. Craig and M. J. Cullum, Eds., vol. 5497, Sep. 2004, pp. 542–552.
- [37] M. S. Quassim and A. F. Attia, "Forecasting the global temperature trend according to the predicted solar activity during the next decades,"

- Mem. Soc. Astron. Italiana*, vol. 76, p. 1030, 2005.
- [38] A.-F. Attia, R. Abdel-Hamid, and M. Quassim, "Prediction of Solar Activity Based on Neuro-Fuzzy Modeling," *Sol. Phys.*, vol. 227, pp. 177–191, Mar. 2005.
- [39] M. Mirmomeni, M. Shafiee, C. Lucas, and B. N. Araabi, "Introducing a new learning method for fuzzy descriptor systems with the aid of spectral analysis to forecast solar activity," *Journal of Atmospheric and Solar-Terrestrial Physics*, vol. 68, pp. 2061–2074, Dec. 2006.
- [40] M. S. Quassim, A.-F. Attia, and H. K. Elminir, "Forecasting the Peak Amplitude of the 24th and 25th Sunspot Cycles and Accompanying Geomagnetic Activity," *Sol. Phys.*, vol. 243, pp. 253–258, Jul. 2007.
- [41] A.-F. Attia, H. A. Ismail, and H. M. Basurah, "A Neuro-Fuzzy modeling for prediction of solar cycles 24 and 25," *Ap&SS*, vol. 344, pp. 5–11, Mar. 2013.
- [42] J.-X. Xie, C.-T. Cheng, K.-W. Chau, and Y.-Z. Pei, "A hybrid adaptive time-delay neural network model for multi-step-ahead prediction of sunspot activity," *International Journal of Environment and Pollution*, vol. 28, no. 3–4, pp. 364–381, 2006.
- [43] G. Maris and A. Oncica, "Solar Cycle 24 Forecasts," *Sun and Geosphere*, vol. 1, no. 1, pp. 8–11, Mar. 2006.
- [44] F. Emmert-Streib and M. Dehmer, "Nonlinear Time Series Prediction Based on a Power-Law Noise Model," *International Journal of Modern Physics C*, vol. 18, pp. 1839–1852, 2007.
- [45] A. S. Pandya, D. R. Kulkarni, and J. C. Parikh, "Study of time series prediction under noisy environment," in *Applications and Science of Artificial Neural Networks III*, ser. Proc. SPIE, S. K. Rogers, Ed., vol. 3077, Apr. 1997, pp. 116–126.
- [46] H. Lundstedt, M. Wik, and P. Wintoft, "Synoptic Solar Magnetic Fields: Explored and Predicted," *AGU Fall Meeting Abstracts*, Dec. 2006.
- [47] M. Wik, "Multiresolution Analysis and Prediction of Solar Magnetic Flux," in *37th COSPAR Scientific Assembly*, ser. COSPAR Meeting, vol. 37, 2008, p. 3467.
- [48] D. Gang and Z. Shi-Sheng, "Sunspot number prediction based on process neural network with time-varying threshold functions [j]," *Acta Physica Sinica*, vol. 2, p. 099, 2007.
- [49] J. Uwamahoro, L.-A. McKinnell, and P. J. Cilliers, "Forecasting solar cycle 24 using neural networks," *Journal of Atmospheric and Solar-Terrestrial Physics*, vol. 71, pp. 569–574, Apr. 2009.
- [50] C. Francile and M. L. Luoni, "Hacia la predicción del Número R de Wolf de manchas solares utilizando Redes Neuronales con retardos temporales," *Boletín de la Asociación Argentina de Astronomía La Plata Argentina*, vol. 53, pp. 241–244, 2010.
- [51] M. A. Parodi, H. A. Ceccatto, R. D. Piacentini, and P. J. García, "Actividad solar del ciclo 23. Predicción del máximo y fase decreciente utilizando redes neuronales," *Boletín de la Asociación Argentina de Astronomía La Plata Argentina*, vol. 43, pp. 23–24, 1999.
- [52] A. Ajabshirizadeh, N. Masoumzadeh Jouzdani, and S. Abbassi, "Neural network prediction of solar cycle 24," *Research in Astronomy and Astrophysics*, vol. 11, pp. 491–496, Apr. 2011.
- [53] A. Ajabshirizadeh and M. Juzdani Nafiseh, "Sunspot Number and Solar Radio Flux Prediction by Artificial Neural Network Method," in *38th COSPAR Scientific Assembly*, ser. COSPAR Meeting, vol. 38, 2010, p. 2.
- [54] S. Chattopadhyay, D. Jhajharia, and G. Chattopadhyay, "Trend estimation and univariate forecast of the sunspot numbers: Development and comparison of ARMA, ARIMA and Autoregressive Neural Network models," *Comptes Rendus Geoscience*, vol. 343, pp. 433–442, Jul. 2011.
- [55] C. Jiang and F. Song, "Sunspot forecasting by using chaotic time-series analysis and narx network," *JCP*, vol. 6, no. 7, pp. 1424–1429, 2011.
- [56] Y. Maslennikova and V. Bochkarev, "Solar activity prediction using artificial neural network and singular spectrum analysis," in *39th COSPAR Scientific Assembly*, ser. COSPAR Meeting, vol. 39, Jul. 2012, p. 1194.
- [57] R. Chandra and M. Zhang, "Cooperative coevolution of elman recurrent neural networks for chaotic time series prediction," *Neurocomput.*, vol. 86, pp. 116–123, Jun. 2012.
- [58] D.-C. Park and D.-M. Woo, "Prediction of sunspot series using bilinear recurrent neural network," in *Information Management and Engineering, 2009. ICIME'09. International Conference on*. IEEE, 2009, pp. 94–98.
- [59] T.-H. Kim, D.-C. Park, D.-M. Woo, W. Huh, C.-H. Yoon, H.-U. Kim, and Y. Lee, "Sunspot series prediction using a multiscale recurrent neural network," in *Signal Processing and Information Technology (ISSPIT), 2010 IEEE International Symposium on*. IEEE, 2010, pp. 399–403.
- [60] J. D. Moghaddam, A. Mosallanezhad, and M. Teshnehlab, "Sunspot prediction by a time delay line recurrent fuzzy neural network using emotional learning," in *Fuzzy Systems (IFSC), 2013 13th Iranian Conference on*. IEEE, 2013, pp. 1–5.
- [61] G. Chattopadhyay and S. Chattopadhyay, "Monthly sunspot number time series analysis and its modeling through autoregressive artificial neural network," *European Physical Journal Plus*, vol. 127, p. 43, Apr. 2012.
- [62] Z.-G. Liu and J. Du, "Sunspot time sequences prediction based on process neural network and quantum particle swarm," in *Multimedia Information Networking and Security (MINES), 2012 Fourth International Conference on*. IEEE, 2012, pp. 233–236.
- [63] A. Gkana and L. Zachilas, "Sunspot numbers: Data analysis, predictions and economic impacts," *Journal of Engineering Science and Technology Review*, vol. 8, no. 1, pp. 79–85, 2015, cited By 1.
- [64] M. Parsapoor, U. Bilstrup, and B. Svensson, "Prediction of solar cycle 24," in *2015 International Joint Conference on Neural Networks, IJCNN 2015, Killarney, Ireland, July 12-17, 2015*. IEEE, 2015, pp. 1–8.
- [65] M. Parsapoor, J. Brooke, and B. Svensson, "A new computational intelligence model for long-term prediction of solar and geomagnetic activity," in *Proceedings of the Twenty-Ninth AAAI Conference on Artificial Intelligence, January 25-30, 2015, Austin, Texas, USA*, B. Bonet and S. Koenig, Eds. AAAI Press, 2015, pp. 4192–4193.
- [66] R. Archana, A. Unnikrishnan, and R. Gopikakumari, "Computation of state space evolution of chaotic systems from time series of output, based on neural network models," *International Journal of Engineering Research and Development*, vol. 2, pp. 49–56, Jul. 2012.
- [67] M. Annunziato, S. Pizzuti, and L. S. Tsimring, "Analysis and prediction of spatio-temporal flame dynamics," in *Proceedings of the IEEE Workshop on Nonlinear Dynamics of Electronic Systems: Catania, Italy, 18-20 May 2000*, G. Setti, R. Rovatti, and G. Mazzini, Eds. World Scientific, 2000, pp. 117–121.
- [68] L. Zachilas and A. Gkana, "On the verge of a grand solar minimum: A second mauerer minimum?" *Solar Physics*, vol. 290, no. 5, pp. 1457–1477, 2015.
- [69] Y. Sun, V. Babovic, and E. S. Chan, "Multi-step-ahead model error prediction using time-delay neural networks combined with chaos theory," *Journal of Hydrology*, vol. 395, no. 1–2, pp. 109–116, 2010.
- [70] S.-C. Huang, P.-J. Chuang, C.-F. Wu, and H.-J. Lai, "Chaos-based support vector regressions for exchange rate forecasting," *Expert Systems with Applications*, vol. 37, no. 12, pp. 8590–8598, 2010.
- [71] R. Battiti, "Using mutual information for selecting features in supervised neural net learning," *IEEE Transactions on Neural Networks*, vol. 5, no. 4, pp. 537–550, Jul. 1994.
- [72] R. J. Frank, N. Davey, and S. P. Hunt, "Time series prediction and neural networks," *Journal of Intelligent and Robotic Systems*, vol. 31, no. 1, pp. 91–103, 2001.
- [73] S. BuHamra, N. Smaoui, and M. Gabr, "The box-jenkins analysis and neural networks: prediction and time series modelling," *Applied Mathematical Modelling*, vol. 27, no. 10, pp. 805–815, 2003.
- [74] R. Chandra and M. Zhang, "Cooperative coevolution of elman recurrent neural networks for chaotic time series prediction," *Neurocomputing*, vol. 86, pp. 116–123, 2012.
- [75] Y. S. Maslennikova and V. V. Bochkarev, "Training algorithm for neuro-fuzzy network based on singular spectrum analysis," *CoRR*, vol. abs/1410.1151, 2014.
- [76] T. Sauter, B. Weitzkamp, and C. Schneider, "Spatio-temporal prediction of snow cover in the black forest mountain range using remote sensing and a recurrent neural network," *International Journal of Climatology*, vol. 30, no. 15, pp. 2330–2341, 2010.
- [77] A. H. Ghaderi, B. Bharani, and H. Jalalkamali, "Embedding dimension as input dimension of artificial neural network: A study on stock prices time series," *International Journal of Modern Physics and Applications*, vol. 1, no. 3, pp. 64–72, 2015.
- [78] D. R. Kulkarni, J. C. Parikh, and A. S. Pandya, "Dynamic Predictions from Time Series Data - An Artificial Neural Network Approach," *International Journal of Modern Physics C*, vol. 8, pp. 1345–1360, 1997.
- [79] Z. Sheng, L. Hong-Xing, G. Dun-Tang, and D. Si-Dan, "Determining the input dimension of a neural network for nonlinear time series prediction," *Chinese Physics*, vol. 12, no. 6, p. 594, 2003.
- [80] P. S. Lucio, F. C. Conde, I. F. A. Cavalcanti, A. I. Serrano, A. M. Ramos, and A. O. Cardoso, "Spatiotemporal monthly rainfall reconstruction via artificial neural network - case study: south of Brazil," *Advances in Geosciences*, vol. 10, pp. 67–76, Apr. 2007.
- [81] P. Maass, T. Kohler, J. Kalden, R. Costa, U. Parlitz, C. Merkwirth, and J. Wichard, *Mathematical Methods for Forecasting Bank Transaction Data*, ser. DFG-Schwerpunktprogramm 1114, Mathematical methods for time series analysis and digital image processing. Zentrum für Technomathematik, 2003.
- [82] G. Simon and M. Verleysen, "High-dimensional delay selection for regression models with mutual information and distance-to-diagonal criteria," *Neurocomput.*, vol. 70, no. 7–9, pp. 1265–1275, Mar. 2007.

- [83] M. Ragulskis and K. Lukoseviciute, "Non-uniform attractor embedding for time series forecasting by fuzzy inference systems," vol. 72, pp. 2618–2626, 12 2009.
- [84] Y. Xia, H. Leung, and H. Chan, "A prediction fusion method for reconstructing spatial temporal dynamics using support vector machines," *IEEE Trans. on Circuits and Systems*, vol. 53-II, pp. 62–66, 2006.
- [85] U. Parlitz and C. Merkwirth, "Nonlinear prediction of spatio-temporal time series," in *ESANN*, 2000.

Eurico Covas Biography text here.

John Doe Biography text here.

Jane Doe Biography text here.