




EUTROPHICATION OF SURFACE WATERS

A4b Seminar: Water resources in a quickly changing world Emmanuel Adeleke. D.











A QUICK INTRODUCTION TO EUTROPHICATION

EUTROPHICATION


When a body of water becomes overly enriched with minerals and nutrients which induces excessive growth of plants & algae






- Natural & slow eutrophication process
- Cultural eutrophication process → Human activities
 - Different ecological & societal effects

- Natural & slow eutrophication process
- Cultural eutrophication process → Human activities
 - Different ecological & societal effects




- Natural & slow eutrophication process
- Cultural eutrophication process → Human activities
 - Different ecological & societal effects

➤ Algal bloom




- Natural & slow eutrophication process
- Cultural eutrophication process → Human activities
 - Different ecological & societal effects

- ➤ Algal bloom
- ➤ Sunlight blockage




- Natural & slow eutrophication process
- Cultural eutrophication process → Human activities
 - Different ecological & societal effects

- ➤ Algal bloom
- ➤ Sunlight blockage
- > Death of aquatic flora




- Natural & slow eutrophication process
- Cultural eutrophication process → Human activities
 - Different ecological & societal effects

- ➤ Algal bloom
- ➤ Sunlight blockage
- Death of aquatic flora
- > Oxygen depletion
- > Formation of dead zones(hypoxia)
- Toxic emissions (CO₂, H₂S, CH₄)




Sources that contribute to nutrient loading

- Point sources:
- > Runoff and leachate from waste disposal system
- Untreated sewage
- Non-point sources:
- > Ploughing in agriculture and development
- > Industrial effluents
- ➤ Urban runoff
- Atmospheric deposition from fossil fuel combustion



Major sources of nutrients in lakes












Conceptual diagram of the transfer, retention and removal zones of nitrogen and phosphorus along the land-sea continuum.



Source: Moal et al. 2018. Eutrophication: A new wine in an old bottle. Science of the Total Environment 651 (2019) I-II

• Is it therefore correct to say that phosphorus is a greater contributor to eutrophication than nitrogen?

- Is it therefore correct to say that phosphorus is a greater contributor to eutrophication than nitrogen?
 - ➤ Does not readily leave the surface?

- Is it therefore correct to say that phosphorus is a greater contributor to eutrophication than nitrogen?
 - ➤ Does not readily leave the surface?
 - > Critical nutrient for lakes. Plant size ~ Quantity of P available


- Is it therefore correct to say that phosphorus is a greater contributor to eutrophication than nitrogen?
 - ➤ Does not readily leave the surface?
 - ➤ Critical nutrient for lakes. Plant size ~ Quantity of P available
 - > P input largely caused by human activity


- Is it therefore correct to say that phosphorus is a greater contributor to eutrophication than nitrogen?
 - ➤ Does not readily leave the surface?
 - ➤ Critical nutrient for lakes. Plant size ~ Quantity of P available
 - > P input largely caused by human activity
 - > Sewage minus P loses algal capability

- Is it therefore correct to say that phosphorus is a greater contributor to eutrophication than nitrogen?
 - > Does not readily leave the surface?
 - ➤ Critical nutrient for lakes. Plant size ~ Quantity of P available
 - > P input largely caused by human activity
 - > Sewage minus P loses algal capability

	Phosphorus	Nitrogen
Cycle	Solid or liquid form	Gas phase
Movement	Less mobile	More mobile
Storage time	Centuries	Decades
Capacity	Retention	Elimination
	Soils and sediments	Atmosphere, waterbodies
Wastewater contains	25-40 mg per liter	6 - 10 mg per liter

EXPERIMENTAL LAKES AREA – CANADA





Schindler et al. 1974

EUTROPHICATION as a global & stubborn problem!

Eutrophication is not a new phenomenon

• Each day, Detroit, Cleveland and 120 other municipalities fill Erie with 1.5 billion gallons of inadequately treated wastes, including nitrates and phosphates. These chemicals act as fertilizer for growths of algae that suck oxygen from the lower depths and rise to the surface as odoriferous green scum. Commercial and game fish ... have nearly vanished ... Weeds proliferate, turning water frontage into swamp. In short, Lake Erie is in danger of dying by suffocation.

The August 1969 issue of Time Magazine

Case Study – The North American Great Lakes

- Lake Superior, Michigan, Ontario, Erie and Huron
- Industrial growth in the 1940s & 1950s resulted in oil pollution
- Eutrophication began in the 1960s as a result of excessive P inputs
- Massive fish die-offs in Erie, Michigan and Ontario.
- 1972 Great Lakes Water Quality Agreement
- Great Lakes Nutrient Initiative with a focus on Erie








Eutrophication at present





Table 2. Major environmental problems occurring in lakes throughout the world

- Low water-level due to the over-use of water from lakes resulting in a pronounced deterioration of water quality and adverse changes in the ecosystems.
- Rapid siltation in lakes and reservoirs caused by accelerated soil erosion resulting from the overuse or misuse of arable and grazing lands and forests within their drainage areas.
- Acidification of lakes due to acid precipitation, resulting in the extinction of fish and the degradation of ecosystems.
- Contamination of the water, sediment and organisms with toxic chemicals originated from agriculture (pesticides) and industrial wastes.
- Eutrophication from inputs of nitrogen and/or phosphorus compounds discharged from industries, agricultural land, homes, urban and road surfaces etc., and resulting in heavy blooms of phytoplankton, deterioration of water quality, and a decrease of biodiversity.
- In extreme cases, the complete collapse of aquatic ecosystems.




Eutrophication at present – drivers



- Direct drivers
 - Energy consumption
 - Fertilizer consumption
 - Land-use conversion
- Indirect drivers
 - Population growth
 - Economic growth
 - Agricultural intensification

>,,Diffuse nitrogen and phosphorus pollutions are now the main drivers of the new wave of eutrophication"

World Hypoxic and Eutrophic Areas



Of the 415 areas around the world, 169 are hypoxic, 13 classified as "systems in recovery" and 233 are eutrophic areas.

- The Baltic Sea
- Laurentian Great Lakes
- Gulf of Mexico
- The Venice Lagoon
- Most lakes & coastal areas in China, Lake Victoria, the Brittany coast, Mediterranean lagoons and





 Goal 14: Conserve and sustainably use the oceans, seas and marine resources for sustainable development





- Goal 14: Conserve and sustainably use the oceans, seas and marine resources for sustainable development
- Target I: Reduce marine & coastal pollution by 2025





- Goal 14: Conserve and sustainably use the oceans, seas and marine resources for sustainable development
- Target I: Reduce marine & coastal pollution by 2025
- Progress as of 2018: Continued deterioration of coastal waters due to pollution and eutrophication.
 - By 2050, coastal eutrophication will increase by 20%

 Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication

- Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication
- Other regions?

- Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication
- Other regions? Lack of reliable data!


- Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication
- Other regions? Lack of reliable data!
- data do not exist or are not publicly available

- Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication
- Other regions? Lack of reliable data!
- > data do not exist or are not publicly available
- inconsistency in terms of parameters measured, indicators used and scale

- Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication
- Other regions? Lack of reliable data!
- > data do not exist or are not publicly available
- inconsistency in terms of parameters measured, indicators used and scale
- most underrepresented region Asia

- Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication
- Other regions? Lack of reliable data!
- > data do not exist or are not publicly available
- inconsistency in terms of parameters measured, indicators used and scale
- most underrepresented region Asia
- as well as Africa, Latin America and the Carribean.

- Recent coastal survey of the US & EU found 78% and 65% of their respective coastal areas to exhibit eutrophication
- Other regions? Lack of reliable data!
- > data do not exist or are not publicly available
- inconsistency in terms of parameters measured, indicators used and scale
- most underrepresented region Asia
- > as well as Africa, Latin America and the Carribean.
- \triangleright absence of detailed nutrient budgets \rightarrow lack of research efforts





ADDRESSING THE PROBLEMS OF EUTROPHICATION

- Limit aquatic fertility
- > Diversion of nutrient-bearing streams
- > Sewage treatment
- Stimulate diseases & parasites among unwanted plants
- Use toxic chemicals
- Biological or tranditional methods

- Limit aquatic fertility
- Diversion of nutrient-bearing streams
- > Sewage treatment
- Stimulate diseases & parasites among unwanted plants
- Use toxic chemicals
- Biological or tranditional methods

NUTRIENTS TO REMOVE

- Phosphorus P / /
- Nitrogen N 🗘
- Carbon C
- +16 other micronutrients e.g. O, H2, Si, K, Mg, S, Cl, Na, Fe, Zn, Cu.

- Limit aquatic fertility
- Diversion of nutrient-bearing streams
- > Sewage treatment
- Stimulate diseases & parasites among unwanted plants
- Use toxic chemicals
- Biological or tranditional methods

NUTRIENTS TO REMOVE

- Nitrogen N 1
- Carbon C
- +16 other micronutrients e.g. O,
 H2, Si, K, Mg, S, Cl, Na, Fe, Zn, Cu.

- Eliminate phosphates in detergents
- Technological advancement
- Control dietary N & P

- Limit aquatic fertility
- Diversion of nutrient-bearing streams
- > Sewage treatment
- Stimulate diseases & parasites among unwanted plants
- Use toxic chemicals
- Biological or traditional methods

ROLE OF GOVERNMENTS

- Grant Funds for Municipal waste treatments
- Research and development
- Enforcement
- Legislation

NUTRIENTS TO REMOVE

- Nitrogen N 1
- Carbon C
- +16 other micronutrients e.g. O,
 H2, Si, K, Mg, S, Cl, Na, Fe, Zn, Cu.

- Eliminate phosphates in detergents
- Technological advancement
- Control dietary N & P

Case Study: Lake Victoria

- dense mats of water hyacinth float on huge surface areas
- most parts show oxygen depletion and reduced transparency
- many haplochromis or "furu" fish species are extinct
- eutrophication controlled by biological methods
- a beetle feeding on water hyacinths
- weevil N. eichhornaiae





45





improved water quality monitoring & assessment of estuarine and coastal waters.



- improved water quality monitoring & assessment of estuarine and coastal waters.
- time series monitoring data



- improved water quality monitoring & assessment of estuarine and coastal waters.
- time series monitoring data
- widespread use & adoption of methods



- improved water quality monitoring & assessment of estuarine and coastal waters.
- time series monitoring data
- widespread use & adoption of methods
- develop credible and consistent proxies for water quality e.g. turbidity, sulfide smell.



- improved water quality monitoring & assessment of estuarine and coastal waters.
- time series monitoring data
- widespread use & adoption of methods
- develop credible and consistent proxies for water quality e.g. turbidity, sulfide smell.
- more decision support tools should be developed
 - watershed models
 - nutrient balance assessments
 - regional and international online information portals
 - nutrient loss estimation tools



- improved water quality monitoring & assessment of estuarine and coastal waters.
- time series monitoring data
- widespread use & adoption of methods
- develop credible and consistent proxies for water quality e.g. turbidity, sulfide smell.
- more decision support tools should be developed
 - watershed models
 - nutrient balance assessments
 - regional and international online information portals
 - nutrient loss estimation tools
- interdisciplinary research



- improved water quality monitoring & assessment of estuarine and coastal waters.
- time series monitoring data
- widespread use & adoption of methods
- develop credible and consistent proxies for water quality e.g. turbidity, sulfide smell
- more decision support tools should be developed
 - watershed models
 - nutrient balance assessments
 - regional and international online information portals
 - nutrient loss estimation tools
- interdisciplinary research
- improving knowledge on eutrophication is the first step in developing robust policy measures to reverse its impacts.

Conclusions

- Eutrophication is a unique problem, not limited spatially or temporally.
- Point sources of phosphates in industrial effluents and sewage were the drivers of eutrophication in the past.
- Non-point source of nitrogen and phosphorus pollutions are the new drivers of eutrophication today
- There is need to improve knowledge & provide data on eutrophic systems

Discussion

- Can you think of other ways by which nutrient loading of surface waters can be controlled?
- How do we provide more globally representative datasets on eutrophic & hypoxic systems?
- Global or regional (national) approach towards nutrient control?
- Best approach to control fertilizer use and stop nutrient runoff from agricultural sources? legislation? enforcement?