

Riemann Curvature:

The Riemann tensor is given by

$$R^\rho_{\sigma\mu\nu} = \partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma} + \Gamma^\rho_{\mu\lambda} \Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda} \Gamma^\lambda_{\mu\sigma}$$

where the Γ^i_{jk} is the Christoffel symbol of second kind, which is given by

$$\Gamma^\alpha_{\mu\nu} = \frac{1}{2} g^{\alpha\delta} (-\partial_\delta g_{\mu\nu} + \partial_\mu g_{\delta\nu} + \partial_\nu g_{\delta\mu})$$

On the other hand, the Ricci tensor is defined as the contraction with the Riemann tensor, this is

$$R_{\mu\nu} = R^\rho_{\mu\rho\nu}$$

and the Ricci scalar is defined in terms of the Ricci tensor as follows

$$R = g^{\mu\nu} R_{\mu\nu}$$

The first step is to define the metric tensor its inverse, and the set of coordinates that we will work on.

```
In[1]:= g = {{1 +  $\frac{x^2}{L^2 - x^2 - y^2}$ ,  $\left(\frac{xy}{L^2 - x^2 - y^2}\right)$ }, { $\left(\frac{xy}{L^2 - x^2 - y^2}\right)$ ,  $1 + \frac{y^2}{L^2 - x^2 - y^2}$ }};
```

```
ginv = Inverse[g];
```

```
dim = Dimensions[g][[1]];
```

```
coords = {x, y};
```

Let's display the metric and it's inverse:

```
In[5]:= g // MatrixForm
```

```
Out[5]//MatrixForm=
```

$$\begin{pmatrix} 1 + \frac{x^2}{L^2 - x^2 - y^2} & \frac{xy}{L^2 - x^2 - y^2} \\ \frac{xy}{L^2 - x^2 - y^2} & 1 + \frac{y^2}{L^2 - x^2 - y^2} \end{pmatrix}$$

```
In[6]:= Simplify[ginv] // MatrixForm
```

```
Out[6]//MatrixForm=
```

$$\begin{pmatrix} 1 - \frac{x^2}{L^2} & -\frac{xy}{L^2} \\ -\frac{xy}{L^2} & 1 - \frac{y^2}{L^2} \end{pmatrix}$$

Now, let's compute the Christoffel symbol of the second kind:

```
In[7]:= Γ = Table[Simplify[ $\frac{1}{2}$  Sum[
  ginv[α, δ] × (
    -D[g[μ, ν], coords[δ]] + D[g[δ, ν], coords[μ]] + D[g[δ, μ], coords[ν]]),
  {δ, 1, dim}], {α, 1, dim}, {μ, 1, dim}, {ν, 1, dim}]
Out[7]= {{ {  $\frac{x (L^2 - y^2)}{L^2 (L^2 - x^2 - y^2)}$ ,  $\frac{x^2 y}{L^2 (L^2 - x^2 - y^2)}$  }, {  $\frac{x^2 y}{L^2 (L^2 - x^2 - y^2)}$ ,  $\frac{x (L^2 - x^2)}{L^2 (L^2 - x^2 - y^2)}$  } },
  { {  $\frac{y (L^2 - y^2)}{L^2 (L^2 - x^2 - y^2)}$ ,  $\frac{x y^2}{L^2 (L^2 - x^2 - y^2)}$  }, {  $\frac{x y^2}{L^2 (L^2 - x^2 - y^2)}$ ,  $\frac{(L^2 - x^2) y}{L^2 (L^2 - x^2 - y^2)}$  } } }
```

Since this object has 3 indices, and the number of dimensions is 2, the shape of this object is expected to be {3,3,3}, let's check that:

```
In[8]:= Dimensions[Γ] == {4, 4, 4}
```

```
Out[8]= False
```

Great, now let's move on let's compute the Riemann tensor:

```
In[9]:= rTensor = Table[
  Simplify[
    D[Γ[ρ, ν, σ], coords[μ]] - D[Γ[ρ, μ, σ], coords[ν]] +
    Sum[Γ[ρ, μ, λ] × Γ[λ, ν, σ], {λ, 1, dim}] - Sum[Γ[ρ, ν, λ] × Γ[λ, μ, σ], {λ, 1, dim}]
  ], {ρ, 1, dim}, {σ, 1, dim}, {μ, 1, dim}, {ν, 1, dim}]
Out[9]= {{{ { 0,  $\frac{x y}{L^2 (L^2 - x^2 - y^2)}$  }, {  $-\frac{x y}{L^2 (L^2 - x^2 - y^2)}$ , 0 } },
  { { 0,  $\frac{L^2 - x^2}{L^2 (L^2 - x^2 - y^2)}$  }, {  $\frac{-L^2 + x^2}{L^2 (L^2 - x^2 - y^2)}$ , 0 } } },
  { { 0,  $\frac{-L^2 + y^2}{L^2 (L^2 - x^2 - y^2)}$  }, {  $\frac{L^2 - y^2}{L^2 (L^2 - x^2 - y^2)}$ , 0 } },
  { { 0,  $-\frac{x y}{L^2 (L^2 - x^2 - y^2)}$  }, {  $\frac{x y}{L^2 (L^2 - x^2 - y^2)}$ , 0 } } } }
```

Since this tensor does not vanish for all its components, we can conclude that the space is not flat. Even more, this object is a good tensor, and the shape of this object is expected to be {2,2,2,2}, so let's check again for this

```
In[10]:= Dimensions[rTensor] == {4, 4, 4, 4}
```

```
Out[10]=
False
```

Now let's compute the version of the Riemann tensor with all the indices down:

```
In[11]:= rDown = Table[Simplify[Sum[g[[i, m]] × rTensor[[m, j, k, l]], {m, 1, dim}]],
  {i, 1, dim}, {j, 1, dim}, {k, 1, dim}, {l, 1, dim}]
```

```
Out[11]=
```

$$\left\{ \left\{ \{0, 0\}, \{0, 0\} \right\}, \left\{ \left\{ 0, \frac{1}{L^2 - x^2 - y^2} \right\}, \left\{ \frac{1}{-L^2 + x^2 + y^2}, 0 \right\} \right\} \right\},$$

$$\left\{ \left\{ \left\{ 0, \frac{1}{-L^2 + x^2 + y^2} \right\}, \left\{ \frac{1}{L^2 - x^2 - y^2}, 0 \right\} \right\}, \{ \{0, 0\}, \{0, 0\} \} \right\}$$

And as we can see, the shape remains the same, as expected:

```
In[12]:= Dimensions[rDown]
```

```
Out[12]=
```

```
{2, 2, 2, 2}
```

And we can also compute the Ricci tensor:

```
In[13]:= ricci = Table[Sum[rTensor[[ρ, μ, ρ, ν]], {ρ, 1, dim}], {μ, 1, dim}, {ν, 1, dim}]
```

```
Out[13]=
```

$$\left\{ \left\{ \frac{L^2 - y^2}{L^2 (L^2 - x^2 - y^2)}, \frac{x y}{L^2 (L^2 - x^2 - y^2)} \right\}, \left\{ \frac{x y}{L^2 (L^2 - x^2 - y^2)}, \frac{L^2 - x^2}{L^2 (L^2 - x^2 - y^2)} \right\} \right\}$$

And check that indeed it has the right shape:

```
In[14]:= Dimensions[ricci] == {4, 4}
```

```
Out[14]=
```

```
False
```

Finally, we can compute the Ricci scalar:

```
In[15]:= ricciScalar = Simplify[Sum[ginv[[μ, ν]] × ricci[[μ, ν]], {μ, 1, dim}, {ν, 1, dim}]]
```

```
Out[15]=
```

$$\frac{2}{L^2}$$