

Illustrative arithmetic: $B = 2, H = 3$; not pilot data

(a) Recipient correctness

		$j = 1$	$j = 2$	$j = 3$	Sequence mean	
					$\frac{0}{\quad\quad\quad} \frac{1}{\quad\quad\quad}$	
$b = 1$	E	1	1	?	$[\frac{2}{3}, 1]$	
	O	1	0	0	$\frac{1}{3}$	
$b = 2$	E	1	1	0	$\frac{2}{3}$	
	O	1	0	?	$[\frac{1}{3}, \frac{2}{3}]$	

$q_E^{\text{obs}} \in [\frac{2}{3}, \frac{5}{6}]$

$q_O^{\text{obs}} \in [\frac{1}{3}, \frac{1}{2}]$

$\delta^{\text{obs}} \in [\frac{1}{6}, \frac{1}{2}]$

Lower bound: $E \text{ ?} \mapsto 0, O \text{ ?} \mapsto 1$

(b) Input + output tokens

		$j = 0$	$j = 1$	$j = 2$	$j = 3$	Total
		donor				
$b = 1$	E	1	1	1	1	4
	O	1	1	1	1	4
$b = 2$	E	2	2	2	2	8
	O	4	4	4	4	16

$S_E = 4 + 8 = 12, \quad S_O = 4 + 16 = 20$

$\widehat{R}_T = 1 - \frac{12}{20} = 40\%$

Block reductions: 0%, 50%

Unweighted mean: 25%