

Cassini's Model

Introduction


In the 17th Century, everyone assumed that the atmosphere terminated abruptly in a smooth surface where it met “the æther,” just like the smooth surface where still water meets the “subtler” medium of air. (The æther was supposed to be subtler still than air; so, why not? The gas laws were not known.) This picture of a homogeneous layer of air surrounding the Earth goes back at least to a work on optics attributed to [Ptolemy](#), which says that “at the surface between air and ether, there is a refraction of the visual ray according to the difference in density between these two media.”

This view had been explicitly adopted by [Kepler](#) in his failed attempt to understand refraction, and it was adopted by Giovanni Domenico Cassini as well.

Unlike Kepler, Cassini knew the [sine law](#) of refraction. And it was a simple exercise in elementary trigonometry for him to apply it to the uniform atmosphere, thus making the first quantitative model for astronomical refraction.

The Model

Cassini's model is so simple that anyone who had trigonometry in high school can follow it. Here's how it works:



Because of the refraction that occurs at **P**, the ray direction outside the atmosphere is different: the ray meets the local vertical there at an angle z_2 , which is bigger than z_1 . If the refractive index of air is n , the law of refraction tells us that

So much for the law of refraction. The nice thing is that we can calculate the sine of z_1 from the sine of the observed zenith distance, z_0 . If you recall the Law of Sines for plane triangles, and apply it to the triangle OPC, you'll see that

But the sine of the angle COP is the same as the sine of its supplement, z_0 . So we have:

or

Now that we have $\sin z_1$, the law of refraction gives us $\sin z_2$:

and consequently

$$z_2 = \arcsin \{ [nR/(R+h)] \sin z_0 \} .$$

But (as you can see from the [diagram](#) above) the refraction, r , is just the difference of these two angles:

$$r = z_2 - z_1 .$$

So, because we already had an expression for $\sin z_1$ a few lines [above](#):

$$r = \arcsin \{ [nR/(R+h)] \sin z_0 \} - \arcsin \{ [R/(R+h)] \sin z_0 \} .$$

That's it! We're done. This is Cassini's refraction model.

Copyright © 2003 – 2006, 2012 Andrew T. Young

Back to the ...

[astronomical refraction page](#)

or the [GF home page](#)

or the [overview page](#)