MAT211: Calculus-II

Lecture Notes

Instructor: Emon Hossain

February 23, 2026

Contents

1 Lecture-01 **2**

2 Lecture-02 **4**

3 Lecture-03 **7**

1 Lecture-01

Question: what did we mean by the derivative $f'(a)$?
Answer: Let U be an open subset of $\mathbb{R}$ and $f : U \to \mathbb{R}$ a function. Then f is differentiable at a, with derivative m, if and only if

$$\lim_{h\to 0} \frac{1}{h}\left(\underbrace{f(a+h)-f(a)}_{\Delta f} - \underbrace{mh}_{(*)}\right) = 0 \tag{1}$$

$(*)$: linear function of $\Delta x = h$.
Question: Why impose the domain U that needs to be an open set?
Answer: Write the answer.
The function mh that multiplies h by the derivative m is thus a linear function of the change in x. This definition emphasizes the idea that a function f is differentiable at a point a if the increment Δf to the function is well approximated by a linear function of the increment h to the variable, this linear function is $f'(a)h$. Let's see how we can confirm that:
Suppose $f(x)$ is our desired function and we want to approximate the function in the neighborhood at $x = a$. For this purpose, we will draw several straight lines through the point $(a, f(a))$ and see which line approximates the function best. Here, the best means, the linear line should approximate the function well at $x = a + h$. Suppose the slope of that linear function is m.



Then,

$$\frac{L(a+h)-L(a)}{a+h-a} = m$$
$$\frac{L(a+h)-f(a)}{h} = m$$
$$L(a+h) = f(a) + mh$$

Now, we want to minimize the error, $E_m(h) = f(a+h) - L(a+h)$. Hence, we need to find the optimal value for m, for which:

$$\frac{E_m(h)}{h} \to 0$$

$$\boxed{\text{optimal } m \iff \lim_{h\to 0} \frac{E_m(h)}{h} = 0}$$

A simple computation show that,

$$\lim_{h\to 0} \frac{E_m(h)}{h} = \lim_{h\to 0} \frac{f(a+h) - f(a) - mh}{h} = 0$$

which is the desired condition 1, we showed in the definition.

2 Lecture-02

Definition 2.1 (Euclidean Vector Space)**.** A Euclidean vector space is a finite-dimensional inner product space over $\mathbb{R}$.

Definition 2.2 (Inner product space)**.** An inner product space is a vector space V over the field F together with an inner product, that is, a map

$$\langle \cdot, \cdot \rangle : V \times V \to F$$

that satisfies the following three properties for all vectors $x, y, z \in V$ and all scalars $a, b \in F$.

1. (Symmetry) $\langle x, y \rangle = \langle y, x \rangle$.
2. (Linearity in the first argument) $\langle ax + by, z \rangle = a\langle x, z \rangle + b\langle y, z \rangle$.
3. (Positive-definiteness) If $x \neq 0$ then, $\langle x, x \rangle > 0$.

Example 2.3. Verify that the dot product function $\langle \cdot, \cdot \rangle : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ is an inner product.

To verify this we must prove the three conditions shown above. Let $u = (a, b)$ and $v = (c, d)$. Then,

1. $\langle u, v \rangle = ac + bd = ca + db = \langle v, u \rangle$.
2. Let $a, b \in \mathbb{R}$ and $u, v, w \in \mathbb{R}^2$ with $u = (u_1, u_2), v = (v_1, v_2), w = (w_1, w_2)$. Then

$$\begin{aligned}
\langle au + bv, w \rangle &= (au_1 + bv_1)w_1 + (au_2 + bv_2)w_2 \\
&= au_1w_1 + bv_1w_1 + au_2w_2 + bv_2w_2 \\
&= a(u_1w_1 + u_2w_2) + b(v_1w_1 + v_2w_2) \\
&= a\langle u, w \rangle + b\langle v, w \rangle.
\end{aligned}$$

3. For any $u = (u_1, u_2)$,

$$\langle u, u \rangle = {u_1}^2 + {u_2}^2.$$

   Since squares of non-zero numbers are positive, we have ${u_1}^2 > 0, {u_2}^2 > 0$ and therefore, $\langle u, u \rangle > 0$.

Definition 2.4 (Informal)**.** Define **norm** of a vector $x \in V$ as

$$||x|| = \sqrt{\langle x, x \rangle}.$$

Definition 2.5 (Metric Space)**.** A metric space is an ordered pair (M, d) where M is a set and d is a **metric** (distance) on M, i.e., a function

$$d : M \times M \to \mathbb{R}$$

satisfying the following axioms for all points $x, y, z \in M$:

1. $d(x, y) \geq 0$;
2. $d(x, y) = 0 \iff x = y$;

3. $d(x, y) = d(y, x)$.

4. $d(x, z) \leq d(x, y) + d(y, z)$.

Example 2.6 (Tax-cab Metric)**.** In $\mathbb{R}^2$, the taxi-cab distance between $p = (p_1, p_2)$ and $q = (q_1, q_2)$ is $|q_1 - p_1| + |q_2 - p_2|$. It is easy to verify that this is a metric.

Definition 2.7 (Open ball)**.** For any $x \in \mathbb{R}^n$ and any $r > 0$, the **open ball** of radius r around x is the subset

$$B_r(x) = \{y \in \mathbb{R}^n \mid |x - y| < r\}.$$



Definition 2.8 (Open sets of $\mathbb{R}^n$)**.** A subset $U \subseteq \mathbb{R}^n$ is open if for every point $x \in U$, there exists $r > 0$ such that the open ball $B_r(x)$ is contained in U.

Example 2.9. We prove that the Cartesian product $(a_1, b_1) \times (a_2, b_2)$ of two open intervals in $\mathbb{R}$ is an open set in $\mathbb{R}^2$ with the standard Euclidean topology.

Proof. Let d denote the Euclidean metric on $\mathbb{R}^2$:

$$d\big((x_1, y_1), (x_2, y_2)\big) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Take an arbitrary point $p = (x, y) \in (a_1, b_1) \times (a_2, b_2)$. Then

$$a_1 < x < b_1, \qquad a_2 < y < b_2.$$

Define the four positive distances to the endpoints:

$$d_1 = x - a_1, \quad d_2 = b_1 - x, \quad d_3 = y - a_2, \quad d_4 = b_2 - y.$$

Let

$$r = \min\{d_1, d_2, d_3, d_4\} > 0.$$

We claim that the open ball $B(p, r) = \{q \in \mathbb{R}^2 : d(p, q) < r\}$ is contained in $(a_1, b_1) \times (a_2, b_2)$.

Indeed, take any point $q = (u, v) \in B(p, r)$. Then

$$|u - x| \leq d(p, q) < r, \qquad |v - y| \leq d(p, q) < r.$$

Hence

$$x - r < u < x + r, \qquad y - r < v < y + r.$$

Because $r \leq x - a_1$ and $r \leq b_1 - x$, we have

$$a_1 \leq x - r < u < x + r \leq b_1,$$

so $u \in (a_1, b_1)$. Similarly, $r \leq y - a_2$ and $r \leq b_2 - y$ imply

$$a_2 \leq y - r < v < y + r \leq b_2,$$

so $v \in (a_2, b_2)$. Consequently $q = (u, v) \in (a_1, b_1) \times (a_2, b_2)$.

Thus every point of the product has an open neighbourhood (an open ball) contained in the product. By definition, $(a_1, b_1) \times (a_2, b_2)$ is open in $\mathbb{R}^2$. $\square$

Definition 2.10 (Closed set)**.** A closed set A is defined by its ability to contain all of its limit points, meaning if a sequence of points $\{x_n\}$ within A converges to a limit x, then that limit x must also be in A.

Definition 2.11 (Sequence)**.** A sequence is a function whose domain is $\mathbb{N}$.

3 Lecture-03

Definition 3.1 (Sequence)**.** A sequence is a function where domain is $\mathbb{N}$.

Example 3.2. For example, let $x_n = \frac{1}{n}$ or $\left\{\frac{1}{n}\right\}_{n=1}^{\infty}$, be a sequence. Here, we write the sequence $f : \mathbb{N} \to \mathbb{R}$, where $f(n) = x_n = \frac{1}{n}$.

Definition 3.3 (Convergence of a Sequence)**.** A sequence of real numbers converges to a real number a if for every number $\varepsilon > 0$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$

$$|a_n - a| < \varepsilon$$

Example 3.4. Take our earlier example 3.2 , prove that $x_n = \left\{\frac{1}{n}\right\}_{n=1}^{\infty}$ converges to 0.

Proof. Let, $\varepsilon > 0$, we choose $N \in \mathbb{N}$ such that $N > 1/\varepsilon$ (which is possible due to the Archimedean property), giving us $\frac{1}{N} < \varepsilon$.

Then, for all $n \geq N$ we have

$$\left|\frac{1}{n} - 0\right| \leq \left|\frac{1}{N} - 0\right| < \varepsilon$$

□

Example 3.5. Consider $x_{n+1} = \frac{1}{2}\left(x_n + \frac{2}{x_n}\right)$, $x_0 > 0$

Show that x_n converges.

Proof. First, note that $x_n >= 0$ for all $n \in \mathbb{N}$. Because, $x_0 > 0$ and for induction, if we assume that $x_n > 0$ then,

$$x_{n+1} = \frac{1}{2}\left(x_n + \frac{2}{x_n}\right) > 0$$

since all the terms in left side is positive.

Then note that, if you start abouve $\sqrt{2}$ you stay above and decrease. Refer to the picture below for an example.



Next, to rigorously proof that the limit exists. We can do so by proving this sequence to be monotone and bounded. We can prove that by testing both cases $0 < x_0 \leq \sqrt{2}$ and $0 < x_0 < \sqrt{2}$. Now, let $(x_n) \to L$, where $L > 0$. Then $(x_{n+1}) \to L$

$$\begin{aligned} L &= \lim_{n\to\infty} x_{n+1} \\ &= \lim_{n\to\infty} \frac{1}{2}\left(x_n + \frac{2}{x_n}\right) \\ &= \lim_{n\to\infty} \frac{x_n}{2} + \frac{1}{x_n} \\ &= \frac{L}{2} + \frac{1}{L} \end{aligned}$$

This gives us,

$$L = \frac{L}{2} + \frac{1}{L} \implies \frac{L}{2} = \frac{1}{L} \implies L = \sqrt{2}$$

Hence, the given sequence converges to $\sqrt{2}$. □

Definition 3.6 (Convergent sequence; limit of sequence for $\mathbb{R}^n$ Space)**.** A sequence $i \to a_i$ of points $\mathbb{R}^n$ converges to $a \in \mathbb{R}^n$ if

$$\forall \varepsilon > 0, \exists M | m > M \implies |a_m - a| < \varepsilon$$

We then call a, the *limit* of the sequence

This also holds for vector a_i.

$$a_m = [(a_m)_1, (a_m)_2, \cdots, (a_m)_n] \to [a_1, a_2, \cdots, a_n] \in \mathbb{R}^n$$

Here, a_1 is the convergent point of sequence $(a_m)_1, (a_{m+1})_1, \cdots$. And so on.

Proposition 3.7 (Convergence in terms of coordinates)**.** *A sequence $m \to a_m$ with $a_m \in \mathbb{R}^n$ convergers to a if and only if each coordinate;* i.e. *for all $1 \leq j \leq n$, the jth coordinate $(a_i)_j)$ of a_i converges to a_j, the jth coordinate of the limit a.*

Proof. Let $\|\cdot\|$ denote the standard Euclidean norm in $\mathbb{R}^n$. Recall the definition of sequence convergence: $a_m \to a$ means that for every $\epsilon > 0$, there exists an integer N such that for all $m \geq N$, $\|a_m - a\| < \epsilon$.

Forward Direction (⇒):
Assume $a_m \to a$. We want to show that for any fixed coordinate $j \in \{1, \ldots, n\}$, the sequence $(a_m)_j \to a_j$.

Let $\epsilon > 0$ be given. Because $a_m \to a$, there exists an integer N such that for all $m \geq N$:

$$\|a_m - a\| < \epsilon$$

Observe that the absolute value of any single coordinate difference is bounded by the Euclidean distance:

$$|(a_m)_j - a_j| = \sqrt{((a_m)_j - a_j)^2} \leq \sqrt{\sum_{k=1}^{n} ((a_m)_k - a_k)^2} = \|a_m - a\|$$

Therefore, for all $m \geq N$, we have:

$$|(a_m)_j - a_j| \leq \|a_m - a\| < \epsilon$$

Since for any $\epsilon > 0$ we found an N that satisfies this condition, $(a_m)_j \to a_j$ for each j.

Reverse Direction ($\Leftarrow$):
Assume that for each $1 \leq j \leq n$, the coordinate sequence $(a_m)_j \to a_j$. We want to show that the vector sequence $a_m \to a$.

Let $\epsilon > 0$ be given. By the definition of convergence for each coordinate, there exists an integer N_j such that for all $m \geq N_j$:

$$|(a_m)_j - a_j| < \frac{\epsilon}{\sqrt{n}}$$

We want this to hold for *all* coordinates at the same time, so we choose the largest threshold: let $N = \max\{N_1, N_2, \ldots, N_n\}$.

For any $m \geq N$, m is greater than or equal to every N_j, so the inequality $|(a_m)_j - a_j| < \frac{\epsilon}{\sqrt{n}}$ holds for all $j = 1, \ldots, n$ simultaneously.

Now, we evaluate the distance between the vectors a_m and a for $m \geq N$:

$$\|a_m - a\| = \sqrt{\sum_{j=1}^{n} ((a_m)_j - a_j)^2} < \sqrt{\sum_{j=1}^{n} \left(\frac{\epsilon}{\sqrt{n}}\right)^2}$$

Simplifying the expression under the square root:

$$\|a_m - a\| < \sqrt{\sum_{j=1}^{n} \frac{\epsilon^2}{n}} = \sqrt{n \cdot \frac{\epsilon^2}{n}} = \sqrt{\epsilon^2} = \epsilon$$

Because for any $\epsilon > 0$ we found an N such that $m \geq N \implies \|a_m - a\| < \epsilon$, the sequence a_m converges to a. $\square$