

MAT092

Emon Hossain¹

¹Lecturer
MNS department
Brac University

LECTURE-02

Lecture Goals

By the end of this lecture, you should be able to:

- Understand the definition of continuity at a point.

- Identify different types of discontinuities.

- Check continuity for polynomials, rational and piecewise functions.

- Understand continuity using ε - δ rigor.

Table of Contents

- 1 Intuition of Continuity
- 2 Basic Examples of Continuity
- 3 Discontinuities
- 4 Continuity on Intervals
- 5 Rigorous Definition: ε - δ Continuity
- 6 Summary

What does continuity mean?

Informally, a function is **continuous at a point** if:

You can draw its graph near that point without lifting your pen.

What does continuity mean?

Informally, a function is **continuous at a point** if:

You can draw its graph near that point without lifting your pen.

Intuitively:

- No holes

- No jumps

- No infinite breaks

Limit-based definition (core idea)

A function $f(x)$ is **continuous at** $x = a$ if:

$$\lim_{x \rightarrow a} f(x) = f(a).$$

Limit-based definition (core idea)

A function $f(x)$ is **continuous at** $x = a$ if:

$$\lim_{x \rightarrow a} f(x) = f(a).$$

This means three things must hold:

- (i) $f(a)$ is defined
- (i) $\lim_{x \rightarrow a} f(x)$ exists
- (i) $\lim_{x \rightarrow a} f(x) = f(a)$

Table of Contents

- 1 Intuition of Continuity
- 2 Basic Examples of Continuity**
- 3 Discontinuities
- 4 Continuity on Intervals
- 5 Rigorous Definition: ε - δ Continuity
- 6 Summary

Example 1: Polynomial functions

Consider:

$$f(x) = 3x^2 - 5x + 1.$$

Example 1: Polynomial functions

Consider:

$$f(x) = 3x^2 - 5x + 1.$$

Solution: Polynomials are continuous for all real numbers.

$$\lim_{x \rightarrow a} f(x) = f(a) \quad \text{for all } a \in \mathbb{R}.$$

Example 1: Polynomial functions

Consider:

$$f(x) = 3x^2 - 5x + 1.$$

Solution: Polynomials are continuous for all real numbers.

$$\lim_{x \rightarrow a} f(x) = f(a) \quad \text{for all } a \in \mathbb{R}.$$

\Rightarrow Continuous everywhere.

Example 2: Rational function (continuous domain)

Consider:

$$f(x) = \frac{x^2 - 1}{x - 1}.$$

Example 2: Rational function (continuous domain)

Consider:

$$f(x) = \frac{x^2 - 1}{x - 1}.$$

Simplify:

$$f(x) = x + 1 \quad (x \neq 1).$$

Example 2: Rational function (continuous domain)

Consider:

$$f(x) = \frac{x^2 - 1}{x - 1}.$$

Simplify:

$$f(x) = x + 1 \quad (x \neq 1).$$

At $x = 1$:

$f(1)$ is undefined.

Example 2: Rational function (continuous domain)

Consider:

$$f(x) = \frac{x^2 - 1}{x - 1}.$$

Simplify:

$$f(x) = x + 1 \quad (x \neq 1).$$

At $x = 1$:

$f(1)$ is undefined.

$\Rightarrow f$ is continuous for $x \neq 1$, but discontinuous at $x = 1$.

(Removable discontinuity)

Example 3: Absolute value

Consider:

$$f(x) = |x|.$$

Example 3: Absolute value

Consider:

$$f(x) = |x|.$$

Check continuity at $x = 0$:

$$\lim_{x \rightarrow 0^-} |x| = 0, \quad \lim_{x \rightarrow 0^+} |x| = 0.$$

Example 3: Absolute value

Consider:

$$f(x) = |x|.$$

Check continuity at $x = 0$:

$$\lim_{x \rightarrow 0^-} |x| = 0, \quad \lim_{x \rightarrow 0^+} |x| = 0.$$

Since $f(0) = 0$,

$$\lim_{x \rightarrow 0} f(x) = f(0).$$

Example 3: Absolute value

Consider:

$$f(x) = |x|.$$

Check continuity at $x = 0$:

$$\lim_{x \rightarrow 0^-} |x| = 0, \quad \lim_{x \rightarrow 0^+} |x| = 0.$$

Since $f(0) = 0$,

$$\lim_{x \rightarrow 0} f(x) = f(0).$$

$\Rightarrow f(x) = |x|$ is continuous everywhere.

Table of Contents

- 1 Intuition of Continuity
- 2 Basic Examples of Continuity
- 3 Discontinuities**
- 4 Continuity on Intervals
- 5 Rigorous Definition: ε - δ Continuity
- 6 Summary

Example 4: Jump discontinuity

Define:

$$f(x) = \begin{cases} 2x, & x < 1, \\ 5, & x \geq 1. \end{cases}$$

Example 4: Jump discontinuity

Define:

$$f(x) = \begin{cases} 2x, & x < 1, \\ 5, & x \geq 1. \end{cases}$$

Compute:

$$\lim_{x \rightarrow 1^-} f(x) = 2, \quad \lim_{x \rightarrow 1^+} f(x) = 5.$$

Example 4: Jump discontinuity

Define:

$$f(x) = \begin{cases} 2x, & x < 1, \\ 5, & x \geq 1. \end{cases}$$

Compute:

$$\lim_{x \rightarrow 1^-} f(x) = 2, \quad \lim_{x \rightarrow 1^+} f(x) = 5.$$

Since left \neq right,

$$\lim_{x \rightarrow 1} f(x) \text{ does not exist.}$$

Example 4: Jump discontinuity

Define:

$$f(x) = \begin{cases} 2x, & x < 1, \\ 5, & x \geq 1. \end{cases}$$

Compute:

$$\lim_{x \rightarrow 1^-} f(x) = 2, \quad \lim_{x \rightarrow 1^+} f(x) = 5.$$

Since left \neq right,

$$\lim_{x \rightarrow 1} f(x) \text{ does not exist.}$$

$\Rightarrow f$ is discontinuous at $x = 1$.

Example 5: Infinite discontinuity

Consider:

$$f(x) = \frac{1}{x}.$$

Example 5: Infinite discontinuity

Consider:

$$f(x) = \frac{1}{x}.$$

At $x = 0$:

$$\lim_{x \rightarrow 0^+} f(x) = +\infty, \quad \lim_{x \rightarrow 0^-} f(x) = -\infty.$$

Example 5: Infinite discontinuity

Consider:

$$f(x) = \frac{1}{x}.$$

At $x = 0$:

$$\lim_{x \rightarrow 0^+} f(x) = +\infty, \quad \lim_{x \rightarrow 0^-} f(x) = -\infty.$$

$\Rightarrow f$ has an infinite discontinuity at $x = 0$.

Example 6: Oscillatory discontinuity

Consider:

$$f(x) = \sin\left(\frac{1}{x}\right).$$

Example 6: Oscillatory discontinuity

Consider:

$$f(x) = \sin\left(\frac{1}{x}\right).$$

As $x \rightarrow 0$, the function oscillates infinitely between -1 and 1 .

Example 6: Oscillatory discontinuity

Consider:

$$f(x) = \sin\left(\frac{1}{x}\right).$$

As $x \rightarrow 0$, the function oscillates infinitely between -1 and 1 .

$\lim_{x \rightarrow 0} f(x)$ does not exist.

Example 6: Oscillatory discontinuity

Consider:

$$f(x) = \sin\left(\frac{1}{x}\right).$$

As $x \rightarrow 0$, the function oscillates infinitely between -1 and 1 .

$$\lim_{x \rightarrow 0} f(x) \text{ does not exist.}$$

$\Rightarrow f$ is discontinuous at $x = 0$.

Table of Contents

- 1 Intuition of Continuity
- 2 Basic Examples of Continuity
- 3 Discontinuities
- 4 Continuity on Intervals**
- 5 Rigorous Definition: ε - δ Continuity
- 6 Summary

Continuity on intervals

A function is:

Continuous on (a, b) if continuous at every point in the interval

Continuous on $[a, b]$ if:

Continuous on (a, b)

Right-continuous at a

Left-continuous at b

Example 7: Endpoint continuity

Consider:

$$f(x) = \sqrt{x}, \quad x \in [0, 4].$$

Example 7: Endpoint continuity

Consider:

$$f(x) = \sqrt{x}, \quad x \in [0, 4].$$

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0 = f(0).$$

Example 7: Endpoint continuity

Consider:

$$f(x) = \sqrt{x}, \quad x \in [0, 4].$$

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0 = f(0).$$

$$\lim_{x \rightarrow 4^-} \sqrt{x} = 2 = f(4).$$

Example 7: Endpoint continuity

Consider:

$$f(x) = \sqrt{x}, \quad x \in [0, 4].$$

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0 = f(0).$$

$$\lim_{x \rightarrow 4^-} \sqrt{x} = 2 = f(4).$$

$\Rightarrow f$ is continuous on $[0, 4]$.

Table of Contents

- 1 Intuition of Continuity
- 2 Basic Examples of Continuity
- 3 Discontinuities
- 4 Continuity on Intervals
- 5 Rigorous Definition: ε - δ Continuity**
- 6 Summary

Formal ε - δ definition

A function f is continuous at $x = a$ if:

For every $\varepsilon > 0$, there exists $\delta > 0$ such that

$$|x - a| < \delta \quad \Rightarrow \quad |f(x) - f(a)| < \varepsilon.$$

Formal ε - δ definition

A function f is continuous at $x = a$ if:

For every $\varepsilon > 0$, there exists $\delta > 0$ such that

$$|x - a| < \delta \quad \Rightarrow \quad |f(x) - f(a)| < \varepsilon.$$

This is exactly the ε - δ definition of:

$$\lim_{x \rightarrow a} f(x) = f(a).$$

Example 8: ε - δ proof (linear function)

Show that $f(x) = 2x + 1$ is continuous at $x = 3$.

Example 8: ε - δ proof (linear function)

Show that $f(x) = 2x + 1$ is continuous at $x = 3$.

We must show:

$$|f(x) - f(3)| < \varepsilon.$$

Example 8: ε - δ proof (linear function)

Show that $f(x) = 2x + 1$ is continuous at $x = 3$.

We must show:

$$|f(x) - f(3)| < \varepsilon.$$

Compute:

$$|2x + 1 - 7| = 2|x - 3|.$$

Example 8: ε - δ proof (linear function)

Show that $f(x) = 2x + 1$ is continuous at $x = 3$.

We must show:

$$|f(x) - f(3)| < \varepsilon.$$

Compute:

$$|2x + 1 - 7| = 2|x - 3|.$$

To ensure $2|x - 3| < \varepsilon$, choose:

$$\boxed{\delta = \frac{\varepsilon}{2}}.$$

Example 8: ε - δ proof (linear function)

Show that $f(x) = 2x + 1$ is continuous at $x = 3$.

We must show:

$$|f(x) - f(3)| < \varepsilon.$$

Compute:

$$|2x + 1 - 7| = 2|x - 3|.$$

To ensure $2|x - 3| < \varepsilon$, choose:

$$\boxed{\delta = \frac{\varepsilon}{2}}.$$

Thus $|x - 3| < \delta$ implies $|f(x) - f(3)| < \varepsilon$.



Example 9: ε - δ continuity of x^2

Show that $f(x) = x^2$ is continuous at $x = 1$.

Example 9: ε - δ continuity of x^2

Show that $f(x) = x^2$ is continuous at $x = 1$.

We want:

$$|x^2 - 1| < \varepsilon.$$

Example 9: ε - δ continuity of x^2

Show that $f(x) = x^2$ is continuous at $x = 1$.

We want:

$$|x^2 - 1| < \varepsilon.$$

Factor:

$$|x^2 - 1| = |x - 1||x + 1|.$$

Example 9: ε - δ continuity of x^2

Show that $f(x) = x^2$ is continuous at $x = 1$.

We want:

$$|x^2 - 1| < \varepsilon.$$

Factor:

$$|x^2 - 1| = |x - 1||x + 1|.$$

Assume $|x - 1| < 1 \Rightarrow x \in (0, 2)$ so $|x + 1| < 3$.

Example 9: ϵ - δ continuity of x^2

Show that $f(x) = x^2$ is continuous at $x = 1$.

We want:

$$|x^2 - 1| < \epsilon.$$

Factor:

$$|x^2 - 1| = |x - 1||x + 1|.$$

Assume $|x - 1| < 1 \Rightarrow x \in (0, 2)$ so $|x + 1| < 3$.

Thus:

$$|x^2 - 1| < 3|x - 1|.$$

Example 9: ε - δ continuity of x^2

Show that $f(x) = x^2$ is continuous at $x = 1$.

We want:

$$|x^2 - 1| < \varepsilon.$$

Factor:

$$|x^2 - 1| = |x - 1||x + 1|.$$

Assume $|x - 1| < 1 \Rightarrow x \in (0, 2)$ so $|x + 1| < 3$.

Thus:

$$|x^2 - 1| < 3|x - 1|.$$

Choose:

$$\delta = \min \left\{ 1, \frac{\varepsilon}{3} \right\}.$$



Table of Contents

- 1 Intuition of Continuity
- 2 Basic Examples of Continuity
- 3 Discontinuities
- 4 Continuity on Intervals
- 5 Rigorous Definition: ε - δ Continuity
- 6 Summary**

Key Takeaways

Continuity means: limit equals function value.

Three checks: existence of $f(a)$, limit, and equality.

Polynomials and absolute values are continuous everywhere.

Rational functions are continuous where denominators $\neq 0$.

ε - δ makes continuity mathematically precise.

Questions?