

# MAT215: Complex Variables And Laplace Transformations

Emon Hossain<sup>1</sup>

<sup>1</sup>Lecturer  
MNS department  
Brac University

LECTURE-02

## Some important formulas

$$2 \sin(A) \cos(B) = \sin(A+B) + \sin(A-B)$$

$$2 \cos(A) \cos(B) = \cos(A+B) + \cos(A-B)$$

$$2 \sin(A) \sin(B) = \cos(A-B) - \cos(A+B)$$

$$\sinh(\theta) = \frac{e^{\theta} - e^{-\theta}}{2}, \cosh(\theta) = \frac{e^{\theta} + e^{-\theta}}{2}$$

$$\int e^{at} \sin(bt) dt = \frac{e^{at}}{a^2 + b^2} (a \sin(bt) - b \cos(bt)) + C$$

$$\int e^{at} \cos(bt) dt = \frac{e^{at}}{a^2 + b^2} (a \cos(bt) + b \sin(bt)) + C$$

# Some important Laplace Transform formulas

- Known formula
- First Translation formula:

$$\mathcal{L}\{e^{at}f(t)\} = \mathcal{L}\{f(t)\}_{s \rightarrow s-a} = F(s-a)$$

- Truncation/Second Translation formula:

$$\mathcal{L}\{f(t-a)u(t-a)\} = e^{-as}F(s)$$

- s-differentiation:

$$\mathcal{L}\{(t)^n f(t)\} = (-1)^n \frac{d^n}{ds^n} F(s)$$

- Partial fraction type
- Convolution type (Not very important)

# Example

## Example

Find the Laplace transform of  $f(t) = t$ .

## Example

Find the Laplace transform of  $f(t) = e^{at}$ .

## Example

- Find the Laplace transform of  $f(t) = 2e^{-2t}$ .
- Find the Laplace transform of  $f(t) = 4t^3 - e^{-1}$ .
- Find the Laplace transform of  $f(t) = (t^2 + 1)^2$ .
- Find the Laplace transform of  $f(t) = \cosh(3t) + \sin(2t)$ .

## Example

- a. Find the Laplace transform of  $f(t) = \cos(at)$ .
- b. Find the Laplace transform of  $f(t) = \sin(at)$ .
- c. Find the Laplace transform of  $f(t) = \sinh(at)$ .
- d. Find the Laplace transform of  $f(t) = \cosh(at)$ .

Find the Laplace transform of  $f(t) = \sin(at)$ .

$$\begin{aligned}\mathcal{L}\{\sin(at)\} &= \int_0^{\infty} e^{-st} \sin(at) dt \\ &= \text{Skip way-01}\end{aligned}$$

## Way-02 (best way)

Find the Laplace transform of  $f(t) = \sin(at)$ .

$$\begin{aligned}\mathcal{L}\{\sin(at)\} &= \int_0^{\infty} e^{-st} \sin(at) dt \\&= \int_0^{\infty} e^{-st} \frac{e^{i(at)} - e^{-i(at)}}{2i} dt \\&= \frac{1}{2i} \left( \int_0^{\infty} e^{-st} e^{i(at)} dt - \int_0^{\infty} e^{-st} e^{-i(at)} dt \right) \\&= \frac{1}{2i} \left[ \frac{1}{s - ia} - \frac{1}{s + ia} \right] \\&= \frac{1}{2i} \frac{2ia}{s^2 + a^2} \\&= \frac{a}{s^2 + a^2}\end{aligned}$$

Find the Laplace transform of  $f(t) = \sin(at)$ .

$$\begin{aligned}\mathcal{L}\{\sin(at)\} &= \int_0^{\infty} e^{-st} \sin(at) dt \\&= \lim_{b \rightarrow \infty} \int_0^b e^{-st} \sin(at) dt \\&= \lim_{b \rightarrow \infty} \left[ \frac{e^{-st}(-a \cos(at) - s \sin(at))}{a^2 + s^2} \right]_0^b \\&= \lim_{b \rightarrow \infty} \left( \frac{e^{-sb}(-a \cos(ab) - s \sin(ab))}{a^2 + s^2} - \frac{-a}{a^2 + s^2} \right) \\&= \frac{a}{a^2 + s^2}\end{aligned}$$



## Example

- Find the Laplace transform of  $f(t) = t^3$ .
- Find the Laplace transform of  $f(t) = t^3 e^{5t}$ .

Hint: Use the first Translation formula:

$$\mathcal{L}\{e^{at}f(t)\} = \mathcal{L}\{f(t)\}_{s \rightarrow s-a} = F(s-a)$$

## Example

Find the Laplace transform of  $f(t) = 5\sin(4t)$ .

Find the Laplace transform of  $f(t) = 5\sin(4t)e^{-3t}$ .

Hint: Use the first Translation formula:

$$\mathcal{L}\{e^{at}f(t)\} = \mathcal{L}\{f(t)\}_{s \rightarrow s-a} = F(s-a)$$

### Example

Find the Laplace transform of  $f(t) = (t+2)^2$ .

Find the Laplace transform of  $f(t) = (t+2)^2 e^t$ .

## Example

Find the Laplace transform of  $f(t) = t \sin(at)$ .

## Example

Find the Laplace transform of  $f(t) = t \sin(at)$ .

$$\begin{aligned}\mathcal{L}\{t \sin(at)\} &= \int_0^{\infty} e^{-st} t \sin(at) dt \\&= \int_0^{\infty} t \underbrace{e^{-st} \sin(at)}_v dt \\&= \left[ t \frac{e^{-st}(-a \cos(at) - s \sin(at))}{a^2 + s^2} \right]_0^{\infty} \\&\quad - \int_0^{\infty} \left[ t \frac{e^{-st}(-a \cos(at) - s \sin(at))}{a^2 + s^2} \right]_0^{\infty} dt \\&= \frac{s}{(s^2 + a^2)} \int_0^{\infty} e^{-st} \sin(at) dt + \frac{a}{(s^2 + a^2)} \int_0^{\infty} e^{-st} \cos(at) dt \\&= \frac{s}{(s^2 + a^2)} \cdot \frac{a}{s^2 + a^2} + \frac{a}{(s^2 + a^2)} \cdot \frac{s}{s^2 + a^2} \\&= \frac{2as}{(s^2 + a^2)^2}\end{aligned}$$

**Solution:**

$$\begin{aligned}\mathcal{L}\{t \sin(at)\} &= \int_0^{\infty} e^{-st} t \sin(at) dt \\&= - \int_0^{\infty} e^{-st} \frac{\partial}{\partial a} (\cos(at)) dt \\&= - \frac{\partial}{\partial a} \int_0^{\infty} e^{-st} \cos(at) dt \\&= - \frac{\partial}{\partial a} \left( \frac{s}{s^2 + a^2} \right) \\&= \frac{2as}{(s^2 + a^2)^2}\end{aligned}$$

## Example

Find the Laplace transform of  $f(t) = 2\sinh(3t) + 5\cosh(4t)$ .

Find the Laplace transform of  $f(t) = e^{2t}[2\sinh(3t) + 5\cosh(4t)]$ .

Find the Laplace transform of  $f(t) = te^{2t}[2\sinh(3t) + 5\cosh(4t)]$