

pt

Stimulated Neutrino Transitions and Rabi Oscillations

Lei Ma

Department of Physics
UNM

January 18, 2017

OUTLINE

1. Introduction

What are Neutrinos
Neutrino Oscillations
Why Do Neutrinos Oscillate

2. Matter Effect

Matter Interaction
MSW Effect
Solar Neutrino Problem
Stimulated Neutrino Oscillations

3. Stimulated Neutrino Flavor Transitions

Hamiltonian and Basis
Rabi Oscillations

4. Jacobi-Anger Expansion

Basis and Formalism

5. Summary

OVERVIEW

pt

100

150

200

250

300

350

Introduction

What are Neutrinos

Neutrino Oscillations

Why Do Neutrinos Oscillate

100

Matter Effect

Stiffened Neutrino Flavor Transitions

150


Jacobi-Anger Expansion

Summary

200


250

WHAT ARE NEUTRINOS?



Beta decay and antineutrino production. Source: Beta_Decay@Wikipedia


WHAT ARE NEUTRINOS?



Fermilab 95-759

Neutrinos are


- fermions,
- electrically neutral,
- light.



Adapted from Olga Mena & Stephen Parke (2004)

Table of elementary particles. Source:
Fermi...


WHAT ARE NEUTRINOS?



Fermilab 95-759

Neutrinos are

- fermions,
- electrically neutral,
- light.



Adapted from Olga Mena & Stephen Parke (2004)


Table of elementary particles. Source:

Fermilab 95-759

WHAT IS NEUTRINO OSCILLATION?


Neutrino Oscillation

Neutrino Flavor Conversion



Neutrino Oscillations

WHAT IS NEUTRINO OSCILLATION?



Probabilities of finding neutrinos to be in each flavor.

WHY DO NEUTRINOS OSCILLATE?

$$i\partial_x |\Psi\rangle = \hat{H} |\Psi\rangle$$

- ▶ Basis: Hamiltonian diagonalized basis/mass basis/propagation basis, $\{|\nu_1\rangle, |\nu_2\rangle\}$.

▶

$$\mathbf{H} = -\frac{\omega_v}{2} \sigma_3, \quad \text{where } \omega_v = \frac{\delta m^2}{2E} = \frac{m_2^2 - m_1^2}{2E}.$$

- ▶ The system can be solved given initial condition of the amplitudes of the two eigenstates $(\langle\nu_1|\Psi(0)\rangle, \langle\nu_2|\Psi(0)\rangle)^T$,

$$\begin{pmatrix} \langle\nu_1|\Psi(x)\rangle \\ \langle\nu_2|\Psi(x)\rangle \end{pmatrix} = \begin{pmatrix} \langle\nu_1|\Psi(0)\rangle \exp(i\omega_v x/2) \\ \langle\nu_2|\Psi(0)\rangle \exp(-i\omega_v x/2) \end{pmatrix}$$

WHY DO NEUTRINOS OSCILLATE?

Flavor basis

Neutrino wave function in flavor basis $\{|\nu_e\rangle, |\nu_\mu\rangle\}$ is related to state in energy basis $\{|\nu_1\rangle, |\nu_2\rangle\}$ through

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \begin{pmatrix} \cos \theta_v & \sin \theta_v \\ -\sin \theta_v & \cos \theta_v \end{pmatrix} \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix}$$

θ_v : vacuum mixing angle

WHY DO NEUTRINOS OSCILLATE?

Flavor basis

Neutrino wave function in flavor basis $\{|\nu_e\rangle, |\nu_\mu\rangle\}$ is related to state in energy basis $\{|\nu_1\rangle, |\nu_2\rangle\}$ through

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \begin{pmatrix} \cos \theta_v & \sin \theta_v \\ -\sin \theta_v & \cos \theta_v \end{pmatrix} \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix}$$

θ_v : vacuum mixing angle

Hamiltonian H

Mass basis

$$\begin{aligned} & \frac{\omega_v}{2} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= -\frac{\omega_v}{2} \sigma_3 \end{aligned}$$

Flavor basis

$$\begin{aligned} & \frac{\omega_v}{2} \begin{pmatrix} -\cos 2\theta_v & \sin 2\theta_v \\ \sin 2\theta_v & \cos 2\theta_v \end{pmatrix} \\ &= \frac{\omega_v}{2} (-\cos 2\theta_v \sigma_3 + \sin 2\theta_v \sigma_1) \end{aligned}$$

NATURE OF NEUTRINO OSCILLATION

pt

100

150

200

250

300

350

Transition Probability

100

$$P(|\nu_e\rangle \rightarrow |\nu_\mu\rangle) = \sin^2(2\theta_v) \sin^2(\omega_v x/2)$$

150


$\omega_v = (m_2^2 - m_1^2)/2E$ determines oscillation wavelength.

► Mixing angle θ_v determines flavor oscillation amplitude.



200

250

OVERVIEW



MATTER INTERACTION



MATTER INTERACTION

pt

Hamiltonian with matter interaction in flavor basis ($\omega_\nu = \delta m^2/2E$):

$$\mathbf{H} = \frac{\omega_\nu}{2} \begin{pmatrix} -\cos 2\theta_\nu & \sin 2\theta_\nu \\ \sin 2\theta_\nu & \cos 2\theta_\nu \end{pmatrix} \pm \sqrt{2} G_F n_e(x) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

• vacuum Hamiltonian

► Matter interaction

MATTER INTERACTION

pt

Hamiltonian with matter interaction in flavor basis ($\omega_\nu = \delta m^2/2E$):

$$\mathbf{H} = \frac{\omega_\nu}{2} (-\cos 2\theta_\nu \sigma_3 + \sin 2\theta_\nu \sigma_1) + \frac{\lambda(x)}{2} \sigma_3$$

► vacuum Hamiltonian

► Matter interaction

► $\lambda(x) = \sqrt{2}G_F n_e(x)$

MSW EFFECT

Hamiltonian in Vacuum

pt

100

150

200

250

300

350

$$\mathbb{H}_{\text{vacuum}} = \frac{\omega_v \cos 2\theta_v}{2} \sigma_3 + \frac{\omega_v \sin 2\theta_v}{2} \sigma_1$$

100

$$\begin{aligned} \mathbf{H} &= \frac{\lambda(x) - \omega_v \cos 2\theta_v}{2} \sigma_3 + \frac{\omega_v \sin 2\theta_v}{2} \sigma_1 \\ &= \frac{\omega_m(x) \cos 2\theta_m(x)}{2} \sigma_3 + \frac{\omega_m(x) \sin 2\theta_m(x)}{2} \sigma_1, \end{aligned}$$

150

where

200

$$\omega_m(x) = \sqrt{(\lambda(x) - \omega_v \cos 2\theta_v)^2 + \omega_v^2 \sin^2 2\theta_v},$$

$$\tan 2\theta_m(x) = \frac{\omega_v \sin 2\theta_v}{\omega_v \cos 2\theta_v - \lambda(x)}.$$

250

MSW EFFECT

pt

100

150

200

250

300

350

Constant matter profile λ_0 as an example,

Significance of θ_m

100

Define matter basis $\{|\nu_L\rangle, |\nu_H\rangle\}$

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \begin{pmatrix} \cos \theta_m & \sin \theta_m \\ -\sin \theta_m & \cos \theta_m \end{pmatrix} \begin{pmatrix} |\nu_L\rangle \\ |\nu_H\rangle \end{pmatrix}$$

150

In matter basis

$$\mathbf{H}_{\text{matter-basis}} = -\frac{\omega_m}{2} \sigma_3$$

200

250

MSW RESONANCE

Hamiltonian with Matter Potential

pt

100

150

200

250

300

350

$$\begin{aligned} H &= \frac{\lambda(x) - \omega_v \cos 2\theta_v}{2} \sigma_3 + \frac{\omega_v \sin 2\theta_v}{2} \sigma_1 \\ &= \frac{\omega_m(x) \cos 2\theta_m(x)}{2} \sigma_3 + \frac{\omega_m(x) \sin 2\theta_m(x)}{2} \sigma_1 \end{aligned}$$

100

$$\tan 2\theta_m(x) = \frac{\omega_v \sin 2\theta_v}{\omega_v \cos 2\theta_v - \lambda(x)}.$$

150

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \begin{pmatrix} \cos \theta_m & \sin \theta_m \\ -\sin \theta_m & \cos \theta_m \end{pmatrix} \begin{pmatrix} |\nu_L\rangle \\ |\nu_H\rangle \end{pmatrix}$$


200

Transition Probability

$$P(|\nu_e\rangle \rightarrow |\nu_\mu\rangle) = \sin^2(2\theta_m) \sin^2(\omega_m x)$$

250

SOLAR NEUTRINO PROBLEM



Chlorine detector (Homestake experiment) results and theory predictions. SNU: 1 event for 10^{36} target atoms per second. Kenneth R. Lang (2010)


MSW EFFECT AND SOLAR NEUTRINOS

pt

$$100 = \frac{\lambda(x) - \omega}{2} \frac{\cos 2\theta_v}{2} - \frac{\omega_v \sin^2 \theta}{2} \tau_1$$

$$\begin{pmatrix} |\nu_L\rangle \\ |\nu_H\rangle \end{pmatrix} = \begin{pmatrix} \cos \theta_m & -\sin \theta_m \\ \sin \theta_m & \cos \theta_m \end{pmatrix} \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix}$$

$$H_{\text{matter-basis}} = -\frac{\omega_m}{2} \sigma_3$$



Yellow bar is the resonance point. Red: $|\nu_e\rangle$. Green: $|\nu_\mu\rangle$. Adapted from Smirnov, 2003.

MSW EFFECT

pt

Suppose $\omega_v = (m_2^2 - m_1^2)/2E < 0$,

$$\mathbf{H} = -\frac{\omega_v}{2} \begin{pmatrix} -\cos 2\theta_v & \sin 2\theta_v \\ \sin 2\theta_v & \cos 2\theta_v \end{pmatrix} + \sqrt{2}G_F n_e(x) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$


↓

$$\mathbf{H} = \left(\frac{-\omega_v}{2} \cos 2\theta_v + \frac{\lambda(x)}{2} \right) \sigma_3 - \frac{\omega_v}{2} \sin 2\theta_v \sigma_1$$

SUPERNOVA MATTER DENSITY PROFILE

Why Do We Care

astrophysical environments: supernovae, accretion disks etc



Supernova shock and turbulence. E. Borriello, et al (2014)

$$\Delta n_e(r) = \sum_n c_n \sin(k_n r + \phi_n)$$

STIMULATED NEUTRINO OSCILLATIONS

Matter Profile

pt

100

150

200

250

300

350

$$\lambda(x) = \lambda_0 + \delta\lambda(x)$$

100 is

Background matter basis: Hamiltonian is diagonalized with only background matter profile λ_0 ,

150

$$H_{\text{background}} = -\frac{\omega_m}{2}\sigma_3.$$

Hamiltonian

200


$$H = \frac{1}{2} (-\omega_m + \delta\lambda(x) \cos 2\theta_m) \sigma_3 - \frac{\delta\lambda(x)}{2} \sin \theta_m \sigma_1.$$

250

STIMULATED NEUTRINO OSCILLATIONS

P. Krastev and A. Smirnov (1989); J. Kneller et al (2013);

pt Patton et al (2014); 150 200 250 300 350



Stimulated oscillations. $\lambda(x) = \lambda_0 + \Lambda \sin(kx)$ with $\hat{x} = \omega_m x$, $\Lambda = 0.1\omega_m$,
 $k = 0.995\omega_m$, $\theta_m = \pi/6$

OVERVIEW

pt

100

150

200

250

300

350

Introduction

Matter Effect

100

Stimulated Neutrino Flavor Transitions

Hamiltonian and Basis

Rabi Oscillations

150

Jacobi-Anger Expansion

Summary

200

250

HAMILTONIAN

pt

Hamiltonian in Background Matter Basis

$$100 \quad \mathbf{H} = \frac{1}{2} (-\omega_m + \delta\lambda(x) \cos 2\theta_m) \sigma_3 - \frac{\delta\lambda(x)}{2} \sin 2\theta_m \sigma_1.$$

Matter profile

$$150 \quad \lambda(x) = \lambda_0 + A \cos(kx),$$

$$200 \quad \mathbf{H} = \frac{1}{2} (-\omega_m + \cos 2\theta_m A \cos(kx)) \sigma_3 - \frac{\sin 2\theta_m}{2} A \cos(kx) \sigma_1.$$

250

RABI OSCILLATIONS

pt

100

150

200

250

300

350

Rabi Oscillation

100


Hamiltonian

$$H = \frac{\omega_0}{2} \sigma_z - \frac{\alpha}{2} \begin{pmatrix} 0 & e^{ikt} \\ e^{-ikt} & 0 \end{pmatrix}$$

$$E_2 = \frac{\omega_0}{2}$$

$$E_1 = -\frac{\omega_0}{2}$$

Incoming light



Frequency : k

200

250

RABI OSCILLATIONS

$$\frac{1}{2} (-\omega_m + \cos 2\theta_m A \cos(kx)) \sigma_3 - \frac{\sin 2\theta_m}{2} A \cos(kx) \sigma_1$$


Rabi Oscillation

Hamiltonian

$$-\frac{\omega_0}{2} \sigma_3 - \frac{\alpha}{2} \begin{pmatrix} 0 & e^{ikt} \\ e^{-ikt} & 0 \end{pmatrix}$$

$$E_2 = \frac{\omega_0}{2}$$

$$E_1 = -\frac{\omega_0}{2}$$



The transition probability from low energy to high energy is

$$P_{1 \rightarrow 2} = \frac{\alpha^2}{\alpha^2 + (\omega_0 - k)^2} \sin^2 \left(\frac{\Omega_R}{2} t \right),$$

where

$$\Omega_R = \sqrt{\alpha^2 + (\omega_0 - k)^2}$$

is Rabi frequency.

RABI OSCILLATIONS


Rabi Oscillations

Hamiltonian

$$-\frac{\hbar\omega_0}{2}\sigma_3 - \frac{\alpha}{2} \begin{pmatrix} 0 & e^{ikt} \\ e^{-ikt} & 0 \end{pmatrix}$$

$$E_2 = \frac{\hbar\omega_0}{2}$$

$$E_1 = -\frac{\hbar\omega_0}{2}$$



Frequency : k


The transition probability from low energy to high energy is

$$P_{1 \rightarrow 2} = \frac{1}{1 + D^2} \sin^2 \left(\frac{\Omega_R}{2} t \right),$$

where

$$D = \left| \frac{\omega_0 - k}{\alpha} \right|.$$

RABI OSCILLATIONS



Amplitude of Rabi oscillations for different driving field frequency k


VISUALIZING RABI OSCILLATIONS

$$-\frac{\omega_0}{2}\sigma_3 - \frac{\alpha}{2} \begin{pmatrix} 0 & e^{ikt} \\ e^{-ikt} & 0 \end{pmatrix}$$

$$-\frac{\omega_0}{2}\sigma_3 - \frac{\alpha}{2} \cos(kt)\sigma_1 + \frac{\alpha}{2} \sin(kt)\sigma_2$$

$$= (\alpha \cos(kt) \quad -\alpha \sin(kt) \quad \omega_0) \cdot \begin{pmatrix} -\sigma_1/2 \\ -\sigma_2/2 \\ -\sigma_3/2 \end{pmatrix}$$

$$= \vec{H} \cdot (-\vec{\sigma}/2)$$



$$D = \left| \frac{\omega_0 - k}{\alpha} \right|$$

is ratio of the energy gap in corotating frame to width of resonance.

INTERFERENCES OF RABI OSCILLATIONS

pt

$$\mathbf{H} = \frac{1}{2} \left(-\omega_m + \cos 2\theta_m A \cos(kx) \right) \sigma_3 - \frac{\sin 2\theta_m}{2} A \cos(kx) \sigma_1$$

$$\rightarrow -\frac{\omega_m}{2} \sigma_3 - \frac{A \sin 2\theta_m}{2} \cos(kx) \sigma_1$$

$$= \begin{pmatrix} 0 \\ 0 \\ \omega_m \end{pmatrix} + \frac{A \sin 2\theta_m}{2} \begin{pmatrix} \cos(kx) \\ \sin(kx) \\ 0 \end{pmatrix} + \frac{A \sin 2\theta_m}{2} \begin{pmatrix} \cos(-kx) \\ \sin(-kx) \\ 0 \end{pmatrix}$$

200 frequencies!

250

INTERFERENCES OF RABI OSCILLATIONS

pt

$$\vec{H} = \begin{pmatrix} 0 \\ 0 \\ \omega_m \end{pmatrix} + \alpha_1 \begin{pmatrix} \cos(k_1 x) \\ -\sin(k_1 x) \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} \cos(k_2 x) \\ -\sin(k_2 x) \\ 0 \end{pmatrix}$$

Changing frame of the second frequency,

$$\vec{H} = \begin{pmatrix} 0 \\ 0 \\ \omega_m - k_2 \end{pmatrix} + \alpha_1 \begin{pmatrix} \cos(k_1 - k_2 x) \\ -\sin(k_1 - k_2 x) \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

Energy gap in this frame becomes the length of the vector

$$\begin{pmatrix} 0 \\ 0 \\ \omega_m - k_2 \end{pmatrix} + \alpha_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

INTERFERENCES OF RABI OSCILLATIONS

$$\vec{H} = \begin{pmatrix} 0 \\ \omega_m \end{pmatrix} + \alpha_1 \begin{pmatrix} \cos(k_1 x) \\ \sin(k_1 x) \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} \cos(k_2 x) \\ \sin(k_2 x) \\ 0 \end{pmatrix}$$

Corotating frame of the second frequency,

$$\vec{H} = \begin{pmatrix} 0 \\ 0 \\ \omega_m - k_2 \end{pmatrix} + \alpha_1 \begin{pmatrix} \cos(k_1 - k_2 x) \\ -\sin(k_1 - k_2 x) \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

Energy gap in this frame becomes the length of the vector

$$\sqrt{(\omega_m - k_2)^2 + \alpha_2^2} \rightarrow \omega_m - k_2 + \frac{1}{2} \frac{\alpha_2^2}{\omega_m - k_2}$$

Relative detuning

$$D' = \left| \frac{\omega_m - k_1}{\alpha_1} + \frac{\alpha_2^2}{2\alpha_1(\omega_m - k_2)} \right|$$





INTERFERENCES OF RABI OSCILLATIONS

$$D' = \left| \frac{\omega_m - k_1}{\alpha_1} + \frac{\alpha_2^2}{2\alpha_1(\omega_m - k_2)} \right|$$

pt

Two driving frequencies k_1 , and k_2 , with amplitude α_1 , and α_2

Destruction effect: $k_1 = \omega_m$, $|\alpha_2| \gg \sqrt{2\omega_m|\alpha_1(k_2 - \omega_m)|} \equiv \alpha_{2,c}$




200
Resonance Destruction

Resonance Survival

250

INTERFERENCES OF RABI OSCILLATIONS




Grid lines: amplitude predicted using $1/(1 + D'^2)$

α_2, k_1 values			
Dashed	dotted	dash-dotted	solid
$10^{-2} \omega_m, 10 \omega_m$	$10^{-2} \omega_m, 10^{-1} \omega_m$	$5.0 \times 10^{-2} \omega_m, 10 \omega_m$	$5 \times 10^{-2} \omega_m, 10^{-1} \omega_m$

RABI FORMULA WORKS

$$\vec{H} = \begin{pmatrix} 0 & \\ 0 & \end{pmatrix} + \frac{A \sin 2\theta_m}{2} \begin{pmatrix} \cos(kx) & \\ -\sin(kx) & 0 \end{pmatrix} + \frac{A \sin 2\theta_m}{2} \begin{pmatrix} \cos(-kx) & \\ -\sin(-kx) & 0 \end{pmatrix}$$



Lines: Rabi formula

Dots, diamonds, triangles, and squares are for $k = \omega_m$, $k = (1 - 2 \times 10^{-5})\omega_m$, and $k = (1 - 10^{-4})\omega_m$ respectively.

$$\alpha_{2,C} = 2\alpha = 2 \times \frac{A \sin 2\theta_m}{2}$$

OVERVIEW

pt

100

150

200

250

300

350

Introduction

Matter Effect

100

Stimulated Neutrino Flavor Transitions

Jan 150
Basis and Formalism

Summary

200

250

INTERFERENCES OF RABI OSCILLATIONS

pt

100

150

200

250

300

350

We have been making approximations.

100

$$\mathbf{H} = \frac{1}{2} \left(-\omega_m + \cos 2\theta_m A \cos(kx) \right) \sigma_3 - \frac{\sin 2\theta_m}{2} A \cos(kx) \sigma_1$$

150

$$\rightarrow -\frac{\omega_m}{2} \sigma_3 - \frac{A \sin 2\theta_m}{2} \cos(kx) \sigma_1$$

We need a better basis.

200

250

RABI BASIS

Hamiltonian in Background Matter Basis

pt

$$\mathbf{H} = \frac{1}{2} (-\omega_m + \delta\lambda(x) \cos 2\theta_m) \sigma_3 - \frac{\delta\lambda(x)}{2} \sin \theta_m \sigma_1.$$

100

A Better Basis

Define Rabi basis $\{|\tilde{\nu}_L\rangle, |\tilde{\nu}_H\rangle\}$ is related to background matter basis $\{|\nu_L\rangle, |\nu_H\rangle\}$ through

150

$$\begin{pmatrix} |\nu_L\rangle \\ |\nu_H\rangle \end{pmatrix} = \begin{pmatrix} e^{-i\eta(x)} & 0 \\ 0 & e^{i\eta(x)} \end{pmatrix} \begin{pmatrix} |\tilde{\nu}_L\rangle \\ |\tilde{\nu}_H\rangle \end{pmatrix},$$

where

200

$$\eta(x) - \eta(0) = \frac{\cos 2\theta_m}{2} \int_0^x \delta\lambda(\tau) d\tau.$$

250

SINGLE FREQUENCY MATTER PROFILE

Matter profile

pt

$$\lambda = \lambda_0 + \frac{2\pi}{k} \cos(kx), \quad 100 \quad 150 \quad 200 \quad 250 \quad 300 \quad 350$$

Hamiltonian in new basis

$$\tilde{H} = -\frac{\omega_m}{2} \sigma_3 - \frac{\delta \lambda(x)}{2} \sin 2\theta_m \begin{pmatrix} 0 & e^{2i\eta(x)} \\ e^{-2i\eta(x)} & 0 \end{pmatrix} = -\frac{\omega_m}{2} \sigma_3 + \begin{pmatrix} 0 & h \\ h^* & 0 \end{pmatrix}$$

Hamiltonian in New Basis

150

$$h \equiv -\frac{\delta \lambda(x)}{2} e^{2i\eta(x)}$$

200

$$= \frac{i}{4} \left[\exp \left(i(k + \omega_m)x + i \cos 2\theta_m \frac{A}{k} \cos(kx) \right) - \exp \left(i(-k + \omega_m)x + i \cos 2\theta_m \frac{A}{k} \cos(kx) \right) \right]$$

250

SINGLE FREQUENCY MATTER PROFILE

Off-diagonal Term in Our System

$$\tilde{H} = -\frac{\omega_m}{2}\sigma_3 + \begin{pmatrix} 0 & h \\ h^* & 0 \end{pmatrix}$$

$$h \propto \left[\exp \left(i(k + \omega_m)x + i \cos 2\theta_m \frac{A}{k} \cos(kx) \right) - \exp \left(i(-k + \omega_m)x + i \cos 2\theta_m \frac{A}{k} \cos(kx) \right) \right]$$

Jacobi-Anger expansion

$$e^{i\beta \cos(kx)} = \sum_{n=-\infty}^{\infty} i^n J_n(\beta) e^{inkx},$$

where $J_n(\beta)$ are Bessel's functions of the first kind.

SINGLE FREQUENCY MATTER PROFILE

pt

100

150

200

250

300

350

Scaled Quantities

100 Characteristic scale: ω_m

► $\hat{A} = A/\omega_m$

► $\hat{k} = k/\omega_m$

150 $\hat{x} = \omega_m x$

► $\hat{h} = h/\omega_m$

200

250

SINGLE FREQUENCY MATTER PROFILE

pt

100

150

200

250

300

350

Rotation Wave Approximation

100 off-diagonal element of Hamiltonian

150

$$\tilde{\mathbf{H}} = \sum_{n=-\infty}^{\infty} \begin{pmatrix} 0 & \frac{1}{2} \hat{B}_n e^{i(n\hat{k}-1)\hat{x}} \\ \frac{1}{2} \hat{B}_n^* e^{-i(n\hat{k}-1)\hat{x}} & 0 \end{pmatrix}$$

where $\hat{B}_n = -(-i)^n n \hat{k} \tan 2\theta_m J_n(\hat{A} \cos 2\theta_m / \hat{k})$.

200

250

SINGLE FREQUENCY MATTER PROFILE

Transition Probability

pt

100

150

200

250

300

350

$$P_{L \rightarrow H}^{(n)} = \frac{|\hat{B}_n|^2}{|\hat{B}_n|^2 + (n\hat{k} - 1)^2} \sin^2 \left(\frac{q^{(n)}}{2} x \right),$$

100

where

$$150 \quad q^{(n)} = \sqrt{|\Gamma^{(n)}|^2 + (n\hat{k} - 1)^2}, \quad \text{frequency of oscillations}$$


$$\Gamma^{(n)} = |\hat{B}_n|, \quad \text{width of resonance } (n\hat{k} \text{ as parameter})$$

Re 200 nance conditions

$$\hat{k} \sim \frac{1}{n}$$


250

SINGLE FREQUENCY MATTER PROFILE



Resonances of different $n = 1/\hat{k}$. Width becomes extremely narrow for high n .

SINGLE FREQUENCY MATTER PROFILE



Resonances of different $n = 1/\hat{k}$. Width becomes extremely narrow for high order.

SINGLE FREQUENCY MATTER PROFILE REVISITED

pt

100

150

200

250

300

350

$$k_1 = \omega_m$$

	D	D'_1	$2\pi\omega_m/\Omega_n$
100			
1	0	-	3.2×10^5
-1	10^5	4.8×10^{-6}	3.1
150			
-	1.1×10^9	2.1×10^{-14}	6.3
-2	3.4×10^9	6.9×10^{-15}	2.1

200

250

SINGLE FREQUENCY MATTER PROFILE REVISITED

pt

100

150

200

250

300

350

$$k_1 = (1 - 2 \times 10^{-5})\omega_m$$

100	n	D	D'_1	$2\pi\omega_m/\Omega_n$
	1	1	-	2.2×10^5
	-1	10^5	1	3.1
150	2	1.1×10^9	1	6.3
	-2	3.4×10^9	1	2.1

200

250

SINGLE FREQUENCY MATTER PROFILE REVISITED

pt

100

150

200

250

300

350


$$k_1 = (1 - 10^{-4})\omega_m$$

z	D	D'_1	$2\pi\omega_m/\Omega_n$
1	5.2	-	6.2×10^4
-1	10^5	5.2	3.1
2	1.1×10^9	5.2	6.3
-2	3.4×10^9	5.2	2.1

200

250

CASTLE WALL MATTER PROFILE



CASTLE WALL MATTER PROFILE

pt

100

150

200


250

300

350

Table: Relative detuning of each
frequ 100

$\{n_1, n_2\}$	D	$D'_{\{1,0\}}$
$\{1, 0\}$	0	-
$\{-1, 0\}$	48	1.0×10^{-2}
$\{0, 1\}$	1.5×10^2	1.1×10^{-3}
$\{2, 0\}$	2.4×10^2	2.0×10^{-4}



200

250

OVERVIEW

pt

100

150

200

250

300

350

Introduction

Mä 100 Effect

Stimulated Neutrino Flavor Transitions

Ja 150 Anger Expansion

Summary

200

250

SUMMARY

pt

100

150

200

250

300

350

- ▶ The fact that neutrino flavor states are not mass states causes vacuum oscillations.
- ▶ MSW resonance happens when matter potential cancels out the vacuum diagonal elements of the Hamiltonian.
- ▶ Even matter profile doesn't match MSW requirement, variation in matter profile can cause resonances.
- ▶ Single frequency perturbations in matter profile is a combination of many Rabi oscillations.
- ▶ Rabi oscillations with two driving fields of different frequencies: large width to destroy the resonance.

200

250

BACKUP SLIDES

pt

100

150

200

250

300

350

100

BACKUP SLIDES

150

200

250

RABI OSCILLATIONS

pt

100

150

200

250

300

350

The coupling strength is calculated as

100

$$\alpha = \langle 1 | \mathbf{d} \cdot \mathbf{E} | 2 \rangle$$

where the electric field is

150

$$\mathbf{E} = \mathbf{E}_0 \sin(kt).$$

and \mathbf{d} is the dipole moment.

200

250

INTERFERENCES OF RABI OSCILLATIONS

$$-\frac{\omega_0}{2}\sigma_3 - \frac{\alpha}{2} \begin{pmatrix} 0 & e^{ikt} \\ e^{-ikt} & 0 \end{pmatrix}$$

pt

100

150

200

250

300

350

$$H = \frac{1}{2} \left(-\omega_m + \cos 2\theta_m A \cos(kx) \right) \sigma_3 - \frac{\sin 2\theta_m}{2} A \cos(kx) \sigma_1$$

$$\rightarrow -\frac{\omega_m}{2} \sigma_3 - \frac{A \sin 2\theta_m}{2} \cos(kx) \sigma_1$$

$$= \frac{\omega_m}{2} \sigma_3 - \frac{A \sin 2\theta_m}{2} \frac{1}{2} \begin{pmatrix} 0 & e^{ikx} \\ e^{-ikx} & 0 \end{pmatrix} - \frac{A \sin 2\theta_m}{2} \frac{1}{2} \begin{pmatrix} 0 & e^{i(-k)x} \\ e^{-i(-k)x} & 0 \end{pmatrix}$$

200

250

PARAMETERS USED FOR VACUUM OSCILLATIONS

pt

100

150

200

250

300

350

100

$$\theta_{12} = 33.36/180\pi; \theta_{13} = 8.66/180\pi; \theta_{23} = 40/180 * \pi; \delta_{cp} = 0;$$

$$m_1^2 = 0.01; m_2^2 = m_1^2 + 0.000079; E = 1\text{MeV}$$

150

200

250

SINGLE FREQUENCY MATTER PROFILE

pt

100

150

200

250

300

350

Why Does It Work?

100

$$J_n(n \operatorname{sech} \alpha) \sim \frac{e^{-n(\alpha - \tanh \alpha)}}{\sqrt{2\pi n \tanh \alpha}}, \quad \text{for large } n$$

\Rightarrow

150

$$\Gamma \propto \hat{B}_n \propto \frac{e^{-n(\alpha - \tanh \alpha)}}{\sqrt{2\pi n \tanh \alpha}}$$

Small perturbation \Rightarrow Small $\hat{A} \Rightarrow$ Large $\alpha \Rightarrow$ Drops fast at large n .

200

250

TWO-FREQUENCY MATTER PROFILE

pt

100

150

200

250

300

350

Matter Profile

100

$$\lambda(x) = \lambda_0 + \delta\lambda(x), \quad \delta\lambda(x) = A_1 \sin(k_1 x) + A_2 \sin(k_2 x).$$

150

200

250

TWO-FREQUENCY MATTER PROFILE $\hat{h} = \sum_{n=-\infty}^{\infty} \frac{1}{2} \hat{B}_n e^{i(n\hat{k}-1)\hat{x}}$,

Hamiltonian Off-diagonal Element

pt

Apply Jacobi-Anger expansion,

$$\hat{h} = \sum_{n_1=-\infty}^{\infty} \sum_{n_2=-\infty}^{\infty} \frac{1}{2} \hat{B}_{n_1, n_2}(\hat{k}_1, \hat{k}_2) e^{i(n_1 \hat{k}_1 + n_2 \hat{k}_2 - 1)\hat{x}},$$

where

$$\begin{aligned} & \hat{B}_{n_1, n_2}(\hat{k}_1, \hat{k}_2) \\ &= -(-i)^{n_1+n_2} (n_1 \hat{k}_1 + n_2 \hat{k}_2) J_{n_1} \left(\frac{\hat{A}_1 \cos 2\theta_m}{\hat{k}_1} \right) J_{n_2} \left(\frac{\hat{A}_2 \cos 2\theta_m}{\hat{k}_2} \right) \end{aligned}$$

Which terms are important?

TWO-FREQUENCY MATTER PROFILE

pt

100

150

200

250

300

350

Resonance Lines

100

There are still resonances, i.e., (almost) zero phases, on lines

$$n_{1,0}\hat{k}_1 + n_{2,0}\hat{k}_2 - 1 = 0$$


150

$\{\hat{k}_1, \hat{k}_2\}$ plane. \Rightarrow Resonance width for each point on resonance lines.

200


250

TWO-FREQUENCY MATTER PROFILE $\hat{h} = \sum_{n_1} \sum_{n_2} \frac{1}{2} \hat{B}_{n_1, n_2} (\hat{k}_1, \hat{k}_2) e^{i(n_1 \hat{k}_1 + n_2 \hat{k}_2 - 1) \hat{x}}$



Density plot of transition amplitudes calculated using only one term out of the whole summation in Hamiltonian. $n_1, n_2 \in [-2, 2]$

TWO-FREQUENCY MATTER PROFILE



Resonance line, distance to resonance, and width

TWO-FREQUENCY MATTER PROFILE

Width

pt

100

150

200

250

300

350

$$\Gamma_2 = \frac{\hat{B}_{n_1, n_2}(\hat{k}_{1, \text{intercept}}, \hat{k}_{2, \text{intercept}})}{\sqrt{n_1^2 + n_2^2}}.$$

100

Distance to Resonance Line

150

$$d = \frac{|n_1 \hat{k}_{10} + n_2 \hat{k}_{20} - 1|}{\sqrt{n_1^2 + n_2^2}}.$$


Distance to Resonance Width Ratio

200


$$Q_2 = \frac{d}{\Gamma_2}.$$

250


TWO-FREQUENCY MATTER PROFILE




TWO-FREQUENCY MATTER PROFILE




TWO-FREQUENCY MATTER PROFILE




TWO-FREQUENCY MATTER PROFILE



TWO-FREQUENCY MATTER PROFILE



TWO-FREQUENCY MATTER PROFILE



BESSEL'S FUNCTION

pt

100

150

200

250

300

350

100

$$J_n(\beta) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m! \Gamma(m+n+1)} \left(\frac{\beta}{2}\right)^{2m+n}$$

150

200

250

REFERENCES I

pt

100

150

200

250

300

350

100

150

200

250