

Student Information

Name : Emre

ID : 2521581

Answer 1

a) Write a context-free grammar for the language $L_1 = \{w | w \in \{a, b\}^* \wedge w \text{ has twice as many b's as a's}\}$.

Let G be the grammar (V, Σ, R, S) for the language L_1 where

$$\begin{aligned} V &= \{S, a, b\}, \\ \Sigma &= \{a, b\}, \\ R &= \{S \rightarrow aSbSb, \\ &\quad S \rightarrow bSaSb, \\ &\quad S \rightarrow bSbSa, \\ &\quad S \rightarrow e\}. \end{aligned}$$

b) Write a context-free grammar for the language $L_2 = \{a^n b^m | m, n \in \mathbb{N} \wedge m \leq n \leq 2m\}$.

Let G be the grammar (V, Σ, R, S) for the language L_2 where

$$\begin{aligned} V &= \{S, a, b\}, \\ \Sigma &= \{a, b\}, \\ R &= \{S \rightarrow aSbSa, \\ &\quad S \rightarrow aSaSb, \\ &\quad S \rightarrow bSaSa, \\ &\quad S \rightarrow bSa, \\ &\quad S \rightarrow aSb, \\ &\quad S \rightarrow e\}. \end{aligned}$$

c) Formally define and draw a PDA that accepts L_1 .

Let $(K, \Sigma, V, \Delta, p, \{q\})$ be the sextuple for the language L_1 where K, Σ, V, R are defined on part a.

Δ contains the following transitions:

- 1) $((p, e, e), (q, S))$
- 2) $((p, e, A), (q, x))$ for each rule $A \rightarrow x$ in R
- 3) $((q, a, a), (q, e))$ for each $a \in \Sigma$

Then,

$$\begin{aligned} \Delta = \{ & ((p, e, e), (q, S)), \\ & ((p, e, S), (q, aSbSa)), \\ & ((p, e, S), (q, bSaSa)), \\ & ((p, e, S), (q, aSaSb)), \\ & ((p, e, S), (q, e)), \\ & ((p, a, a), (q, e)), \\ & ((p, b, b), (q, e)) \}. \end{aligned}$$

d) Write a context-free grammar for the language $L_3 = L_1 \cup L_2$. Let $G_1 = (V_1, \Sigma_1, R_1, S_1)$ for the language L_1 and $G_2 = (V_2, \Sigma_2, R_2, S_2)$ for the language L_2 .

Then $G_3 = (V_1 \cup V_2 \cup \{S\}, \Sigma_1 \cup \Sigma_2, R_1 \cup R_2 \cup \{S \rightarrow S_1, S \rightarrow S_2\})$ for the language L_3 .

$$\begin{aligned}
V_3 &= \{S, S_1, S_2, a, b\} \\
\Sigma &= \{a, b\} \\
R &= \{S \rightarrow S_1, \\
&\quad S \rightarrow S_2, \\
&\quad S_1 \rightarrow aS_1bS_1b \\
&\quad S_1 \rightarrow bS_1aS_1b \\
&\quad S_1 \rightarrow bS_1bS_1a \\
&\quad S_1 \rightarrow e \\
&\quad S_2 \rightarrow aS_2bS_2a \\
&\quad S_2 \rightarrow aS_2aS_2b \\
&\quad S_2 \rightarrow bS_2aS_2a \\
&\quad S_2 \rightarrow bS_2a \\
&\quad S_2 \rightarrow aS_2b \\
&\quad S_2 \rightarrow e \\
&\quad \}.
\end{aligned}$$

Answer 2

a)

b)

Answer 3

a)

b)