Centre No.					Pape	r Refer	ence			Surname	Initial(s)
Candidate No.			6	6	6	8	/	0	1	Signature	

Paner Reference(s)

6668/01

Edexcel GCE

Further Pure Mathematics FP2 Advanced/Advanced Subsidiary

Friday 21 June 2013 – Morning

Time: 1 hour 30 minutes

ľ	Ma	terials	rec	uired	for	examination
-						

Mathematical Formulae (Pink)

Items included with question papers

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer for each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 9 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy. ©2013 Pearson Education Ltd.

W850/R6668/57570 5/5/5/5/



Turn over

Total



Examiner's use only Team Leader's use only

$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} + x \frac{\mathrm{d}y}{\mathrm{d}x} = 2\cos x$$

(a) Find $\frac{d^3y}{dx^3}$ in terms of x, $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$.

At x = 0, y = 1 and $\frac{dy}{dx} = 3$

(b) Find the value of $\frac{d^3y}{dx^3}$ at x = 0

(1)

(c) Express y as a series in ascending powers of x, up to and including the term in x^3 .

(3)

(3)

Question 1 continued	



Leave blank

- **2.** (a) Sketch, on the same axes,
 - $(i) \quad y = |2x 3|$
 - (ii) $y = 4 x^2$

(3)

(b) Find the set of values of x for which

$$4 - x^2 > \left| 2x - 3 \right|$$

(6)

estion 2 continued	



Leave blank

3.

$$f(x) = \ln(1 + \sin kx)$$

where k is a constant, $x \in \mathbb{R}$ and $-\frac{\pi}{2} < kx < \frac{3\pi}{2}$

(a) Find f'(x)

(2)

(b) Show that $f''(x) = \frac{-k^2}{1 + \sin kx}$

(3)

(c) Find the Maclaurin series of f(x), in ascending powers of x, up to and including the term in x^3 .

(4)

estion 3 continued		



estion 3 continued	

Question 3 continued	Lea blar
Question 5 continued	
	Q3
(Total 9 marks)	



4.	Find the general solution of the differential equation

$$x\frac{\mathrm{d}y}{\mathrm{d}x} + (1 + x\cot x)y = \sin x, \qquad 0 < x < \pi$$

giving your answer in the form $y = f(x)$.	

(9)

	Leave blank
Question 4 continued	



estion 4 continued		
		-
		-
		-

Question 4 continued	Leav blanl
	Q4
(Total 9 marks)	



Leave	
blank	

5. (a) Express $\frac{2}{r(r+1)(r+2)}$ in partial fractions.

(3)

(b) Using your answer to part (a) and the method of differences, show that

$$\sum_{r=1}^{n} \frac{2}{r(r+1)(r+2)} = \frac{n(n+3)}{2(n+1)(n+2)}$$
(4)

	I
Question 5 continued	l t



estion 5 continued		

Question 5 continued	Le bla
	Q5



6.	Solve the equation $z^5 = -16\sqrt{3} + 16i$	
	giving your answers in the form $r(\cos \theta + i \sin \theta)$, where $r > 0$ and $-\pi < \theta < \pi$.	(8

nestion 6 continued	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	-
	_
	_
	_
	_
	_
	-
	_
	_



Leave	
blank	

7. (a) Find the value of the constant λ for which $y = \lambda x e^{2x}$ is a particular integral of the differential equation

$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} - 4y = 6\mathrm{e}^{2x}$$

(4)

(b) Hence, or otherwise, find the general solution of the differential equation

$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} - 4y = 6\mathrm{e}^{2x}$$

(3)

nestion 7 continued	

Leave blank

- **8.** A complex number z is represented by the point P on an Argand diagram.
 - (a) Given that |z| = 1, sketch the locus of P.

(1)

The transformation T from the z-plane to the w-plane is given by

$$w = \frac{z + 7i}{z - 2i}$$

(b) Show that T maps |z| = 1 onto a circle in the w-plane.

(5)

(c) Show that this circle has its centre at w = -5 and find its radius.

(2)

	Leave blank
Question 8 continued	Dialik



estion 8 continued		

stion 8 continued	

9.




Figure 1

Figure 1 shows a sketch of the curves given by the polar equations

$$r = 1$$
 and $r = 2 - 2 \sin \theta$

(a) Find the coordinates of the points where the curves intersect.

(3)

Leave blank

The region S, between the curves, for which r < 1 and for which $r < 2 - 2 \sin \theta$, is shown shaded in Figure 1.

(b) Find, by integration, the area of the shaded region S, giving your answer in the form $a\pi + b\sqrt{3}$, where a and b are rational numbers.

•	\mathbf{o}	
	XΙ	
	\mathbf{v}	

	Leave
Question 9 continued	Otalik

