


Week 4: *Centrality & Variability*

 EMSE 4575: Exploratory Data Analysis

 John Paul Helveston

 September 21, 2022

Quiz solution

Tip of the week:
`theme_set()`


Add "global" settings to all plots

```
library(knitr)
library(tidyverse)
library(here)
knitr::opts_chunk$set(
  warning = FALSE,
  message = FALSE,
  comment = "#>",
  fig.path = "figs/", # Plot save path
  fig.width = 7.252, # Plot dimensions
  fig.height = 4,
  fig.retina = 3 # Better plot resolution
)


theme_set(theme_bw(base_size = 20)) # Set theme for all ggplots
```

```
ggplot(mtcars) +  
  geom_point(aes(x = mpg, y = hp))
```

Default theme



theme_bw(base_size = 20)



Week 4: *Centrality & Variability*

1. Data Types

2. Measures of Centrality & Variability

BREAK

3. Visualizing Centrality & Variability

4. Relationships Between 2 Variables

5. Exploratory Data Analysis

Week 4: *Centrality & Variability*

1. Data Types

2. Measures of Centrality & Variability

BREAK

3. Visualizing Centrality & Variability

4. Relationships Between 2 Variables

5. Exploratory Data Analysis

24,901

Earth's circumference at the equator:
24,901 miles

Types of Data

Categorical

Subdivide things into *groups*

- What type?
- Which category?

Numerical

Measure things with numbers

- How many?
- How much?

Categorical (discrete) variables

Nominal

- Order doesn't matter
- Differ in "name" (nominal) only

e.g. **country** in TB case data:

```
#> # A tibble: 6 × 4
#>   country      year  cases population
#>   <chr>      <dbl> <dbl>      <dbl>
#> 1 Afghanistan 1999     745   19987071
#> 2 Afghanistan 2000    2666   20595360
#> 3 Brazil      1999   37737   172006362
#> 4 Brazil      2000   80488   174504898
#> 5 China       1999  212258  1272915272
#> 6 China       2000  213766  1280428583
```

Ordinal

- Order matters
- Distance between units not equal

e.g.: **Placement** 2017 Boston marathon:

```
#> # A tibble: 6 × 3
#>   Placement `Official Time` Name
#>   <dbl> <time>      <chr>
#> 1         1 02:09:37 Kirui, Geo
#> 2         2 02:09:58 Rupp, Gale
#> 3         3 02:10:28 Osako, Sug
#> 4         4 02:12:08 Biwott, Sh
#> 5         5 02:12:35 Chebet, Wi
#> 6         6 02:12:45 Abdirahman
```

Numerical data

Interval

- Numerical scale with arbitrary starting point
- No "0" point
- Can't say "x" is double "y"

e.g.: **temp** in Beaver data

```
#>   day time  temp activ
#> 1 346  840 36.33     0
#> 2 346  850 36.34     0
#> 3 346  900 36.35     0
#> 4 346  910 36.42     0
#> 5 346  920 36.55     0
#> 6 346  930 36.69     0
```

Ratio

- Has a "0" point
- Can be described as percentages
- Can say "x" is double "y"

e.g.: **height** & **speed** in wildlife impacts

```
#> # A tibble: 6 × 3
#>   incident_date      height speed
#>   <dtm>          <dbl> <dbl>
#> 1 2018-12-31 00:00:00    700   200
#> 2 2018-12-27 00:00:00    600   145
#> 3 2018-12-23 00:00:00     0   130
#> 4 2018-12-22 00:00:00    500   160
#> 5 2018-12-21 00:00:00    100   150
#> 6 2018-12-18 00:00:00   4500   250
```

Key Questions

Categorical

Numerical

Does the order matter?

Is there a "baseline"?

Yes	No
Ordinal	Nominal

Yes	No
Ratio	Interval

Be careful of how variables are encoded!

When numbers are categories

- "Dummy coding": e.g., `passedTest = 1` or `0`)
- "North", "South", "East", "West" = 1, 2, 3, 4

When ratio data are discrete (i.e. counts)

- Number of eggs in a carton, heart beats per minute, etc.
- Continuous variables measured discretely (e.g. age)

Time

- As *ordinal* categories: "Jan.", "Feb.", "Mar.", etc.
- As *interval* scale: "Jan. 1", "Jan. 2", "Jan. 3", etc.
- As *ratio* scale: "30 sec", "60 sec", "70 sec", etc.

Quick practice: What's the data type?

Decide [here](#) (link also in #classroom)

```
wildlife_impacts %>%  
  filter(!is.na(cost_repairs_infl_adj)) %>%  
  select(incident_date, time_of_day, species, cost_repairs_infl_adj) %>%  
  head()
```

```
#> # A tibble: 6 × 4  
#>   incident_date    time_of_day species      cost_repairs_infl_adj  
#>   <dtm>          <chr>      <chr>          <dbl>  
#> 1 2018-10-25 00:00:00 Day      Unknown bird - large      1000  
#> 2 2018-09-05 00:00:00 <NA>     Unknown bird - medium     200  
#> 3 2018-08-09 00:00:00 Day      Semipalmated sandpiper  10000  
#> 4 2018-06-24 00:00:00 Day      Unknown bird - large  100000  
#> 5 2018-02-18 00:00:00 Day      Rough-legged hawk      20000  
#> 6 2018-01-05 00:00:00 Night     Brant                  487000
```


Week 4: *Centrality & Variability*

1. Data Types

2. Measures of Centrality & Variability

BREAK

3. Visualizing Centrality & Variability

4. Relationships Between 2 Variables

5. Exploratory Data Analysis

Summary Measures:

This week: **Centrality** & **Variability**

Next week: **Correlation**

Centrality (a.k.a. The "Average" Value)

A single number representing the *middle* of a set of numbers

Mean: $\frac{\text{Sum of values}}{\# \text{ of values}}$

Median: "Middle" value (50% of data above & below)

Mode: Most frequent value (usually for categorical data)


Mean isn't always the "best" choice

```
wildlife_impacts %>%  
  filter(! is.na(height)) %>%  
  summarise(  
    mean = mean(height),  
    median = median(height))
```

```
#> # A tibble: 1 × 2  
#>   mean median  
#>   <dbl> <dbl>  
#> 1  984.    50
```

Percent of data below mean:


```
#> [1] "73.9%"
```



Beware the "flaw of averages"

What happened to the statistician
that crossed a river with an average
depth of 3 feet?

...he drowned



Variability ("Spread")

Standard deviation: distribution of values relative to the mean

$$s = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N-1}}$$

Interquartile range (IQR): $Q_3 - Q_1$ (middle 50% of data)

Range: max - min

Example: Days to ship

Complaints are coming in about orders shipped from warehouse B, so you collect some data:

daysToShip

```
#>   order warehouseA warehouseB
#> 1      1          3          1
#> 2      2          3          1
#> 3      3          3          1
#> 4      4          4          3
#> 5      5          4          3
#> 6      6          4          4
#> 7      7          5          5
#> 8      8          5          5
#> 9      9          5          5
#> 10    10          5          6
#> 11    11          5          7
#> 12    12          5         10
```

Here, **averages** are misleading:

```
daysToShip %>%
  gather(warehouse, days, warehouseA:warehouseB) %>%
  group_by(warehouse) %>%
  summarise(
    mean    = mean(days),
    median  = median(days))
```

```
#> # A tibble: 2 × 3
#>   warehouse    mean median
#>   <chr>      <dbl>  <dbl>
#> 1 warehouseA  4.25    4.5
#> 2 warehouseB  4.25    4.5
```

Example: Days to ship

Complaints are coming in about orders shipped from warehouse B, so you collect some data:

daysToShip


```
#>   order warehouseA warehouseB
#> 1      1          3          1
#> 2      2          3          1
#> 3      3          3          1
#> 4      4          4          3
#> 5      5          4          3
#> 6      6          4          4
#> 7      7          5          5
#> 8      8          5          5
#> 9      9          5          5
#> 10    10          5          6
#> 11    11          5          7
#> 12    12          5         10
```

Variability reveals difference in days to ship:

```
daysToShip %>%
  gather(warehouse, days, warehouseA:warehouseB) %>%
  group_by(warehouse) %>%
  summarise(
    mean    = mean(days),
    median  = median(days),
    range   = max(days) - min(days),
    sd      = sd(days))
```



```
#> # A tibble: 2 × 5
#>   warehouse    mean median  range    sd
#>   <chr>      <dbl>  <dbl> <dbl> <dbl>
#> 1 warehouseA  4.25    4.5    2  0.866
#> 2 warehouseB  4.25    4.5    9  2.70
```


Example: Days to ship



Interpreting the standard deviation

$$s = \sqrt{\frac{\sum_{i=1}^N (x_i - \bar{x})^2}{N-1}}$$



Outliers



Mean & Standard Deviation are sensitive to outliers

Outliers: $Q_1 - 1.5IQR$ or $Q_3 + 1.5IQR$

Extreme values: $Q_1 - 3IQR$ or $Q_3 + 3IQR$

```
data1 <- c(3,3,4,5,5,6,6,7,8,9)
```

- Mean: 5.6
- Standard Deviation: 2.01
- Median: 5.5
- IQR: 2.5

```
data2 <- c(3,3,4,5,5,6,6,7,8,20)
```

- Mean: 6.7
- Standard Deviation: 4.95
- Median: 5.5
- IQR: 2.5

Robust statistics for continuous data (less sensitive to outliers)

Centrality: Use *median* rather than *mean*

Variability: Use *IQR* rather than *standard deviation*

Practice with summary measurements

1) Read in the following data sets:

- `milk_production.csv`
- `lotr_words.csv`

2) For each variable in each data set, if possible, summarize its

1. **Centrality**

2. **Variability**

Break!

Stand up, Move around, Stretch!

05 : 00

Week 4: *Centrality & Variability*

1. Data Types

2. Measures of Centrality & Variability

BREAK

3. Visualizing Centrality & Variability


4. Relationships Between 2 Variables

5. Exploratory Data Analysis

"Visualizing data helps us think"

	A		B		C		D	
	x	y	x	y	x	y	x	y
	10	8.04	10	9.14	10	7.46	8	6.58
	8	6.95	8	8.14	8	6.77	8	5.76
	13	7.58	13	8.74	13	12.74	8	7.71
	9	8.81	9	8.77	9	7.11	8	8.84
	11	8.33	11	9.26	11	7.81	8	8.47
	14	9.96	14	8.1	14	8.84	8	7.04
	6	7.24	6	6.13	6	6.08	8	5.25
	4	4.26	4	3.1	4	5.39	19	12.5
	12	10.84	12	9.13	12	8.15	8	5.56
	7	4.82	7	7.26	7	6.42	8	7.91
	5	5.68	5	4.74	5	5.73	8	6.89
Sum:	99	82.51	99	82.51	99	82.5	99	82.51
Mean:	9	7.5	9	7.5	9	7.5	9	7.5
St. Dev:	3.3	2	3.3	2	3.3	2	3.3	2

Anscombe's Quartet



The data *type* determines
how to summarize it

Nominal (Categorical)

Measures:

- Frequency counts / Proportions

Charts:

- Bars

Ordinal (Categorical)

Measures:

- Frequency counts / Proportions
- **Centrality:** Median, Mode
- **Variability:** IQR

Charts:

- Bars

Numerical (Continuous)

Measures:

- **Centrality:** Mean, median
- **Variability:** Range, standard deviation, IQR

Charts:

- Histogram
- Boxplot


Summarizing **Nominal** data

Summarize with counts / percentages Visualize with bars

```
wildlife_impacts %>%  
  count(operator, sort = TRUE) %>%  
  mutate(p = n / sum(n))
```

```
#> # A tibble: 4 × 3  
#>   operator          n      p  
#>   <chr>        <int> <dbl>  
#> 1 SOUTHWEST AIRLINES 17970 0.315  
#> 2 UNITED AIRLINES   15116 0.265  
#> 3 AMERICAN AIRLINES 14887 0.261  
#> 4 DELTA AIR LINES   9005 0.158
```

```
wildlife_impacts %>%  
  count(operator, sort = TRUE) %>%  
  ggplot() +  
    geom_col(aes(x = n, y = reorder(operator, n)),  
             width = 0.7) +  
    labs(x = "Count", y = "Operator")
```



Summarizing **Ordinal** data


Summarize: Counts / percentages

```
wildlife_impacts %>%  
  count(incident_month, sort = TRUE) %>%  
  mutate(p = n / sum(n))
```

```
#> # A tibble: 12 × 3  
#>   incident_month     n     p  
#>   <dbl> <int> <dbl>  
#> 1         9 7980 0.140  
#> 2        10 7754 0.136  
#> 3         8 7104 0.125  
#> 4         5 6161 0.108  
#> 5         7 6133 0.108  
#> 6         6 4541 0.0797  
#> 7         4 4490 0.0788  
#> 8        11 4191 0.0736  
#> 9         3 2678 0.0470  
#> 10        12 2303 0.0404  
#> 11         1 1951 0.0342  
#> 12         2 1692 0.0297
```

Visualize: Bars

```
wildlife_impacts %>%  
  count(incident_month, sort = TRUE) %>%  
  ggplot() +  
    geom_col(aes(x = as.factor(incident_month),  
                 y = n), width = 0.7) +  
    labs(x = "Incident month")
```




Summarizing **continuous** variables

Histograms:

- Skewness
- Number of modes

Boxplots:

- Outliers
- Comparing variables



Histogram: Identify Skewness & # of Modes

Summarise:

Mean, median, sd, range, & IQR:


```
summary(wildlife_impacts$height)
```

```
#>      Min. 1st Qu.  Median    Mean  
#>      0.0     0.0    50.0   983.8
```

Visualize:

Histogram (identify skewness & modes)


```
ggplot(wildlife_impacts) +  
  geom_histogram(aes(x = height), bins = 50) +  
  labs(x = 'Height (ft)', y = 'Count')
```



Histogram: Identify Skewness & # of Modes


Height

```
ggplot(wildlife_impacts) +  
  geom_histogram(aes(x = height), bins = 50)  
  labs(x = 'Height (ft)', y = 'Count')
```



Speed


```
ggplot(wildlife_impacts) +  
  geom_histogram(aes(x = speed), bins = 50)  
  labs(x = 'speed (mph)', y = 'Count')
```



Boxplot: Identify outliers


Height

```
ggplot(wildlife_impacts) +  
  geom_boxplot(aes(x = height)) +  
  labs(x = 'Height (ft)', y = NULL)
```




Speed

```
ggplot(wildlife_impacts) +  
  geom_boxplot(aes(x = speed)) +  
  labs(x = 'Speed (mph)', y = NULL)
```




Histogram

- Skewness
- Modes



Boxplot

- Outliers



Practicing visual summaries

1) Read in the following data sets:

- `faithful.csv`
- `marathon.csv`

2) Summarize the following variables using an appropriate chart (bar chart, histogram, and / or boxplot):

- faithful: `eruptions`
- faithful: `waiting`
- marathon: `Age`
- marathon: `State`
- marathon: `Country`
- marathon: ``Official Time``

Week 4: *Centrality & Variability*

1. Data Types

2. Measures of Centrality & Variability

BREAK

3. Visualizing Centrality & Variability

4. Relationships Between 2 Variables

5. Exploratory Data Analysis

Two **Categorical** Variables

Summarize with a table of counts

```
wildlife_impacts %>%  
  count(operator, time_of_day)
```

```
#> # A tibble: 20 × 3  
#>   operator      time_of_day      n  
#>   <chr>      <chr>      <int>  
#> 1 AMERICAN AIRLINES Dawn        458  
#> 2 AMERICAN AIRLINES Day         7809  
#> 3 AMERICAN AIRLINES Dusk         584  
#> 4 AMERICAN AIRLINES Night        3710  
#> 5 AMERICAN AIRLINES <NA>        2326  
#> 6 DELTA AIR LINES Dawn         267  
#> 7 DELTA AIR LINES Day         4846  
#> 8 DELTA AIR LINES Dusk         353  
#> 9 DELTA AIR LINES Night        2090  
#> 10 DELTA AIR LINES <NA>        1449  
#> 11 SOUTHWEST AIRLINES Dawn         394  
#> 12 SOUTHWEST AIRLINES Day         9109
```

Two **Categorical** Variables

Convert to "wide" format with `pivot_wider()` to make it easier to compare values


```
wildlife_impacts %>%  
  count(operator, time_of_day) %>%  
  pivot_wider(names_from = time_of_day, values_from = n)
```

```
#> # A tibble: 4 × 6  
#>   operator      Dawn    Day   Dusk  Night  `NA`  
#>   <chr>      <int> <int> <int> <int> <int>  
#> 1 AMERICAN AIRLINES    458  7809   584  3710  2326  
#> 2 DELTA AIR LINES     267  4846   353  2090  1449  
#> 3 SOUTHWEST AIRLINES   394  9109   599  5425  2443  
#> 4 UNITED AIRLINES     151  3359   181  1510  9915
```

Two **Categorical** Variables

Visualize with bars:
map **fill** to denote 2nd categorical var


```
wildlife_impacts %>%  
  count(operator, time_of_day) %>%  
  ggplot() +  
  geom_col(  
    aes(  
      x = n,  
      y = reorder(operator, n),  
      fill = reorder(time_of_day, n)  
    ),  
    width = 0.7,  
    position = 'dodge') +  
  theme(legend.position = "bottom") +  
  labs(  
    fill = "Time of day",  
    y = "Airline"  
  )
```



Two **Continuous** Variables

Visualize with scatterplot - looking for *clustering* and/or *correlational* relationship


```
ggplot(wildlife_impacts) +  
  geom_point(  
    aes(  
      x = speed,  
      y = height  
    ),  
    size = 0.5) +  
  labs(  
    x = 'Speed (mph)',  
    y = 'Height (f)'  
  )
```



One **Continuous**, One **Categorical**

Visualize with **boxplot**

```
ggplot(wildlife_impacts) +  
  geom_boxplot(  
    aes(  
      x = speed,  
      y = operator)  
    ) +  
  labs(  
    x = 'Speed (mph)',  
    y = 'Airline'  
  )
```



Practice with visualizing *relationships*

1) Read in the following data sets:

- `marathon.csv`
- `wildlife_impacts.csv`

2) Visualize the *relationships* between the following variables using an appropriate chart (bar plots, scatterplots, and / or box plots):

- marathon: `Age` & `Official Time`
- marathon: `Country` & `Official Time`
- wildlife_impacts: `state` & `operator`

Week 4: *Centrality & Variability*

1. Data Types

2. Measures of Centrality & Variability

BREAK

3. Visualizing Centrality & Variability

4. Relationships Between 2 Variables

5. **Exploratory Data Analysis**

Exploratory Analysis

Goal: **Form** hypotheses.

Improves quality of **questions**.

(do this in THIS class)

Confirmatory Analysis

Goal: **Test** hypotheses.

Improves quality of **answers**.

(do this in your stats classes)


Don't be Icarus



"Far better an approximate answer to the *right* question, which is often vague, than an exact answer to the *wrong* question, which can always be made precise."

— John Tukey

**EDA is an iterative process to help you
understand your data and ask better questions**



Visualizing variation

Ask yourself:

- What type of **variation** occurs within my variables?
- What type of **covariation** occurs between my variables?

Check out [these guides](#)

Variation		Covariation	
		Categorical Y	Continuous Y
Categorical X	Categorical	Heatmap or Count	Boxplot
	Continuous	Boxplot (with coord_flip)	Scatterplot (many to one) line chart (one to one)

Practice doing EDA

- 1) Read in the `candy_rankings.csv` data sets
- 2) Preview the data, note the data types and what each variable is.
- 3) Visualize (at least) three *relationships* between two variables (guided by a question) using an appropriate chart:
 - Bar chart
 - Scatterplot
 - Boxplot

Start thinking about research questions

Writing a research question

Follow [these guidelines](#) - your question should be:

- **Clear:** your audience can easily understand its purpose without additional explanation.
- **Focused:** it is narrow enough that it can be addressed thoroughly with the data available and within the limits of the final project report.
- **Concise:** it is expressed in the fewest possible words.
- **Complex:** it is not answerable with a simple "yes" or "no," but rather requires synthesis and analysis of data.
- **Arguable:** its potential answers are open to debate rather than accepted facts (do others care about it?)

Writing a research question

Look at examples: See the [Example Projects Page](#) page

Start now!

Mini Project 1: Due next week (9/27)

Project Proposal: Due in two weeks (10/04)