



The Crouzeix-Raviart Finite Element Method for a Nonconforming Formulation of the Rudin-Osher-Fatemi Model Problem

Enrico Bergmann

Humboldt-Universität zu Berlin

June 16, 2021

Table of Contents

① Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

② Primal-Dual Iteration

③ Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Table of Contents

1 Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

2 Primal-Dual Iteration

3 Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Let U be an open subset of \mathbb{R}^d . A function $v \in L^1(U)$ is a function of bounded variation iff

$$|v|_{\text{BV}(U)} := \sup_{\substack{\phi \in C_c^1(U; \mathbb{R}^d) \\ \|\phi\|_{L^\infty(U)} \leq 1}} \int_U v \operatorname{div}(\phi) \, dx < \infty.$$

The space of all such functions is denoted by $\text{BV}(U)$.

Let U be an open subset of \mathbb{R}^d . A function $v \in L^1(U)$ is a function of bounded variation iff

$$|v|_{\text{BV}(U)} := \sup_{\substack{\phi \in C_c^1(U; \mathbb{R}^d) \\ \|\phi\|_{L^\infty(U)} \leq 1}} \int_U v \operatorname{div}(\phi) \, dx < \infty.$$

The space of all such functions is denoted by $\text{BV}(U)$. It is a Banach space equipped with the norm $\|\bullet\|_{\text{BV}(U)} := \|\bullet\|_{L^1(U)} + |\bullet|_{\text{BV}(U)}$.

Let U be an open subset of \mathbb{R}^d . A function $v \in L^1(U)$ is a function of bounded variation iff

$$|v|_{\text{BV}(U)} := \sup_{\substack{\phi \in C_c^1(U; \mathbb{R}^d) \\ \|\phi\|_{L^\infty(U)} \leq 1}} \int_U v \operatorname{div}(\phi) \, dx < \infty.$$

The space of all such functions is denoted by $\text{BV}(U)$. It is a Banach space equipped with the norm $\|\bullet\|_{\text{BV}(U)} := \|\bullet\|_{L^1(U)} + |\bullet|_{\text{BV}(U)}$.

We have $W^{1,1}(\Omega) \subset \text{BV}(\Omega)$ with $\|v\|_{\text{BV}(\Omega)} = \|v\|_{W^{1,1}(\Omega)}$ for all $v \in W^{1,1}(\Omega)$.

Hedy Attouch, Giuseppe Buttazzo, and Gérard Michaille.

Variational Analysis in Sobolev and BV Spaces. Applications to PDEs and Optimization. Second Edition. Vol. 17.

MOS-SIAM Series on Optimization. Philadelphia: Society for Industrial and Applied Mathematics, Mathematical Optimization Society, 2014. ISBN: 978-1-611973-47-1

Lawrence C. Evans and Ronald F. Gariepy. **Measure Theory and Fine Properties of Functions.** CRC Press, 1992. ISBN: 0-8493-7157-0

Table of Contents

1 Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

2 Primal-Dual Iteration

3 Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Let $\Omega \subset \mathbb{R}^2$ be a bounded polygonal Lipschitz domain.

Rudin-Osher-Fatemi (ROF) model problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $g \in L^2(\Omega)$ minimize the functional

$$I(v) := |v|_{BV(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

amongst all $v \in BV(\Omega) \cap L^2(\Omega)$.

Let $\Omega \subset \mathbb{R}^2$ be a bounded polygonal Lipschitz domain.

Rudin-Osher-Fatemi (ROF) model problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $g \in L^2(\Omega)$ minimize the functional

$$I(v) := |v|_{\text{BV}(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

Leonid I. Rudin, Stanley Osher, and Emad Fatemi. “Nonlinear total variation based noise removal algorithms”. In: **Physica D: Nonlinear Phenomena**. Vol. 60. 1-4. 1992, pp. 259–268. DOI: [10.1016/0167-2789\(92\)90242-F](https://doi.org/10.1016/0167-2789(92)90242-F). URL: [https://doi.org/10.1016/0167-2789\(92\)90242-F](https://doi.org/10.1016/0167-2789(92)90242-F)

Let $\Omega \subset \mathbb{R}^2$ be a bounded polygonal Lipschitz domain.

Rudin-Osher-Fatemi (ROF) model problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $g \in L^2(\Omega)$ minimize the functional

$$I(v) := |v|_{\text{BV}(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

Leonid I. Rudin, Stanley Osher, and Emad Fatemi. “Nonlinear total variation based noise removal algorithms”. In: **Physica D: Nonlinear Phenomena**. Vol. 60. 1-4. 1992, pp. 259–268. DOI: 10.1016/0167-2789(92)90242-F. URL: [https://doi.org/10.1016/0167-2789\(92\)90242-F](https://doi.org/10.1016/0167-2789(92)90242-F)

Let $\Omega \subset \mathbb{R}^2$ be a bounded polygonal Lipschitz domain.

Rudin-Osher-Fatemi (ROF) model problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $g \in L^2(\Omega)$ minimize the functional

$$I(v) := |v|_{\text{BV}(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

Leonid I. Rudin, Stanley Osher, and Emad Fatemi. “Nonlinear total variation based noise removal algorithms”. In: **Physica D: Nonlinear Phenomena**. Vol. 60. 1-4. 1992, pp. 259–268. DOI: 10.1016/0167-2789(92)90242-F. URL: [https://doi.org/10.1016/0167-2789\(92\)90242-F](https://doi.org/10.1016/0167-2789(92)90242-F)

Original picture⁰



⁰<https://homepages.cae.wisc.edu/~ece533/images/cameraman.tif>

Original picture⁰



Input signal



The input signal was created by adding AWGN with a SNR of 20 to the original picture.

⁰<https://homepages.cae.wisc.edu/~ece533/images/cameraman.tif>

$$I(v) := |v|_{BV(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

Original picture



Input signal



$$I(v) := |v|_{BV(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

Original picture



Input signal




$$\alpha = 10^5$$

$$I(v) := |v|_{BV(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

Original picture



Input signal



$\alpha = 10^3$




$\alpha = 10^5$

$$I(v) := |v|_{\text{BV}(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

Original picture



Input signal



$\alpha = 10^3$



$\alpha = 10^4$



$\alpha = 10^5$

Pascal Getreuer. “Rudin-Osher-Fatemi Total Variation Denoising using Split Bregman”. In: **Image Processing On Line** 2 (2012), pp. 74–95. URL: <https://doi.org/10.5201/ipol.2012.g-tvd>

Table of Contents

1 Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

2 Primal-Dual Iteration

3 Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Continuous problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $f \in L^2(\Omega)$ minimize the functional

$$E(v) := \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f v \, dx$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

Continuous problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $f \in L^2(\Omega)$ minimize the functional

$$E(v) := \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f v \, dx$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

Continuous problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $f \in L^2(\Omega)$ minimize the functional

$$E(v) := \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f v \, dx$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

For $f = \alpha g$ the functional E has the same minimizers as

$$I(v) = |v|_{\text{BV}(\Omega)} + \frac{\alpha}{2} \|v - g\|_{L^2(\Omega)}^2$$

in $\{v \in \text{BV}(\Omega) \cap L^2(\Omega) \mid \|v\|_{L^1(\partial\Omega)} = 0\}$.

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u \in \text{BV}(\Omega) \cap L^2(\Omega)$ for $E(v) = \frac{\alpha}{2}\|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} fv \, dx$ amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u \in \text{BV}(\Omega) \cap L^2(\Omega)$ for $E(v) = \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f v \, dx$ amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$.

Lemma

Let $v \in \text{BV}(\Omega)$. For all $x \in \mathbb{R}^d$, define

$$\tilde{v}(x) := \begin{cases} v(x) & \text{if } x \in \Omega, \\ 0 & \text{if } x \in \mathbb{R}^d \setminus \overline{\Omega}. \end{cases}$$

Then $\tilde{v} \in \text{BV}(\mathbb{R}^d)$ and $|\tilde{v}|_{\text{BV}(\mathbb{R}^d)} = |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)}$.

Let U be an open subset of \mathbb{R}^d .

Definition (Weak convergence in $BV(U)$)

Let $(v_n)_{n \in \mathbb{N}} \subset BV(U)$ and $v \in BV(U)$ with $v_n \rightarrow v$ in $L^1(U)$ as $n \rightarrow \infty$. Then $(v_n)_{n \in \mathbb{N}}$ converges weakly to v in $BV(U)$ iff, for all $\phi \in C_0(U; \mathbb{R}^d)$, it holds

$$\int_U v_n \operatorname{div}(\phi) \, dx \rightarrow \int_U v \operatorname{div}(\phi) \, dx \quad \text{as } n \rightarrow \infty.$$

We write $v_n \rightharpoonup v$ as $n \rightarrow \infty$.

Theorem

Let $v \in L^1(U)$ and $(v_n)_{n \in \mathbb{N}} \subset BV(U)$ with $\sup_{n \in \mathbb{N}} |v_n|_{BV(U)} < \infty$ and $v_n \rightarrow v$ in $L^1(U)$ as $n \rightarrow \infty$. Then $v \in BV(U)$ and $|v|_{BV(U)} \leq \liminf_{n \rightarrow \infty} |v_n|_{BV(U)}$. Furthermore, $v_n \rightarrow v$ in $BV(U)$.

Theorem

Let $v \in L^1(U)$ and $(v_n)_{n \in \mathbb{N}} \subset BV(U)$ with $\sup_{n \in \mathbb{N}} |v_n|_{BV(U)} < \infty$ and $v_n \rightarrow v$ in $L^1(U)$ as $n \rightarrow \infty$. Then $v \in BV(U)$ and $|v|_{BV(U)} \leq \liminf_{n \rightarrow \infty} |v_n|_{BV(U)}$. Furthermore, $v_n \rightarrow v$ in $BV(U)$.

Let U be a bounded Lipschitz domain.

Theorem

Let $(v_n)_{n \in \mathbb{N}} \subset BV(U)$ be bounded. Then there exists some subsequence $(v_{n_k})_{k \in \mathbb{N}}$ of $(v_n)_{n \in \mathbb{N}}$ and $v \in BV(U)$ such that $v_{n_k} \rightarrow v$ in $L^1(U)$ as $k \rightarrow \infty$.

Theorem (Stability)

Let $f_1, f_2 \in L^2(\Omega)$. For $\ell \in \{1, 2\}$, let $u_\ell \in \text{BV}(\Omega) \cap L^2(\Omega)$ minimize

$$E_\ell(v) := \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f_\ell v \, dx$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$. Then

$$\|u_1 - u_2\| \leq \frac{1}{\alpha} \|f_1 - f_2\|.$$

Theorem (Stability)

Let $f_1, f_2 \in L^2(\Omega)$. For $\ell \in \{1, 2\}$, let $u_\ell \in \text{BV}(\Omega) \cap L^2(\Omega)$ minimize

$$E_\ell(v) := \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f_\ell v \, dx$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$. Then

$$\|u_1 - u_2\| \leq \frac{1}{\alpha} \|f_1 - f_2\|.$$

Sören Bartels. **Numerical Methods for Nonlinear Partial Differential Equations**. Vol. 47. Springer Series in Computational Mathematics. Springer International Publishing, 2015. ISBN: 978-3-319-13796-4. DOI: 10.1007/978-3-319-13797-1, Chapter 10, p. 297-319.

Theorem (Stability)

Let $f_1, f_2 \in L^2(\Omega)$. For $\ell \in \{1, 2\}$, let $u_\ell \in \text{BV}(\Omega) \cap L^2(\Omega)$ minimize

$$E_\ell(v) := \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f_\ell v \, dx$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$. Then

$$\|u_1 - u_2\| \leq \frac{1}{\alpha} \|f_1 - f_2\|.$$

Sören Bartels. **Numerical Methods for Nonlinear Partial Differential Equations**. Vol. 47. Springer Series in Computational Mathematics. Springer International Publishing, 2015. ISBN: 978-3-319-13796-4. DOI: 10.1007/978-3-319-13797-1, Chapter 10, p. 297-319.

Theorem (Stability)

Let $f_1, f_2 \in L^2(\Omega)$. For $\ell \in \{1, 2\}$, let $u_\ell \in \text{BV}(\Omega) \cap L^2(\Omega)$ minimize

$$E_\ell(v) := \frac{\alpha}{2} \|v\|^2 + |v|_{\text{BV}(\Omega)} + \|v\|_{L^1(\partial\Omega)} - \int_{\Omega} f_\ell v \, dx$$

amongst all $v \in \text{BV}(\Omega) \cap L^2(\Omega)$. Then

$$\|u_1 - u_2\| \leq \frac{1}{\alpha} \|f_1 - f_2\|.$$

Sören Bartels. **Numerical Methods for Nonlinear Partial Differential Equations**. Vol. 47. Springer Series in Computational Mathematics. Springer International Publishing, 2015. ISBN: 978-3-319-13796-4. DOI: 10.1007/978-3-319-13797-1, Chapter 10, p. 297-319.

Table of Contents

1 Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

2 Primal-Dual Iteration

3 Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Let \mathcal{T} be a regular triangulation of Ω .

For all $v_{\text{CR}} \in \text{CR}^1(\mathcal{T})$,

$$|v_{\text{CR}}|_{\text{BV}(\Omega)} = \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} + \sum_{F \in \mathcal{E}(\Omega)} \|[v_{\text{CR}}]_F\|_{L^1(F)}.$$

In particular, $\text{CR}^1(\mathcal{T}) \subset \text{BV}(\Omega)$.

$$E(v_{\text{CR}}) = \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + |v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$$

$$|v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} = \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} + \sum_{F \in \mathcal{E}} \|[v_{\text{CR}}]_F\|_{L^1(F)}$$

$$E(v_{\text{CR}}) = \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + |v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$$

$$|v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} = \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} + \sum_{F \in \mathcal{E}} \|[v_{\text{CR}}]_F\|_{L^1(F)}$$

$$E(v_{\text{CR}}) = \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + |v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$$

$$|v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} = \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} + \sum_{F \in \mathcal{E}} \|[v_{\text{CR}}]_F\|_{L^1(F)}$$

Discrete problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $f \in L^2(\Omega)$ minimize the functional

$$E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$$

amongst all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

$$E(v_{\text{CR}}) = \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + |v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$$

$$|v_{\text{CR}}|_{\text{BV}(\Omega)} + \|v_{\text{CR}}\|_{L^1(\partial\Omega)} = \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} + \sum_{F \in \mathcal{E}} \| [v_{\text{CR}}]_F \|_{L^1(F)}$$

Discrete problem

For a parameter $\alpha \in \mathbb{R}_+$ and an input signal $f \in L^2(\Omega)$ minimize the functional

$$E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$$

amongst all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ for $E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$ amongst all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ for
 $E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$ *amongst all*
 $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ for $E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$ amongst all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

Let $K := \{\Lambda \in L^\infty(\Omega; \mathbb{R}^2) \mid |\Lambda(\bullet)| \leq 1 \text{ a.e. in } \Omega\}$ and, for all $(v_{\text{CR}}, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$,

$$L(v_{\text{CR}}, \Lambda_0) := \int_{\Omega} \Lambda_0 \cdot \nabla_{\text{NC}} v_{\text{CR}} \, dx + \frac{\alpha}{2} \|v_{\text{CR}}\|^2 - \int_{\Omega} f v_{\text{CR}} \, dx - I_K(\Lambda_0).$$

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ for $E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$ amongst all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

Let $K := \{\Lambda \in L^\infty(\Omega; \mathbb{R}^2) \mid |\Lambda(\bullet)| \leq 1 \text{ a.e. in } \Omega\}$ and, for all $(v_{\text{CR}}, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$,

$$L(v_{\text{CR}}, \Lambda_0) := \int_{\Omega} \Lambda_0 \cdot \nabla_{\text{NC}} v_{\text{CR}} \, dx + \frac{\alpha}{2} \|v_{\text{CR}}\|^2 - \int_{\Omega} f v_{\text{CR}} \, dx - I_K(\Lambda_0).$$

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ for $E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$ amongst all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

Let $K := \{\Lambda \in L^\infty(\Omega; \mathbb{R}^2) \mid |\Lambda(\bullet)| \leq 1 \text{ a.e. in } \Omega\}$ and, for all $(v_{\text{CR}}, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$,

$$L(v_{\text{CR}}, \Lambda_0) := \int_{\Omega} \Lambda_0 \cdot \nabla_{\text{NC}} v_{\text{CR}} \, dx + \frac{\alpha}{2} \|v_{\text{CR}}\|^2 - \int_{\Omega} f v_{\text{CR}} \, dx - I_K(\Lambda_0).$$

Theorem (Existence and uniqueness of a minimizer)

There exists a unique minimizer $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ for $E_{\text{NC}}(v_{\text{CR}}) := \frac{\alpha}{2} \|v_{\text{CR}}\|^2 + \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \int_{\Omega} f v_{\text{CR}} \, dx$ amongst all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$.

Let $K := \{\Lambda \in L^\infty(\Omega; \mathbb{R}^2) \mid |\Lambda(\bullet)| \leq 1 \text{ a.e. in } \Omega\}$ and, for all $(v_{\text{CR}}, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$,

$$L(v_{\text{CR}}, \Lambda_0) := \int_{\Omega} \Lambda_0 \cdot \nabla_{\text{NC}} v_{\text{CR}} \, dx + \frac{\alpha}{2} \|v_{\text{CR}}\|^2 - \int_{\Omega} f v_{\text{CR}} \, dx - I_K(\Lambda_0).$$

Minimax problem

Find $(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$ such that

$$L(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) = \inf_{v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})} \sup_{\Lambda_0 \in P_0(\mathcal{T}; \mathbb{R}^2)} L(v_{\text{CR}}, \Lambda_0).$$

$$L(v_{\text{CR}}, \Lambda_0) := \int_{\Omega} \Lambda_0 \cdot \nabla_{\text{NC}} v_{\text{CR}} \, dx + \frac{\alpha}{2} \|v_{\text{CR}}\|^2 - \int_{\Omega} f v_{\text{CR}} \, dx - I_K(\Lambda_0)$$

Minimax problem

Find $(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$ such that

$$L(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) = \inf_{v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})} \sup_{\Lambda_0 \in P_0(\mathcal{T}; \mathbb{R}^2)} L(v_{\text{CR}}, \Lambda_0).$$

$$L(v_{\text{CR}}, \Lambda_0) := \int_{\Omega} \Lambda_0 \cdot \nabla_{\text{NC}} v_{\text{CR}} \, dx + \frac{\alpha}{2} \|v_{\text{CR}}\|^2 - \int_{\Omega} f v_{\text{CR}} \, dx - I_K(\Lambda_0)$$

Minimax problem

Find $(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$ such that

$$L(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) = \inf_{v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})} \sup_{\Lambda_0 \in P_0(\mathcal{T}; \mathbb{R}^2)} L(v_{\text{CR}}, \Lambda_0).$$

This problem has a solution

$$(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) \in \text{CR}_0^1(\mathcal{T}) \times (P_0(\mathcal{T}; \mathbb{R}^2) \cap K).$$

$$L(v_{\text{CR}}, \Lambda_0) := \int_{\Omega} \Lambda_0 \cdot \nabla_{\text{NC}} v_{\text{CR}} \, dx + \frac{\alpha}{2} \|v_{\text{CR}}\|^2 - \int_{\Omega} f v_{\text{CR}} \, dx - I_K(\Lambda_0)$$

Minimax problem

Find $(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \mathbb{R}^2)$ such that

$$L(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) = \inf_{v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})} \sup_{\Lambda_0 \in P_0(\mathcal{T}; \mathbb{R}^2)} L(v_{\text{CR}}, \Lambda_0).$$

This problem has a solution

$$(\tilde{u}_{\text{CR}}, \bar{\Lambda}_0) \in \text{CR}_0^1(\mathcal{T}) \times (P_0(\mathcal{T}; \mathbb{R}^2) \cap K).$$

R. Tyrrell Rockafellar. **Convex Analysis**. New Jersey: Princeton University Press, 1970. ISBN: 0-691-08069-0

Theorem (Equivalent characterizations)

For a function $\tilde{u}_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ the following statements are equivalent.

- (i) \tilde{u}_{CR} solves the discrete problem.
- (ii) There exists $\bar{\Lambda}_0 \in P_0(\mathcal{T}; \mathbb{R}^2)$ with $|\bar{\Lambda}_0(\bullet)| \leq 1$ a.e. in Ω s.t.

$$\bar{\Lambda}_0(\bullet) \cdot \nabla_{\text{NC}} \tilde{u}_{\text{CR}}(\bullet) = |\nabla_{\text{NC}} \tilde{u}_{\text{CR}}(\bullet)| \quad \text{a.e. in } \Omega$$

and

$$(\bar{\Lambda}_0, \nabla_{\text{NC}} v_{\text{CR}}) = (f - \alpha \tilde{u}_{\text{CR}}, v_{\text{CR}}) \quad \text{for all } v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T}).$$

- (iii) For all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$,

$$(f - \alpha \tilde{u}_{\text{CR}}, v_{\text{CR}} - \tilde{u}_{\text{CR}}) \leq \|\nabla_{\text{NC}} v_{\text{CR}}\|_{L^1(\Omega)} - \|\nabla_{\text{NC}} \tilde{u}_{\text{CR}}\|_{L^1(\Omega)}.$$

Table of Contents

1 Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

2 Primal-Dual Iteration

3 Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}}),$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}}), \tau > 0$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(u_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ *for* u_j ,

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(u_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ *for* u_j , *and set*

$$v_j := \frac{u_j - u_{j-1}}{\tau}.$$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(u_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ for u_j , and set

$$v_j := \frac{u_j - u_{j-1}}{\tau}.$$

Output: Sequence $(u_j, \Lambda_j)_{j \in \mathbb{N}}$ in $\text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(u_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ for u_j , and set

$$v_j := \frac{u_j - u_{j-1}}{\tau}.$$

Output: Sequence $(u_j, \Lambda_j)_{j \in \mathbb{N}}$ in $\text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(\textcolor{red}{u}_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ for u_j , and set

$$v_j := \frac{u_j - u_{j-1}}{\tau}.$$

Output: Sequence $(u_j, \Lambda_j)_{j \in \mathbb{N}}$ in $\text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(u_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ for u_j , and set

$$v_j := \frac{u_j - u_{j-1}}{\tau}.$$

Output: Sequence $(u_j, \Lambda_j)_{j \in \mathbb{N}}$ in $\text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(u_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ for u_j , and set

$$v_j := \frac{u_j - u_{j-1}}{\tau}.$$

Output: Sequence $(u_j, \Lambda_j)_{j \in \mathbb{N}}$ in $\text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$

Algorithm (Primal-dual iteration)

Input: $(u_0, \Lambda_0) \in \text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$, $\tau > 0$, $\varepsilon_{\text{stop}} > 0$

Initialize $v_0 := 0$ in $\text{CR}_0^1(\mathcal{T})$.

for $j = 1, 2, \dots$

$$\tilde{u}_j := u_{j-1} + \tau v_{j-1}, \quad \Lambda_j := \frac{\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j}{\max \{1, |\Lambda_{j-1} + \tau \nabla_{\text{NC}} \tilde{u}_j|\}},$$

solve

$$\frac{1}{\tau} a_{\text{NC}}(u_j, \bullet) + \alpha(u_j, \bullet) = \frac{1}{\tau} a_{\text{NC}}(u_{j-1}, \bullet) + (f, \bullet) - (\Lambda_j, \nabla_{\text{NC}} \bullet)$$

in $\text{CR}_0^1(\mathcal{T})$ for u_j , and set

$$v_j := \frac{u_j - u_{j-1}}{\tau}. \text{ Terminate iteration if } \|v_j\| < \varepsilon_{\text{stop}}.$$

Output: Sequence $(u_j, \Lambda_j)_{j \in \mathbb{N}}$ in $\text{CR}_0^1(\mathcal{T}) \times P_0(\mathcal{T}; \overline{B_{\mathbb{R}^2}})$

Theorem (Convergence of the primal-dual iteration)

Let $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ solve the discrete problem, $\bar{\Lambda}_0 \in P_0(\mathcal{T}; \mathbb{R}^2)$ satisfy $|\bar{\Lambda}_0(\bullet)| \leq 1$ a.e. in Ω as well as

$$\bar{\Lambda}_0(\bullet) \cdot \nabla_{\text{NC}} u_{\text{CR}}(\bullet) = |\nabla_{\text{NC}} u_{\text{CR}}(\bullet)| \quad \text{a.e. in } \Omega$$

and

$$(\bar{\Lambda}_0, \nabla_{\text{NC}} v_{\text{CR}}) = (f - \alpha u_{\text{CR}}, v_{\text{CR}}) \quad \text{for all } v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T}),$$

and $\tau \in (0, 1]$. Then the iterates $(u_j)_{j \in \mathbb{N}}$ of the primal-dual iteration converge to u_{CR} in $L^2(\Omega)$.

Theorem (Convergence of the primal-dual iteration)

Let $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ solve the discrete problem, $\bar{\Lambda}_0 \in P_0(\mathcal{T}; \mathbb{R}^2)$ satisfy $|\bar{\Lambda}_0(\bullet)| \leq 1$ a.e. in Ω as well as

$$\bar{\Lambda}_0(\bullet) \cdot \nabla_{\text{NC}} u_{\text{CR}}(\bullet) = |\nabla_{\text{NC}} u_{\text{CR}}(\bullet)| \quad \text{a.e. in } \Omega$$

and

$$(\bar{\Lambda}_0, \nabla_{\text{NC}} v_{\text{CR}}) = (f - \alpha u_{\text{CR}}, v_{\text{CR}}) \quad \text{for all } v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T}),$$

and $\tau \in (0, 1]$. Then the iterates $(u_j)_{j \in \mathbb{N}}$ of the primal-dual iteration converge to u_{CR} in $L^2(\Omega)$.

For all $J \in \mathbb{N}$,

$$\sum_{j=1}^J \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Theorem (Convergence of the primal-dual iteration)

Let $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ solve the discrete problem, $\bar{\Lambda}_0 \in P_0(\mathcal{T}; \mathbb{R}^2)$ satisfy $|\bar{\Lambda}_0(\bullet)| \leq 1$ a.e. in Ω as well as

$$\bar{\Lambda}_0(\bullet) \cdot \nabla_{\text{NC}} u_{\text{CR}}(\bullet) = |\nabla_{\text{NC}} u_{\text{CR}}(\bullet)| \quad \text{a.e. in } \Omega$$

and

$$(\bar{\Lambda}_0, \nabla_{\text{NC}} v_{\text{CR}}) = (f - \alpha u_{\text{CR}}, v_{\text{CR}}) \quad \text{for all } v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T}),$$

and $\tau \in (0, 1]$. Then the iterates $(u_j)_{j \in \mathbb{N}}$ of the primal-dual iteration converge to u_{CR} in $L^2(\Omega)$.

For all $J \in \mathbb{N}$,

$$\sum_{j=1}^J \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Theorem (Convergence of the primal-dual iteration)

Let $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ solve the discrete problem, $\bar{\Lambda}_0 \in P_0(\mathcal{T}; \mathbb{R}^2)$ satisfy $|\bar{\Lambda}_0(\bullet)| \leq 1$ a.e. in Ω as well as

$$\bar{\Lambda}_0(\bullet) \cdot \nabla_{\text{NC}} u_{\text{CR}}(\bullet) = |\nabla_{\text{NC}} u_{\text{CR}}(\bullet)| \quad \text{a.e. in } \Omega$$

and

$$(\bar{\Lambda}_0, \nabla_{\text{NC}} v_{\text{CR}}) = (f - \alpha u_{\text{CR}}, v_{\text{CR}}) \quad \text{for all } v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T}),$$

and $\tau \in (0, 1]$. Then the iterates $(u_j)_{j \in \mathbb{N}}$ of the primal-dual iteration converge to u_{CR} in $L^2(\Omega)$.

$$\sum_{j=1}^{\infty} \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Table of Contents

① Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

② Primal-Dual Iteration

③ Numerical Examples

Settings


Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator


Convergence of the Iteration

Let $\Omega = (-1, 1)^2$. Define $f \in H_0^1(\Omega)$ by $f(x) = \tilde{f}(|x|)$ for all $x \in \Omega$ with

$$\tilde{f}(r) := \begin{cases} \alpha - 12(2 - 9r) & \text{if } 0 \leq r \leq \frac{1}{6}, \\ 6r\alpha - \frac{1}{r} & \text{if } \frac{1}{6} \leq r \leq \frac{1}{3}, \\ 2\alpha + 6\pi \sin(\pi(6r - 2)) - \frac{1}{r} \cos(\pi(6r - 2)) & \text{if } \frac{1}{3} \leq r \leq \frac{1}{2}, \\ \alpha(5 - 6r) + \frac{1}{r} & \text{if } \frac{1}{2} \leq r \leq \frac{5}{6}, \\ -3\pi \sin(\pi(6r - 5)) + \frac{1 + \cos(\pi(6r - 5))}{2r} & \text{if } \frac{5}{6} \leq r \leq 1. \end{cases}$$




f for $\alpha = 1$




f for $\alpha = 1$ along the axes

Then the solution to the continuous problem with input signal f is given by $u \in H_0^1(\Omega)$ defined by $u(x) = \tilde{u}(|x|)$ for all $x \in \Omega$ with

$$\tilde{u}(r) := \begin{cases} 1 & \text{if } 0 \leq r \leq \frac{1}{6}, \\ 6r & \text{if } \frac{1}{6} \leq r \leq \frac{1}{3}, \\ 2 & \text{if } \frac{1}{3} \leq r \leq \frac{1}{2}, \\ 5 - 6r & \text{if } \frac{1}{2} \leq r \leq \frac{5}{6}, \\ 0 & \text{if } \frac{5}{6} \leq r. \end{cases}$$




u




u along the axes

Then the solution to the continuous problem with input signal f is given by $u \in H_0^1(\Omega)$ defined by $u(x) = \tilde{u}(|x|)$ for all $x \in \Omega$ with

$$\tilde{u}(r) := \begin{cases} 1 & \text{if } 0 \leq r \leq \frac{1}{6}, \\ 6r & \text{if } \frac{1}{6} \leq r \leq \frac{1}{3}, \\ 2 & \text{if } \frac{1}{3} \leq r \leq \frac{1}{2}, \\ 5 - 6r & \text{if } \frac{1}{2} \leq r \leq \frac{5}{6}, \\ 0 & \text{if } \frac{5}{6} \leq r. \end{cases}$$



u



u along the axes

It holds $E(u) \approx -2.058034062391$.


For $\alpha = 10000$ let the input signal represent the grayscale of an image in $[0, 1]^{256 \times 256}$ multiplied with α scaled to the domain $\Omega = (0, 1)^2$.



Image cameraman


Initial Triangulations for the Input Signals

Input signal f



$$\Omega = (-1, 1)^2$$

Input signal cameraman



$$\Omega = (0, 1)^2$$

Table of Contents

1 Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

2 Primal-Dual Iteration

3 Numerical Examples

Settings


Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Choice of τ


For the rest of the presentation (unless otherwise specified) let the bulk parameter be $\theta = 0.5$, and $\varepsilon_{\text{stop}} = 10^{-4}$.



Input signal f

Choice of τ


For the rest of the presentation (unless otherwise specified) let the bulk parameter be $\theta = 0.5$, and $\varepsilon_{\text{stop}} = 10^{-4}$.



Input signal cameraman

Choice of τ

For the rest of the presentation (unless otherwise specified) let the bulk parameter be $\theta = 0.5$, and $\varepsilon_{\text{stop}} = 10^{-4}$.



Input signal f

Conclusion and Hypothesis

For the rest of the presentation $\tau = 1$.

Conclusion and Hypothesis

For the rest of the presentation $\tau = 1$.

Convergence of the iterates of the primal-dual iteration to the discrete solution u_{CR} followed from

$$\sum_{j=1}^{\infty} \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Conclusion and Hypothesis

For the rest of the presentation $\tau = 1$.

Convergence of the iterates of the primal-dual iteration to the discrete solution u_{CR} followed from

$$\sum_{j=1}^{\infty} \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Conclusion and Hypothesis

For the rest of the presentation $\tau = 1$.

Convergence of the iterates of the primal-dual iteration to the discrete solution u_{CR} followed from

$$\sum_{j=1}^{\infty} \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Settings with $\tau = 1.2$ and no convergence were observed.

Conclusion and Hypothesis

For the rest of the presentation $\tau = 1$.

Convergence of the iterates of the primal-dual iteration to the discrete solution u_{CR} followed from

$$\sum_{j=1}^{\infty} \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Settings with $\tau = 1.2$ and no convergence were observed.

Conclusion and Hypothesis

For the rest of the presentation $\tau = 1$.

Convergence of the iterates of the primal-dual iteration to the discrete solution u_{CR} followed from

$$\sum_{j=1}^{\infty} \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Settings with $\tau = 1.2$ and no convergence were observed.

For $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$, define $J_1 : \text{CR}_0^1(\mathcal{T}) \rightarrow P_1(\mathcal{T}) \cap C_0(\Omega)$ by

$$J_1 v_{\text{CR}}(z) := |\mathcal{T}(z)|^{-1} \sum_{T \in \mathcal{T}(z)} v_{\text{CR}}|_T(z) \quad \text{for all } z \in \mathcal{N}(\Omega).$$

Conclusion and Hypothesis

For the rest of the presentation $\tau = 1$.

Convergence of the iterates of the primal-dual iteration to the discrete solution u_{CR} followed from

$$\sum_{j=1}^{\infty} \|u_{\text{CR}} - u_j\|^2 \leq \frac{1}{2\alpha\tau} (\|u_{\text{CR}} - u_0\|_{\text{NC}}^2 + \|\bar{\Lambda}_0 - \Lambda_0\|^2).$$

Settings with $\tau = 1.2$ and no convergence were observed.


For $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$, define $J_1 : \text{CR}_0^1(\mathcal{T}) \rightarrow P_1(\mathcal{T}) \cap C_0(\Omega)$ by

$$J_1 v_{\text{CR}}(z) := |\mathcal{T}(z)|^{-1} \sum_{T \in \mathcal{T}(z)} v_{\text{CR}}|_T(z) \quad \text{for all } z \in \mathcal{N}(\Omega).$$

Use $\hat{u}_0 := J_1 u_{\text{CR}, \mathcal{T}} \in P_1(\mathcal{T}) \cap C_0(\Omega) \subseteq P_1(\hat{\mathcal{T}}) \cap C_0(\Omega) \subseteq \text{CR}_0^1(\hat{\mathcal{T}})$ as input for the iteration on the refined triangulation $\hat{\mathcal{T}}$.


Choice of $\varepsilon_{\text{stop}}$

With $\tau = 1$ the stopping criterion reads $\|u_j - u_{j-1}\|_{\text{NC}} < \varepsilon_{\text{stop}}$.



Choice of $\varepsilon_{\text{stop}}$

With $\tau = 1$ the stopping criterion reads $\|u_j - u_{j-1}\|_{\text{NC}} < \varepsilon_{\text{stop}}$.



For the rest of the presentation $\varepsilon_{\text{stop}} = 10^{-4}$.

Table of Contents

① Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

② Primal-Dual Iteration

③ Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

TODO translate and rewrite

Theorem

Sei Ω konvex, $f \in H_0^1(\Omega)$ das Eingangssignal für mit Lösung $u \in H_0^1(\Omega)$ und minimaler Energie $E(u)$ sowie für mit Lösung $u_{\text{CR}} \in \text{CR}_0^1(\Omega)$ und minimaler Energie $E_{\text{NC}}(u_{\text{CR}})$. Dann gilt

$$E_{\text{NC}}(u_{\text{CR}}) + \frac{\alpha}{2} \|u - u_{\text{CR}}\|^2 - \frac{\kappa_{\text{CR}}}{\alpha} \|h_{\mathcal{T}}(f - \alpha u_{\text{CR}})\| \|\nabla f\| \leq E(u).$$

Dabei ist die Konstante $\kappa_{\text{CR}} := \sqrt{1/48 + 1/j_{1,1}^2}$ mit der kleinsten positiven Nullstelle $j_{1,1}$ der Bessel-Funktion erster Art. Insbesondere gilt dann für

$$E_{\text{GLEB}} := E_{\text{NC}}(u_{\text{CR}}) - \frac{\kappa_{\text{CR}}}{\alpha} \|h_{\mathcal{T}}(f - \alpha u_{\text{CR}})\| \|\nabla f\|, \quad (1)$$

dass $E_{\text{NC}}(u_{\text{CR}}) \geq E_{\text{GLEB}}$ und $E(u) \geq E_{\text{GLEB}}$.

TODO translate and rewrite, leave out the d and make it 2. Say we choose $\gamma = 1$ because we want to refine towards the discontinuities

Definition (Verfeinerungsindikator)

Für $d \in \mathbb{N}$ (in dieser Arbeit stets $d = 2$) und $0 < \gamma \leq 1$ definieren wir für alle $T \in \mathcal{T}$ und $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$ die Funktionen

$$\eta_{\text{V}}(T) := |T|^{2/d} \|f - \alpha u_{\text{CR}}\|_{L^2(T)}^2 \quad \text{und}$$

$$\eta_{\text{J}}(T) := |T|^{\gamma/d} \sum_{F \in \mathcal{E}(T)} \| [u_{\text{CR}}]_F \|_{L^1(F)}.$$

Damit definieren wir den Verfeinerungsindikator $\eta := \sum_{T \in \mathcal{T}} \eta(T)$, wobei

$$\eta(T) := \eta_{\text{V}}(T) + \eta_{\text{J}}(T) \quad \text{für alle } T \in \mathcal{T}.$$

cameraman triangulation figures to show the effect of the refinement indicator

convergence graphs error and refinement indicator and probably its
volume and jump contributions
adaptive and uniform
maybe also plot refinement indicator and its contributions for
cameraman
say expected rates from bartels and compare to them

plot differences between energies and E_{gleb}
plot the differences between the discrete energies and the exact
energy in the graph with all E_{gleb} differences
note that $E_{\text{nc}} - E_{\text{gleb}}$ and $E - E_{\text{gleb}}$ don't differ much obviously
because $E_{\text{nc}} - E$ tends to zero (as seen in the plots)
show strict convexity theorem here to show that it holds (even
'better' than the theorem guarantees)

show error, η (without contributions) together with the differences of the energies and EGLEB

Table of Contents

① Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

② Primal-Dual Iteration

③ Numerical Examples

Settings


Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

TODO

think about where to position this topic wise



previous frame cont.

show that the energy converges from above to the exact energy (not necessarily monotonically, i.e. choose some example where it simply converges and one example where there are oscillations, i.e. two pictures here

also mention that it converges to something slightly below the exact energy, i.e. choose the pictures accordingly, maybe even plot the error between the energies of the iterates and the exact energy

THAT MEANS choose one picture from f where the exact energy can be seen and maybe one from the cameraman

mention that during the afem loop we will see, that this undershooting decrease (as one would expect, the discrete energy converges to the exact energy)

show the result of an iteration (plot of u_{Approx} next to u_{Exact})
(along the axes as well) and show cameraman grayscale plot
definitely mention degrees of freedom if only one level is shown!!!

End slide with some picture maybe or just end on the last slide in experiments (with might be a plot with everything in it, which would be nice

Table of Contents

① Recapitulation

Functions of Bounded Variation

Rudin-Osher-Fatemi Model Problem

Continuous Problem

Discrete Problem

② Primal-Dual Iteration

③ Numerical Examples

Settings

Choice of Parameters

Guaranteed lower Energy Bound and Refinement Indicator

Convergence of the Iteration

Appendix

L2 Sprünge vielleicht auswerten (bleiben sie konstant. . . , if we consider them, it becomes conforming
die L2 Sprung entwicklung einiger experiment (iteration auswerten, iteration selbst und Afem loop insgesamt). bleiben sicherlich konstant oder sowas

different norm for termination criteria comparison (energy difference not good because oscillations, everything else (don't show L2 error squared) is similar, just different height

compare times without preallocating and with (inform about the extreme improvement of performance

Let $u_P : [0, \infty) \rightarrow \mathbb{R}$ with $u_P(r) = 0$ for $r \geq 1$, and, for all $x \in \Omega$, $u(x) = u_P(|x|)$.

Let $u_P : [0, \infty) \rightarrow \mathbb{R}$ with $u_P(r) = 0$ for $r \geq 1$, and, for all $x \in \Omega$, $u(x) = u_P(|x|)$. Furthermore, assume the existence of $\partial_r u_P$ a.e. in $[0, \infty)$, the existence of the derivative of

$$\operatorname{sgn}(\partial_r u_P(r)) := \begin{cases} -1 & \text{für } \partial_r u_P(r) < 0, \\ x \in [0, 1] & \text{für } \partial_r u_P(r) = 0, \\ 1 & \text{für } \partial_r u_P(r) > 0. \end{cases}$$

a.e. in $[0, \infty)$, and that $\operatorname{sgn}(\partial_r u_P(r))/r \rightarrow 0$ as $r \rightarrow 0$.

Let $u_P : [0, \infty) \rightarrow \mathbb{R}$ with $u_P(r) = 0$ for $r \geq 1$, and, for all $x \in \Omega$, $u(x) = u_P(|x|)$. Furthermore, assume the existence of $\partial_r u_P$ a.e. in $[0, \infty)$, the existence of the derivative of

$$\operatorname{sgn}(\partial_r u_P(r)) := \begin{cases} -1 & \text{für } \partial_r u_P(r) < 0, \\ x \in [0, 1] & \text{für } \partial_r u_P(r) = 0, \\ 1 & \text{für } \partial_r u_P(r) > 0. \end{cases}$$

a.e. in $[0, \infty)$, and that $\operatorname{sgn}(\partial_r u_P(r))/r \rightarrow 0$ as $r \rightarrow 0$. For all $r \in [0, \infty)$, define

$$f_P(r) := \alpha u_P(r) - \partial_r (\operatorname{sgn}(\partial_r u_P(r))) - \frac{\operatorname{sgn}(\partial_r u_P(r))}{r}$$

Let $u_P : [0, \infty) \rightarrow \mathbb{R}$ with $u_P(r) = 0$ for $r \geq 1$, and, for all $x \in \Omega$, $u(x) = u_P(|x|)$. Furthermore, assume the existence of $\partial_r u_P$ a.e. in $[0, \infty)$, the existence of the derivative of

$$\operatorname{sgn}(\partial_r u_P(r)) := \begin{cases} -1 & \text{für } \partial_r u_P(r) < 0, \\ x \in [0, 1] & \text{für } \partial_r u_P(r) = 0, \\ 1 & \text{für } \partial_r u_P(r) > 0. \end{cases}$$

a.e. in $[0, \infty)$, and that $\operatorname{sgn}(\partial_r u_P(r))/r \rightarrow 0$ as $r \rightarrow 0$. For all $r \in [0, \infty)$, define

$$f_P(r) := \alpha u_P(r) - \partial_r (\operatorname{sgn}(\partial_r u_P(r))) - \frac{\operatorname{sgn}(\partial_r u_P(r))}{r}$$

Then u solves the continuous problem on $\Omega \supseteq \{w \in \mathbb{R}^2 \mid |w| \leq 1\}$ if the input signal is $f(x) := f_P(|x|)$.