

Convergent Science Physical Oncology



TOPICAL REVIEW

Cancer as a disorder of patterning information: computational and biophysical perspectives on the cancer problem

Douglas Moore^{1,3}, Sara I Walker^{1,2} and Michael Levin^{3,4}

¹ BEYOND Center for Fundamental Concepts in Science Arizona State University, Tempe, AZ, United States of America

² School of Earth and Space Exploration, Arizona State University, Tempe, AZ, United States of America

³ Biology Department, Allen Discovery Center at Tufts, Tufts University, Medford, MA, United States of America

⁴ Author to whom any correspondence should be addressed.

E-mail: michael.levin@tufts.edu

Keywords: bioelectricity, ion channel, gap junction, morphogenetic field

Abstract

The current paradigm views cancer as arising clonally from a degradation of genetic information in single cells. A complementary perspective, originating at the dawn of modern developmental biology, is that cancer is the result of a system disorder of algorithms that normally orchestrate individual cell activities toward specific anatomical structures and away from tumorigenesis. A view of cancer as a disease of geometry focuses on the pathways that allow cells to cooperate to build and maintain large-scale anatomical patterning. Cancer may result when cells stop maintaining higher-order structures and reduce the boundary of their computational selves to a single-cell level, reverting to a unicellular lifestyle in which the rest of the organism is merely part of the environment at the expense of which all living things survive. While this view has been widely discussed, little progress has been made in providing a quantitative, mechanistic framework within which this perspective's specific and unique implications for treatment strategies can be tested and biomedically exploited. Here, we highlight two recent areas of progress which may facilitate much-needed progress on the cancer problem. First, we review the roles that endogenous bioelectrical networks, operating across many tissues *in vivo*, play as a medium of information processing in tumor suppression, progression, and reprogramming. Second, we provide a primer to the development of computational theory and tools for quantifying the information and causal control structures in cancer and other complex biological systems. Rigorous mathematical formalisms now exist to measure and analyze the extent to which 'a whole is more than the sum of its parts', applications of which could lead to new strategies for cancer reprogramming. Here, we review the basic landscape of these related subfields, and sketch specific ways in which a synthesis of novel integrative biophysics and mathematical analysis may contribute to novel ways to understand and address cancer *in vivo*.

1. Introduction

Cancer is well-recognized not only as an extremely pernicious medical problem, but also as a group of phenomena deeply connected to the fundamental questions of evolution, multicellularity, pattern (dis)regulation, and the interplay between the genome and the environment [1, 2]. Two fundamental paradigms currently divide the field. One is that cancer results from disorders of genetics: cancer cells are fundamentally broken due to genomic mutation or instability, and their clonal progeny form tumors and metastases. This view is sometimes called the SMT, *somatic mutation theory* [3–5]. A competing

perspective is that of cancer as a circuit disease—a disorder of the complex biophysical, transcriptional, and epigenetic dynamics that regulate cellular state and the ability of cells to cooperate *in vivo* [6–8]. Under this view (sometimes called the tissue organization field theory or TOFT, [9, 10]), cancer is akin to a traffic jam [11]—the problem is not a permanent discrete alteration within a founder cell and all of its clonal descendants, but an undesirable stable attractor in the complex network of controls that normally guides anatomical homeostasis [12].

The relative merits of the two models have been discussed extensively [10, 13–17]. We do not rehash the arguments here, nor do we claim that cancer is a

monolithic phenomenon that is entirely explained by either model in all cases. Nevertheless, there is widespread agreement that fundamental advances in the basic understanding of cancer are necessary to break the stalemate in cancer medicine. Here, we focus on the under-appreciated developmental perspective because it holds potential to generate such fundamental advances. We review the classical and recent data suggesting a complementary top-down strategy for control of the cancer process, and outline an emerging set of quantitative approaches that we believe are an ideal formalism from which to functionally address this conceptualization of the cancer problem at the bench.

This view places the locus of cancer at a multicellular level—specifically, a failure of the mechanisms that orchestrate individual cell behaviors toward large-scale anatomical outcomes: the organism’s target morphology [18]. This is important, because this view makes unique predictions that could overcome the increasingly acknowledged gap between the amount of effort spent on molecular biology and the modest gains in cancer medicine to date [19]. For example, the SMT model implies that cancer cells are irrevocably damaged and must be killed to address cancer outcomes. In contrast, TOFT suggests rebooting/reprogramming strategies that could avoid toxic chemotherapy and compensatory proliferation by shifting control networks back into the regime that normally instructs their morphogenetic functions within a complex organism. In the SMT model, each cancer cell must be addressed individually, lest it initiate relapse. TOFT suggests that diagnostics and intervention may include a non-local, tissue-scale component involving long-range integration of cell states and communication into global (dis)order. Finally, SMT and TOFT make opposing predictions about the relationship between induction of regenerative repair and carcinogenic susceptibility (discussed below).

Indeed, an anatomical framework is what is used to diagnose cancer clinically, and is the context in which SMT-derived or other therapies are evaluated. Nevertheless, the developmental perspective receives less attention because the successes of mechanistic molecular genetics (genomics, high throughput sequencing) make it easy to amass increasingly high-resolution subcellular data comparing cells within and across tumors, making it tempting to search for the source of transformation within the massive volumes of genomic data. The alternative requires a much more difficult journey to understanding how the default program of unicellular life is normally harnessed in the context of morphogenesis, regeneration, and tumor suppression towards creation and maintenance of a complex body. Below, we first review some little-known data that support a top-down perspective and functional importance of anatomical homeostasis [20] for the cancer problem. We then discuss two essential aspects of a new approach: mathematical theory applicable to higher-order biological function, and

an emerging set of biophysical control mechanisms that can be effectively targeted by interventions motivated by the theoretical work.

Crucially, a full understanding of the cancer problem will involve not only the mechanisms of cellular control, but also the algorithms that guide their communication and other activities at a population level. We propose a research program that includes investigating how cell networks operate as well as the development of new quantitative theory that describes the control policies and information-processing executed by multicellular tissues. This latter point is essential if the field is to move beyond the qualitative idea of target morphology as the missing control factor when cells go rogue, toward specific models that make testable, quantitative predictions to identify novel intervention strategies for biomedicine. In the next section, we introduce the unacquainted reader to basic concepts of information theory as applied to biological systems. How do cell networks process information, and how can cancer biologists acquire data that can be analyzed using these methods? We then go on to review how endogenous bioelectric circuits help regulate large-scale growth and form (section 3), and how information theory may yield insights into their control processes (section 4). We conclude with a discussion of cancer as a disease of global physiological dynamics (section 5), and how this may be viewed from the perspective of information theory as a breakdown or reduction of information integration (section 6).

2. Information in biological systems

The cancer problem has been considered from many different modeling perspectives ranging from the standard mechanistic models [21–25], to applications of population dynamics [26–29], evolutionary game theory [30–32], and models of swarm dynamics [33, 34]. However, these methods generally only consider one or two levels of organization in describing the dynamics of the systems they consider, and almost exclusively view the lowest level as giving rise to the higher levels. This ultimately endows the lowest level of organization with the most causal power, and rejects the possibility that, as complex systems, cells and collectives exhibit behavior and function that cannot necessarily be better described or deduced based solely on the dynamics of smaller and smaller components [35]. Moreover, the above approaches largely do not address the loss of patterning or the goal-seeking behavior of cancer systems, which we argue in are its defining characteristics and offer tractable points for intervention.

An alternative approach is to take a more wholistic perspective, treating the system as composed of many levels of organization and inquiring about the interaction and information flow between them. We propose an ‘*information-theoretic toolkit*’ which may

be utilized to refine and quantify how the various levels of organization of a system interact and ultimately exert influence upon one another. Such a quantitative approach will be essential in the coming decades to begin to understand the measurements that cells and tissues make *in vivo*, which include measurements of distant cells' properties and likely measurements of properties that do not exist at the level of single cells or molecules (e.g. organ size, topology, or position).

Information theory has gained widespread application in the biological sciences, from understanding the information content of genomes [36–38], to identifying individuality [39], and patterns of information processing in brain tissue [40–42]. It has even been suggested by Smith that without drawing on informational analogies, the discoveries made in molecular biology over the last 60 years would have been accomplished at a slower pace, or perhaps not even at all [43]. However, information theory has so far seen less application to the problem of cancer (see e.g. [44, 45] for exceptions) than other areas of biological research despite the fact that information is a way to quantify patterning and structure, precisely the features that we argue break down during oncogenesis making it a disease of pattern dysregulation.

Modern formulations of information theory began with the seminal work of Claude Shannon in 1948 who defined a quantity now termed *Shannon entropy* (H),

$$H(X) = - \sum_{i=1}^N p(x_i) \log p(x_i), \quad (1)$$

where X is a finite random variable, which can take on one of N different states, $\{x_1, x_2, \dots, x_N\}$ each with probability $p(x_i)$. Shannon entropy is often informally described as the expected degree of surprise at learning the outcome of an event: the more unlikely the event, the more surprised you feel at learning that it happened. Equation (1) forms the core of information theory. To quantify shared information due to communication, Shannon introduced the concepts of sender and receiver, described by two random variables X and Y , respectively. The idea is to determine how strongly events drawn from each are correlated as, upon communication, the state of sender and receiver should become correlated. This can be thought of as inquiring as to how much information variable X carries about variable Y , and vice versa. One can then ask, by making observations of X , how much can be learned about Y ? To quantify this notion of information, Shannon defined *mutual information* as

$$I(X, Y) = \sum_{i=1}^{N_x} \sum_{j=1}^{N_y} p(x_i, y_j) \log \frac{p(x_i, y_j)}{p(x_i)p(y_j)}, \quad (2)$$

where $p(x, y)$ denotes the probability of observing X and Y in the states x and y , respectively. Intuitively, mutual information precisely measures how much the distribution of X and Y jointly differs from the distribution of each independently. That is,

equation (2) can be thought of as the entropy shared between X and Y .

As a conceptual example of the role mutual information plays in biological systems, consider two cells communicating via cell potentials and inter- and intracellular ion concentrations. The expectation is that the internal state of a cell is in some way correlated with the pumping of ions in and out of the cell and thereby with the cell's membrane potential. They should therefore have non-zero mutual information. When neighboring cells observe changes in intercellular ion concentrations, via changes in their membrane potentials, they respond mechanistically. Other examples of the application of mutual information in biological modeling include quantifying the correlation between the information content of the genomes and their environment [46], between molecules within the cell [47] and between early life replicators [48] and their environment.

It should be noted that mutual information is not directional: it is not concerned with which event preceded the other and therefore is not a measure of causation. In biomedical research, we are often interested in the idea of *intervention* rather than correlation. For example, in cancer therapy a small molecule drug might be used to block a cell surface receptor, or a key point in a chemical pathway within the cell, inactivating some downstream process, such as the production of cellular growth factors [49, 50]. Two cellular growth factors could be the output of some of the same series of steps and therefore would be correlated even if they do not causally interact. The goal of an information-based approach is to uncover the structure of integrated healthy tissue and how this structure breaks down because of cancer progression. While correlations do not directly imply causation, they nonetheless resolve features of the non-linear causal structure of a complex system. We discuss below under what circumstances this could also be useful to identifying targets for therapy. The next section reviews the biological foundation for such an information-based approach.

3. Cancer as a disorder of pattern regulation

3.1. Dynamic pattern control as anatomical homeostasis: the healthy state

The existence of multicellular organisms requires that individual cells abandon their normal unicellular behavior patterns; they must specialize morphologically and behaviorally, reduce or abandon their own proliferative tendencies, and cooperate to build and continuously repair complex large-scale anatomical structures (as occurs in embryogenesis and regenerative repair/remodeling, respectively). A process that subjugates individual agents' needs towards group-level goals will inevitably suffer occasional breakdowns; cancer is in a sense, the cost of doing business as a metazoan [51]. A related issue




Figure 1. Morphogenetic field underlies large-scale pattern control and remodeling. (A) Substructures and cells in an organism *in vivo* continuously receive information from the entire body; this information makes up the morphogenetic field (mediated by gradients of secreted factors, physical forces (stresses and strains), bioelectric states, and extracellular matrix components). The morphogenetic field is what allows the coordination of cell activities toward correct patterning during embryogenesis and regeneration, and what allows structures to resist dysmorphologies such as cancer and aging. (B) Amphibian models reveal large-scale (non-local) aspects of the morphogenetic field: tails transplanted to the flank slowly remodel into limbs—structures more appropriate to that anatomical location. Importantly, even the tail tip cells (whose local environment, a tail, is correct) remodel into fingers, illustrating the integrated large-scale anatomical surveillance and remodeling of pattern homeostasis. (C) The view of cancer as a disease of geometry is supported by data showing that when provided with strong patterning cues (in this case, when implanted into a developing embryo), the cells contribute to normal morphogenesis, not tumors. (D) The same is true in regeneration, where initiation of regenerative patterning of a new limb is able to reprogram limb tumors in that region. Artwork in panels (B)–(D) produced by Jeremy Guay of Peregrine Creative. Panel (A) used with permission from [478] John Wiley & Sons. © 2013 Wiley Periodicals, Inc. Panel (B) reproduced from [182] with permission of The Royal Society of Chemistry.

for future investigation is: what does a unicellular organism's cancer look like? Is it possible for a paramecium to enter a cancer state? The SMT model might predict novel phenotypes (increased proliferation rate? increased motility?) after DNA damage to typical tumor suppressor genes. The TOFT model would instead predict that default unicellular behavior is the cancer phenotype, and that further cancerization of a unicellular animal might be impossible (although it might predict interesting effects of TOFT-inspired cancer therapies upon normal unicellular organisms).

The antithesis of cancer is the remarkable fidelity of pattern homeostasis observed in biological organisms—maintenance of the same body over hundreds of years in some long-lived species. How embryos, which begin as one stem cell (the fertilized egg), reliably build a complex stereotypic anatomy is a major unanswered puzzle, but it is clear that a complex morphogenetic field of chemical, biomechanical, and bioelectrical signals impinges on cells and provides instructive influence *in vivo* (figure 1(A)). However, development is just the first manifestation of a more fundamental

property of living things: pattern homeostasis [18, 52]. The problem of reliably generating a specific anatomy is difficult enough, but at least the feed-forward development of a zygote into a predictable outcome gives some hope that complexity theory will explain the eventual emergence of a specific structural endpoint. However, beyond forming a single well-defined structure, many biological systems have the ability to *repair* their complex shape if it is damaged in unpredictable ways [53], and remodel themselves to fit specific anatomical criteria (figure 1(B)). Many early embryos, when cut in half, or combined, will reorganize, and give rise to normal organisms [54–56]. Even in adulthood, some organisms (such as salamanders) can regenerate amputated limbs, jaws, hearts, tails, and portions of the brain [57, 58]. Masters of regeneration, such as planaria [59, 60], have also solved the aging problem—achieving body-wide immortality through continuous regeneration (while individual cells senesce and die). Importantly, planaria reproduce by fission, bypassing the Weissmann barrier [61] and accumulating mutation at a high rate [62], despite a remarkably rock-solid developmental and regenerative morphology.

The problem before us is that of closed loop morphostasis: maintaining a complex shape despite deformations that cannot be known in advance [63]. How do systems detect when their correct shape has been altered by injury or disease, compute what steps to take to restore their correct target morphology, and decide when to stop growing (after their anatomical goal has been reached)? Regenerating animals rebuild precisely what is missing—no more, no less—in the face of external injuries they could not have anticipated. When half of a planarian's head is removed, a perfect match is restored, of the right size and orientation, and then growth stops. Such systems process information about their current state and its deviation from the target morphology, perform computations about what should be done next, and make decisions about what to grow and when to stop [64, 65]. Importantly, these decisions are global. When planaria are bisected, the two blastemas form a head and tail, respectively, even though the cells on either side of the cut were adjacent neighbors. A tail grafted to the flank of a salamander slowly remodels to a structure more appropriate to its new location [66], illustrating shape homeostasis towards a normal amphibian body plan (figure 1(B)). Even the tail tip cells (which are in their expected location, at the tip of a tail) slowly become fingers. Similarly, planarian tumors disappear when regeneration is induced even at the opposite end of the animal [67]. This all shows that the remodeling is driven, at least in part, by non-local information.

The common feature in these examples is that individual cell behaviors are guided toward *large-scale anatomical setpoints*—goal states of system-level homeostatic processes that harness cell activity toward a specific morphological outcome. This requires long-range coordination among cells, and significant amounts of signaling that enables sensing of local and long-range states, whether this takes place through a series of neighbor interactions or via a long-range signaling modality. It is easy to see that cells that cannot perceive the morphogenetic field of organizing cues [18, 68–70], or that end up in locales with a local low-point in the morphogenetic field, would be unable to coordinate with others and revert to single-cell behavior. In such a case, cells would proliferate, consume resources, migrate, and evolve, without regard for the whole of the organism, as observed in cancer.

The distinction between cell-level mechanisms and global patterning needs is clearly observed in an old experiment showing a kind of 'implementation independence' of morphogenesis. Kidney tubules in the newt possess a constant size, while cell size can vary drastically with ploidy [71]. The tubule pattern (a macroscopic goal state) can be implemented by diverse underlying molecular mechanisms such as cell-to-cell interactions (when there are many small cells) or cytoskeletal bending that curls one cell around itself, to make a tubule (when cells are very large). Such discordance between macroscopic pattern and underly-

ing molecular mechanisms is one way to conceptualize the often-bewildering lack of one-to-one match between molecular marker states and cancer outcomes. A key puzzle for future research is to clarify how cells can sense metrics not defined at the individual cell level (e.g. size, position, orientation, and symmetry of a given organ), and how these macroscopic states can serve functional determinants of phenotype. A quantitative view of systems in which global states have causal power and dominate the functional roles of molecular states is discussed below in 5.

3.2. Cancer as a disease of anatomical homeostasis

Importantly, cancer is not an imposition of runaway growth—instead, it is a lifting of normal constraints and order provided by the host. 'No cancer exhibits any trait which cannot be found in some normal tissue as the expression of normal genomic activity; no cancer grows faster than an embryo nor is any cancer cell more invasive than a macrophage nor are cancer cell lines more immortal than are germ lines' [72, 73]. Indeed, because growth and disorder is the default condition, continuous input from the host is required to keep tissue organization. One mechanism used to coordinate cell activity away from cancer is communication via the nervous system. Tumors are readily induced by denervation in salivary organ and alimentary canal in cockroach [74, 75] and in mammalian skin [76]. Similarly, tumors are chemically induced more easily in denervated rabbit ears as compared with contralateral controls bearing normal innervation [76]; the same has been observed in sarcomas implanted into normal or denervated frog limbs [77]. The ability of the host to regulate local cell activity in accordance to global patterning needs can be generalized as the influence of a '*morphogenetic field*'—the sum total of signaling cues impinging upon all cells *in vivo*.

Cancerous failure of morphostasis can occur because a morphogenetic field is missing, altered, or not successfully perceived (all three of which can occur due to genetic or physiological state change). Cells in dispersed monolayer culture are several orders of magnitude more sensitive to chemical carcinogenesis than are organized tissues within an intact organism [78], and placing normal primary mammalian cells in culture results in the appearance of cells with malignant potential [78–81]. Chick embryos infected with the v-Src virus exhibit no malignant phenotype, but the same cells in culture undergo massive transformation [82]. A number of recent papers stress the suppressive nature of signals from neighboring tissues [83–86]. Consistently, re-establishing appropriate interactions of human cancer cells with the microenvironment and normal neighbors underlies the observed reversion of malignant phenotype in a number of cell lines [87–89].

Consistent with the need for cell-to-cell interactions in suppressing cancer are data showing that tumorigenesis is promoted when cells are isolated from

their neighbors (and thus from the morphogenetic guidance they would otherwise receive) by physical barriers. Implanting into connective tissue of the rat rectangles of inert plastic, metal foil, or glass coverslips induces sarcomas when the material is $>1\text{ cm}^2$. If the material is perforated, the incidence is reduced, and the effect does not occur with powders of the same material (which actually increases surface area, ruling out chemical induction or genetic damage mechanisms) [90–93]. These kind of data clearly underscore the importance of environment [94], not only intrinsic state.

More recent data have focused attention on interruption of cell-to-cell communication via ions and other small molecules through electrical synapses known as gap junctions (GJs)—aqueous channels made of connexin proteins that allow physiological signaling directly between the cytoplasmic interiors of docking adjacent cells [95–100]. For example, Connexin32-deficient mice have a 25-fold increased incidence of spontaneous liver tumors [101]. Gap junctional isolation is known to be a tumor-promoting agent [100, 102–107], although there are counterexamples [108]. Active GJ communication allows cells to make sophisticated decisions comparing relative levels of specific compounds between themselves and their neighbors [109] and thus can underlie the transmission of physiological patterning signals [110–120].

Anticipating recent discoveries of the importance of gap-junction cell-to-cell communication for planarian regenerative patterning [119, 121], in 1965 Seiler-Aspang described planarian experiments in which a carcinogen led to formation of many head teratomas with irregular nerves and un-oriented eyes, concluding that ‘the *cell-isolating* action of the carcinogen prevents formation of a single morphogenetic field and leads to the establishment of several separated fields of reduced dimensions’ [67]. Thus, tumors could also represent establishment of a local ‘subfield’—a fragmentation of the host’s morphogenetic field such that integration with the host body plan is lost. Unlike normal somatic tissues, which remodel when transplanted into foreign locations [66, 122, 123], the histopathological structure of metastasis reflects the tissue of origin, not of their destination [16], confirming an inability to respond to neighboring signals such as positional information and remodeling cues.

One of the key predictions of a view of cancer as a disease of geometry is that it should be reversible. If it results from the lack of morphogenetic influences upon cells, then contexts in which very strong patterning cues are being imposed should have the power to reverse or reprogram neoplasm. The morphogenetic field ought to be the most active and accessible during embryogenesis. It is thus not surprising that despite considerable malignancy and aneuploidy, tumor cells introduced into wild-type embryos become integrated (figure 1(C)) as normal tissue [124–134]. Human metastatic melanoma cells injected into zebrafish embryos

acquire a non-neoplastic phenotype, but form tumors when injected into zebrafish *after* organogenesis [135, 136]. Likewise, implanted sarcoma progressed in 80% of adult rats but only in 6.4% of rat embryos. Similar data have been recently shown for chick and other kinds of embryos that are able to tame aggressive cancer cells when these are implanted [136–139]. Cancer normalization can occur cell-autonomously [140], or induced by communication from other cells, such as the mammary stroma [124, 141–146]. Indeed the embryonic field present in the blastocyst can normalize several types of cancer cells including those isolated from embryonic carcinoma, leukemia and neuroblastoma [134], although the limits of this normalization process (with respect to large-scale chromosome aberrations found in some tumors) remain to be probed fully. Thus, active patterning signals can normalize cancer (over-ride genetic defects and reboot cell behavior programs); this is a finding that is not predicted by the cell-level view of cancer as embryos have high levels of many growth factors that could be expected to potentiate tumor growth (and do, in experimental contexts such as cell culture which is devoid of large-scale patterning structure).

It has been long known that regeneration and cancer are closely related [147–152]. Highly-regenerative organisms are resistant to carcinogenesis and indeed activating regenerative response (figure 1(D)) can normalize existing tumors [147–150, 152–156], although this does not always occur [157]. The inverse relationship between regeneration and cancer susceptibility [72, 158] is more compatible with the importance of morphogenetic field guidance than with cancer risk associated with the presence of highly-active, undifferentiated cells [68]. Mammalian liver regeneration can overcome cancer—early nodules initiated by carcinogens are remodeled to normal-appearing liver [159, 160], hepatocarcinoma cells can be normalized by injection into wild-type liver [161, 162], and over 95% of nascent tumor sites remodel into normal tissue by the highly-regenerative liver [163–165]. Clinically however, this capability is tempered by the fact that many patients have a damaged liver (Hepatitis C or other burden which induces cirrhosis and may prevent regenerative response). In zebrafish brain regeneration, a remarkable degree of aneuploidy does not lead to cancer, while amphibian limb regeneration can likewise normalize tumors [6, 151, 166], showing that in some cases, an active patterning program can trump chromosomal damage [167].

What kind of data are conveyed by the morphogenetic field? Information conveyed by the morphogenetic field includes positional coordinates. It has long been known that mismatches in positional identity, in grafting experiments, causes supernumerary structures to form [168–170]—the so-called ‘rule of normal neighbors’, in which tissues grow as needed to intercalate new structures to keep continuous positional information gradients. In the cancer field, this

is observed in differences of anterior-posterior axial positions to cancer susceptibility, and in the ability of tumors to form from the juxtaposition of normal but ectopic tissues. Tumors grow on posterior regions of *Triturus* less readily than they do on anterior regions [67], and numerous such differences are observed in human tumors as well [171–176]. Despite lack of DNA damage or cytotoxic chemical stressor, transplantation of rat testis to the spleen induces formation of interstitial cell tumors [177], while normal rat ovary tissue put into normal rat spleen results in malignant neoplasm [178]. Implantation of early embryos (which organize their own field of signals) under the kidney capsule of an adult makes transplantable malignant teratomas despite a lack of any infective, chemical, or radiation initiator to cause genetic damage [16], while normal adult *Xenopus* kidney implanted in the non-amputated forelimbs of recently-metamorphosed larvae will make lymphosarcomas as well as accessory limb structures [179]. Implantation of mouse embryos into adults causes teratocarcinomas [180], possibly due to an interference between the host and implanted embryo's morphogenetic field signals.

The morphogenetic field is mediated by a number of physical modalities, including the well-known gradients of chemical morphogens (secreted factors), extracellular matrix, physical forces (tension and compression), and bioelectric cell states. Addressing cancer from the perspective of large-scale coordination cues requires familiarity with the properties of mechanisms that underlie long-range pattern control. Non-neural bioelectricity is precisely such a set of mechanisms, and is increasingly implicated in cancer by molecular studies [181]. Most importantly, the known ability of bioelectric networks to process global information and implement networks with multiple causal layers (e.g. goal-directed activity mediated by brain bioelectric circuits) make it an ideal nexus from which to understand and manipulate cancer as an error of global organization [20, 182]. Thus, in prelude to the discussion of mathematical theory germane to top-down approaches to cancer, we first review a set of important emerging mechanisms that provide plentiful fodder—both quantitative and conceptual data—for computational analyses to the control of cancer as a defection from normal developmental cues.

3.3. Bioelectric regulators of growth and form: a new locus of intervention in cancer

As in the brain, somatic endogenous ion currents, plasma membrane resting potentials (V_{mem}), and electric fields are produced by the activity of ion channel and pump proteins across cell membranes. One difference from action potentials is their temporal dynamics, which are slower and continuous (not discrete millisecond spiking activity as in the CNS). The tissue-wide gradients thus produced, and the networks of endogenous bioelectric states present across many cell types (figure 2) serve as instructive determinants


of cell behavior [183, 184], and—more importantly, of collective dynamics during pattern regulation. Due to length limitations, here we omit discussion of other biophysical phenomena including transepithelial electric fields [185, 186], ultraweak photon emission [187, 188], and coherent AC electromagnetic fields [189, 190], all of which also have considerable relevance for the cancer problem. Likewise, the many interesting studies of applied field effects in the cancer context are discussed in several excellent recent reviews [191–194]. Here we focus on distributions of V_{mem} or membrane potential [195], as a mediator of information processing in cellular networks. Due to their central role in pattern homeostasis, these bioelectric patterns are both an important target for neoplastic dysregulation and biomedical strategies seeking to restore appropriate dynamics between cells and body-wide patterning fields.

3.4. Bioelectric states control cell behavior

Bioelectric properties of cells and the electrical states of cells in the microenvironment control several key behaviors of relevance to cancer (reviewed in [186, 196–202]). For example, electric fields generated by ion pumping across epithelia serve as migration cues for cellular galvanotaxis [203–207]—an important guidance modality for cell movement within the host. Cell shape changes, such as increased arborization, are also driven by endogenous electric fields and changes in V_{mem} [208–212]. Together, migration and shape properties are key elements of successful metastasis [213, 214].

Resting potentials determine differentiation state and proliferation; generally, a depolarized state is indicative of plastic, undifferentiated cells (e.g. stem cells), while differentiation is caused by increase of negative V_{mem} [215]. Functional control of cell state by changes in V_{mem} has been observed in many kinds of stem and progenitor cells [216–226], including adult human mesenchymal stem cells [222, 227], which can be kept stem-like despite the presence of chemical differentiation factors by forced depolarization, and in induced pluripotent stem cells [228]. Even mature CNS neurons can be made to re-enter mitosis by sustained depolarization [229, 230], revealing the power of transmembrane potential to regulate proliferative potential in adult somatic cells.

Changes in V_{mem} of cells, such as cancer-associated depolarization, can trigger transcriptional changes by (1) regulating the movement of morphogens such as serotonin, calcium, and inositol triphosphate through gap junctions [116, 117, 231–235], (2) controlling the import/export of small signaling molecules such as butyrate across membrane exchangers [208, 235–238], and (3) modulating the activity level of phosphatases such as PTEN [239–242]. Together, these transduction mechanisms convert an essentially biophysical state change into secondary messenger events that impact on transcriptional and epigenetic regulation of loci



such as NODAL or SLUG which are important intermediate effectors for the cancer phenotype [238, 243].

3.5. Spatio-temporal gradients of V_{mem} are instructive cues maintaining healthy anatomy

Importantly, relatively recent development of state-of-the-art tools for the detection and experimental manipulation of biophysical signals in multicellular patterning contexts [244–247] has revealed how distributions of voltage gradients mediate positional information, organ identity of large cell groups, and initiation signals for complex developmental modules such as tail or limb regeneration. Using a combination of fluorescent voltage-reporter dyes to characterize spatial V_{mem} distributions and functional studies using targeted misexpression of well-characterized ion transporters to specifically modify those gradients *in vivo*, instructive signaling roles of transmembrane voltage gradients have been identified in embryogenesis and regeneration, adding to the list of such roles identified in classical work that utilized functional physiology [248, 249]. A number of recent molecular studies using unbiased approaches in human syndromes and non-human model systems have identified a range of ion channels, gap junctions, and ion pumps in developmental and regenerative morphogenesis of the face, brain, heart, appendages,

growth of the cerebellum [250–258] and many other structures [259–271]. These physiological states are ideal inputs for the computational analysis delineated in the next section.

During early frog development, the redistribution of maternal potassium channels and proton pumps in early blastomeres results in a V_{mem} gradient that determines asymmetric expression of key transcription factors [243, 272, 273]. Later, during craniofacial patterning, the positions of the eyes [274] and other elements of the face [275] are determined by a regionalization of naïve ectoderm into distinct domains of hyperpolarized cells—a bioelectric pre-pattern or scaffold for the nascent structures. These voltage gradients regulate the expression of genes like *Frizzled*, and artificially altering this pattern by misexpression of specific ion channels and pumps is sufficient to perturb normal craniofacial anatomy and to reprogram tissues far away from the head to form properly-patterned eyes [274]. Importantly, in such cases, as in the cancer phenotypes discussed below, it is really the V_{mem} that is the necessary and sufficient factor for inducing specific shape change—it does not matter which ion translocator protein is involved, or what ion species is used: a given voltage change, no matter how it is produced, activates specific downstream events.

During regeneration of flatworms, the patterning activity of adult stem cells (neoblasts) is regulated by gap junctional connectivity and a set of proton and potassium flows [119, 121, 276]. By regulation of apoptotic remodeling and downstream activity of genes such as *Wnt11* [277], the physiological gradient determines the anatomy of the organs built after injury. In vertebrates, where electric fields were long ago implicated in limb regeneration [278–285], recent experiments showed that driving proton and sodium fluxes can force complete tail [286, 287] or limb [288] regeneration in a range of non-regenerative conditions. The mechanisms involve guidance of innervation into the stump, activation of blastema genes such as *MSX1*, *Notch*, *Delta*, *BMP2*, and *BMP4*, and induction of cell proliferation in the wound mesenchyme.

With respect to wound healing, inhibition is known to be a tumor promoting agent [289–291], elegant molecular genetic experiments have now revealed some of the elements underlying endogenous electric field-mediated cell migrations. Epithelial wound closure involves Integrin Beta-4 (ITGB4), Cyclic AMP, betaphosphatidylinositol-3-OH kinase- γ (PI(3)K γ) and phosphatase and tensin homolog (PTEN) [197, 292–295]. Having seen that endogenous electric fields and V_{mem} gradients play an instructive role in normal patterning, what is the evidence that dysregulation of bioelectrical communication can underlie the cancer phenotype?

3.6. Ion channels are oncogenes and important drug targets

The view that cancer is a developmental disorder predicts that molecular mechanisms known to be important mediators of the morphogenetic field would be involved in tumorigenesis. Indeed, there is mounting evidence that the bioelectric cues that establish normal pattern can go awry and result in cancerous growth (reviewed in [199, 296, 297]). The function of ion channels is involved in the self-sufficiency in growth signals, insensitivity to anti-growth signals, evasion of programmed cell death, limitless replicative potential, sustained angiogenesis, and tissue invasion and metastasis [200]. Ion channels, pumps, and gap junctions are now recognized as oncogenes [199], predictive markers [200], and an important set of targets for new cancer drugs that seek to modulate cell behavior by tweaking electrical controls of proliferation or metastatic behavior [298]. Importantly however, onco-channel misregulation occurs not only through mutations in channel genes but also by changes in the rich network of events that implement post-translational gating of wild-type ion channels.

Tumor cells differ from untransformed cells in terms of the type of ion channels and pumps they express and in the resulting membrane potential of the cells [299–308]. Some channel levels are thus used as markers, such as the K2P channel TREK-1 and the

sodium channel NaV in prostate cancer [309, 310], and the TRPM1 channel in melanoma [311, 312]. Tumor cells' membrane voltage is often determined by a different transporter than that of normal cells and it has been suggested that this gives the cells a selective advantage [303].

The function of ion translocators, such as voltage-gated K⁺ channels [313, 314] and Cl[−] channels [315], controls the proliferation rate of a number of cells that often form tumors [316–327] or leukemia [328]. ERG is particularly involved in cell growth signals [303, 329–333], and is implicated in transformation of prostate epithelium [334]. Also implicated are 2-pore channels such as KCNK9 [335, 336], and voltage-gated sodium channels, being definitive oncogenes—necessary and sufficient for a transformed phenotype [337]. Metastatic potential correlates with voltage-gated inward sodium current and it has been suggested that some sodium channels may be oncofetal genes, encoding signals that are active during the rapid and autonomous growth of tumors and embryos [310, 338–341].

Migration of cells including B-16 melanoma is dependent on K⁺ channels [342]. The voltage-gated sodium channels (VGSCs) potentiate breast cancer metastasis [341], and indeed the involvement of NaV in the galvanotaxis that allows prostate and breast cancer cells to move across vessel lumens [213, 337, 343–349] is one of the leading stories on ion channels in cancer. Highly up-regulated activities of NaV confers on cancer cells directional motility and invasive characteristics via Ca²⁺ and pH-sensitive cytoskeletal remodeling processes which facilitate metastasis [337, 341]. Certain channelopathies result in syndromes associated with cancer such as the lung cancer seen in Lambert–Eaton syndrome [350], and the tumors present in Beckwith–Wiedemann syndrome, which is caused by abnormal imprinting of a voltage-gated potassium channel [351–353].

The involvement of ion channels in transformation, growth control, and metastasis has led to efforts to develop potassium, chloride, and sodium channel and pump modulators as clinical agents for ovarian [354], breast [355], and prostate [356] cancer [298, 357]. Unbiased drug screens for inhibitors of cancer stem cells have implicated potassium ionophores such as salinomycin [358], and tetraethylammonium (TEA, a potassium channel blocker), was found to suppress colony formation in endometrial cancer cells while withdrawal of TEA resulted in a significant enhancement of tumorigenesis [359]. Known cancer-fighting agents are now beginning to be recognized as ion channel modulators, for example resveratrol [360–362]. Indeed, drugs that target membrane voltage-generating transporters have already shown clinical promise in cancer [363].

Which ion channels should be targeted by therapeutic drugs? In an important sense, focusing on the channel gene or protein may be missing the bigger picture. In the current literature, ion translocators are




Figure 3. Cancer: detecting its bioelectric signature and suppressing tumors by managing bioelectric state. Microinjection of human oncogene mRNA (A) induces tumor-like structures (ITLS) in *Xenopus* embryos (A'). Soaking in voltage-reporting fluorescent dye reveals the voltage depolarization that occurs at the earliest stages of tumorigenesis (B). Remarkably, tumors normally induced by Xrel3 fused to a fluorescent red protein (C) can be suppressed by misexpression of hyperpolarizing channels such as Kv1.5, even when the hyperpolarizing region is at a considerable distance from the oncogene (visualized by its red fluorescence). Panels C and D reproduced from [398]. CC BY-NC 3.0.

usually treated as single genes or proteins responsible for a specific cell behavior (metastasis, hyperproliferation, etc)—a cell-level view that neglects the fact that numerous channels and pumps contribute to the V_{mem} gradients that mediate large-scale patterning cues [202, 364]. The true impact of bioelectricity in cancer will only occur when we understand and target the storage of patterning information in *physiological* networks that is misprocessed in cancer [365, 366]; such networks are dynamical systems with complex feedback between the post-translational gating of many different channel and pump proteins.

3.7. A unique bioelectric signature reveals incipient cancer

A variety of bioelectric properties have been used as detection modalities for tumors; these capitalize on cancer cells' distinct electrical impedance [367–381] or ion content [382, 383]. Zeta potential is also associated with cancer; for example asbestos fibers and sheets of positively-charged materials (but not powders of the same material) induce tumors, probably by acting as a capacitor for bioelectric potential, the positive side corresponding to the electron sink existing at a wound (reviewed in [384]). In this section, we focus on depolarized V_{mem} , which has been suggested to correspond to the cancer state [215, 385–387].

One way to probe the physiology of the effects of canonical mammalian oncogenes (Gli1, Xrel3 and KRASG12D) and a mutant tumor suppressors ($p53^{\text{Trp}248}$) *in vivo* is to misexpress them in *Xenopus* and zebrafish embryos [388–391], which induces the formation of tumor like structures (ITLS). ITLS's thus form as a result of genetic interference with signaling pathways altered in several cancer types including basal cell carcinoma, lung cancer, leukemia, melanoma, and rhabdomyosarcoma [392–395]. Examination of injected animals using fluorescence reporters of V_{mem} [396] revealed unique depolarization of tumors (and increased sodium content) (figures 3(A) and (B)) compared to healthy surrounding tissues [237, 397]. Moreover, depolarization foci are present in oncogene-expressing, pre-neoplastic cells that are yet to undergo transformation or show any morphological phenotype. Such depolarized foci, while present in only 19–30% of oncogene-injected embryos (depending on oncogene used), predict tumor formation with 50–56% success rate (15–21% false negatives). For comparison, prostate specific antigen (PSA) level in the serum, when used as a biomarker for prostate cancer, has ~29% predictive value [399, 400].

The existing data reveal local bioelectric disruptions that correlate with incipient cancer and could be exploited as a diagnostic modality for prediction of

tumor sites and margins. In addition, it is imperative to establish molecular models in which to investigate the fact that transplanted or chemically-induced tumors can be detected by aberrant voltmeter readings taken at locations far away from the tumor [8, 401–406]. In the meantime, work has progressed to show that voltage states are not merely a marker that correlates with cancer, but are a functional driver of the process. These bioelectric readings, together with the information analyses discussed in sections 4 and 6, may enable both, highly predictive new cancer biomarkers, and a unique description of the functional aspects of biophysical regulators of cancer progression.

3.8. Bioelectric modulation can induce global metastatic phenotype

Can bioelectric disruption alone, in the absence of classic carcinogens or mutagenic compounds, induce a cancer phenotype? Genetic or pharmacological targeting of a specific cell population (sparse but ubiquitous cells expressing the GlyR channel) in an amphibian model revealed a remarkable phenotype: over-proliferation, increased migration, and drastic arborization of melanocytes (pigment cells). By transiently depolarizing GlyR-expressing cells *in vivo*, an entirely different cell type (melanocytes) underwent a metastatic-like conversion, turning on expression of genes such as Sox10 and SLUG [209]. The melanocytes acquired a dendritic morphology, upregulated mitotic activity, and invaded blood vessels and soft tissues like the neural tube lumen and brain. In addition to melanocytes, disorganization and ectopic growth of blood vessels was also observed [397], but otherwise the tadpoles were remarkably normal in terms of overall growth and development. Importantly, the same exact effect was induced by *any* method of depolarization, including by the movement of chloride, sodium, potassium, or hydrogen ions—truly an effect initiated by V_{mem} depolarization, not any specific gene product or chemical ion species, and a metastatic phenotype induced by a purely physiological perturbation.

Most interesting was the non-cell-autonomous nature of the effect: the cells that acquired a melanoma-like phenotype were not the cells whose resting potential was changed (the GlyR-bearing cells were thus called ‘instructor cells’) [208, 209]. Indeed, only a small number of instructor cells had to be depolarized in order to induce the hyperpigmentation phenotype (which is all-or-none within any individual animal). How was the communication between these two cell types mediated? A suppression screen testing the several known methods of transducing voltage change into transcriptional cascades revealed that the serotonin transporter SERT, which powers uptake or efflux of serotonin depending on resting potential, was involved (much like in the bioelectric regulation of left-right patterning); blockade of SERT could rescue the hyperpigmentation effect, and direct applica-

tion of serotonin could trigger a similar phenotype. This pathway, comprising bioelectric dynamics and downstream neurotransmitter responses (all outside the nervous system) has now been extensively analyzed by machine-learning approaches to assist human scientists in rationally modifying its complex dynamics *in vivo* [397, 407, 408].

3.9. Hyperpolarization inhibits oncogene-induced tumorigenesis

The above data show that depolarization of V_{mem} can, itself, activate a metastatic-like phenotype. What role could V_{mem} play in carcinogenesis induced by genetic perturbation? Well-characterized channels such as inward rectifying potassium channel (Kir 4.1) and constitutively open glycine-gated chloride channel mutant (GlyRF99A) can be used to generate strong hyperpolarizing currents [409, 410]. When GlyRF99A was co-injected with the Xrel3 oncogene in *Xenopus* larvae [237], it significantly suppressed the incidence of ITLS formation induced by oncogene alone (figures 3(C) and (D)). Florescent tags on the oncogene protein revealed that hyperpolarization could prevent the formation of tumor-like structures despite very robust expression of oncogene in cells. The use of several different hyperpolarizing channels based on Cl^- and K^+ confirmed that suppression of neoplastic transformation was due to V_{mem} hyperpolarization, as opposed to ion-specific or scaffolding functions of the ion channel proteins. Consistent with this, data in rats showed that the ion channel modulator drug ivermectin can likewise modulate the effectiveness of carcinogenic compounds [411].

More recent work has used light to trigger optogenetic ion channels in prevention and normalization of human oncogene-induced tumors [412]. Crucially, a dissection of the spatial relationship between the cells expressing the oncogene and the cells expressing hyperpolarizing channels revealed *long-range* signaling mediated by electrical synapses known as gap junctions, and bacterial-derived butyrate [413]. Thus, whether or not a tumor formed on one side of the animal depended on whether or not cells on the opposite sides were connected electrically, demonstrating the importance of distributed decision-making in the appearance of tumors at a given site and the importance of not restricting attention to the tumor site itself.

These results must be translated into mammalian models, and tested in a variety of different types of cancer in order to determine in what cases mutational burden has to be addressed before cure, and in which cases patterning signals might be sufficient to permanently solve the problem regardless of the DNA state. A crucial component of this effort is to understand the information-bearing signals that keep cells, both normal and mutant, under control [7, 414]. Study of these signals must take place both at the bench (unraveling the biochemical and biophysical mediators of pattern




Figure 4. A toy example of how to apply active information or transfer entropy. (a) V_{mem} time series data for two cells X and Y. (b) The time series is binned per some specification. While continuous time series can be handled in principle, binning is more common. (c) The binned time series is broken up into histories, $x_n^{(k)}$ and y_n , and futures, x_{n+1} , and used to estimate probabilities, e.g. $p(x_n^{(k)}, x_{n+1})$, $p(x_n^{(k)}, y_n, x_{n+1})$, etc... (d) The active information and transfer entropy are then computed based on the probabilities.

control mechanisms) and *in silico* (establishing rigorous analytical frameworks for quantifying and decoding specific control policies operating in cell collectives). Advances in this area will also help identify key tipping points in physiological and genetic state space that reveal when a system can and cannot be brought back to normal by specific interventions. Having seen the relevance of patterned bioelectric states for both normal development and cancer, we next describe the information dynamics approach to information theory as a quantitative toolset for understanding information processing in development and cancer.

4. Information dynamics

Information dynamics is an extension of information theory that permits quantifying the component operations of computation within dynamical systems by analyzing their time series data [415]. It has utility in decoding the ‘logic of life’ [416] as one can make direct analogy with the more intuitive computational concepts of information storage and processing. In information dynamics, these concepts are quantified by active information (AI) and transfer entropy (TE), respectively. Figure 4 provides a schematic description of how to apply AI and TE to time series data, e.g. V_{mem} for two cell membranes.

4.1. Information storage

In an isolated mechanical system with no noise, past states are good predictors of future states. However, in the messy and noisy cellular environment, the past states of a process within a cellular network, or an individual cell within a tissue, will in general be

inadequate for guiding prediction of future states due to strong nonlinear coupling between cellular components. Local active information storage (LAIS) quantifies the degree to which uncertainty about the future of an agent X is reduced by knowledge of its own past states [417]. Formally, local active information storage for an agent X is the local (unaveraged) mutual information between its semi-infinite past $x_n^{(k)}$ (as $k \rightarrow \infty$) and its next state x_{n+1} at time $t + 1$:

$$a_X(n+1) = \lim_{k \rightarrow \infty} \log \frac{p(x_n^{(k)}, x_{n+1})}{p(x_n^{(k)})p(x_{n+1})}, \quad (3)$$

where $x_n^{(k)} = \{x_{n-k+1}, x_{n-k+2}, \dots, x_n\}$ is a vector of k previous states of X. For real-world data, time series are finite, and local active information is often approximated by defining a history length k shorter than the dynamical time scales of interest. Averaging equation (3) gives the *active information* for a dynamical process [417]:

$$A_X(k) = \sum_{n=k}^{N-1} p(x_n^{(k)}, x_{n+1}) \log \frac{p(x_n^{(k)}, x_{n+1})}{p(x_n^{(k)})p(x_{n+1})}. \quad (4)$$

Average and local active information have been used to understand information storage in biological systems as diverse as swarm behavior [418–420], gene regulatory networks [421], neural information processing [40, 41], and to evolve artificial information processing systems [415]. Application to neural information processing is particularly relevant to the cancer problem since the measurements of voltage potential tracking V_{mem} across populations of cells is similar in data content to that acquired by fMRI of brain tissue. In particular, LAIS has been applied to

voltage sensitive dye imaging from brain tissue from area 18 of the cat visual cortex [40]. LAIS was shown to reflect functional properties, such as the response to stimulus, and the surprise on change of stimulus (despite the locally random nature of the stimulus).

As discussed in section 3.4, tumor cells' V_{mem} is typically determined by different transporters than normal cells [303] and no single channel dominates the global dynamics of large-scale patterning. Applied to physiological networks of cell cultures containing cancer, LAIS might separate cells with the cancer phenotype from healthy cells. It could be used to identify functional interfaces of the intra and inter-cellular interactions of cancer and normal cells, given that functional control of cell state changes is known to be driven by changes in V_{mem} [215]. Since healthy cells are expected to act 'more collectively' within a tissue we might expect LAIS to be, on average, lower for healthy cells than for cancerous cells that have gone rogue and act independently. For cancer, an interesting question is whether this information-theoretic approach would pick out behavior that was largely uncorrelated (e.g. random behavior of the cancer phenotype) or if cancer cells are predictable based on their own local past history.

4.2. Information processing

Information processing in information dynamics is quantified using Schreiber's transfer entropy (TE) [422], a directional measure of information transfer between random variables. The information transferred from a source node Y to a target node X may be defined as the reduction in uncertainty provided by the source about the next state of the target, above the reduction in uncertainty due to knowledge of the past of the target. Formally, TE from Y to X is the mutual information between the previous state of the source y_n , and the next state of the target x_{n+1} , conditioned on k previous states of target, $x_n^{(k)}$:


$$T_{Y \rightarrow X}(k) = \sum_{n=k}^{N-1} p(x_n^{(k)}, y_n, x_{n+1}) \log \frac{p(x_{n+1}|x_n^{(k)}, y_n)}{p(x_{n+1}|x_n^{(k)})}. \quad (5)$$

TE naturally exhibits an asymmetry between the source and target nodes. This asymmetry lends a directionality to the measure, and as such TE can be utilized to measure 'information flows', which cannot be captured by symmetric measures such as mutual information [422]. One can view the transfer entropy as conditional mutual information between a target and source, conditioned on the past states of the target. The basic idea is that the prediction of the next value for a given variable at step n , given knowledge of its past k states, could be improved by taking account of the state of other variables at step n . If that is the case, then in some sense information has been transferred between the two variables since there is a directed correlation. If the correlation is high, it implies that knowledge of the other source variable leads to better prediction of the state of the destination variable at step n .

In neuroscience, transfer entropy is used to uncover effective connectivity [423, 424]. It has been used to evaluate information transfer between auditory cortical neurons [42], and on simulated data for cortical interactions [425], electroencephalogram (EEG) data [426], and fMRI data [427]. In the case of cancer, TE could be used to determine effective connectivity of cancer and healthy tissue, and infer if these represent distinct patterns in information flow, related to whether the behavior of cancer cells is somehow coordinated (non-zero TE) within the population of cancer cells, or if instead behavior of cancer cells is coordinated with the surrounding healthy tissue. Multivariate analysis can be implemented to infer additional structure by assuming that X and Y in equation (5) no longer represent individual elements in a system, but instead are collections of sets of elements. For example, one could measure the directional information transfer between a tumor and surrounding tissue, or between cell types with different phenotypes. Inter-regional information transfer has, for example, been measured in neural fMRI data [428] to infer the informational structure of visuo-motor tracking in the brain. Importantly, metrics from computational neuroscience have begun to be applied to the problem of developmental/regenerative patterning [65, 182, 429], and the emerging field of primitive cognition suggests a research program to understand the information processing/computational capacity of tumor cells versus normal cells, and tumors versus normal organs.

4.3. Information processing, storage and drug targets in gene regulatory networks

Recently, information dynamics analysis was utilized to study the informational structure of gene regulatory networks using Boolean models [421, 430, 431]. Both active information and transfer entropy were applied to study cell cycle regulation of the fission yeast *Schizosaccharomyces Pombe* [421]. In a Boolean network model, genes are represented by nodes (or equivalently in this abstract representation, their protein product) and can take on a value of '1' or '0' corresponding to whether the gene is expressed or not, respectively [432]. In the case of the fission yeast cell cycle Boolean model, the network consists of nine nodes, shown in figure 5(A). The biologically functional attractor, corresponding to the stationary G1 resting phase of the cell, dominates the attractor landscape: for this network 73% of all states will eventually converge to this single state [433]. The four nodes highlighted in figure 5(A) are identified as a 'control kernel' that regulates the function of the entire network. Pinning the state of control nodes to their value in the primary attractor corresponding to the G1 resting phase increases this to 100% convergence, such that every possible trajectory terminates on the primary attractor. Control kernels were also discovered in a number of other Boolean network models for gene regulatory networks [434]. A survey of the control



nodes of these regulatory networks reveals that drug targets are statistically overrepresented [434].

Interestingly, the active information and transfer entropy also indicate a special role for control nodes in the correlational structure of the fission yeast cell cycle network. A scaling relation for TE between pairs of nodes was compared to the TE scaling of Erdős-Rényi random networks and scale-free random networks by Kim *et al* [421], shown in figure 5(B). The analysis shows that the biological network is quite distinct from random in terms of amount of ‘information processed’ as quantified by TE. Important for discussion here is that the most distinctive regime for the real fission yeast model ($> 2\sigma$ deviation from either the Erdős-Rényi or scale-free random networks, highlighted between red dashed lines in figure 5(B)) is dominated by information transfer either to control nodes from the rest of the network, or from the rest of the network to control nodes (see [421], figure 5). That is, the control nodes act as hubs for information processing in the network. They additionally play a dominant role in information storage, with higher values of active information than the majority of other nodes in the network [430]. While this provides only one case study, it opens the possibility that informational analyses of time series data of gene expression might provide a non-interventionist approach to identifying drug targets that regulate cellular behavior. Much work remains must be done to detail how information is organized through the pattern of gene-expression states in both healthy and cancerous cells, and in turn whether this kind of approach can indeed provide a novel platform for identifying targets for therapy.

Cancer involves processes guided by changes in gradients of membrane potential (V_{mem}) and also by gene

regulatory networks within cells. Part of the power of information-based approaches is that they are agnostic to the details of the problem to which they are applied. Thus, information-based approaches could provide a unified framework for addressing the onset of and progression of cancer as it occurs both within a cell through changes to the attractor landscapes of the gene regulatory networks of cancer cells [435] and to the neural-like pattern memories encoded as attractors in the state space of bioelectric networks. Analyzing multiple levels of information dynamics in tandem could provide greater insights into the complexity of the cancer disease.

5. Global physiological dynamics underlie cancer: information, not molecules

‘For those who believe in the simplification and rationalization of the cancer process, the actual course of research on the molecular basis of cancer has been largely disappointing. Rather than revealing a small number of genetic and biochemical determinants operating within cancer cells, molecular analyses of human cancers have revealed a bewilderingly complex array of such factors’ [436]. It is now appreciated that the essence of cancer may not be in specific driver genes but in the dynamics of cells traversing state spaces and shifting between different attractors [12, 437]. While these state spaces are commonly thought of in terms of transcription (gene-regulatory networks), the data on bioelectricity in cancer suggests that another important concept may be the physiological state space [438].

The existing data in developmental bioelectrics reveal that the decision of any given region to develop

a tumor or not depends on the bioelectric state of far-off regions of the body, and the combined activity of many ion translocators that interact with, and partially compensate for, each other to establish a specific spatio-temporal voltage profile. Thus, predicting and normalizing cancer may be best achieved not by local, gene-focused approaches, but rather a focus on information (which bioelectrical networks are superbly suited for processing). What do cells measure, remember, and use for decision-making as they shift between normal and neoplastic state?

One way to think about cancer is as a shift of the border between self and environment [439, 440]—a contraction of the functional integration of the self-organizing system [441, 442]. In contrast to normal cells, which work for the good of a macroscopic host (the combined ‘self’), cancer cells have reduced their computational boundary to their own surface: to each cancer cell, the whole body is just ‘the environment’, at the expense of which all living things reproduce and survive. Cancer may be not be an increase of selfishness, but rather an unchanged self-interest coupled to a shift (reduction) of the internal self-model [65, 443–445]. This view challenges us to explore models that rely not on selfishness-based evolutionary dynamics but rather on the computational aspects of what cells and cell collectives measure and how they form internal models of self and environment [445–447].

It is also possible that the right level of description for some tumors is not the single cancer cell but the tumor, which may embody not only a *loss* of normal organ patterning dominating individual cell dynamics, but also a coherent, goal-seeking subsystem that exists outside the morphospace of healthy somatic organs. Tumors may not be just aggregates of replicating selfish neoplastic cells but complex living entities composed of numerous cell types that work together to acquire nutrients, survive, and evade the efforts of an environment that is trying to kill them [448, 449]. Perhaps, a tumor is, in some practical sense, an independent colonial organism [450] with its own (primitive) morphogenetic field, and a self-boundary that is more than unicellular but reduced from the whole body to that of a much smaller structure. Such a view is suggested by a number of findings. First, histological analysis indicates that tumors can indeed be regarded as complex tissues with a distinct internal organization [79, 451]. Tumors reproduce themselves via metastasis, and execute many adaptive strategies (such as up-regulating multi-drug resistance proteins in the face of chemotherapy) to preserve their homeostasis and existence—just as organisms within an ecological niche do [452–454]. Much like organisms maintaining morphostasis, tumors maintain their identity during massive cell turnover during selection for founder cells resistant to chemotherapy drugs [455]. Recent work describes the highly malignant brain tumor as an ‘opportunistic, self-organizing, and adaptive complex dynamic biosystem’ [456], which is also a great

description of an embryo; proper characterization of the essential principles predictive of the properties of tumor invasion makes uses of concepts such as least resistance, most permission, and highest attraction—these are systems-level, goal-directed elements that are very compatible with the conceptual modeling techniques suggested for understanding embryogenesis and regeneration of whole organisms [20, 182]. Much analysis and experimentation will be needed to reveal whether a tumor and its activity in the body is best modeled as the activity of unicellular organisms, emergent swarm dynamics of the tumor ‘colony’, or a cognitive structure embodied in a kind of group cognition of the tumor ‘organoid’ [457–460].

Next, we consider computational strategies that can be applied to sharpen, quantify, and make rigorous the above-mentioned qualitative arguments about the most effective (causally potent) level of organization at which to attack the cancer problem and the information flow that exists in normal and abnormal morphogenetic field scenarios.

6. Integration and information

We argue in the previous section that cancer may be the result of a retracted notion of ‘self’, a contraction of the functional integration of a self-organizing system of cells [441, 442]. One important aspect is whether a tissue acts collectively as a single entity, or whether distinct regions take-on independent function—it is the latter case that we might expect to give rise to cancer. Information theory offers attractive methods for precisely quantifying such notions of integration. We review two information-theoretic formalisms that attempt to capture various aspects of integration: integrated information theory (IIT) [461, 462] and integrated spatiotemporal patterns (ISTP) [463].

6.1. Integrated information theory

One promising direction is provided by the concept of integrated information. Integrated information theory (IIT) is one of the most widely discussed measures of integration. IIT quantifies how much information a system generates as a whole, above and beyond its individual parts [464]. While IIT was originally developed as a measure of consciousness (which clearly must have some ‘wholeness’), its applicability is broad. We discuss here two measures from IIT: effective information (EI) and *integrated information* (φ). Unlike in information dynamics, both EI and φ characterize entire network *states*, rather than individual nodes, and they do not generally require time series data to compute.

Effective information quantifies the information generated by a system when it enters a state G . More formally, the effective information for each realized state of a system, given by $EI(G)$, is calculated as the relative entropy of the *a posteriori repertoire* with respect to the *a priori repertoire*:

$$\text{EI}(G') = H(p^{\max}(G)) - H(p(G \rightarrow G')). \quad (6)$$

The *a priori repertoire* is defined as the maximum entropy distribution, $p^{\max}(G)$, where all system states are treated as equally likely. The *a posteriori repertoire*, $p(G \rightarrow G')$, is defined as the repertoire of possible states that could have led to the state G' . In other words, $\text{EI}(G')$ measures how much knowledge of how the system behaves reduces the uncertainty about the possible states that might have preceded G' . EI is then used to calculate *integrated information*, which quantifies by how much the ‘sum is more than the parts’. φ is quantified as the information generated by a system when it enters a particular state G' , as compared to sum of information generated independently by its parts. More specifically, φ can be calculated as follows: (1) divide the system entering a state G' into distinctive parts and calculate EI for each part, (2) compute the difference between the sum of EIs from every part and EI of the whole system, (3) repeat the first two steps with all possible partitions. φ is then the minimum difference between EI from the whole system and the sum of EIs for its parts (we refer readers to [461, 462, 464] for more details on calculating φ). If $\varphi(G') > 0$, then the causal structure of network generates more information as a whole, than as a set of independent parts when it enters the state G' . For $\varphi(G') = 0$, there exist connections in the system that can be removed without loss of information. φ can capture how much a system is ‘more than its parts’ suggesting that it is the right sort-of measure for understanding when a collective is really a single entity. In major transitions we might expect systems to become more integrated as the transition proceeds and in cancer we should expect the combined system of healthy and cancerous tissue to become less integrated as the disease progresses.

Integrated information theory has been used to study the structure of the fission yeast cell cycle network, and the network as a whole has been found to maintain integration through the sequence of states corresponding to the phases of the cell cycle [465]. A subgraph of the network in figure 3(A), known as a ‘backbone motif’, was also analyzed [466]. This backbone motif is the minimal sub-network of the original fission yeast cell cycle Boolean network capable of the same function as the original network, namely it can reproduce the sequence of states for each phase of the cycle. IIT analysis of this reduced network demonstrated that while it could produce the same function, it did not maintain integration through each step of the cycle. Extrapolating to the case of cancer, one can separate function from integration, and loss of the latter does not immediately imply loss of the former. However, due to the rapid increase of the number of partitions necessary to calculate φ as system size increases, IIT is impractical to calculate for systems larger than about ten elements (roughly the size of the fission yeast network). Other measures that approximate φ , or quantify different aspects of integration

[467], will be necessary for application to the cancer problem.

6.2. Agency and integrated spatiotemporal patterns

Colloquially, integrated information theory (IIT) focuses on whether a system is greater than the sum of its parts. It attempts, in a rigorous way, to make clear whether the system can be subdivided, and to quantify how much ‘more’ the system is as a whole. While this concept of integration is powerful, it says little about how the system in question is integrated *with its environment*. An alternative approach to assessing integration, integrated spatiotemporal patterns (ISTP) [463], offers a less assertive take on integration that can account for environmental context in a more flexible way. Specifically, ISTP attempts to provide a quantitative notion of *agency*. Since describing cancer requires not just understanding how it behaves as a collective, but also how it interacts with its environment, ISTP is a promising tool for understanding cancer dynamics.

One intriguing possibility is that tumorigenesis amounts to increased agency of collective and/or constituent cells. Phenomena such as metastasis and angiogenesis lend credence to this hypothesis, though few experiments address this possibility directly. The degree to which the cancer cells are collectively integrated could, based on known relationships between gap junction connectivity and proliferation [468, 469], translate into more aggressive disorders. This section lays out how ISTP could prove experimentally valuable in quantifying the agency of tissues, with cancers as an example of interest.

The notion of agency is notoriously difficult to define much less rigorously quantify. However, there are properties that are commonly accepted as requirements for agency: *individuality*, *perception*, and *action* [470]. Individuality expresses the idea of an ‘entity’, something to which perception and action can be attributed. This generally requires a larger context than the individual itself, e.g. a tumor is defined in the context of a tissue or a body, an organism in the context of an environment, etc... Within the realm of information theory, individuality is typically quantified in terms of either mutual information or relative entropy between the constituent parts of the individual and the environmental setting. More formally, individuality can be expressed in terms of the *integration* of a subsystem with its environmental context—low integration suggests that the entire system is well understood in terms of how the individual interacts with its environmental context (and vice versa). Given such an individual, *perception* amounts to the individual’s interactions with its environment and *action* is the subsystem’s response to external stimuli.

Within the context of a given tissue, each cell has some degree of agency—it is an individual that perceives changes in its environment and responds accordingly. However, tissues themselves are reasonably

homogeneous and each cell is functionally similar, if not equivalent, to every other cell. What makes cancer cells distinct is that they break this homogeneity and take on a different function; they become distinguishable from their surrounding tissue. What's more they begin to decouple from their environment, which arguably allows for metastasis and greater proliferation, and makes them easier to culture [471–473]. The expectation is that they become less integrated with their surroundings and take on greater agency than their neighboring cells. However, this is not to say that one should expect that all tumors themselves are integrated structures. This said, the most prolific or metastatic scenarios show signs of cooperative behavior or swarm-like behavior, and as such may be identified in terms of their increased degree of agency and integration.

The concept of *integrated spatiotemporal patterns* was introduced in the work of Biehl et al [463] to make explicit the notion of agency in terms of spatiotemporal integration. The authors build upon the works of Krakauer et al [39] and Friston [474] to provide a framework which can represent agents. Biehl et al began by expressing the structure of discrete, finite-time dynamical systems in terms of dynamical Bayesian networks [463]. These directed, acyclic networks are composed of nodes, each associated with a finite random variable, and directed edges signifying correlational dependence. Essentially, each node represents some aspect of the system, e.g. a chemical concentration, the expression of a gene, the expressed phenotype of a cell, etc, and the edges represent conditional dependence between those variables, e.g. phenotypes may be conditionally dependent on gene expression. As noted by Ay and Polani [475], Bayesian networks have temporal structure; the nodes can be partitioned into sets $\{V_0, V_1, \dots\}$ such that if a node $A \in V_{t+1}$ then there exists a node $B \in V_t$ with an edge from B to A . In other words, the nodes in the network can be broken into *time slices* where nodes in subsequent time slices have correlational dependence on nodes in the previous time slice.

Given such a Bayesian network, a spatiotemporal pattern is defined to be a collection of nodes in the network together with specific values for those nodes [463]. For example, given a network $G \rightarrow P$ which represents the correlational dependence of a phenotype P on the expression of some gene G , then a spatiotemporal pattern may be $x_O = \{1 \in G, 0 \in P\}$ representing the pattern 'the gene is expressed while the particular phenotype is absent'. Such a pattern has associated with it a probability $p_O(x_O)$ of occurrence. In a practical situation, this probability would be approximated by the observed frequency of the pattern relative to every expressible pattern of those nodes, here G and P .

The integration of a pattern can then be assessed in terms of local mutual information, first introduced by Fano and Hawkins (1961) [476],

$$mi(X_1 = x_1, \dots, X_n = x_n) = \log \frac{p(x_1, \dots, x_n)}{\prod_{i=1}^n p(x_i)} \quad (7)$$

which is an unaveraged, multivariate form of the mutual information discussed in equation (2)). Unlike mutual information, local mutual information can take on negative values. This signifies that observing the random variables X_1, \dots, X_n in the states x_1, \dots, x_n together is less likely than would be expected if the variables were fully independent. The evidence for integration for a spatiotemporal pattern x_O relative to a partitioning π of the pattern is then defined by

$$mi_\pi(x_O) = \begin{cases} 0 & \text{if } p_O(x_O) = 0 \\ \log \frac{p_O(x_O)}{\prod_{Q \in \pi} p_Q(x_Q)} & \text{otherwise.} \end{cases} \quad (8)$$

The pattern x_O is *integrated* if and only if $mi_\pi(x_O) > 0$ for every partitioning π of x_O . In other words, the pattern is integrated if considering all the components of the system jointly is always better than assuming that they are independent, regardless of how you group those components.

Continuing the example above, there is only one proper partitioning of the pattern $x_O = \{1 \in G, 0 \in P\}$, namely $\pi = \{\{1 \in G\}, \{0 \in P\}\}$. Using equation (8), the evidence for integration of x_O relative to π is


$$mi_\pi(x_O) = \begin{cases} 0 & \text{if } p_O(x_O) = 0 \\ \log \frac{p(G=1, P=0)}{p(G=1)p(P=0)} & \text{otherwise.} \end{cases}$$

In this case, if $p(G=1, P=0) > p(G=1)p(P=0)$, then we would say that the pattern x_O is integrated. As such, from the point of view of the ISTP formalism, x_O would represent an agent in the dynamical system $G \rightarrow P$.

The simplest approach to applying this formalism to real systems proceeds as follows.

1. Identify the degrees of freedom of interest, e.g. phenotypes, genes, chemical concentrations, etc...
2. Observe the state of each degree of freedom over a sufficiently long period.
3. Construct an approximation of the system's Bayesian network from the time-series data.
4. Assess the integration of each spatiotemporal pattern in the system using equation (8).

Unfortunately, the systems of interest to cancer biology would likely consist of hundreds or thousands of degrees of freedom. Some of these degrees of freedom may be binary, e.g. a gene is expressed or suppressed, but most would likely admit many more



states, e.g. chemical concentrations or cell membrane potentials. Identifying agents using the formalism thus described would prove computationally challenging, if not impossible, due to fact that the number of partitionings of a spatiotemporal pattern grows faster than exponentially with the number of nodes in the pattern. To overcome this limitation, Biehl *et al* suggest only computing the evidence for integration in the finest partitioning (the partitioning in which each node is in its own partition). This approximation is computationally tractable for all but the most pathological cases. Using this approach, Biehl *et al* present preliminary evidence that ISTP can accommodate metabolism, motility and counterfactual variation as intended [463].

As tools in the cancer biologist's toolkit, IIT and ISTP offer powerful new ways of understanding how cancer develops and progresses over time. While measures themselves are likely computationally intractable,

approximations exist that will make analyzing the systems of interest possible [463, 467]. The concept of integration may provide insight into phenomena such as metastasis and angiogenesis which could provide targets for treatment. The hypothesis that cancer virulence is related to its degree of integration, internally and with its environment, can be assessed head on with these measures. Establishing such a connection could provide new biomedical techniques for treating cancer, possibly even reverting the cancer state altogether.

7. Conclusion

Over the last several decades we have made tremendous advances in understanding the molecular constituents of living matter. Nonetheless, we are still far from a general understanding of what life is, or how the millions of components of a cell orchestrate to produce the properties we associate with the living state.

The phenomena of complex organ regeneration, and of developmental reprogramming of cancer, challenge our field to augment the increasing resolution of cellular and molecular analysis with the equally necessary integration that explains the information content of (not only mechanisms involved in) higher levels of organization (figure 6). In cancer, many of these properties fail in ways that still preserve features necessary to persist as a living system. Cancer thus represents a functioning example of a living alternative that of healthy tissue, one that is operating with nearly identical molecular components, but very different growth and patterning dynamics. This is perhaps one reason the problem of cancer has been so intractable: reductionist approaches that focus on molecular components alone are hard pressed to distinguish between a cancerous and healthy cell as they share much of the same molecular biology, albeit differentially expressed. Treating cancer as a disease therefore cannot be disentangled from understanding more fundamental aspects of living organization, including what leads to the robustness of the living state, and what alternative forms this robust state of order might take on.

These properties are perhaps best characterized by adopting an informational [416, 431] perspective. Practical advances in this field will require quantitative analyses of patterning information processing in tractable animal models. Key open questions include: (1) determining (perhaps using integrated information metrics) the organizational level at which decisions in cancer are made—which types of tumors are their own integrated entity and which are simply reversions of somatic cells to a unicellular past, and using tools of connectionist computational neuroscience to understand the representation of anatomical pattern states in bioelectrical properties of tissues (figure 6), (2) identifying effective ways to couple cells back to the morphogenetic field (perhaps through the development of gap junction opener drugs), (3) using physiological reporter modalities (such as voltage-sensitive dyes) to detect pre-cancer signatures of incipient patterning failures *in vivo*, and (4) identifying ionoceuticals (blends of the many available and already human-approved ion channel drugs) that can directly mimic specific patterning cues to induce normalization (transition existing successes in amphibia to mammalian models). Recent technologies in interrogating cell perception space [477] suggest numerous approaches to quantify information processed within and among cell collectives, which could be applied to bioelectric signaling to better understand this important layer of the morphogenetic field and then impose states that increase whole-body integrated pattern dynamics. Of course, interventions derived from this perspective will have to be integrated into a biomedical program that functions despite the real-life stressors to which human patients are continuously exposed.

Here we have demonstrated how novel empirical data on cancer progress can be combined with new insights from information-based approaches to the characterization of cancers to provide new insights into the cancer problem, and hopefully new pathways for diagnosis and treatment.

Acknowledgments

ML is grateful to many members of the cancer and developmental biology communities for discussions of these topics. This paper is dedicated to the memory of Panagiotis ‘Takis’ Tsonis, a brilliant, versatile developmental biologist who contributed significantly to the cancer/regeneration interface. ML, SIW, and DM are supported by the Templeton Foundation (TWCF0089/AB55 and TWCF0140). ML also gratefully acknowledges an Allen Discovery Center award from The Paul G Allen Frontiers Group, and support by the G Harold and Leila Y Mathers Charitable Foundation.

ORCID iDs

Douglas G. Moore  <https://orcid.org/0000-0003-4003-4320>

Sara I Walker  <https://orcid.org/0000-0001-5779-2772>

Michael Levin  <https://orcid.org/0000-0001-7292-8084>

References

- [1] Johnston R N, Pai S B and Pai R B 1992 The origin of the cancer cell: oncogeny reverses phylogeny *Biochem. Cell Biol.* **70** 831–4
- [2] Huang S 2011 On the intrinsic inevitability of cancer: from foetal to fatal attraction *Semin. Cancer Biol.* **21** 183–99
- [3] Vaux D L 2011 In defense of the somatic mutation theory of cancer *Bioessays* **33** 341–3
- [4] Bjerkvig R, Tysnes B B, Aboody K S, Najbauer J and Terzis A J 2005 Opinion: the origin of the cancer stem cell: current controversies and new insights *Nat. Rev. Cancer* **5** 899–904
- [5] Hanahan D and Weinberg R A 2000 The hallmarks of cancer *Cell* **100** 57–70
- [6] Waddington C H 1935 Cancer and the theory of organisers *Nature* **135** 606–8
- [7] Rubin H 1985 Cancer as a dynamic developmental disorder *Cancer Res.* **45** 2935–42 (PMID: 3891078)
- [8] Burr H S 1940 Biologic organization and the cancer problem *Yale J. Biol. Med.* **12** 277–82 (PMCID: PMC2602176)
- [9] Soto A M and Sonnenschein C 2011 The tissue organization field theory of cancer: a testable replacement for the somatic mutation theory *Bioessays* **33** 332–40
- [10] Sonnenschein C, Soto A M, Rangarajan A and Kulkarni P 2014 Competing views on cancer *J. Biosci.* **39** 281–302
- [11] Smithers D W 1962 An attack on cytologism *Lancet* **1** 493–9
- [12] Huang S, Ernberg I and Kauffman S 2009 Cancer attractors: a systems view of tumors from a gene network dynamics and developmental perspective *Semin. Cell Dev. Biol.* **20** 869–76
- [13] Bizzarri M and Cucina A 2016 SMT and TOFT: why and how they are opposite and incompatible paradigms *Acta Biotheor.* **64** 221–39
- [14] Bizzarri M, Cucina A, Conti F and D’Anselmi F 2008 Beyond the oncogene paradigm: understanding complexity in cancerogenesis *Acta Biotheor.* **56** 173–96
- [15] Tarin D 2012 Clinical and biological implications of the tumor microenvironment *Cancer Microenviron.* **5** 95–112

- [16] Tarin D 2011 Cell and tissue interactions in carcinogenesis and metastasis and their clinical significance *Semin. Cancer Biol.* **21** 72–82
- [17] Soto A M and Sonnenschein C 2014 One hundred years of somatic mutation theory of carcinogenesis: is it time to switch? *Bioessays* **36** 118–20
- [18] Levin M 2012 Morphogenetic fields in embryogenesis, regeneration and cancer: non-local control of complex patterning *Biosystems* **109** 243–61
- [19] Siegel R L, Miller K D and Jemal A 2016 Cancer statistics *Cancer J. Clin.* **66** 7–30
- [20] Pezzulo G and Levin M 2016 Top-down models in biology: explanation and control of complex living systems above the molecular level *J. R. Soc. Interface* **13** 20160555
- [21] Song Y, Wang Y, Tong C, Xi H, Zhao X, Wang Y and Chen L 2017 A unified model of the hierarchical and stochastic theories of gastric cancer *Br. J. Cancer* **116** 973–89
- [22] Torquato S 2011 Toward an Ising model of cancer and beyond *Phys. Biol.* **8** 15017
- [23] Biava P M, Basevi M, Biggiero L, Borgonovo A, Borgonovo E and Burigana F 2011 Cancer cell reprogramming: stem cell differentiation stage factors and an agent based model to optimize cancer treatment *Curr. Pharm. Biotechnol.* **12** 231–42
- [24] Byrne H M 2010 Dissecting cancer through mathematics: from the cell to the animal model *Nat. Rev. Cancer* **10** 221–30
- [25] Hadi M and Marashi S-A 2014 Reconstruction of a generic metabolic network model of cancer cells *Mol. Biosyst.* **10** 3014–21
- [26] Johnston M D, Edwards C M, Bodmer W F, Maini P K and Chapman S J 2007 Mathematical modeling of cell population dynamics in the colonic crypt and in colorectal cancer *Proc. Natl Acad. Sci.* **104** 4008–13
- [27] la Porta C A M, Zapperi S and Sethna J P 2012 Senescent cells in growing tumors: Population dynamics and cancer stem cells *PLoS Comput. Biol.* **8** e1002316
- [28] Garner A L, Lau Y Y, Jordan D W, Uhler M D and Gilgenbach R M 2006 Implications of a simple mathematical model to cancer cell population dynamics *Cell Prolif.* **39** 15–28
- [29] Zhou D, Wu D, Li Z, Qian M and Zhang M Q 2013 Population dynamics of cancer cells with cell state conversions, *Quant. Biol.* **1** 201–8
- [30] Cleveland C, Liao D and Austin R 2012 Physics of cancer propagation: a game theory perspective *AIP Adv.* **2** 011202
- [31] Pacheco J M, Santos F C and Dingli D 2014 The ecology of cancer from an evolutionary game theory perspective *Interface Focus* **4** 20140019
- [32] Dingli D, Chalub F A C C, Santos F C, Van Segbroeck S and Pacheco J M 2009 Cancer phenotype as the outcome of an evolutionary game between normal and malignant cells *Br. J. Cancer* **101** 1130–6
- [33] Tarabichi M, Antoniou A, Saiselet M, Pita J M, Andry G, Dumont J E, Detours V and Maenhaut C 2013 Systems biology of cancer: entropy, disorder, and selection-driven evolution to independence, invasion and ‘swarm intelligence’ *Cancer Metastasis Rev.* **32** 403–21
- [34] Rosenfeld S 2013 Critical junction: nonlinear dynamics, swarm intelligence and cancer research 2013 *IEEE Symp. on Computational Intelligence in Bioinformatics and Computational Biology* pp 206–11
- [35] Anderson P W et al 1972 More is different *Science* **177** 393–6
- [36] Yockey H P 1956 An application of information theory to the physics of tissue damage *Radiat. Res.* **5** 146–55
- [37] Yockey H P 1974 An application of information theory to the central dogma and the sequence hypothesis *J. Theor. Biol.* **46** 369–406
- [38] Adami C and Cerf N J 2000 Physical complexity of symbolic sequences *Phys. D Nonlinear Phenom.* **137** 62–9
- [39] Krakauer D, Bertschinger N, Olbrich E, Ay N and Flack J C 2014 The information theory of individuality (arXiv:1412.2447)
- [40] Wibral M, Lizier J T, Vögler S, Priesemann V and Galuske R 2014 Local active information storage as a tool to understand distributed neural information processing *Front. Neuroinform.* **8** 1
- [41] Wibral M, Lizier J T and Priesemann V 2014 How to measure local active information storage in neural systems 2014 *8th Conf. of the European Study Group on Cardiovascular Oscillations* pp 131–2
- [42] Gourevitch B and Eggermont J J 2007 Evaluating information transfer between auditory cortical neurons *J. Neurophysiol.* **97** 2533–43
- [43] Smith J M 2000 The concept of information in biology *Phil. Sci.* **67** 177–94
- [44] Gatenby R A and Frieden B R 2002 Application of information theory and extreme physical information to carcinogenesis *Cancer Res.* **62** 3675–84
- [45] Gatenby R A and Frieden B R 2004 Information dynamics in carcinogenesis and tumor growth *Mutat. Res. Mol. Mech. Mutagen.* **568** 259–73
- [46] Mirmomeni M, Punch W F and Adami C 2014 Is information a selectable trait? (arXiv:1408.3651)
- [47] Adami C 2004 Information theory in molecular biology *Phys. Life Rev.* **1** 3–22
- [48] Mathis C, Bhattacharya T and Walker S I 2015 The emergence of life as a first order phase transition *Astrobiology* **17** 266–76
- [49] Sia D, Alsinet C, Newell P and Villanueva A 2014 VEGF signaling in cancer treatment *Curr. Pharm. Des.* **20** 2834–42
- [50] Bartolotti M, Franceschi E and Brandes A A 2012 EGF receptor tyrosine kinase inhibitors in the treatment of brain metastases from non-small-cell lung cancer *Expert Rev. Anticancer Ther.* **12** 1429–35
- [51] Davies P C W and Lineweaver C H 2011 Cancer tumors as Metazoa 1.0: tapping genes of ancient ancestors *Phys. Biol.* **8** 15001
- [52] Ingber D E and Levin M 2007 What lies at the interface of regenerative medicine and developmental biology? *Development* **134** 2541–7
- [53] Birnbaum K D and Alvarado A S 2008 Slicing across kingdoms: regeneration in plants and animals *Cell* **132** 697–710
- [54] Tarkowski A K 1961 Mouse chimaeras developed from fused eggs *Nature* **190** 857–60
- [55] Mintz B 1971 Genetic mosaicism *in vivo*: development and disease in allophenic mice *Fed. Proc.* **30** 935–43
- [56] Takano K, Ito Y, Obata S, Oinuma T, Komazaki S, Nakamura H and Asashima M 2007 Heart formation and left-right asymmetry in separated right and left embryos of a newt *Int. J. Dev. Biol.* **51** 265–72
- [57] Baddour J A, Sousounis K and Tsonis P A 2012 Organ repair and regeneration: an overview *Birth Defects Res. C* **96** 1–29
- [58] McCusker C and Gardiner D M 2011 The axolotl model for regeneration and aging research: a mini-review *Gerontology* **57** 565–71
- [59] Lobo D, Beane W S and Levin M 2012 Modeling planarian regeneration: a primer for reverse-engineering the worm *PLoS Comput Biol.* **8** e1002481
- [60] Gentile L, Cebria F and Bartscherer K 2011 The planarian flatworm: an *in vivo* model for stem cell biology and nervous system regeneration *Dis. Models Mech.* **4** 12–9
- [61] Neuhof M, Levin M and Rechavi O 2016 Vertically- and horizontally-transmitted memories—the fading boundaries between regeneration and inheritance in planaria *Biol. Open* **5** 1177–88
- [62] Nishimura O, Hosoda K, Kawaguchi E, Yazawa S, Hayashi T, Inoue T, Umesono Y and Agata K 2015 Unusually large number of mutations in asexually reproducing clonal planarian *dugesia japonica* *PLoS One* **10** e0143525
- [63] Levin M 2011 The wisdom of the body: future techniques and approaches to morphogenetic fields in regenerative medicine, developmental biology and cancer *Regen. Med.* **6** 667–73
- [64] Chandebois R 1980 Cell sociology and the problem of automation in the development of pluricellular animals *Acta Biotheor.* **29** 1–35
- [65] Friston K, Levin M, Sengupta B and Pezzulo G 2015 Knowing one’s place: a free-energy approach to pattern regulation *J. R. Soc. Interface* **12** 20141383
- [66] Farinella-Ferruzza N 1956 The transformation of a tail into a limb after xenoplastic transformation *Experientia* **15** 304–5

- [67] Seilern-Aspang F and Kratochwill L 1965 Relation between regeneration and tumor growth *Regeneration in Animals and Related Problems* (Amsterdam: North-Holland) pp 452–73
- [68] Bizzarri M, Cucina A, Biava P M, Proietti S, D'Anselmi F, Dinicola S, Pasqualato A and Lisi E 2011 Embryonic morphogenetic field induces phenotypic reversion in cancer cells *Curr. Pharm. Biotechnol.* **12** 243–53
- [69] Morozova N and Shubin M 2013 The geometry of morphogenesis and the morphogenetic field concept *Pattern Formation in Morphogenesis* vol 15, ed V Capasso et al (Berlin: Springer) pp 255–82
- [70] Belousov L V, Opitz J M and Gilbert S F 1997 Life of Alexander G. Gurwitsch and his relevant contribution to the theory of morphogenetic fields *Int. J. Dev. Biol.* **41** 771–9
- [71] Fankhauser G 1945 Maintenance of normal structure in heteroploid salamander larvae, through compensation of changes in cell size by adjustment of cell number and cell shape *J. Exp. Zool.* **100** 445–55
- [72] Prehn R T 1997 Regeneration versus neoplastic growth *Carcinogenesis* **18** 1439–44
- [73] Prehn R T 1994 Cancers beget mutations versus mutations beget cancers *Cancer Res.* **54** 5296–300
- [74] Scharrer B 1953 Insect tumors induced by nerve severance: incidence and mortality *Cancer Res.* **13** 73–6
- [75] Scharrer B and Lochhead M S 1950 Tumors in the invertebrates: a review *Cancer Res.* **10** 403–19
- [76] Pawlowski A and Weddell G 1967 Induction of tumours in denervated skin *Nature* **213** 1234–6
- [77] Outzen H C, Custer R P and Prehn R T 1976 Influence of regenerative capacity and innervation on oncogenesis in the adult frog (*Rana pipiens*) *J. Natl. Cancer Inst.* **57** 79–84
- [78] Parodi S and Brambilla G 1977 Relationships between mutation and transformation frequencies in mammalian cells treated 'in vitro' with chemical carcinogens *Mutat. Res.* **47** 53–74
- [79] Clark W H Jr 1995 The nature of cancer: morphogenesis and progressive (self)-disorganization in neoplastic development and progression *Acta Oncol.* **34** 3–21
- [80] Kraemer P M, Travis G L, Ray F A and Cram L S 1983 Spontaneous neoplastic evolution of Chinese hamster cells in culture: multistep progression of phenotype *Cancer Res.* **43** 4822–7
- [81] Cram L S, Bartholdi M F, Ray F A, Travis G L and Kraemer P M 1983 Spontaneous neoplastic evolution of Chinese hamster cells in culture: multistep progression of karyotype *Cancer Res.* **43** 4828–37
- [82] Dolberg D S and Bissell M J 1984 Inability of Rous sarcoma virus to cause sarcomas in the avian embryo *Nature* **309** 552–6
- [83] Baker S G, Soto A M, Sonnenschein C, Cappuccio A, Potter J D and Kramer B S 2009 Plausibility of stromal initiation of epithelial cancers without a mutation in the epithelium: a computer simulation of morphostats *BMC Cancer* **9** 89
- [84] Potter J D 2007 Morphogens, morphostats, microarchitecture and malignancy *Nat. Rev. Cancer* **7** 464–74
- [85] Potter J D 2001 Morphostats: a missing concept in cancer biology *Cancer Epidemiol. Biomarkers Prev.* **10** 161–70
- [86] Soto A M and Sonnenschein C 2004 The somatic mutation theory of cancer: growing problems with the paradigm? *Bioessays* **26** 1097–107
- [87] Bissell M J, Radisky D C, Rizki A, Weaver V M and Petersen O W 2002 The organizing principle: microenvironmental influences in the normal and malignant breast *Differentiation* **70** 537–46
- [88] Barcellos-Hoff M H 2001 It takes a tissue to make a tumor: epigenetics, cancer and the microenvironment *J. Mammary Gland Biol. Neoplasia* **6** 213–21
- [89] Wang F, Weaver V M, Petersen O W, Larabell C A, Dedhar S, Briand P, Lupu R and Bissell M J 1998 Reciprocal interactions between beta1-integrin and epidermal growth factor receptor in three-dimensional basement membrane breast cultures: a different perspective in epithelial biology *Proc. Natl Acad. Sci. USA* **95** 14821–6
- [90] Oppenheimer B S, Oppenheimer E T and Stout A P 1952 Sarcomas induced in rodents by imbedding various plastic films *Proc. Soc. Exp. Biol. Med.* **79** 366–9
- [91] Bischoff F and Bryson G 1964 Carcinogenesis through solid state surfaces *Prog. Exp. Tumor Res.* **5** 85–133
- [92] Oppenheimer B S, Oppenheimer E T and Stout A P 1953 Carcinogenic effect of imbedding various plastic films in rats and mice *Surg. Forum* **4** 672–6
- [93] Oppenheimer B S, Oppenheimer E T, Stout A P and Danishefsky I 1953 Malignant tumors resulting from embedding plastics in rodents *Science* **118** 305–6
- [94] Bissell M J and Hines W C 2011 Why don't we get more cancer? A proposed role of the microenvironment in restraining cancer progression *Nat. Med.* **17** 320–9
- [95] Wade M H, Trosko J E and Schindler M 1986 A fluorescence photobleaching assay of gap junction-mediated communication between human cells *Science* **232** 525–8
- [96] Ruch R J and Trosko J E 2001 Gap-junction communication in chemical carcinogenesis *Drug Metab. Rev.* **33** 117–24
- [97] Dufloot-Dancer A, Mesnil M and Yamasaki H 1997 Dominant-negative abrogation of connexin-mediated cell growth control by mutant connexin genes *Oncogene* **15** 2151–8
- [98] Yamasaki H, Krutovskikh V, Mesnil M, Tanaka T, Zaidan-Dagli M L and Omori Y 1999 Role of connexin (gap junction) genes in cell growth control and carcinogenesis *C. R. Acad. Sci. III* **322** 151–9
- [99] Omori Y, Zaidan Dagli M L, Yamakage K and Yamasaki H 2001 Involvement of gap junctions in tumor suppression: analysis of genetically-manipulated mice *Mutat. Res.* **477** 191–6
- [100] Yamasaki H, Mesnil M, Omori Y, Mironov N and Krutovskikh V 1995 Intercellular communication and carcinogenesis *Mutat. Res.* **333** 181–8
- [101] Temme A, Buchmann A, Gabriel H D, Nelles E, Schwarz M and Willecke K 1997 High incidence of spontaneous and chemically induced liver tumors in mice deficient for connexin32 *Curr. Biol.* **7** 713–6
- [102] Loewenstein W R 1980 Junctional cell-to-cell communication and growth control *Ann. N. Y. Acad. Sci.* **339** 39–45
- [103] Loewenstein W R 1979 Junctional intercellular communication and the control of growth *Biochim. Biophys. Acta* **560** 1–659
- [104] Loewenstein W R 1969 Transfer of information through cell junctions and growth control *Proc. Can. Cancer Conf.* **8** 162–70
- [105] Loewenstein W R and Kanno Y 1966 Intercellular communication and the control of tissue growth: lack of communication between cancer cells *Nature* **209** 1248–9
- [106] Rose B, Mehta P P and Loewenstein W R 1993 Gap-junction protein gene suppresses tumorigenicity *Carcinogenesis* **14** 1073–5
- [107] Mesnil M, Crespin S, Avanzo J L and Zaidan-Dagli M L 2005 Defective gap junctional intercellular communication in the carcinogenic process *Biochim. Biophys. Acta* **1719** 125–45
- [108] Stoletov K, Strnadel J, Zardouzan E, Momiyama M, Park F D, Kelber J A, Pizzo D P, Hoffman R, Vandenberg S R and Klemke R L 2013 Role of connexins in metastatic breast cancer and melanoma brain colonization *J. Cell Sci.* **126** 904–13
- [109] Moreno E 2008 Is cell competition relevant to cancer? *Nat. Rev. Cancer* **8** 141–7
- [110] Dobrowolski R, Hertig G, Lechner H, Worsdorfer P, Wulf V, Dicke N, Eckert D, Bauer R, Schorle H and Willecke K 2009 Loss of connexin43-mediated gap junctional coupling in the mesenchyme of limb buds leads to altered expression of morphogens in mice *Hum. Mol. Genet.* **18** 2899–911
- [111] Jin E J, Lee S Y, Jung J C, Bang O S and Kang S S 2008 TGF-beta3 inhibits chondrogenesis of cultured chick leg bud mesenchymal cells via downregulation of connexin 43 and integrin beta4 *J. Cell Physiol.* **214** 345–53
- [112] Pizard A, Burgon P G, Paul D L, Bruneau B G, Seidman C E and Seidman J G 2005 Connexin 40, a target of transcription factor tbx5, patterns wrist, digits and sternum *Mol. Cell Biol.* **25** 5073–83

- [113] Levin M and Mercola M 2000 Expression of connexin 30 in *Xenopus* embryos and its involvement in hatching gland function *Dev. Dyn.* **219** 96–101
- [114] Lecanda F, Warlow P M, Sheikh S, Furlan F, Steinberg T H and Civitelli R 2000 Connexin43 deficiency causes delayed ossification, craniofacial abnormalities and osteoblast dysfunction *J. Cell Biol.* **151** 931–44
- [115] Warner A 1999 Interactions between growth factors and gap junctional communication in developing systems *Novartis Found. Symp.* **219** 60–72
- [116] Levin M and Mercola M 1998 Gap junctions are involved in the early generation of left-right asymmetry *Dev. Biol.* **203** 90–105
- [117] Levin M and Mercola M 1999 Gap junction-mediated transfer of left-right patterning signals in the early chick blastoderm is upstream of Shh asymmetry in the node *Development* **126** 4703–14
- [118] Levin M 2007 Gap junctional communication in morphogenesis *Prog. Biophys. Mol. Biol.* **94** 186–206
- [119] Oviedo N J, Morokuma J, Walentek P, Kema I P, Gu M B, Ahn J M, Hwang J S, Gojobori T and Levin M 2010 Long-range neural and gap junction protein-mediated cues control polarity during planarian regeneration *Dev. Biol.* **339** 188–99
- [120] Schiffmann Y 2008 The Turing-Child energy field as a driver of early mammalian development *Prog. Biophys. Mol. Biol.* **98** 107–17
- [121] Nogi T and Levin M 2005 Characterization of innexin gene expression and functional roles of gap-junctional communication in planarian regeneration *Dev. Biol.* **287** 314–35
- [122] Holtfreter J 1955 Transformation of a tail into a limb or gill-like structures *J. Exp. Zool.* **129** 623–48
- [123] Farinella-Ferruzza N 1953 Risultati di trapianti di bottone codale di urodeli su anuri e vice versa *Riv. Biol.* **45** 523–7
- [124] Mintz B and Illmensee K 1975 Normal genetically mosaic mice produced from malignant teratocarcinoma cells *Proc. Natl Acad. Sci. USA* **72** 3585–9
- [125] Illmensee K and Mintz B 1976 Totipotency and normal differentiation of single teratocarcinoma cells cloned by injection into blastocysts *Proc. Natl Acad. Sci. USA* **73** 549–53
- [126] Astigiano S, Damonte P, Fossati S, Boni L and Barbieri O 2005 Fate of embryonal carcinoma cells injected into postimplantation mouse embryos *Differentiation* **73** 484–90
- [127] Webb C G, Gootwine E and Sachs L 1984 Developmental potential of myeloid leukemia cells injected into midgestation embryos *Dev. Biol.* **101** 221–4
- [128] Li L, Connelly M C, Wetmore C, Curran T and Morgan J I 2003 Mouse embryos cloned from brain tumors *Cancer Res.* **63** 2733–6
- [129] Pierce G B and Wells R S 1982 The blastocyst in control of embryonal carcinoma *Prog. Clin. Biol. Res.* **85** 593–600
- [130] Pierce G B, Arechaga J and Wells R S 1986 Embryonic control of cancer *Prog. Clin. Biol. Res.* **226** 67–77
- [131] Podesta A H, Mullins J, Pierce G B and Wells R S 1984 The neurula stage mouse embryo in control of neuroblastoma *Proc. Natl Acad. Sci. USA* **81** 7608–11
- [132] Pierce G B, Aguilar D, Hood G and Wells R S 1984 Trophoblast in control of murine embryonal carcinoma *Cancer Res.* **44** 3987–96
- [133] Pierce G B 1983 The cancer cell and its control by the embryo. Rous-Whipple award lecture *Am. J. Pathol.* **113** 117–24
- [134] Pierce G B, Pantazis C G, Caldwell J E and Wells R S 1982 Specificity of the control of tumor formation by the blastocyst *Cancer Res.* **42** 1082–7
- [135] Haldi M, Ton C, Seng W L and McGrath P 2006 Human melanoma cells transplanted into zebrafish proliferate, migrate, produce melanin, form masses and stimulate angiogenesis in zebrafish *Angiogenesis* **9** 139–51
- [136] Lee L M, Seftor E A, Bonde G, Cornell R A and Hendrix M J 2005 The fate of human malignant melanoma cells transplanted into zebrafish embryos: assessment of migration and cell division in the absence of tumor formation *Dev. Dyn.* **233** 1560–70
- [137] Kasemeier-Kulesa J C, Teddy J M, Postovit L M, Seftor E A, Seftor R E, Hendrix M J and Kulesa P M 2008 Reprogramming multipotent tumor cells with the embryonic neural crest microenvironment *Dev. Dyn.* **237** 2657–66
- [138] Hendrix M J, Seftor E A, Seftor R E, Kasemeier-Kulesa J, Kulesa P M and Postovit L M 2007 Reprogramming metastatic tumour cells with embryonic microenvironments *Nat. Rev. Cancer* **7** 246–55
- [139] Kulesa P M, Kasemeier-Kulesa J C, Teddy J M, Margaryan N V, Seftor E A, Seftor R E and Hendrix M J 2006 Reprogramming metastatic melanoma cells to assume a neural crest cell-like phenotype in an embryonic microenvironment *Proc. Natl Acad. Sci. USA* **103** 3752–7
- [140] Hochedlinger K, Blleloch R, Brennan C, Yamada Y, Kim M, Chin L and Jaenisch R 2004 Reprogramming of a melanoma genome by nuclear transplantation *Genes Dev.* **18** 1875–85
- [141] Kenny P A and Bissell M J 2003 Tumor reversion: correction of malignant behavior by microenvironmental cues *Int. J. Cancer* **107** 688–95
- [142] Zhang W, Remenyik E, Zelterman D, Brash D E and Wikonkal N M 2001 Escaping the stem cell compartment: sustained UVB exposure allows p53-mutant keratinocytes to colonize adjacent epidermal proliferating units without incurring additional mutations *Proc. Natl. Acad. Sci. USA* **98** 13948–53
- [143] Chao D L, Eck J T, Brash D E, Maley C C and Luebeck E G 2008 Preneoplastic lesion growth driven by the death of adjacent normal stem cells *Proc. Natl Acad. Sci. USA* **105** 15034–9
- [144] Maffini M V, Calabro J M, Soto A M and Sonnenschein C 2005 Stromal regulation of neoplastic development: age-dependent normalization of neoplastic mammary cells by mammary stroma *Am. J. Pathol.* **167** 1405–10
- [145] Bussard K M, Boulanger C A, Booth B W, Bruno R D and Smith G H 2010 Reprogramming human cancer cells in the mouse mammary gland *Cancer Res.* **70** 6336–43
- [146] Booth B W, Boulanger C A, Anderson L H and Smith G H 2011 The normal mammary microenvironment suppresses the tumorigenic phenotype of mouse mammary tumor virus-neu-transformed mammary tumor cells *Oncogene* **30** 679–89
- [147] Tsonis P A 1983 Effects of carcinogens on regenerating and non-regenerating limbs in amphibia (review) *Anticancer Res.* **3** 195–202
- [148] Wolsky A 1978 Regeneration and cancer *Growth* **42** 425–6
- [149] Donaldson D J and Mason J M 1975 Cancer-related aspects of regeneration research: a review *Growth* **39** 475–96
- [150] Ruben L N, Balls M and Stevens J 1966 Cancer and super-regeneration in *Triturus viridescens* limbs *Experientia* **22** 260–1
- [151] Rose S M and Wallingford H M 1948 Transformation of renal tumors of frogs to normal tissues in regenerating limbs of salamanders *Science* **107** 1948
- [152] Brookes J P 1998 Regeneration and cancer *Biochim. Biophys. Acta* **1377** M1–11
- [153] Tsonis P A and Eguchi G 1981 Carcinogens on regeneration. Effects of N-methyl-N'-nitro-N-nitrosoguanidine and 4-nitroquinoline-1-oxide on limb regeneration in adult newts *Differentiation* **20** 52–60
- [154] Pizzarello D J and Wolsky A 1966 Carcinogenesis and regeneration in newts *Experientia* **22** 387–8
- [155] Okamoto M 1997 Simultaneous demonstration of lens regeneration from dorsal iris and tumour production from ventral iris in the same newt eye after carcinogen administration *Differentiation* **61** 285–92
- [156] Zilakos N P, Zafiratos C S and Parchment R E 1996 Stage-dependent genetically-based deformities of the regenerating newt limb from 4-nitroquinoline-N-oxide mutagenesis: potential embryonic regulation of cancer *Differentiation* **60** 67–74
- [157] Sheremetieva E A 1965 Spontaneous melanoma in regenerating tails of axolotls *J. Exp. Zool.* **158** 101–21
- [158] Breedis C 1952 Induction of accessory limbs and of sarcoma in the Newt (*Triturus viridescens*) with carcinogenic substances *Cancer Res.* **12** 861–6

- [159] Farber E 1984 Pre-cancerous steps in carcinogenesis. Their physiological adaptive nature *Biochim. Biophys. Acta* **738** 171–80
- [160] Farber E 1984 The multistep nature of cancer development *Cancer Res.* **44** 4217–23
- [161] Coleman W B, Wennerberg A E, Smith G J and Grisham J W 1993 Regulation of the differentiation of diploid and some aneuploid rat liver epithelial (stemlike) cells by the hepatic microenvironment *Am. J. Pathol.* **142** 1373–82
- [162] McCullough K D, Coleman W B, Ricketts S L, Wilson J W, Smith G J and Grisham J W 1998 Plasticity of the neoplastic phenotype *in vivo* is regulated by epigenetic factors *Proc. Natl Acad. Sci. USA* **95** 15333–8
- [163] Tatematsu M, Nagamine Y and Farber E 1983 Redifferentiation as a basis for remodeling of carcinogen-induced hepatocyte nodules to normal appearing liver *Cancer Res.* **43** 5049–58
- [164] Ogawa K, Solt D B and Farber E 1980 Phenotypic diversity as an early property of putative preneoplastic hepatocyte populations in liver carcinogenesis *Cancer Res.* **40** 725–33
- [165] Enomoto K and Farber E 1982 Kinetics of phenotypic maturation of remodeling of hyperplastic nodules during liver carcinogenesis *Cancer Res.* **42** 2330–5
- [166] Needham J 1936 New advances in the chemistry and biology of organized growth: (section of pathology) *Proc. R. Soc. Med.* **29** 1577–626
- [167] Zupanc G K 2008 Adult neurogenesis and neuronal regeneration in the brain of teleost fish *J. Physiol. Paris.* **102** 357–73
- [168] Mitternath J E 1981 The rule of normal neighbors: a hypothesis for morphogenetic pattern regulation *Dev. Biol.* **88** 15–26
- [169] Bryant P J and Simpson P 1984 Intrinsic and extrinsic control of growth in developing organs *Q. Rev. Biol.* **59** 387–415
- [170] French V 1980 Positional information around the segments of the cockroach leg *J. Embryol. Exp. Morphol.* **59** 281–313
- [171] Auerbach R, Morrissey L W, Kubai L and Sidky Y A 1978 Regional differences in tumor growth: studies of the vascular system *Int. J. Cancer* **22** 40–6
- [172] Auerbach R, Morrissey L W and Sidky Y A 1978 Gradients in tumour growth *Nature* **274** 697–9
- [173] Auerbach R, Morrissey L W and Sidky Y A 1978 Regional differences in the incidence and growth of mouse tumors following intradermal or subcutaneous inoculation *Cancer Res.* **38** 1739–44
- [174] Auerbach R and Auerbach W 1982 Regional differences in the growth of normal and neoplastic cells *Science* **215** 127–34
- [175] Kubai L and Auerbach R 1980 Regional differences in the growth of skin transplants *Transplantation* **30** 128–31
- [176] Morrissey L W, Sidky Y A and Auerbach R 1980 Regional differences in the growth of tumor cells injected intraperitoneally into syngeneic adult mice *Cancer Res.* **40** 2197–201
- [177] Biskind M S and Biskind G R 1945 Tumor of rat testis produced by heterotransplantation of infantile testis to spleen of adult castrate *Proc. Soc. Exp. Biol. Med.* **59** 4–8
- [178] Biskind M S and Biskind G S 1944 Development of tumors in the rat ovary after transplantation into the spleen *Proc. Soc. Exp. Biol. Med.* **55** 176–9
- [179] Balls M and Ruben L N 1964 Variation in the response of xenopus laevis to normal tissue homografts *Dev. Biol.* **10** 92–104
- [180] Stevens L C 1970 The development of transplantable teratocarcinomas from intratesticular grafts of pre- and postimplantation mouse embryos *Dev. Biol.* **21** 364–82
- [181] Bates E 2015 Ion channels in development and cancer *Annu. Rev. Cell Dev. Biol.* **31** 231–47
- [182] Pezzulo G and Levin M 2015 Re-membering the body: applications of computational neuroscience to the top-down control of regeneration of limbs and other complex organs *Integr. Biol.* **7** 1487–517
- [183] Borgens R B, Robinson K R, Venable J W and McGinnis M E 1989 *Electric Fields in Vertebrate Repair* (New York: Alan R. Liss)
- [184] Lund E J 1947 *Bioelectric Fields and Growth* (Austin, TX: University of Texas Press)
- [185] Pullar C E 2011 *The Physiology of Bioelectricity in Development, Tissue Regeneration and Cancer* (Boca Raton, FL: CRC Press)
- [186] McCaig C D, Song B and Rajnicek A M 2009 Electrical dimensions in cell science *J. Cell Sci.* **122** 4267–76
- [187] Popp F A 2009 Cancer growth and its inhibition in terms of coherence *Electromagn. Biol. Med.* **28** 53–60
- [188] Takeda M, Kobayashi M, Takayama M, Suzuki S, Ishida T, Ohnuki K, Moriya T and Ohuchi N 2004 Biophoton detection as a novel technique for cancer imaging *Cancer Sci.* **95** 656–61
- [189] Pokorný J, Hasek J, Vanis J and Jelinek F 2008 Biophysical aspects of cancer-electromagnetic mechanism *Indian J. Exp. Biol.* **46** 310–21
- [190] Frohlich H 1978 Coherent electric vibrations in biological systems and the cancer problem *IEEE Trans. Microw. Theory Tech.* **26** 613–8
- [191] Funk R H, Monsees T and Ozkucur N 2009 Electromagnetic effects—from cell biology to medicine *Prog. Histochem. Cytochem.* **43** 177–264
- [192] Cifra M, Fields J Z and Farhadi A 2011 Electromagnetic cellular interactions *Prog. Biophys. Mol. Biol.* **105** 223–46
- [193] Wang E T and Zhao M 2010 Regulation of tissue repair and regeneration by electric fields *Chin. J. Traumatol.* **13** 55–61
- [194] Zhao M 2009 Electrical fields in wound healing—an overriding signal that directs cell migration *Semin. Cell Dev. Biol.* **20** 674–82
- [195] Levin M 2012 Molecular bioelectricity in developmental biology: new tools and recent discoveries: control of cell behavior and pattern formation by transmembrane potential gradients *Bioessays* **34** 205–17
- [196] Levin M 2009 Bioelectric mechanisms in regeneration: Unique aspects and future perspectives *Semin. Cell Dev. Biol.* **20** 543–56
- [197] Rajnicek A M, Foubister L E and McCaig C D 2006 Growth cone steering by a physiological electric field requires dynamic microtubules, microfilaments and Rac-mediated filopodial asymmetry *J. Cell Sci.* **119** 1736–45
- [198] Levin M and Stevenson C G 2012 Regulation of cell behavior and tissue patterning by bioelectrical signals: challenges and opportunities for biomedical engineering *Annu. Rev. Biomed. Eng.* **14** 295–323
- [199] Becchetti A 2011 Ion channels and transporters in cancer. 1. Ion channels and cell proliferation in cancer *Am. J. Physiol. Cell Physiol.* **301** C255–65
- [200] Prevarskaya N, Skryma R and Shuba Y 2010 Ion channels and the hallmarks of cancer *Trends Mol. Med.* **16** 107–21
- [201] Pardo L A, Contreras-Jurado C, Zientkowska M, Alves F and Stuhmer W 2005 Role of voltage-gated potassium channels in cancer *J. Membr. Biol.* **205** 115–24
- [202] Kunzelmann K 2005 Ion channels and cancer *J. Membr. Biol.* **205** 159–73
- [203] Stock C, Ludwig F T, Hanley P J and Schwab A 2013 Roles of ion transport in control of cell motility *Compr. Physiol.* **3** 59–119
- [204] Sun Y S, Peng S W, Lin K H and Cheng J Y 2012 Electrotaxis of lung cancer cells in ordered three-dimensional scaffolds *Biomicrofluidics* **6** 14102–14
- [205] Yan X, Han J, Zhang Z, Wang J, Cheng Q, Gao K, Ni Y and Wang Y 2009 Lung cancer A549 cells migrate directionally in DC electric fields with polarized and activated EGFRs *Bioelectromagnetics* **30** 29–35
- [206] Pu J, McCaig C D, Cao L, Zhao Z, Segall J E and Zhao M 2007 EGF receptor signalling is essential for electric-field-directed migration of breast cancer cells *J. Cell Sci.* **120** 3395–403
- [207] Yao L, Pandit A, Yao S and McCaig C D 2001 Electric field-guided neuron migration: a novel approach in neurogenesis *Tissue Eng. B* **17** 143–53
- [208] Blackiston D J, Adams D S, Lemire J M, Lobikin M and Levin M 2011 Transmembrane potential of GlyCl-expressing instructor cells induces a neoplastic-like conversion of melanocytes via a serotonergic pathway *Dis. Model. Mech.* **4** 67–85

- [209] Morokuma J, Blackiston D J, Adams D S, Seeböhm G, Trimmer B and Levin M 2008 Modulation of potassium channel function confers a hyperproliferative invasive phenotype on embryonic stem cells *Proc. Natl Acad. Sci. USA* **105** 16608–13
- [210] Pu J and Zhao M 2005 Golgi polarization in a strong electric field *J. Cell Sci.* **118** 1117–28
- [211] Harris A K, Pryer N K and Paydarfar D 1990 Effects of electric fields on fibroblast contractility and cytoskeleton *J. Exp. Zool.* **253** 163–76
- [212] Baumann M 2001 Early stage shape change of human erythrocytes after application of electric field pulses *Mol. Membr. Biol.* **18** 153–60
- [213] Brackenbury W J, Djamgoz M B and Isom L L 2008 An emerging role for voltage-gated Na⁺ channels in cellular migration: regulation of central nervous system development and potentiation of invasive cancers *Neuroscientist* **14** 571–83
- [214] Isbilen B, Fraser S P and Djamgoz M B 2006 Docosahexaenoic acid (omega-3) blocks voltage-gated sodium channel activity and migration of MDA-MB-231 human breast cancer cells *Int. J. Biochem. Cell Biol.* **38** 2173–82
- [215] Binggeli R and Weinstein R C 1986 Membrane potentials and sodium channels: hypotheses for growth regulation and cancer formation based on changes in sodium channels and gap junctions *J. Theor. Biol.* **123** 377–401
- [216] Root C M, Velazquez-Ulloa N A, Monsalve G C, Minakova E and Spitzer N C 2008 Embryonically expressed GABA and glutamate drive electrical activity regulating neurotransmitter specification *J. Neurosci.* **28** 4777–84
- [217] Pineda R H, Svoboda K R, Wright M A, Taylor A D, Novak A E, Gamse J T, Eisen J S and Ribera A B 2006 Knockdown of Nav1.6a Na⁺ channels affects zebrafish motoneuron development *Development* **133** 3827–36
- [218] Svoboda K R, Linares A E and Ribera A B 2001 Activity regulates programmed cell death of zebrafish Rohon-Beard neurons *Development* **128** 3511–20
- [219] Messenger N J and Warner A E 2000 Primary neuronal differentiation in *Xenopus* embryos is linked to the beta(3) subunit of the sodium pump *Dev. Biol.* **220** 168–82
- [220] Ng S Y, Chin C H, Lau Y T, Luo J, Wong C K, Bian Z X and Tsang S Y 2010 Role of voltage-gated potassium channels in the fate determination of embryonic stem cells *J. Cell Physiol.* **224** 165–77
- [221] van Vliet P, de Boer T P, van der Heyden M A, El Tamer M K, Sluijter J P, Doevendans P A and Goumans M J 2010 Hyperpolarization induces differentiation in human cardiomyocyte progenitor cells *Stem Cell Rev.* **6** 178–85
- [222] Lange C, Prenninger S, Knuckles P, Taylor V, Levin M and Calegari F 2011 The H(+) vacuolar ATPase maintains neural stem cells in the developing mouse cortex *Stem Cells Dev.* **20** 843–50
- [223] Liu J H, Bijlenga P, Fischer-Lougheed J, Occhiodoro T, Kaelin A, Bader C R and Bernheim L 1998 Role of an inward rectifier K⁺ current and of hyperpolarization in human myoblast fusion *J. Physiol.* **510** 467–76
- [224] Occhiodoro T, Bernheim L, Liu J H, Bijlenga P, Sinnreich M, Bader C R and Fischer-Lougheed J 1998 Cloning of a human ether-a-go-go potassium channel expressed in myoblasts at the onset of fusion *FEBS Lett.* **434** 177–82
- [225] Bijlenga P, Occhiodoro T, Liu J H, Bader C R, Bernheim L and Fischer-Lougheed J 1998 An ether-a-go-go K⁺ current, Ih-eag, contributes to the hyperpolarization of human fusion-competent myoblasts *J. Physiol.* **512** 317–23
- [226] Hinard V, Belin D, König S, Bader C R and Bernheim L 2008 Initiation of human myoblast differentiation via dephosphorylation of Kir2.1 K⁺ channels at tyrosine 242 *Development* **135** 859–67
- [227] Sundelacruz S, Levin M and Kaplan D L 2008 Membrane potential controls adipogenic and osteogenic differentiation of mesenchymal stem cells *PLoS One* **3** e3737
- [228] Jiang P, Rushing S, Kong C W, Fu J, Lieu D K, Chan C, Deng W and Li R 2009 Electrophysiological properties of human induced pluripotent stem cells *Am. J. Physiol. Cell Physiol.* **298** C486–95
- [229] Stillwell E F, Cone C M and Cone C D 1973 Stimulation of DNA synthesis in CNS neurones by sustained depolarisation *Nat. New Biol.* **246** 110–1
- [230] Cone C D 1980 Ionically mediated induction of mitogenesis in CNS neurons *Ann. N. Y. Acad. Sci.* **339** 115–31
- [231] Saez J C, Connor J A, Spray D C and Bennett M V L 1989 Hepatocyte gap junctions are permeable to the second messenger, inositol 1,4,5-trisphosphate and to calcium ions *Proc. Natl. Acad. Sci.* **86** 2708–12
- [232] Toyofuku T, Yabuki M, Otsu K, Kuzuya T, Hori M and Tada M 1998 Intercellular calcium signaling via gap junction in connexin-43-transfected cells *J. Biol. Chem.* **273** 1519–28
- [233] Stains J P and Civitelli R 2004 Gap junctions regulate extracellular signal-regulated kinase (ERK) signaling to affect gene transcription *J. Mol. Biol. Cell* **16** 64–72
- [234] Palacios-Prado N and Bukauskas F F 2012 Modulation of metabolic communication through gap junction channels by transjunctional voltage; synergistic and antagonistic effects of gating and ionophoresis *Biochim. Biophys. Acta* **1818** 1884–94
- [235] Fukumoto T, Kema I P and Levin M 2005 Serotonin signaling is a very early step in patterning of the left-right axis in chick and frog embryos *Curr. Biol.* **15** 794–803
- [236] Fukumoto T, Blakely R and Levin M 2005 Serotonin transporter function is an early step in left-right patterning in chick and frog embryos *Dev. Neurosci.* **27** 349–63
- [237] Chernet B T and Levin M 2013 Transmembrane voltage potential is an essential cellular parameter for the detection and control of tumor development in a *Xenopus* model *Dis. Model. Mech.* **6** 595–607
- [238] Tseng A S and Levin M 2012 Transducing bioelectric signals into epigenetic pathways during tadpole tail regeneration *Anat. Rec.* **295** 1541–51
- [239] Worby C A and Dixon J E 2005 Phosphoinositide phosphatases: emerging roles as voltage sensors? *Mol. Interv.* **5** 274–7
- [240] Murata Y, Iwasaki H, Sasaki M, Inaba K and Okamura Y 2005 Phosphoinositide phosphatase activity coupled to an intrinsic voltage sensor *Nature* **435** 1239–43
- [241] Okamura Y and Dixon J E 2011 Voltage-sensing phosphatase: its molecular relationship with PTEN *Physiol.* **26** 6–13
- [242] Lacroix J, Halaszovich C R, Schreiber D N, Leitner M G, Bezanilla F, Oliver D and Villalba-Galea C A 2011 Controlling the activity of a phosphatase and tensin homolog (PTEN) by membrane potential *J. Biol. Chem.* **286** 17945–53
- [243] Carneiro K, Donnet C, Rejtar T, Karger B L, Barisone G A, Diaz E, Kortagere S, Lemire J M and Levin M 2011 Histone deacetylase activity is necessary for left-right patterning during vertebrate development *BMC Dev. Biol.* **11** 29
- [244] Adams D S and Levin M 2013 Endogenous voltage gradients as mediators of cell-cell communication: strategies for investigating bioelectrical signals during pattern formation *Cell Tissue Res.* **352** 95–122
- [245] Adams D S 2008 A new tool for tissue engineers: ions as regulators of morphogenesis during development and regeneration *Tissue Eng. A* **14** 1461–8
- [246] Levi M 2007 Large-scale biophysics: ion flows and regeneration *Trends Cell Biol.* **17** 262–71
- [247] Mutoh H, Perron A, Akemann W, Iwamoto Y and Knopfel T 2011 Optogenetic monitoring of membrane potentials *Exp. Physiol.* **96** 13–8
- [248] Hotary K B and Robinson K R 1992 Evidence of a role for endogenous electrical fields in chick embryo development *Development* **114** 985–96
- [249] Shi R and Borgens R B 1995 Three-dimensional gradients of voltage during development of the nervous system as invisible coordinates for the establishment of embryonic pattern *Dev. Dyn.* **202** 101–14
- [250] Rakic P and Sidman R L 1973 Sequence of developmental abnormalities leading to granule cell deficit in cerebellar cortex of weaver mutant mice *J. Comp. Neurol.* **152** 103–32
- [251] Rakic P and Sidman R L 1973 Weaver mutant mouse cerebellum: defective neuronal migration secondary to abnormality of Bergmann glia *Proc. Natl Acad. Sci. USA* **70** 240–4

- [252] Hatten M E, Liem R K and Mason C A 1986 Weaver mouse cerebellar granule neurons fail to migrate on wild-type astroglial processes *in vitro* *J. Neurosci.* **6** 2676–83
- [253] Patil N, Cox D R, Bhat D, Faham M, Myers R M and Peterson A S 1995 A potassium channel mutation in weaver mice implicates membrane excitability in granule cell differentiation *Nat. Genet.* **11** 126–9
- [254] Civitelli R 2008 Cell-cell communication in the osteoblast/osteocyte lineage *Arch. Biochem. Biophys.* **473** 188–92
- [255] Chopra S S, Stroud D M, Watanabe H, Bennett J S, Burns C G, Wells K S, Yang T, Zhong T P and Roden D M 2010 Voltage-gated sodium channels are required for heart development in zebrafish *Circ. Res.* **106** 1342–50
- [256] Teng G Q, Zhao X, Lees-Miller J P, Quinn F R, Li P, Rancourt D E, London B, Cross J C and Duff H J 2008 Homozygous missense N629D hERG (KCNH2) potassium channel mutation causes developmental defects in the right ventricle and its outflow tract and embryonic lethality *Circ. Res.* **103** 1483–91
- [257] Uzun S, Gokce S and Wagner K 2005 Cystic fibrosis transmembrane conductance regulator gene mutations in infertile males with congenital bilateral absence of the vas deferens *Tohoku J. Exp. Med.* **207** 279–85
- [258] Wilschanski M, Dupuis A, Ellis L, Jarvi K, Zielenski J, Tullis E, Martin S, Corey M, Tsui L C and Durie P 2006 Mutations in the cystic fibrosis transmembrane regulator gene and *in vivo* transepithelial potentials *Am. J. Respir. Crit. Care Med.* **174** 787–94
- [259] Rock J R, Futtner C R and Harfe B D 2008 The transmembrane protein TMEM16A is required for normal development of the murine trachea *Dev. Biol.* **321** 141–9
- [260] Iwashita M, Watanabe M, Ishii M, Chen T, Johnson S L, Kurachi Y, Okada N and Kondo S 2006 Pigment pattern in jaguar/obelix zebrafish is caused by a Kir7.1 mutation: implications for the regulation of melanosome movement *PLoS Genet.* **2** e197
- [261] Watanabe M, Iwashita M, Ishii M, Kurachi Y, Kawakami A, Kondo S and Okada N 2006 Spot pattern of leopard Danio is caused by mutation in the zebrafish connexin41.8 gene *EMBO Rep.* **7** 893–7
- [262] Hoptak-Solga A D, Nielsen S, Jain I, Thummel R, Hyde D R and Iovine M K 2008 Connexin43 (GJA1) is required in the population of dividing cells during fin regeneration *Dev. Biol.* **317** 541–8
- [263] Gloyn A L *et al* 2004 Activating mutations in the gene encoding the ATP-sensitive potassium-channel subunit Kir6.2 and permanent neonatal diabetes *New Engl. J. Med.* **350** 1838–49
- [264] Culiat C T, Stubbs L J, Woychik R P, Russell L B, Johnson D K and Rinchik E M 1995 Deficiency of the beta 3 subunit of the type A gamma-aminobutyric acid receptor causes cleft palate in mice *Nat. Genet.* **11** 344–6
- [265] Wee E L and Zimmerman E F 1985 GABA uptake in embryonic palate mesenchymal cells of two mouse strains *Neurochem. Res.* **10** 1673–88
- [266] Miller R P and Becker B A 1975 Teratogenicity of oral diazepam and diphenylhydantoin in mice *Toxicol. Appl. Pharmacol.* **32** 53–61
- [267] Homanics G E *et al* 1997 Mice devoid of gamma-aminobutyrate type A receptor beta3 subunit have epilepsy, cleft palate and hypersensitive behavior *Proc. Natl Acad. Sci. USA* **94** 4143–8
- [268] Bendahhou S, Donaldson M R, Plaster N M, Tristani-Firouzi M, Fu Y H and Ptacek L J 2003 Defective potassium channel Kir2.1 trafficking underlies Andersen–Tawil syndrome *J. Biol. Chem.* **278** 51779–85
- [269] Dahal G R, Rawson J, Gassaway B, Kwok B, Tong Y, Ptacek L J and Bates E 2012 An inwardly rectifying K⁺ channel is required for patterning *Development* **139** 3653–64
- [270] Debeer P, Van Esch H, Huysmans C, Pijkels E, De Smet L, Van de Ven W, Devriendt K and Fryns J P 2005 Novel GJA1 mutations in patients with oculo-dento-digital dysplasia (ODDD) *Eur. J. Med. Genet.* **48** 377–87
- [271] Pizzuti A, Flex E, Mingarelli R, Salpietro C, Zelante L and Dallapiccola B 2004 A homozygous GJA1 gene mutation causes a Hallermann–Streiff/ODDD spectrum phenotype *Hum. Mutat.* **23** 286–6
- [272] Adams D S, Robinson K R, Fukumoto T, Yuan S, Albertson R C, Yelick P, Kuo L, McSweeney M and Levin M 2006 Early, H⁺-V-ATPase-dependent proton flux is necessary for consistent left-right patterning of non-mammalian vertebrates *Development* **133** 1657–71
- [273] Vandenberg L N, Lemire J M and Levin M 2012 Serotonin has early, cilia-independent roles in *Xenopus* left-right patterning *Dis. Model. Mech.* **6** 261–8
- [274] Pai V P, Aw S, Shomrat T, Lemire J M and Levin M 2012 Transmembrane voltage potential controls embryonic eye patterning in *Xenopus laevis* *Development* **139** 313–23
- [275] Vandenberg L N, Morrie R D and Adams D S 2011 V-ATPase-dependent ectodermal voltage and pH regionalization are required for craniofacial morphogenesis *Dev. Dyn.* **240** 1889–904
- [276] Beane W S, Morokuma J, Adams D S and Levin M 2011 A chemical genetics approach reveals H,K-ATPase-mediated membrane voltage is required for planarian head regeneration *Chem. Biol.* **18** 77–89
- [277] Beane W S, Morokuma J, Lemire J M and Levin M 2013 Bioelectric signaling regulates head and organ size during planarian regeneration *Development* **140** 313–22
- [278] Bodemer C W 1964 Evocation of regrowth phenomena in anuran limbs by electrical stimulation of the nerve supply *Anat. Rec.* **148** 441–57
- [279] Becker R O 1961 The bioelectric factors in amphibian-limb regeneration *J. Bone Jt. Surg. Am.* **43A** 643–56
- [280] Becker R O and Spadaro J A 1972 Electrical stimulation of partial limb regeneration in mammals *Bull. N. Y. Acad. Med.* **48** 627–41
- [281] Borgens R B 1984 Are limb development and limb regeneration both initiated by an integumentary wounding? A hypothesis *Differentiation* **28** 87–93
- [282] Borgens R B, Rouleau M F and DeLanney L E 1983 A steady efflux of ionic current predicts hind limb development in the axolotl *J. Exp. Zool.* **228** 491–503
- [283] Borgens R B, Venable J W Jr and Jaffe L F 1979 Reduction of sodium dependent stump currents disturbs urodele limb regeneration *J. Exp. Zool.* **209** 377–86
- [284] Borgens R B, Venable J W Jr and Jaffe L F 1977 Bioelectricity and regeneration: large currents leave the stumps of regenerating newt limbs *Proc. Natl Acad. Sci. USA* **74** 4528–32
- [285] Borgens R B, Venable J W Jr and Jaffe L F 1977 Bioelectricity and regeneration. I. Initiation of frog limb regeneration by minute currents *J. Exp. Zool.* **200** 403–16
- [286] Adams D S, Masi A and Levin M 2007 H⁺ pump-dependent changes in membrane voltage are an early mechanism necessary and sufficient to induce *Xenopus* tail regeneration *Development* **134** 1323–35
- [287] Tseng A S, Beane W S, Lemire J M, Masi A and Levin M 2010 Induction of vertebrate regeneration by a transient sodium current *J. Neurosci.* **30** 13192–200
- [288] Tseng A and Levin M 2013 Cracking the bioelectric code: probing endogenous ionic controls of pattern formation *Commun. Integr. Biol.* **6** 1–8
- [289] Riss J *et al* 2006 Cancers as wounds that do not heal: differences and similarities between renal regeneration/repair and renal cell carcinoma *Cancer Res.* **66** 7216–24
- [290] Kenny P A, Nelson C M and Bissell M J 2006 The ecology of tumors: by perturbing the microenvironment, wounds and infection may be key to tumor development *Scientist* **20** 30
- [291] Dolberg D S, Hollingsworth R, Hertle M and Bissell M J 1985 Wounding and its role in RSV-mediated tumor formation *Science* **230** 676–8
- [292] Zhao M *et al* 2006 Electrical signals control wound healing through phosphatidylinositol-3-OH kinase-gamma and PTEN *Nature* **442** 457–60
- [293] Rajnicek A M, Foubister L E and McCaig C D 2007 Prioritising guidance cues: directional migration induced by

- substratum contours and electrical gradients is controlled by a rho/cdc42 switch *Dev. Biol.* **312** 448–60
- [294] Pullar C E, Baier B S, Kariya Y, Russell A J, Horst B A, Marinkovich M P and Isseroff R R 2006 Beta4 integrin and epidermal growth factor coordinately regulate electric field-mediated directional migration via Rac1 *Mol. Biol. Cell* **17** 4925–35
- [295] Pullar C E and Isseroff R R 2005 Cyclic AMP mediates keratinocyte directional migration in an electric field *J. Cell Sci.* **118** 2023–34
- [296] Huber S M 2013 Oncochannels *Cell Calcium* **53** 241–55
- [297] Arcangeli A 2011 Ion channels and transporters in cancer. 3. Ion channels in the tumor cell-microenvironment cross talk *Am. J. Physiol. Cell Physiol.* **301** C762–71
- [298] Arcangeli A, Crociani O, Lastraioli E, Masi A, Pillozzi S and Becchetti A 2009 Targeting ion channels in cancer: a novel frontier in antineoplastic therapy *Curr. Med. Chem.* **16** 66–93
- [299] Farias L M et al 2004 Ether a go-go potassium channels as human cervical cancer markers *Cancer Res.* **64** 6996–7001
- [300] Killion J J 1984 Electrical properties of normal and transformed mammalian cells *Biophys. J.* **45** 523–8
- [301] Martinez-Zaguilan R and Gillies R J 1992 A plasma membrane V-type H(+) -ATPase may contribute to elevated intracellular pH (pHin) in some human tumor cells *Ann. N. Y. Acad. Sci.* **671** 478–80
- [302] Martinez-Zaguilan R, Lynch R M, Martinez G M and Gillies R J 1993 Vacuolar-type H(+) -ATPases are functionally expressed in plasma membranes of human tumor cells *Am. J. Physiol.* **265** C1015–29
- [303] Bianchi L et al 1998 HERG encodes a K⁺ current highly conserved in tumors of different histogenesis: a selective advantage for cancer cells? *Cancer Res.* **58** 815–22
- [304] Arcangeli A, Bianchi L, Becchetti A, Faravelli L, Coronello M, Mini E, Olivetto M and Wanke E 1995 A novel inward-rectifying K⁺ current with a cell-cycle dependence governs the resting potential of mammalian neuroblastoma cells *J. Physiol.* **489** 455–71
- [305] McLean L A, Roscoe J, Jorgensen N K, Gorin F A and Cala P M 2000 Malignant gliomas display altered pH regulation by NHE1 compared with nontransformed astrocytes *Am. J. Physiol. Cell Physiol.* **278** C676–88
- [306] Wissenbach U, Niemeyer B A, Fixemer T, Schneidewind A, Trost C, Cavalie A, Reus K, Meese E, Bonkhoff H and Flockerzi V 2001 Expression of CaT-like, a novel calcium-selective channel, correlates with the malignancy of prostate cancer *J. Biol. Chem.* **276** 19461–8
- [307] Wissenbach U, Niemeyer B, Himmerkus N, Fixemer T, Bonkhoff H and Flockerzi V 2004 TRPV6 and prostate cancer: cancer growth beyond the prostate correlates with increased TRPV6 Ca²⁺ channel expression *Biochem. Biophys. Res. Commun.* **322** 1359–63
- [308] Marino A A, Iliev I G, Schwalke M A, Gonzalez E, Marler K C and Flanagan C A 1994 Association between cell membrane potential and breast cancer *Tumour Biol.* **15** 82–9
- [309] Voloshyna I, Besana A, Castillo M, Matos T, Weinstein I B, Mansukhani M, Robinson R B, Cordon-Cardo C and Feinmark S J 2008 TREK-1 is a novel molecular target in prostate cancer *Cancer Res.* **68** 1197–203
- [310] Diss J K, Stewart D, Pani F, Foster C S, Walker M M, Patel A and Djamgoz M B 2005 A potential novel marker for human prostate cancer: voltage-gated sodium channel expression in vivo *Prostate Cancer Prostatic Dis.* **8** 266–73
- [311] Duncan L M, Deeds J, Cronin F E, Donovan M, Sober A J, Kauffman M and McCarthy J J 2001 Melastatin expression and prognosis in cutaneous malignant melanoma *J. Clin. Oncol.* **19** 568–76
- [312] Zhiqi S, Soltani M H, Bhat K M, Sangha N, Fang D, Hunter J J and Setaluri V 2004 Human melastatin 1 (TRPM1) is regulated by MITF and produces multiple polypeptide isoforms in melanocytes and melanoma *Melanoma Res.* **14** 509–16
- [313] Conti M 2004 Targeting K⁺ channels for cancer therapy *J. Exp. Ther. Oncol.* **4** 161–6
- [314] Fraser S P, Grimes J A and Djamgoz M B 2000 Effects of voltage-gated ion channel modulators on rat prostatic cancer cell proliferation: comparison of strongly and weakly metastatic cell lines *Prostate* **44** 61–76
- [315] Schlichter L C, Sakellaropoulos G, Ballyk B, Pennefather P S and Phipps D J 1996 Properties of K⁺ and Cl[−] channels and their involvement in proliferation of rat microglial cells *Glia* **17** 225–36
- [316] Schuldiner S and Rozengurt E 1982 Na⁺/H⁺ antiport in Swiss 3T3 cells: mitogenic stimulation leads to cytoplasmic alkalinization *Proc. Natl Acad. Sci. USA* **79** 7778–82
- [317] Lopez-Rivas A, Adelberg E A and Rozengurt E 1982 Intracellular K⁺ and the mitogenic response of 3T3 cells to peptide factors in serum-free medium *Proc. Natl Acad. Sci. USA* **79** 6275–9
- [318] Burns C P and Rozengurt E 1984 Extracellular Na⁺ and initiation of DNA synthesis: role of intracellular pH and K⁺ *J. Cell Biol.* **98** 1082–9
- [319] Pillozzi S et al 2002 HERG potassium channels are constitutively expressed in primary human acute myeloid leukemias and regulate cell proliferation of normal and leukemic hemopoietic progenitors *Leukemia* **16** 1791–8
- [320] Wohlrab D, Wohlrab J and Markwardt F 2000 Electrophysiological characterization of human keratinocytes using the patch-clamp technique *Exp. Dermatol.* **9** 219–23
- [321] Dalle-Lucca S L, Dalle-Lucca J J, Borges A C, Ihara S S and Paiva T B 2000 Abnormal proliferative response of the carotid artery of spontaneously hypertensive rats after angioplasty may be related to the depolarized state of its smooth muscle cells *Braz. J. Med. Biol. Res.* **33** 919–27
- [322] Wohlrab D and Hein W 2000 Der Einfluss von Ionenkanalmodulatoren auf das Membranpotential humaner Chondrozyten *Orthopade* **29** 80–4
- [323] MacFarlane S N and Sontheimer H 2000 Changes in ion channel expression accompany cell cycle progression of spinal cord astrocytes *Glia* **30** 39–48
- [324] Kamleiter M, Hanemann C O, Kluwe L, Rosenbaum C, Wosch S, Mautner V F, Muller H W and Grafe P 1998 Voltage-dependent membrane currents of cultured human neurofibromatosis type 2 Schwann cells *Glia* **24** 313–22
- [325] Wang S, Melkounian Z, Woodfork K A, Cather C, Davidson A G, Wonderlin W F and Strobl J S 1998 Evidence for an early G1 ionic event necessary for cell cycle progression and survival in the MCF-7 human breast carcinoma cell line *J. Cell. Physiol.* **176** 456–64
- [326] Knutson P, Ghiani C A, Zhou J M, Gallo V and McBain C J 1997 K⁺ channel expression and cell proliferation are regulated by intracellular sodium and membrane depolarization in oligodendrocyte progenitor cells *J. Neurosci.* **17** 2669–82
- [327] Puro D G, Roberge F and Chan C C 1989 Retinal glial cell proliferation and ion channels: a possible link *Invest. Ophthalmol. Vis. Sci.* **30** 521–9
- [328] Banderali U, Belke D, Singh A, Jayanthan A, Giles W R and Narendran A 2011 Curcumin blocks Kv11.1 (erg) potassium current and slows proliferation in the infant acute monocytic leukemia cell line THP-1 *Cell. Physiol. Biochem.* **28** 1169–80
- [329] Lin H, Xiao J, Luo X, Wang H, Gao H, Yang B and Wang Z 2007 Overexpression HERG K⁺ channel gene mediates cell-growth signals on activation of oncoproteins SP1 and NF-KB and inactivation of tumor suppressor Nkx3.1 *J. Cell. Physiol.* **212** 137–47
- [330] Wang H, Zhang Y, Cao L, Han H, Wang J, Yang B, Nattel S and Wang Z 2002 HERG K⁺ channel, a regulator of tumor cell apoptosis and proliferation *Cancer Res.* **62** 4843–8
- [331] Lastraioli E et al 2004 HERG1 gene and HERG1 protein are overexpressed in colorectal cancers and regulate cell invasion of tumor cells *Cancer Res.* **64** 606–11
- [332] Wang Z 2004 Roles of K⁺ channels in regulating tumour cell proliferation and apoptosis *Pflugers Arch.* **448** 274–86
- [333] Arcangeli A 2005 Expression and role of hERG channels in cancer cells *Novartis Found Symp.* **266** 224–5
- [334] Klezovitch O, Risk M, Coleman I, Lucas J M, Null M, True L D, Nelson P S and Vasioukhin V 2008 A causal role for ERG in

- neoplastic transformation of prostate epithelium *Proc. Natl Acad. Sci. USA* **105** 2105–10
- [335] Mu D *et al* 2003 Genomic amplification and oncogenic properties of the KCNK9 potassium channel gene *Cancer Cell* **3** 297–302
- [336] Pei L, Wisner O, Slavin A, Mu D, Powers S, Jan L Y and Hoey T 2003 Oncogenic potential of TASK3 (Kcnk9) depends on K⁺ channel function *Proc. Natl Acad. Sci. USA* **100** 7803–7
- [337] Bennett E S, Smith B A and Harper J M 2004 Voltage-gated Na⁺ channels confer invasive properties on human prostate cancer cells *Pflugers Arch.* **447** 908–14
- [338] Brackenbury W J and Djamgoz M B 2006 Activity-dependent regulation of voltage-gated Na⁺ channel expression in Mat-LyLu rat prostate cancer cell line *J. Physiol.* **573** 343–56
- [339] Onganer P U, Seckl M J and Djamgoz M B 2005 Neuronal characteristics of small-cell lung cancer *Br. J. Cancer* **93** 1197–201
- [340] Onganer P U and Djamgoz M B 2005 Small-cell lung cancer (human): potentiation of endocytic membrane activity by voltage-gated Na(+) channel expression *in vitro J. Membr. Biol.* **204** 67–75
- [341] Fraser S P *et al* 2005 Voltage-gated sodium channel expression and potentiation of human breast cancer metastasis *Clin. Cancer Res.* **11** 5381–9
- [342] Liu L Y, Hu C L, Ma L J, Zhang Z H and Mei Y A 2004 ET-1 inhibits B-16 murine melanoma cell migration by decreasing K(+) currents *Cell Motil. Cytoskeleton.* **58** 127–36
- [343] Mycielska M E and Djamgoz M B 2004 Cellular mechanisms of direct-current electric field effects: galvanotaxis and metastatic disease *J. Cell Sci.* **117** 1631–9
- [344] Fraser S P, Ozerlat-Gunduz I, Onkal R, Diss J K, Latchman D S and Djamgoz M B 2010 Estrogen and non-genomic upregulation of voltage-gated Na(+) channel activity in MDA-MB-231 human breast cancer cells: role in adhesion *J. Cell. Physiol.* **224** 527–39
- [345] Onkal R, Mattis J H, Fraser S P, Diss J K, Shao D, Okuse K and Djamgoz M B 2008 Alternative splicing of Nav1.5: an electrophysiological comparison of ‘neonatal’ and ‘adult’ isoforms and critical involvement of a lysine residue *J. Cell Physiol.* **216** 716–26
- [346] Diss J K, Fraser S P, Walker M M, Patel A, Latchman D S and Djamgoz M B 2008 Beta-subunits of voltage-gated sodium channels in human prostate cancer: quantitative *in vitro* and *in vivo* analyses of mRNA expression *Prostate Cancer Prostatic Dis.* **11** 325–33
- [347] Ding Y, Brackenbury W J, Onganer P U, Montano X, Porter L M, Bates L F and Djamgoz M B 2008 Epidermal growth factor upregulates motility of Mat-LyLu rat prostate cancer cells partially via voltage-gated Na⁺ channel activity *J. Cell Physiol.* **215** 77–81
- [348] Uysal-Onganer P and Djamgoz M B 2007 Epidermal growth factor potentiates *in vitro* metastatic behaviour of human prostate cancer PC-3M cells: involvement of voltage-gated sodium channel *Mol. Cancer* **6** 76
- [349] Brackenbury W J and Djamgoz M B 2007 Nerve growth factor enhances voltage-gated Na⁺ channel activity and transwell migration in Mat-LyLu rat prostate cancer cell line *J. Cell Physiol.* **210** 602–8
- [350] Takamori M 1999 An autoimmune channelopathy associated with cancer: Lambert–Eaton myasthenic syndrome *Intern. Med.* **38** 86–96
- [351] Weksberg R *et al* 2002 Discordant KCNQ1OT1 imprinting in sets of monozygotic twins discordant for Beckwith–Wiedemann syndrome *Hum. Mol. Genet.* **11** 1317–25
- [352] Weksberg R *et al* 2001 Tumor development in the Beckwith–Wiedemann syndrome is associated with a variety of constitutional molecular 11p15 alterations including imprinting defects of KCNQ1OT1 *Hum. Mol. Genet.* **10** 2989–3000
- [353] Lee M P, Hu R J, Johnson L A and Feinberg A P 1997 Human KVLQT1 gene shows tissue-specific imprinting and encompasses Beckwith–Wiedemann syndrome chromosomal rearrangements *Nat. Genet.* **15** 181–5
- [354] Frede J, Fraser S P, Oskay-Ozcelik G, Hong Y, Ioana Braicu E, Sehouli J, Gabra H and Djamgoz M B 2013 Ovarian cancer: ion channel and aquaporin expression as novel targets of clinical potential *Eur. J. Cancer* **49** 2331–44
- [355] Speyer C L, Smith J S, Banda M, DeVries J A, Mekani T and Gorski D H 2012 Metabotropic glutamate receptor-1: a potential therapeutic target for the treatment of breast cancer *Breast Cancer Res. Treat.* **132** 565–73
- [356] Yildirim S, Altun S, Gumushan H, Patel A and Djamgoz M B 2012 Voltage-gated sodium channel activity promotes prostate cancer metastasis *in vivo Cancer Lett.* **323** 58–61
- [357] Spugnini E P, Citro G and Fais S 2010 Proton pump inhibitors as anti vacuolar-ATPases drugs: a novel anticancer strategy *J. Exp. Clin. Cancer Res.* **29** 44
- [358] Gupta P B, Onder T T, Jiang G, Tao K, Kuperwasser C, Weinberg R A and Lander E S 2009 Identification of selective inhibitors of cancer stem cells by high-throughput screening *Cell* **138** 645–59
- [359] Schickling B M, Aykin-Burns N, Leslie K K, Spitz D R and Korovkina V P 2011 An inhibitor of K⁺ channels modulates human endometrial tumor-initiating cells *Cancer Cell Int.* **11** 25
- [360] Fraser S P, Peters A, Fleming-Jones S, Mukhey D and Djamgoz M B 2014 Resveratrol: inhibitory effects on metastatic cell behaviors and voltage-gated Na(+) channel activity in rat prostate cancer *in vitro Nutrition Cancer* **66** 1047–58
- [361] Yang S, Yu B, Sui Y J, Zhang Y F, Wang X, Hou S G, Ma T H and Yang H 2013 CFTR chloride channel is a molecular target of the natural cancer preventive agent resveratrol *Pharmazie* **68** 772–6
- [362] Shen Y A, Lin C H, Chi W H, Wang C Y, Hsieh Y T, Wei Y H and Chen Y J 2013 Resveratrol impedes the stemness, epithelial-mesenchymal transition and metabolic reprogramming of cancer stem cells in nasopharyngeal carcinoma through p53 activation *Evid. Based. Complement. Alternat. Med.* **2013** 590393
- [363] Kometiani P, Liu L and Askari A 2005 Digitalis-induced signaling by Na⁺/K⁺-ATPase in human breast cancer cells *Mol. Pharmacol.* **67** 929–36
- [364] Blackiston D J, McLaughlin K A and Levin M 2009 Bioelectric controls of cell proliferation: ion channels, membrane voltage and the cell cycle *Cell Cycle* **8** 3519–28
- [365] Rubin H 1992 Mechanisms for enduring biological change *Am. J. Physiol.* **262** L111–3
- [366] Rubin H 1990 On the nature of enduring modifications induced in cells and organisms *Am. J. Physiol.* **258** L19–24
- [367] Stojadinovic A *et al* 2005 Electrical impedance scanning for the early detection of breast cancer in young women: preliminary results of a multicenter prospective clinical trial *J. Clin. Oncol.* **23** 2703–15
- [368] Bauer J, Capra S and Davies P S 2005 Estimation of total body water from foot-to-foot bioelectrical impedance analysis in patients with cancer cachexia—agreement between three prediction methods and deuterium oxide dilution *J. Hum. Nutrition Diet.* **18** 295–300
- [369] Aberg P, Geladi P, Nicander I, Hansson J, Holmgren U and Ollmar S 2005 Non-invasive and microinvasive electrical impedance spectra of skin cancer—a comparison between two techniques *Ski. Res. Technol.* **11** 281–6
- [370] Gupta D, Lis C G, Dahlk S L, Vashi P G, Grutsch J F and Lammersfeld C A 2004 Bioelectrical impedance phase angle as a prognostic indicator in advanced pancreatic cancer *Br. J. Nutrition* **92** 957–62
- [371] Gupta D, Lammersfeld C A, Burrows J L, Dahlk S L, Vashi P G, Grutsch J F, Hoffman S and Lis C G 2004 Bioelectrical impedance phase angle in clinical practice: implications for prognosis in advanced colorectal cancer *Am. J. Clin. Nutrition* **80** 1634–8
- [372] Gatta L 2004 The clinical role of extracellular bioimpedance tomography (Gastro-Mida(x)) in the diagnosis of colorectal diseases *Minerva Med.* **95** 541–56
- [373] Aberg P, Nicander I, Hansson J, Geladi P, Holmgren U and Ollmar S 2004 Skin cancer identification using

- multifrequency electrical impedance—a potential screening tool *IEEE Trans. Biomed. Eng.* **51** 2097–102
- [374] Toso S, Piccoli A, Gusella M, Menon D, Bononi A, Crepaldi G and Ferrazzi E 2000 Altered tissue electric properties in lung cancer patients as detected by bioelectric impedance vector analysis *Nutrition* **16** 120–4
- [375] Konishi Y, Morimoto T, Kinouchi Y, Iritani T and Monden Y 1995 Electrical properties of extracted rat liver tissue *Res. Exp. Med.* **195** 183–92
- [376] Kimura S, Morimoto T, Uyama T, Monden Y, Kinouchi Y and Iritani T 1994 Application of electrical impedance analysis for diagnosis of a pulmonary mass *Chest* **105** 1679–82
- [377] Morimoto T, Kimura S, Konishi Y, Komaki K, Uyama T, Monden Y, Kinouchi Y and Iritani T 1993 A study of the electrical bio-impedance of tumors *J. Invest. Surg.* **6** 25–32
- [378] Morimoto T, Kinouchi Y, Iritani T, Kimura S, Konishi Y, Mitsuyama N, Komaki K and Monden Y 1990 Measurement of the electrical bio-impedance of breast tumors *Eur. Surg. Res.* **22** 86–92
- [379] Davies R J, Joseph R, Asbun H and Sedwitz M 1989 Detection of the cancer-prone colon, using transepithelial impedance analysis *Arch. Surg.* **124** 480–4
- [380] Cuzick J et al 1998 Electropotential measurements as a new diagnostic modality for breast cancer *Lancet* **352** 359–63
- [381] Faupel M, Vanel D, Barth V, Davies R, Fentiman I S, Holland R, Lamarque J L, Sacchini V and Schreer I 1997 Electropotential evaluation as a new technique for diagnosing breast lesions *Eur. J. Radiol.* **24** 33–8
- [382] Moyer M P, Moyer R C and Waite M R 1982 A survey of intracellular Na⁺ and K⁺ of various normal, transformed and tumor cells *J. Cell. Physiol.* **113** 129–33
- [383] Ouwerkerk R, Jacobs M A, Macura K J, Wolff A C, Stearns V, Mezban S D, Khouri N F, Bluemke D A and Bottomley P A 2007 Elevated tissue sodium concentration in malignant breast lesions detected with non-invasive ²³Na MRI *Breast Cancer Res. Treat.* **106** 151–60
- [384] Beech J A 1997 Bioelectric potential gradients may initiate cell cycling: ELF and zeta potential gradients may mimic this effect *Bioelectromagnetics* **18** 341–8
- [385] Cone C D 1971 Unified theory on the basic mechanism of normal mitotic control and oncogenesis *J. Theor. Biol.* **30** 151–81
- [386] Cone C D 1974 The role of the surface electrical transmembrane potential in normal and malignant mitogenesis *Ann. N. Y. Acad. Sci.* **238** 420–35
- [387] Binggeli R and Cameron I L 1980 Cellular potentials of normal and cancerous fibroblasts and hepatocytes *Cancer Res.* **40** 1830–5
- [388] Dahmane N, Lee J, Robins P, Heller P and Ruiz i Altaba A 1997 Activation of the transcription factor Gli1 and the Sonic hedgehog signalling pathway in skin tumours *Nature* **389** 876–81
- [389] Wallingford J B, Seufert D W, Virta V C and Vize P D 1997 p53 activity is essential for normal development in *Xenopus* *Curr. Biol.* **7** 747–57
- [390] Yang S, Lockwood A, Hollett P, Ford R and Kao K 1998 Overexpression of a novel *Xenopus* rel mRNA gene induces tumors in early embryos *J. Biol. Chem.* **273** 13746–52
- [391] Le X, Langenau D M, Keefe M D, Kutok J L, Neuberger D S and Zon L I 2007 Heat shock-inducible Cre/Lox approaches to induce diverse types of tumors and hyperplasia in transgenic zebrafish *Proc. Natl Acad. Sci. USA* **104** 9410–5
- [392] Stratton M R, Cooper C S, Gusterson B A and Fisher C 1989 Detection of point mutations in N-ras and K-ras genes of human embryonal rhabdomyosarcomas using oligonucleotide probes and the polymerase chain reaction *Cancer Res.* **49** 6324–7
- [393] Gilmore T D, Kalaitzidis D, Liang M C and Starczynowski D T 2004 The c-Rel transcription factor and B-cell proliferation: a deal with the devil *Oncogene* **23** 2275–86
- [394] McNulty S E, Del Rosario R, Cen D, Meyskens F L and Yang S 2004 Comparative expression of NF- κ B proteins in melanocytes of normal skin versus benign intradermal naevus and human metastatic melanoma biopsies *Pigment Cell Res.* **17** 173–80
- [395] Clement V, Sanchez P, de Tribolet N, Radovanovic I and Ruiz i Altaba A 2007 HEDGEHOG-GLI1 signaling regulates human glioma growth, cancer stem cell self-renewal and tumorigenicity *Curr. Biol.* **17** 165–72
- [396] Wolff C, Fuks B and Chatelain P 2003 Comparative study of membrane potential-sensitive fluorescent probes and their use in ion channel screening assays *J. Biomol. Screen* **8** 533–43
- [397] Lobikin M, Chernet B, Lobo D and Levin M 2012 Resting potential, oncogene-induced tumorigenesis and metastasis: the bioelectric basis of cancer *in vivo Phys. Biol.* **9** 65002
- [398] Chernet B T and Levin M 2014 Transmembrane voltage potential of somatic cells controls oncogene-mediated tumorigenesis at long-range *Oncotarget* **5** 3287–306
- [399] Thompson I M et al 2004 Prevalence of prostate cancer among men with a prostate-specific antigen level ≤ 4.0 ng per milliliter *New Engl. J. Med.* **350** 2239–46
- [400] Smith D S, Humphrey P A and Catalona W J 1997 The early detection of prostate carcinoma with prostate specific antigen: the Washington University experience *Cancer* **80** 1852–6
- [401] Burr H S, Strong L C and Smith G M 1938 Bioelectric correlates of methylcholanthrene-induced tumors in mice *Yale J. Biol. Med.* **10** 539–44
- [402] Burr H S 1952 Electrometrics of atypical growth *Yale J. Biol. Med.* **25** 67–75
- [403] Burr H S 1941 Changes in the field properties of mice with transplanted tumors *Yale J. Biol. Med.* **13** 783–8
- [404] Nordenstrom B E 1994 The paradigm of biologically closed electric circuits (BCEC) and the formation of an International Association (IABC) for BCEC systems *Eur. J. Surg. Suppl.* **574** 7–23
- [405] Nordenstrom B, Naslund I, Nordenstrom J and Wersall P 1994 *Proc. of the IABC International-Association-for-Biologically-Closed-Electric-Circuits (BCEC) in Medicine and Biology—Main Topic—Activation of BCEC-Channels for Electrochemical Therapy (ECT) of Cancer—Foreword (Stockholm, 12–15 September 1993)* (*Eur. J. Surg.*) p 5
- [406] Nordenstrom B E 1992 Impact of biologically closed electric circuits (BCEC) on structure and function *Integr. Physiol. Behav. Sci.* **27** 285–303
- [407] Lobikin M, Lobo D, Blackiston D J, Martyniuk C J, Tkachenko E and Levin M 2015 Serotonergic regulation of melanocyte conversion: a bioelectrically regulated network for stochastic all-or-none hyperpigmentation *Sci. Signal* **8** ra99
- [408] Lobo D, Lobikin M and Levin M 2017 Discovering novel phenotypes with automatically inferred dynamic models: a partial melanocyte conversion in *Xenopus* *Sci. Rep.* **7** 41339
- [409] Vafa B, Lewis T M, Cunningham A M, Jacques P, Lynch J W and Schofield P R 1999 Identification of a new ligand binding domain in the $\alpha 1$ subunit of the inhibitory glycine receptor *J. Neurochem.* **73** 2158–66
- [410] Aw S, Adams D S, Qiu D and Levin M 2008 H,K-ATPase protein localization and Kir4.1 function reveal concordance of three axes during early determination of left-right asymmetry *Mech. Dev.* **125** 353–72
- [411] O'Connor P J, MacNaught F, Butler W H, Cooper D P, Margison G P and Povey A C 2001 Increased pathology incidence in the forestomach of rats maintained on a diet containing ivermectin and given a single dose of N-methyl-N1-nitro-N-nitrosoguanidine *J. Environ. Pathol. Toxicol. Oncol.* **20** 223–7
- [412] Chernet B T, Adams D S, Lobikin M and Levin M 2016 Use of genetically encoded, light-gated ion translocators to control tumorigenesis *Oncotarget* **7** 19575–88
- [413] Chernet B T, Fields C and Levin M 2015 Long-range gap junctional signaling controls oncogene-mediated tumorigenesis in *Xenopus laevis* embryos *Front. Physiol.* **5** 519
- [414] Rubin H 2006 What keeps cells in tissues behaving normally in the face of myriad mutations? *BioEssays* **28** 515–24
- [415] Prokopenko M, Gerasimov V and Tanev I 2008 Evolving spatiotemporal coordination in a modular robotic system *From Animals to Animals 9: 9th Int. Conf. on Simulation*

- of *Adaptive Behavior*, SAB 2006, Proc. (Rome, Italy, 25-29 September 2006) ed S Nolfi et al (Berlin: Springer) pp 558–69
- [416] Nurse P 2008 Life, logic and information *Nature* **454** 424–6
- [417] Lizier J T, Prokopenko M and Zomaya A Y 2012 Local measures of information storage in complex distributed computation *Inf. Sci.* **208** 39–54
- [418] Wang X R, Miller J M, Lizier J T, Prokopenko M and Rossi L F 2011 Measuring information storage and transfer in swarms *Proc. Elev. Eur. Conf. Synth. Simul. Living Syst. (ECAL 2011)* pp 838–45
- [419] Wang X R, Miller J M, Lizier J T, Prokopenko M and Rossi L F 2012 Quantifying and tracing information cascades in swarms *PLoS One* **7** e40084
- [420] Miller J M, Wang X R, Lizier J T, Prokopenko M and Rossi L F 2014 Measuring information dynamics in swarms *Guided Self-Organization: Inception* ed M Prokopenko (Berlin: Springer) pp 343–64
- [421] Kim H, Davies P and Walker S I 2015 New scaling relation for information transfer in biological networks *J. R. Soc. Interface* **12** 20150944
- [422] Schreiber T 2011 Measuring information transfer *Phys. Rev. Lett.* **85** 461–4
- [423] Vicente R, Wibral M, Lindner M and Pipa G 2011 Transfer entropy—a model-free measure of effective connectivity for the neurosciences *J. Comput. Neurosci.* **30** 45–67
- [424] Nigam S et al 2016 Rich-club organization in effective connectivity among cortical neurons *J. Neurosci.* **36** 670–84
- [425] Honey C J, Kotter R, Breakspear M and Sporns O 2007 Network structure of cerebral cortex shapes functional connectivity on multiple time scales *Proc. Natl Acad. Sci.* **104** 10240–5
- [426] Chávez M, Martinier J and Le Van Quyen M 2003 Statistical assessment of nonlinear causality: application to epileptic EEG signals *J. Neurosci. Methods* **124** 113–28
- [427] Hinrichs H, Heinze H J and Schoenfeld M A 2006 Causal visual interactions as revealed by an information theoretic measure and fMRI *Neuroimage* **31** 1051–60
- [428] Lizier J T, Heinze H J, Horstmann A, Haynes J-D and Prokopenko M 2011 Multivariate information-theoretic measures reveal directed information structure and task relevant changes in fMRI connectivity *J. Comput. Neurosci.* **30** 85–107
- [429] Grossberg S 1978 Communication, memory and development *Prog. Theor. Biol.* **5** 183–232
- [430] Walker S I, Kim H and Davies P C W 2016 The informational architecture of the cell *Phil. Trans. R. Soc. A* **374** 20150057
- [431] Davies P C W and Walker S I 2016 The hidden simplicity of biology *Rep. Prog. Phys.* **79** 102601
- [432] Kauffman S A 1969 Metabolic stability and epigenesis in randomly constructed genetic nets *J. Theor. Biol.* **22** 437–67
- [433] Davidich M I and Bornholdt S 2007 Boolean network model predicts cell cycle sequence of fission yeast *PLoS One* **3** e1672
- [434] Kim J, Park S-M and Cho K-H 2013 Discovery of a kernel for controlling biomolecular regulatory networks *Sci. Rep.* **3** 2223
- [435] Choi M, Shi J, Jung S H, Chen X and Cho K-H 2012 Attractor landscape analysis reveals feedback loops in the p53 network that control the cellular response to DNA damage *Sci. Signal* **5** ra83
- [436] Hahn W C and Weinberg R A 2002 Rules for making human tumor cells *New Engl. J. Med.* **347** 1593–603
- [437] Huang S and Ingber D E 2006 A non-genetic basis for cancer progression and metastasis: self-organizing attractors in cell regulatory networks *Breast Dis.* **26** 27–54
- [438] Sullivan K G, Emmons-Bell M and Levin M 2016 Physiological inputs regulate species-specific anatomy during embryogenesis and regeneration *Commun. Integr. Biol.* **9** e192733
- [439] Matsuno K 2014 Self-identities and durability of biosystems via their abstracting capacity *Biosystems* **120** 31–4
- [440] Pattee H H 2001 The physics of symbols: bridging the epistemic cut *Biosystems* **60** 5–21
- [441] Rosen R 1985 *Anticipatory Systems: Philosophical, Mathematical and Methodological Foundations* (Oxford: Pergamon)
- [442] Bertschinger N, Olbrich E, Ay N and Jost J 2008 Autonomy: an information theoretic perspective *Biosystems* **91** 331–45
- [443] Duthen Y, Luga H, Lassabe N, Cussat-Blanc S, Breton T and Pascale J 2010 An introduction to the bio-logic of artificial creatures *Intell. Comput. Graph.* **321** 1–23
- [444] Schmidhuber J 2007 Prototype resilient, self-modeling robots *Science* **316** 688
- [445] Bongard J, Zykov V and Lipson H 2006 Resilient machines through continuous self-modeling *Science* **314** 1118–21
- [446] Pattee H H 1989 The measurement problem in artificial world models *Biosystems* **23** 281–9
- [447] Pattee H H 2012 Cell psychology: an evolutionary approach to the symbol-matter problem *LAWS, LANGUAGE and LIFE: Howard Pattee's Classic Papers on the Physics of Symbols with Contemporary Commentary* (Dordrecht: Springer) pp 165–79
- [448] Tarin D 2013 Role of the host stroma in cancer and its therapeutic significance *Cancer Metastasis Rev.* **32** 553–66
- [449] Egeblad M, Nakasone E S and Werb Z 2010 Tumors as organs: complex tissues that interface with the entire organism *Dev. Cell* **18** 884–901
- [450] Vincent M 2012 Cancer: a de-repression of a default survival program common to all cells? A life-history perspective on the nature of cancer *Bioessays* **34** 72–82
- [451] Dean M 1998 Cancer as a complex developmental disorder—nineteenth Cornelius P. Rhoads Memorial Award Lecture *Cancer Res.* **58** 5633–6
- [452] Chabner B A and Roberts T G Jr 2005 Timeline: chemotherapy and the war on cancer *Nat. Rev. Cancer* **5** 65–72
- [453] Nooter K and Herweijer H 1991 Multidrug resistance (mdr) genes in human cancer *Br. J. Cancer* **63** 663–9
- [454] Krishna R and Mayer L D 2000 Multidrug resistance (MDR) in cancer. Mechanisms, reversal using modulators of MDR and the role of MDR modulators in influencing the pharmacokinetics of anticancer drugs *Eur. J. Pharm. Sci.* **11** 265–83
- [455] Shah N P, Skaggs B J, Branford S, Hughes T P, Nicoll J M, Paquette R L and Sawyers C L 2007 Sequential ABL kinase inhibitor therapy selects for compound drug-resistant BCR-ABL mutations with altered oncogenic potency *J. Clin. Invest.* **117** 2562–9
- [456] Deisboeck T S, Berens M E, Kansal A R, Torquato S, Stemmer-Rachamimov A O and Chiocca E A 2001 Pattern of self-organization in tumour systems: complex growth dynamics in a novel brain tumour spheroid model *Cell Prolif.* **34** 115–34
- [457] Guttal V and Couzin I D 2010 Social interactions, information use and the evolution of collective migration *Proc. Natl Acad. Sci. USA* **107** 16172–7
- [458] Deisboeck T S and Couzin I D 2009 Collective behavior in cancer cell populations *Bioessays* **31** 190–7
- [459] Couzin I D 2009 Collective cognition in animal groups *Trends Cogn. Sci.* **13** 36–43
- [460] Couzin I D, Krause J, James R, Ruxton G D and Franks N R 2002 Collective memory and spatial sorting in animal groups *J. Theor. Biol.* **218** 1–11
- [461] Tononi G 2008 Consciousness as integrated information: a provisional manifesto *Biol. Bull.* **215** 216–42
- [462] Oizumi M, Albantakis L and Tononi G 2014 From the phenomenology to the mechanisms of consciousness: integrated information theory 3.0 *PLoS Comput. Biol.* **10** e1003588
- [463] Biehl M, Ikegami T and Polani D 2016 Towards information based spatiotemporal patterns as a foundation for agent representation in dynamical systems *Proc. of the Artificial Life Conf. 2016* pp 722–9
- [464] Hoel E P, Albantakis L and Tononi G 2013 Quantifying causal emergence shows that macro can beat micro *Proc. Natl Acad. Sci.* **110** 19790–5
- [465] Marshall W, Kim H, Walker S I, Tononi G and Albantakis L 2017 How causal analysis can reveal autonomy in biological systems *Phil. Trans. A submitted*
- [466] Wang G, Du C, Chen H, Simha R, Rong Y, Xiao Y and Zeng C 2010 Process-based network decomposition reveals backbone motif structure *Proc. Natl Acad. Sci. USA* **107** 10478–83
- [467] Tegmark M 2016 Improved measures of integrated information *PLOS Comput. Biol.* **12** e1005123

- [468] McDonough W S, Johansson A, Joffe H, Giese A and Berens M E 1999 Gap junction intercellular communication in gliomas is inversely related to cell motility *Int. J. Dev. Neurosci.* **17** 601–11
- [469] Oliveira R, Christov C, Guillamo J S, De Boüard S, Palfi S, Venance L, Tardy M and Peschanski M 2005 Contribution of gap junctional communication between tumor cells and astroglia to the invasion of the brain parenchyma by human glioblastomas *BMC Cell Biol.* **6** 7
- [470] Barandiaran X E, Di Paolo E and Rohde M 2009 Defining agency: individuality, normativity, asymmetry and spatio-temporality in action *Adapt. Behav.* **17** 367–86
- [471] Katakura Y, Alam S and Shirahata S 1998 Immortalization by gene transfection *Methods Cell Biol.* **57** 69–91
- [472] Smith J R, Ning Y and Pereira-Smith O M 1992 Why are transformed cells immortal? Is the process reversible? *Am. J. Clin. Nutr.* **55** 1215S–21S
- [473] Noble M, Groves A K, Ataliotis P and Jat P S 1992 From chance to choice in the generation of neural cell lines *Brain Pathol.* **2** 39–46
- [474] Friston K 2008 Life as we know it *J. R. Soc. Interface* **10** 20130475
- [475] Ay N and Polani D 2008 Information flows in causal networks *Adv. Complex Syst.* **11** 17–41
- [476] Fano R M and Hawkins D 1961 Transmission of information: a statistical theory of communications *Am. J. Phys.* **29** 793–4
- [477] Bugaj L J, O'Donoghue G P, Lim W A, O'Donoghue G P and Lim W A 2017 Interrogating cellular perception and decision making with optogenetic tools *J. Cell Biol.* **216** 25LP-28
- [478] Levin M 2013 Reprogramming cells and tissue patterning via bioelectrical pathways: molecular mechanisms and biomedical opportunities *Wiley Interdiscip. Rev. Syst. Biol. Med.* **5** 657–76