

AP Calculus Homework Four – Differentiation

2.3 Implicit Differentiation; 2.4 Estimating a Derivative; 2.5 Derivative of the Inverse of a Function

1. Find $\frac{dy}{dx}$

$$a) (x)' + [\cos(x+y)]' = (0)' \quad 1+y' = \frac{1}{\sin(x+y)}$$

$$\Rightarrow 1 - \sin(x+y)(x'+y') = 0 \Rightarrow \therefore y' = \csc(x+y) - 1$$

(a) $x + \cos(x+y) = 0$

$$b) (\sin x)' - (\cos y)' - (2)' = 0 \Rightarrow \cos x + \sin y \cdot y' = 0$$

$$\therefore y' = -\frac{\cos x}{\sin y}$$

(b) $\sin x - \cos y - 2 = 0$

(c) $3x^2 - 2xy + 5y^2 = 1$

$$c) 3(x^2)' - 2(xy)' + 5(y^2)' = (1)'$$

$$\Rightarrow 3(2x) - 2(x'y + xy') + 5(2y \cdot y') = 0$$

$$\Rightarrow 6x - 2y - 2xy' + 10yy' = 0 \Rightarrow y' = \frac{2y - 6x}{10y - 2x} = \frac{y - 3x}{5y - x}$$

(d) $\sin(xy) = x$

$$y + xy' = \sec(xy)$$

$$\therefore y' = \frac{\sec(xy) - y}{x}$$

(e) $\begin{cases} x = t - \sin t \\ y = 1 - \cos t \end{cases}$

$$e) \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{(1 - \cos t)'}{(t - \sin t)'} = \frac{\sin t}{1 - \cos t}$$

(f) $\begin{cases} x = 1 - e^{-t} \\ y = t + e^{-t} \end{cases}$

$$f) \frac{dy}{dx} = \frac{(t + e^{-t})'}{(1 - e^{-t})'} = \frac{1 - e^{-t}}{e^{-t}} = e^t - 1$$

(g) $\begin{cases} x = \frac{1}{1-t} \\ y = 1 - \ln(1-t) \end{cases}$

$$g) \frac{dy}{dx} = \frac{[1 - \ln(1-t)]'}{(\frac{1}{1-t})'} = \frac{\frac{1}{1-t}}{\frac{1}{(1-t)^2}} = 1-t$$

2. Find $\frac{d^2y}{dx^2}$

(a) $x^2 + y^2 = 25$

$$a) (x^2)' + (y^2)' = (25)' \Rightarrow 2x + 2y \cdot y' = 0 \Rightarrow y' = -\frac{x}{y}$$

$$y'' = -\frac{x'y - xy'}{y^2} = \frac{x(-\frac{x}{y}) - y}{y^2} = \frac{-x^2 - y^2}{y^3} = \frac{-25}{y^3}$$

(b) $\begin{cases} x = t^2 - 1 \\ y = t^4 - 2t^3 \end{cases}$

$$b) \frac{dy}{dx} = \frac{(t^4 - 2t^3)'}{(t^2 - 1)'} = \frac{4t^3 - 6t^2}{2t} = 2t^2 - 3t; \quad \frac{d^2y}{dx^2} = \frac{\frac{d}{dt}(\frac{dy}{dx})}{\frac{dx}{dt}} = \frac{(2t^2 - 3t)'}{2t} = \frac{4t - 3}{2t} = 2 - \frac{3}{2t}$$

(c) $\begin{cases} x = \cos t \\ y = \cos 2t \end{cases}$

$$c) \frac{dy}{dx} = \frac{(\cos 2t)'}{(\cos t)'} = \frac{-\sin 2t (2)}{-\sin t} = \frac{4 \sin t \cos t}{\sin t} = 4 \cos t$$

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dt}(\frac{dy}{dx})}{\frac{dx}{dt}} = \frac{(4 \cos t)'}{(\cos t)'} = \frac{-4 \sin t}{-\sin t} = 4$$

3. In the following eight sub questions, differentiable functions f and g have the values shown in the table.

x	f	f'	g	g'
0	2	1	5	-4
1	3	2	3	-3
2	5	3	1	-2
3	10	4	0	-1

- (a) If $A = f + 2g$, find $A'(3)$

$$A'(x) = f'(x) + 2g'(x)$$

$$\therefore A'(3) = f'(3) + 2g'(3) = 4 + 2(-1) = 2$$

- (c) If $D = \frac{1}{g}$, find $D'(1)$

$$D'(x) = -\frac{1}{(g(x))^2} \cdot g'(x) = -\frac{-3}{3^2} = \frac{1}{3}$$

$$\therefore D' = (g^{-1})' = (-1)g^{-2} \cdot g'$$

- (e) If $K(x) = \left(\frac{f}{g}\right)(x)$, find $K'(0)$

$$K'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

$$K'(0) = \frac{f'(0)g(0) - f(0)g'(0)}{(g(0))^2} = \frac{(1)(5) - (2)(-4)}{5^2} = \frac{13}{25}$$

- (g) If $P(x) = f(x^3)$, find $P'(1)$

$$\therefore P'(x) = f'(x^3)(x^3)' = f'(x^3)(3x^2)$$

$$\therefore P'(1) = f'(1^3)(3(1)^2) = f'(1) \cdot 3 = 2(3) = 6$$

- (b) If $B = f \times g$, find $B'(2)$

$$B'(x) = f'(x)g(x) + f(x)g'(x)$$

$$= (3)(1) + (5)(-2) = -7$$

- (d) If $H(x) = \sqrt{f(x)}$, find $H'(3)$

$$H'(x) = \frac{1}{2}(f(x))^{-\frac{1}{2}}f'(x)$$

$$\therefore H'(3) = \frac{1}{2}[f(3)]^{-\frac{1}{2}}f'(3) = \frac{1}{2}(10)^{-\frac{1}{2}}(4) = \frac{2}{\sqrt{10}} = \frac{\sqrt{10}}{5}$$

- (f) If $M(x) = f(g(x))$, find $M'(1)$

$$M'(x) = f'(g(x))g'(x) \quad \therefore M'(1) = f'(g(1))g'(1)$$

$$= f'(3)(-3)$$

$$= 4(-3) = -12$$

- (h) If $S(x) = f^{-1}(x)$, find $S'(3)$

$$\therefore S'(x) = \frac{1}{f'(x)} \quad \therefore S'(3) = \frac{1}{f'(1)}$$

$$\text{where } x = f(1)$$

$$= \frac{1}{2}$$

4. From the values of f shown in the table below, estimate $f'(2)$.

x	1.92	1.94	1.96	1.98	2.00
$f(x)$	6.00	5.00	4.40	4.10	4.00


$$f'(2) = \lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2} \approx \frac{f(1.98) - f(2)}{1.98 - 2} = \frac{4.10 - 4.00}{-0.02} = \frac{0.1}{-0.02} = -5$$

5. Using the values shown in the table in Q7, estimate $(f^{-1})'(4)$.

$$(f^{-1})'(4) = \frac{1}{f'(2)} \approx \frac{1}{-5}$$

$$\therefore \text{If } b = f(a), \text{ then } (f^{-1})'(b) = \frac{1}{f'(a)}$$

$$y = x^2 - 8x + 10 = x^2 - 2(4)x + 4^2 - 4^2 + 10 = (x-4)^2 - 6$$



6. The “left half” of the parabola defined by $y = x^2 - 8x + 10$ for $x \leq 4$ is a one-to-one function; therefore its inverse is also a function. Call that inverse g . Find $g'(3)$.

$$\text{Let } x^2 - 8x + 10 = 3 \Rightarrow x^2 - 8x + 7 = 0 \Rightarrow (x-1)(x-7) = 0$$

$$\Rightarrow x = 1 \text{ or } x = 7 \quad \because x \leq 4 \quad \therefore x = 1.$$

$$\text{and } f'(x) = (x^2 - 8x + 10)' = 2x - 8; \quad f'(1) = 2(1) - 8 = -6.$$

$$\therefore g'(3) = \frac{1}{f'(1)} = \frac{1}{-6}$$

7. At how many points on the interval $[a, e]$ does the function graphed satisfy the Mean Value Theorem?

