Enhanced dissipation for passive scalars

8 November 2026

Contents

1 Overview **1**
1.1 Map of the wiki . . . 1

I Motivation and history **1**

2 Why it matters **1**

3 History of the problem **1**
3.1 Origins . . . 2
3.2 Spectral methods . . . 2
3.3 Hypocoercivity . . . 2
3.4 Stochastic methods . . . 2
3.5 Hörmander theory . . . 2

4 Recent literature **2**

II Techniques through examples **2**

5 Couette flow **2**
5.1 Setting . . . 2
5.2 The result . . . 2
5.3 One example, every technique . . . 2
5.3.1 Explicit solution . . . 3
5.3.2 Spectral methods . . . 3
5.3.3 Hypocoercivity . . . 3
5.3.4 Stochastic methods . . . 3
5.3.5 Hörmander theory . . . 3
5.4 What this example teaches . . . 3

6 Poiseuille flow **3**
6.1 Setting . . . 3
6.2 The result . . . 3
6.3 What changes compared with Couette flow . . . 3
6.3.1 Explicit solution . . . 4
6.3.2 Spectral methods . . . 4
6.3.3 Hypocoercivity . . . 4
6.3.4 Stochastic methods . . . 4
6.3.5 Hörmander theory . . . 4

III Techniques and general results **4**

7 Spectral methods **4**
7.1 The method . . . 4
7.2 General results . . . 4
7.3 Limitations and comparison . . . 4

8 Hypocoercivity **4**
8.1 The method . . . 5
8.2 General results . . . 5
8.3 Limitations and comparison . . . 5

9 Stochastic methods **5**
9.1 The method . . . 5
9.2 General results . . . 5
9.3 Limitations and comparison . . . 5

10 Hörmander theory **5**
10.1 The method . . . 5
10.2 General results . . . 5
10.3 Limitations and comparison . . . 6

IV Extensions and recent advances **6**

11 State of the art **6**

12 Related results **6**
12.1 Entries . . . 6
12.2 How an entry moves . . . 6

13 Open problems **6**
13.1 Problems . . . 6

14 Raw data **7**
14.1 Entries . . . 7

V Toolkit **7**

15 Notation **7**
15.1 Notation . . . 7
15.2 Definitions . . . 7

16 Standard analysis **7**
16.1 Inequalities . . . 7
16.2 Functional analysis . . . 7

17 Enhanced-dissipation tools **8**
17.1 Tools . . . 8

18 Numerical methods **8**
18.1 How a record works . . . 8
18.2 Records . . . 8
18.2.1 Heat versus shear on the torus . . . 8
18.2.2 Alternating shears with random phases . . . 12
18.3 Solvers . . . 13

19 Lean formalization **13**
19.1 Certificates . . . 13
19.2 How to add a certificate . . . 13

VI About this wiki **13**

20 How to contribute **13**
20.1 Roles . . . 14
20.2 Small corrections . . . 14
20.3 Mathematical contributions . . . 14
20.4 The TeX format . . . 15
20.5 Anatomy of a result . . . 15
20.5.1 Specimen . . . 15
20.6 Labels . . . 16
20.7 Pictures, videos and numerics . . . 16
20.8 Lean . . . 16
20.9 Discussion and conduct . . . 16
20.10Formulas in the wiki . . . 17
20.11The edition . . . 17

21 Contributors **17**
21.1 Editorial board . . . 17
21.2 Associate editors . . . 17
21.3 Reviewers . . . 17
21.4 Community . . . 18

22 Provenance and AI use **18**
22.1 Rules . . . 18
22.2 This version . . . 18
22.3 AI-generated documents . . . 18
22.4 How a result becomes checked . . . 18

23 Literature **19**
23.1 Orientation . . . 19
23.2 General mechanisms and rate limitations . . . 19
23.3 Smooth stationary shears . . . 19
23.4 Circular and Hamiltonian flows and Taylor dispersion . . . 20
23.5 Time dependent shears and localized estimates . . . 20
23.6 Deterministic mixing constructions . . . 21
23.7 Rough velocities and generalized diffusion . . . 21
23.8 Random stirring and transport noise . . . 21
23.9 Inviscid mixing anomalous dissipation and further lower bounds . . . 21

1 Overview

Discuss on Zulip: #overview, topic “Overview”.

[A short introduction to the topic, written as continuous prose: the phenomenon, the equation, why it matters, and what the reader will find in this wiki.]

$$\partial_t f + u \cdot \nabla f = \nu \Delta f, \qquad \nabla \cdot u = 0.$$

Video: `heat-vs-shear.mp4`, shown in the online edition.
Left: the heat equation. Right: the same initial datum carried by the shear flow $u = (\sin y, 0)$, with the same diffusivity. Bottom: the size of the two solutions in time, on a logarithmic scale. [numerics]

Video: `alternating-shears.mp4`, shown in the online edition.
Left: the shear flow $u = (\sin y, 0)$. Right: shears that alternate, horizontal and vertical, each acting for 3 units of time, with random phases. Same initial datum $f_0 = \cos x$, same diffusivity $\nu = 10^{-4}$; both velocity fields have size at most 1. Bottom: the size of the solutions in time, on a logarithmic scale. [numerics]

1.1 Map of the wiki

Diagram: `wiki-map.svg`, shown in the online edition.
The parts of the wiki; click a name to open its page.

Part I
Motivation and history

2 Why it matters

Page team: to be named · Status: skeleton
Discuss on Zulip: #overview, topic “Why it matters”.

[Why the problem matters, in mathematics and in applications, written as the introduction of a paper.]

3 History of the problem

Page team: to be named · Status: skeleton
Discuss on Zulip: #overview, topic “History of the problem”.

[A short account of the progress; then the history technique by technique.]

3.1 Origins

[Before the mathematical theory.]

3.2 Spectral methods

[Contributions and dates; the general theory is on Spectral methods.]

3.3 Hypocoercivity

[Contributions and dates; the general theory is on Hypocoercivity.]

3.4 Stochastic methods

[Contributions and dates; the general theory is on Stochastic methods.]

3.5 Hörmander theory

[Contributions and dates; the general theory is on Hörmander theory.]

4 Recent literature

Page team: to be named · Status: skeleton
Discuss on Zulip: #overview, topic "Recent literature".

[The latest papers, with what each contributes; their results are collected on State of the art and Related results.]

Part II
Techniques through examples

5 Couette flow

Page team: to be named · Status: skeleton
Discuss on Zulip: #examples, topic "Couette flow".

[Lead: what this example shows, in three lines.]

5.1 Setting

[The flow $u = (y, 0)$ on $\mathbb{T} \times \mathbb{R}$: the equation, the Fourier modes, what decays.]

5.2 The result

Theorem 1 (enhanced dissipation for Couette flow)**.** *[The statement.]*

Lean: not yet formalized.

Idea. [The key mechanism in a few lines.]

Proof architecture. [The proof as a short chain of steps, and its diagram.]

5.3 One example, every technique

Each proof below is written for this example only, to show the mechanism; the general results, with complete proofs, are in the part *Techniques and general results*.

5.3.1 Explicit solution

Proof by the explicit solution. [The proof.] □

5.3.2 Spectral methods

Proof by spectral methods. [The proof.] □

General theory: Spectral methods.

5.3.3 Hypocoercivity

Proof by hypocoercivity. [The proof.] □

General theory: Hypocoercivity.

5.3.4 Stochastic methods

Proof by stochastic methods. [The proof.] □

General theory: Stochastic methods.

5.3.5 Hörmander theory

Proof by Hörmander theory. [The proof.] □

General theory: Hörmander theory.

5.4 What this example teaches

[What the example shows, and what the general theory keeps of it.]

6 Poiseuille flow

Page team: to be named · Status: skeleton
Discuss on Zulip: #examples, topic "Poiseuille flow".

[Lead: what changes compared with Couette flow, and why.]

6.1 Setting

[The flow, the domain, the critical point, what decays.]

6.2 The result

Theorem 2 (enhanced dissipation for Poiseuille flow)**.** *[The statement.]*

Lean: not yet formalized.

Idea. [The key mechanism in a few lines.]

Proof architecture. [The proof as a short chain of steps, and its diagram.]

6.3 What changes compared with Couette flow

For each technique: what carries over from Couette flow, what is lost, and the proof for this example.

6.3.1 Explicit solution

[What is lost.]

6.3.2 Spectral methods

Proof by spectral methods. [The proof.] □

6.3.3 Hypocoercivity

Proof by hypocoercivity. [The proof.] □

6.3.4 Stochastic methods

Proof by stochastic methods. [The proof.] □

6.3.5 Hörmander theory

Proof by Hörmander theory. [The proof.] □

Part III
Techniques and general results

7 Spectral methods

Page team: to be named · Status: skeleton
Discuss on Zulip: #techniques, topic "Spectral methods".

[Lead: what the method gives, and under which hypotheses.]

7.1 The method

[The key mechanism, in a few lines.]
Worked example: Couette flow. What changes with a critical point: Poiseuille flow.

7.2 General results

[The theorems, each written as a result (statement, Lean banner, idea, proof architecture, complete proof; see How to contribute); lemmas of general use go to the Toolkit *(Standard analysis, Enhanced-dissipation tools).]*

7.3 Limitations and comparison

[What the method does not give, and how it compares with the other techniques.]

8 Hypocoercivity

Page team: to be named · Status: skeleton
Discuss on Zulip: #techniques, topic "Hypocoercivity".

[Lead: what the method gives, and under which hypotheses.]

8.1 The method

[The key mechanism, in a few lines.]
Worked example: Couette flow. What changes with a critical point: Poiseuille flow.

8.2 General results

[The theorems, each written as a result (statement, Lean banner, idea, proof architecture, complete proof; see How to contribute); lemmas of general use go to the Toolkit *(Standard analysis, Enhanced-dissipation tools).]*

8.3 Limitations and comparison

[What the method does not give, and how it compares with the other techniques.]

9 Stochastic methods

Page team: to be named · Status: skeleton
Discuss on Zulip: #techniques, topic "Stochastic methods".

[Lead: what the method gives, and under which hypotheses.]

9.1 The method

[The key mechanism, in a few lines.]

Worked example: Couette flow. What changes with a critical point: Poiseuille flow.

9.2 General results

[The theorems, each written as a result (statement, Lean banner, idea, proof architecture, complete proof; see How to contribute); lemmas of general use go to the Toolkit *(Standard analysis, Enhanced-dissipation tools).]*

9.3 Limitations and comparison

[What the method does not give, and how it compares with the other techniques.]

10 Hörmander theory

Page team: to be named · Status: skeleton
Discuss on Zulip: #techniques, topic "Hörmander theory".

[Lead: what the method gives, and under which hypotheses.]

10.1 The method

[The key mechanism, in a few lines.]

Worked example: Couette flow. What changes with a critical point: Poiseuille flow.

10.2 General results

[The theorems, each written as a result (statement, Lean banner, idea, proof architecture, complete proof; see How to contribute); lemmas of general use go to the Toolkit *(Standard analysis, Enhanced-dissipation tools).]*

10.3 Limitations and comparison

[What the method does not give, and how it compares with the other techniques.]

Part IV
Extensions and recent advances

11 State of the art

Page team: to be named · Status: skeleton

Discuss on Zulip: #advances, topic “State of the art”.

[The strongest results available to date, in one table.]

Flow class	Domain	Dissipation time	Sharp	Source	Label
—	—	—	—	—	—

12 Related results

Page team: to be named · Status: skeleton
Discuss on Zulip: #advances, topic “Related results”.

Results related to enhanced dissipation and not yet part of the established theory. Each entry gives the statement, a label ([published], [preprint, to be verified] or [raw, unchecked]), the date it was added, its source, and where it would go in the established theory.

12.1 Entries

[First entries.]

12.2 How an entry moves

1. An entry keeps its label until its status changes: raw data checked and posted becomes [preprint, to be verified], a refereed result [published].
2. The editorial board moves established results into the part *Techniques and general results*, or into a new page, with a synthesis proof.

13 Open problems

Page team: to be named · Status: skeleton
Discuss on Zulip: #advances, topic “Open problems”.

Open problems, each with what is known, the expected answer, and the main difficulty.

13.1 Problems

[First problems.]

14 Raw data

Page team: to be named · Status: skeleton
Discuss on Zulip: #advances, topic “Raw data”.

AI-generated proofs, numerical observations and sketches, labelled [raw, unchecked] until a human has checked them. AI-generated text is quoted verbatim, with its source and date; how an entry moves on is explained on Related results.

14.1 Entries

[First entries.]

Part V
Toolkit

15 Notation

Page team: to be named · Status: skeleton
Discuss on Zulip: #toolkit, topic “Notation”.

The notation and the definitions used throughout the wiki.

15.1 Notation

[The scalar, the velocity field, the domains, the Fourier modes, the norms.]

15.2 Definitions

[Dissipation time, enhanced dissipation, and the related notions used in the wiki.]

16 Standard analysis

Page team: to be named · Status: skeleton
Discuss on Zulip: #toolkit, topic “Standard analysis”.

Classical facts of analysis used by the proofs. A statement comes to the Toolkit when several proofs use it, or when it holds in more generality than one result; a step used by one proof only is a claim inside that proof. Each statement is given once, with its proof or a reference, and the proofs elsewhere link to it.

16.1 Inequalities

[Poincaré and interpolation inequalities, and the other inequalities the proofs use.]

16.2 Functional analysis

[Semigroups, spectral theory, and the other facts the proofs use.]

17 Enhanced-dissipation tools

Page team: to be named · Status: skeleton
Discuss on Zulip: #toolkit, topic “Enhanced-dissipation tools”.

Tools specific to enhanced dissipation. A statement comes to the Toolkit when several proofs use it, or when it holds in more generality than one result; a step used by one proof only is a claim inside that proof. Each statement is given once, with its proof or a reference, and the proofs elsewhere link to it.

17.1 Tools

[For instance spectral-gap and resolvent lemmas, and the link between mixing and enhanced dissipation.]

18 Numerical methods

Page team: to be named · Status: skeleton, one record
Discuss on Zulip: #numerics, topic “Numerical methods”.

Every computed picture and video of this wiki has a record on this page: what was computed, how, with which parameters and checks, and the code that produced it.

18.1 How a record works


- A record is a section with an anchor `num-...`; the picture or video links to it with the parameter `record` of Media.
- It gives the equation, the parameters, the method, the checks, the code and how to run it, and its status.
- The code is in the folder `numerics/` of the repository of the wiki, one folder per record, with the output of its checks.

18.2 Records

18.2.1 Heat versus shear on the torus

Produces	the video on the Overview (`heat-vs-shear.mp4`) and the check figure below
Equation	$\partial_t f + \sin y\, \partial_x f = \nu \Delta f$ on $[0, 2\pi)^2$, periodic, and the heat equation $\partial_t f = \nu \Delta f$; same datum $f_0 = \cos x$
Parameters	$\nu = 10^{-3}$, times $0 \le t \le 100$; video of 20 s at 24 frames per second, time running uniformly (5 time units per second), last frame held 1.5 s
Method	one Fourier mode in x: $f = \mathrm{Re}(g(t,y)e^{ix})$ with $\partial_t g = -i \sin y\, g + \nu(\partial_y^2 - 1)g$; 512 Fourier modes in y; Strang splitting with time step 0.01, transport and diffusion steps solved exactly; heat equation solved exactly
Checks	with 1024 modes and time step 0.005 the norms agree to a relative $6 \cdot 10^{-8}$; the energy balance $\|g(t)\|^2 + 2\nu \int_0^t \|(\partial_y g, g)\|^2 ds = \|g(0)\|^2$ holds to $3 \cdot 10^{-7}$; decay rate of the norm on $[50, 100]$: 0.0167, against $\nu^{1/2} \approx 0.0316$
Code	numerics/heat-vs-shear/heat_vs_shear.py in the repository, also below (Python 3, NumPy, Matplotlib, ffmpeg); `python3 heat_vs_shear.py -out ../../wiki/media`, about a minute
Status	computed by Claude (AI) on 8 November 2026; not yet checked by a human [demo text]

Figure: `heat-vs-shear-check.png`, shown in the online edition.



Left: the norms of the two runs and of the heat equation. Right: the relative difference of the two runs and the energy-balance residual.

Code.

```
#!/usr/bin/env python3
"""Heat equation versus advection-diffusion by a shear flow on the torus [0, 2pi)^2.

    d_t f + sin(y) d_x f = nu (d_xx + d_yy) f,      f(0, x, y) = cos x,

compared with the heat equation (the same without the transport term), whose solution is
exp(-nu t) cos x. The initial datum has zero average in x. It is a single Fourier mode in x,
so f(t, x, y) = Re(g(t, y) e^{ix}) with

    d_t g = -i sin(y) g + nu (d_yy - 1) g,      g(0, y) = 1,

solved by a Fourier method in y with Strang splitting: the transport step is the exact
multiplication by exp(-i sin(y) dt), the diffusion step is exact in Fourier variables.
The L2 norm reported is ||f(t)|| / ||f(0)||, that is the root mean square of |g(t, .)|.

Checks (written to heat-vs-shear-check.txt and heat-vs-shear-check.png):
  * the run is repeated with twice the Fourier modes and half the time step, and the norms are
    compared;
  * energy balance: ||g(t)||^2 + 2 nu int_0^t ||(d_y g, g)||^2 ds = ||g(0)||^2 (means over y).

The video lasts --duration seconds; frame s in [0, 1] shows t = T (a s + (1 - a) s^2), a = --
    slow (default 1:
time runs uniformly; a < 1 slows down the start); the last frame is held --hold seconds.

Outputs, in --out: heat-vs-shear.mp4 (video), heat-vs-shear-poster.png (one frame of it),
heat-vs-shear-check.png and heat-vs-shear-check.txt. Needs NumPy, Matplotlib and ffmpeg
(with the libx264 encoder). Run:  python3 heat_vs_shear.py --out ../../wiki/media

Drafted by Claude (AI) on 8 November 2026 for the wiki "Enhanced dissipation for passive
scalars"; record: page Numerical methods, section num-heat-vs-shear.
"""
import argparse
import os
import shutil
import subprocess
import tempfile

import numpy as np
import matplotlib

matplotlib.use("Agg")
import matplotlib.pyplot as plt

HEAT_COLOR = "#b4532a"
SHEAR_COLOR = "#1f4e79"


def solve(nu, T, N, dt, frame_times=()):
    """Times, norms ||f(t)||/||f(0)||, energy-balance residuals, and g(t, y) at frame_times."""
    y = 2 * np.pi * np.arange(N) / N
    eta = np.fft.fftfreq(N, d=1.0 / N)              # integer wavenumbers in y
    symbol = eta ** 2 + 1.0                          # symbol of -(d_yy - 1): mode k = 1 in x
    half = np.exp(-nu * symbol * dt / 2)
    transport = np.exp(-1j * np.sin(y) * dt)
    ghat = np.fft.fft(np.ones(N, dtype=complex))
    steps = int(round(T / dt))
    times = dt * np.arange(steps + 1)
    sq = np.empty(steps + 1)                         # mean over y of |g|^2
    dissip = np.empty(steps + 1)                     # mean over y of |d_y g|^2 + |g|^2
    frames, targets, j = [], list(frame_times), 0
    for n in range(steps + 1):
        power = np.abs(ghat) ** 2 / N ** 2           # Parseval: means over y
        sq[n] = power.sum()
```

```
        dissip[n] = (symbol * power).sum()
        while j < len(targets) and targets[j] <= times[n] + dt / 2:
            frames.append(np.fft.ifft(ghat))
            j += 1
        if n == steps:
            break
        ghat *= half
        ghat = np.fft.fft(np.fft.ifft(ghat) * transport) * half
    integral = np.concatenate([[0.0], np.cumsum((dissip[1:] + dissip[:-1]) / 2) * dt])
    residual = np.abs(sq + 2 * nu * integral - 1.0)
    return times, np.sqrt(sq), residual, frames, y


def field(g, nx):
    """f(x, y) = Re(g(y) e^{ix}) on an nx-by-nx grid (rows: y, columns: x)."""
    step = max(1, len(g) // nx)
    x = 2 * np.pi * np.arange(nx) / nx
    return np.real(g[::step][:, None] * np.exp(1j * x)[None, :])


def draw_frame(path, t, g, nu, times, norms, heat, nx, ymin):
    x = 2 * np.pi * np.arange(nx) / nx
    f_heat = np.exp(-nu * t) * np.cos(x)[None, :] * np.ones((nx, 1))
    f_shear = field(g, nx)
    fig = plt.figure(figsize=(11.2, 7.2), dpi=100)
    gs = fig.add_gridspec(2, 2, height_ratios=[3.6, 1.15], hspace=0.3, wspace=0.06,
                          left=0.08, right=0.97, top=0.845, bottom=0.095)
    for col, (data, title) in enumerate([(f_heat, "Heat equation:  $u = 0$"),
                                         (f_shear, r"Shear flow:  $u = (\sin y,\, 0)$")]):
        ax = fig.add_subplot(gs[0, col])
        ax.imshow(data, origin="lower", extent=[0, 2 * np.pi, 0, 2 * np.pi], cmap="RdBu_r",
                  vmin=-1, vmax=1, interpolation="bilinear")
        ax.set_title(title, fontsize=17, pad=8)
        ax.set_xticks([]); ax.set_yticks([])
        ax.set_aspect("equal")
    fig.suptitle(r"Same initial datum $f_0 = \cos x$, same diffusivity $\nu = 10^{-3}$"
                 + "        t = %d" % round(t), fontsize=17, y=0.975)
    ax = fig.add_subplot(gs[1, :])
    k = np.searchsorted(times, t, side="right")
    ax.semilogy(times, heat, color=HEAT_COLOR, alpha=0.18, lw=2)
    ax.semilogy(times, norms, color=SHEAR_COLOR, alpha=0.18, lw=2)
    ax.semilogy(times[:k], heat[:k], color=HEAT_COLOR, lw=2.4, label="heat equation")
    ax.semilogy(times[:k], norms[:k], color=SHEAR_COLOR, lw=2.4, label="shear flow")
    ax.axvline(t, color="#888888", lw=1)
    ax.set_xlim(0, times[-1]); ax.set_ylim(ymin, 1.5)
    ax.set_yticks([10.0 ** e for e in range(int(np.log10(ymin)), 1)])
    ax.set_xlabel("time  t", fontsize=13)
    ax.set_ylabel(r"$\|f(t)\|\,/\,\|f_0\|$", fontsize=14)
    ax.tick_params(labelsize=11)
    ax.legend(loc="lower left", fontsize=12, frameon=False, ncol=2)
    fig.savefig(path)
    plt.close(fig)


def main():
    ap = argparse.ArgumentParser(description=__doc__.split("\n")[0])
    ap.add_argument("--out", default=".", help="folder for the video and the figures")
    ap.add_argument("--nu", type=float, default=1e-3)
    ap.add_argument("--T", type=float, default=100.0)
    ap.add_argument("--N", type=int, default=512, help="Fourier modes in y")
    ap.add_argument("--dt", type=float, default=0.01)
    ap.add_argument("--duration", type=float, default=20.0, help="length of the video in seconds
    ")
```

```
    ap.add_argument("--fps", type=int, default=24)
    ap.add_argument("--slow", type=float, default=1.0, help="speed of time at the start,
    relative to the mean (1: uniform)")
    ap.add_argument("--hold", type=float, default=1.5, help="seconds the last frame is held")
    ap.add_argument("--nx", type=int, default=256, help="pixels per side of each panel")
    a = ap.parse_args()
    os.makedirs(a.out, exist_ok=True)
    s = np.linspace(0, 1, int(round(a.duration * a.fps)))
    frame_times = a.T * (a.slow * s + (1 - a.slow) * s ** 2)
    times, norms, residual, frames, y = solve(a.nu, a.T, a.N, a.dt, frame_times)
    heat = np.exp(-a.nu * times)
    # check: the same run with N doubled and dt halved
    times2, norms2, residual2, _, _ = solve(a.nu, a.T, 2 * a.N, a.dt / 2)
    diff = np.abs(norms2[::2] - norms) / norms2[::2]
    ymin = 10 ** np.floor(np.log10(norms.min()))
    with open(os.path.join(a.out, "heat-vs-shear-check.txt"), "w") as fh:
        fh.write("nu=%g T=%g N=%d dt=%g frames=%d fps=%d\n" % (a.nu, a.T, a.N, a.dt, len(
    frame_times), a.fps))
        fh.write("final norm: shear %.4e, heat %.4e\n" % (norms[-1], heat[-1]))
        fh.write("max relative difference with (2N, dt/2): %.2e\n" % diff.max())
        fh.write("max energy-balance residual: %.2e (2N, dt/2: %.2e)\n" % (residual.max(),
    residual2.max()))
        fh.write("decay rate of the shear norm on [T/2, T]: %.4f; nu^(1/2) = %.4f\n" % (
            np.log(norms[len(norms) // 2] / norms[-1]) / (times[-1] - times[len(times) // 2]),
    np.sqrt(a.nu)))
    # check figure
    fig, (ax1, ax2) = plt.subplots(1, 2, figsize=(11, 4), dpi=100)
    ax1.semilogy(times, heat, color=HEAT_COLOR, lw=2, label="heat equation (exact)")
    ax1.semilogy(times, norms, color=SHEAR_COLOR, lw=2, label="shear flow, N=%d, dt=%g" % (a.N,
    a.dt))
    ax1.semilogy(times2, norms2, color="k", lw=1, ls="--", label="shear flow, N=%d, dt=%g" % (2
    * a.N, a.dt / 2))
    ax1.set_xlabel("time t"); ax1.set_ylabel("norm ratio"); ax1.legend(frameon=False, fontsize
    =9)
    ax1.set_title("Norms (log scale)")
    ax2.semilogy(times[1:], diff[1:] + 1e-16, color=SHEAR_COLOR, lw=1.5, label="relative
    difference of the two runs")
    ax2.semilogy(times[1:], residual[1:] + 1e-16, color="#6b7280", lw=1.5, label="energy-balance
     residual")
    ax2.set_xlabel("time t"); ax2.legend(frameon=False, fontsize=9); ax2.set_title("Checks")
    fig.tight_layout()
    fig.savefig(os.path.join(a.out, "heat-vs-shear-check.png"))
    plt.close(fig)
    # video
    with tempfile.TemporaryDirectory() as tmp:
        for i, (t, g) in enumerate(zip(frame_times, frames)):
            draw_frame(os.path.join(tmp, "f%04d.png" % i), t, g, a.nu, times, norms, heat, a.nx,
     ymin)
        poster = min(range(len(frame_times)), key=lambda i: abs(frame_times[i] - 0.2 * a.T))
        shutil.copyfile(os.path.join(tmp, "f%04d.png" % poster), os.path.join(a.out, "heat-vs-
    shear-poster.png"))
        subprocess.run(["ffmpeg", "-y", "-loglevel", "error", "-framerate", str(a.fps),
                        "-i", os.path.join(tmp, "f%04d.png"), "-vf", "tpad=stop_mode=clone:
    stop_duration=%g" % a.hold,
                        "-c:v", "libx264", "-pix_fmt", "yuv420p",
                        "-crf", "26", "-preset", "slow", "-movflags", "+faststart",
                        os.path.join(a.out, "heat-vs-shear.mp4")], check=True)
    print(open(os.path.join(a.out, "heat-vs-shear-check.txt")).read().strip())


if __name__ == "__main__":
    main()
```

18.2.2 Alternating shears with random phases

Produces the second video on the Overview (`alternating-shears.mp4`) and the check figure below

Equation $\partial_t f + u \cdot \nabla f = \nu \Delta f$ on $[0, 2\pi)^2$, periodic, with $f_0 = \cos x$ and the alternating shears of Pierrehumbert's model: $u = (\sin(y + \varphi_n), 0)$ for $6n \le t < 6n + 3$ and $u = (0, \sin(x + \psi_n))$ for $6n + 3 \le t < 6n + 6$, the phases independent and uniform on $[0, 2\pi)$; compared with the steady shear $u = (\sin y, 0)$ and with the heat equation. Both velocity fields have size at most 1.

Parameters $\nu = 10^{-4}$, each shear acting for a time 3, times $0 \le t \le 60$, phases from the random generator of NumPy with seed 2026; video of 20 s at 24 frames per second, time running uniformly (3 time units per second), last frame held 1.5 s

Method Fourier pseudo-spectral method on a 512×512 grid; Strang splitting with time step 0.0125, the transport steps (each row or column shifted, through its Fourier coefficients) and the diffusion steps solved exactly; the steady shear as in the record above


Checks with 1024×1024 modes and time step 0.00625 the norms agree to a relative $1.2 \cdot 10^{-6}$; the energy balance $\|f(t)\|^2 + 2\nu \int_0^t \|\nabla f\|^2 ds = \|f_0\|^2$ holds to $3.5 \cdot 10^{-6}$; time for the norm to halve, for $\nu = 10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}$: alternating shears 8.8, 16.3, 23.9, 31.7 (the same with twice the modes), an increase of about 7.6 for each factor 10 in ν; steady shear 8.6, 20.0, 43.9, 94.9; heat equation $\ln 2/\nu$

Context Exponential mixing for randomly shifted alternating shears is proved by Blumenthal, Coti Zelati and Gvalani (2023), and dissipation on the time scale $|\log \nu|$, with high probability, by Cooperman, Iyer and Son (2025); whether the parameters used here satisfy their hypotheses has not been checked

Code numerics/alternating-shears/alternating_shears.py in the repository (Python 3, NumPy, SciPy, Matplotlib, ffmpeg); `python3 alternating_shears.py -out ../../wiki/media`, about 15 minutes

Status computed by Claude (AI) on 8 November 2026; not yet checked by a human [demo text]

Figure: `alternating-shears-check.png`, shown in the online edition.



Left: the norms for $\nu = 10^{-4}$, with two resolutions for the alternating shears. Right: the time for the norm to halve, against $1/\nu$, logarithmic scales.

18.3 Solvers

[The methods used across the records, with their strengths and limits.]

19 Lean formalization

Page team: to be named · Status: skeleton
Discuss on Zulip: #lean, topic "Lean formalization".

A result of this wiki can carry a machine-checked proof in Lean, its *certificate*. A small banner under each theorem says where it stands. A result not yet formalized:
Lean: not yet formalized.

A formalized result, with a link to its certificate:
Lean: formalized.

19.1 Certificates

Result	Status	Certificate	Lean and Mathlib versions
—	—	—	—

19.2 How to add a certificate

[What a certificate contains (repository, commit, versions, the statement as formalized, the axioms used), and how it is checked before its banner turns green.]

Part VI
About this wiki

20 How to contribute

Page team: Michele Dolce · Status: proposal, to be agreed with the editorial board · Text: [demo text]
Discuss on Zulip: #contributing, topic "How to contribute".

This wiki is written from the bottom up, the way a journal works: members of the community propose changes, proposals are discussed and reviewed in the open, an associate editor decides, and accepted changes are written into the wiki. The roles, and who holds them, are on Contributors. [demo text]

20.1 Roles

- **Editorial board**: the scope and the structure of the wiki; appoints the associate editors; settles discussions that end without consensus.
- **Associate editors**: one or more per part, following its Zulip channel. They run the discussion of each proposal, ask reviewers, decide, and write accepted changes into the wiki.
- **Reviewers**: check a proposal at the request of an associate editor, in the open, in its Zulip topic.
- **Community**: propose and discuss. New members are endorsed by an editor or by a member at least two years past the PhD.

The first line of each page names its team. As the wiki grows, busy parts and pages get more associate editors and reviewers; the roles stay the same. On the wiki, everyone with a role is a *member* (reads every page and writes on talk pages); the editorial board and the associate editors are also *editors* (edit articles and templates); administrators give these rights.

20.2 Small corrections

Typos, broken links, a missing reference: members of the wiki leave a note on the talk page of the page (tab *Discussion*, then *Add topic*); others write in the Zulip channel *contributing*, naming the page. The editors go through these notes regularly and make the changes. For now the wiki is open only to invited members.

20.3 Mathematical contributions

A new proof, a shorter argument, a correction to a statement, a new result or a new page goes through Zulip.

1. Download the TeX source of the page: link *TeX source* under the title of every page of the edition, or *Open in Overleaf*. It is a standard LaTeX article that compiles as it is. Change what you want and check that it compiles; then keep only the blocks you changed: a section, or a statement or a proof with its label.

2. Open a topic in the Zulip channel of the part, titled *Proposal: short title*. Attach the file and say in a few lines what it changes, why, and which sources it uses.

3. For two weeks the community and the reviewers discuss the proposal in its topic.

4. Within a week after that, the associate editor decides: accepted, to be revised, or declined. If the discussion ended without consensus, the editorial board decides.

5. A script converts the accepted file to the wiki's format, shows the editor exactly what changes on the page, and flags new citations and links. The associate editor writes the change into the wiki, links the Zulip topic in the edit summary, and the page credits its author.

A proposal changes only the blocks it contains. A paper that is not yet on Literature is proposed on its own, as a small correction of that page.

20.4 The TeX format

The TeX source of a page is a standard LaTeX article: one file that compiles with pdflatex and nothing else (on Overleaf, or with TeXShop, TeXstudio, ...; typeset twice for the citations). Its preamble is the same for every page and is never changed; a contribution defines no macros, since the wiki has none. Sections and statements are numbered in the PDF only.

- Statements: theorem, proposition, lemma, corollary, conjecture, definition, example, remark, claim. The optional argument is the name, the label is the anchor: `\begin{theorem}[name]\label{thm-name}`. Labels start with thm-, prp-, lem-, cor-, def- or clm-.

- After a statement: `\lean{}` (the Lean banner), the environments idea and architecture, and proofs with an optional title and a label, as in `\begin{proof}[Proof by hypocoercivity]\label{proof-name}`. A claim and its proof sit inside the proof that uses them.

- Formulas: `$...$`, `\[ ... \]` and align*; no numbered equations and no labels on equations; none of the commands that the wiki's renderer rejects (the script lists them).

- Citations: `\citet{Key}` gives *Author (year)* and `\citep{Key}` gives *(Author, year)*, with the keys of Literature. The reference list at the end of the file is generated from the reference catalogue; a new citation is added to it when the proposal is accepted.

- Links: on the same page, `\hyperref[thm-name]{the theorem above}` rather than `\ref`, since the wiki has no numbers; to another page, `\wikilink{Page\#anchor}{text}`.
- Labels: `\heuristic{}`, `\numerics{}`, `\published{}`, `\preprint{}`, `\raw{}`.
- A new block goes after an existing one: put the line `%% after: Setting` (a section title or a label) just before it.

20.5 Anatomy of a result

A result is a statement followed by its levels of detail, each folded until the reader opens it:

Statement	in the simplest form, constants hidden
Lean	a one-line banner: formalized or not yet
Idea	the key mechanism in a few lines
Architecture	the proof as a short chain of steps, later with a diagram
Proof	the complete proof, in steps; a statement used only in this proof is a *claim*, with its own folded proof inside

A statement that several proofs use, or that holds in more generality than one result, goes to the *Toolkit* (Standard analysis or Enhanced-dissipation tools) and is linked from the proofs. A short proof is written in one piece. The specimen below shows the boxes at work; open the folds to see the nesting.

20.5.1 Specimen

Theorem 3 (specimen)**.** *The statement, in its simplest form.*

Lean: not yet formalized.

Idea. The key mechanism in a few lines, and the heuristic scaling, labelled [heuristic].

Proof architecture. 1. Step 1 reduces the theorem to a claim.

2. Step 2 proves the claim.
3. Step 3 concludes with a lemma of the toolkit.

Proof. **Step 1.** The reduction.

Claim (specimen)**.** *A statement used only in this proof.*

Proof of the claim. The proof of the claim, folded inside the proof that uses it. □

Step 2. The conclusion, with a lemma of the toolkit.
Sources. This proof follows ... □

20.6 Labels

- [demo text]: text drafted by Claude (AI), until a member of the page team replaces or approves it.
- [heuristic]: a non-rigorous explanation, kept apart from what is proved.
- [numerics]: a computation, with its record on Numerical methods.
- [published], [preprint, to be verified], [raw, unchecked]: the status of an entry on Related results and Raw data: refereed; posted but not refereed; neither posted nor checked by a human (an AI-generated proof, a numerical observation, a sketch).
- Grey quote boxes: AI-generated text, quoted verbatim with its source and date.

20.7 Pictures, videos and numerics

A picture, a video or a diagram is one line in the wiki, the template Media; in TeX, `\media{video}{file}{caption}`. The files are kept in the folder `wiki/media` of the project; the edition shows them, the wiki shows a box with the caption. Every computed picture or video has a record on Numerical methods, with the method, the parameters, the checks and the code.

20.8 Lean

Under each theorem a one-line banner says whether it is formalized: grey while it is not, green with a link to the certificate once it is. Certificates are listed on Lean formalization.

20.9 Discussion and conduct

Discussion takes place on Zulip: one channel per part, one topic per page, and one topic per proposal. The line under the title of each page links to its topic; talk pages on the wiki are for small corrections.

Part	Zulip channel
Overview, Motivation and history	overview
Techniques through examples	examples
Techniques and general results	techniques
Extensions and recent advances	advances
Toolkit	toolkit
Numerical methods	numerics
Lean formalization	lean
About this wiki	contributing

Discuss the mathematics, not the people. Associate editors moderate their channel: they can hide messages and, after a warning, ask the editorial board to remove a member; their decisions can be appealed to the board. The Zulip is not set up yet; once it is, its address goes into Template:Zulip host and the links appear on every page.

20.10 Formulas in the wiki

For editors who write the wiki's own format.

- Inline formulas need `display="inline"`, displayed ones `display="block"`; each formula is written on one line. The renderer rejects `\newcommand`, `\label`, `\eqref` and `\substack`. No formulas in headings.
- Citations link to the Literature page by the key of the reference catalogue: `[[Literature#BedrossianCotiZelati2017|Bedrossian and Coti Zelati (2017)]]`. The section *References* of each page is generated from the catalogue: add the key to its comment line rather than editing the list.
- To link to a result, use the `id` of its box as the anchor, as in `Couette flow#thm-couette`.

20.11 The edition

A static edition of this wiki, with the pictures, the videos and the TeX source of every page, is built from the wiki's text by a script and never edited by hand; its address will be added here.

21 Contributors

Page team: editorial board · Status: to be filled
Discuss on Zulip: #contributing, topic “Contributors”.

The people who write and check this wiki, by role. How they work together is described in How to contribute.

21.1 Editorial board

Decides the scope and the structure of the wiki, appoints the associate editors, and settles a discussion that ends without consensus.
[Names and roles.]

21.2 Associate editors

Each part of the wiki has at least one associate editor, who follows its Zulip channel, runs the discussion of every proposal there, asks reviewers, decides, and writes accepted changes into the wiki. A busy part or page can have several.
[Names, with their parts.]

21.3 Reviewers

Asked by an associate editor to check a proposal: the mathematics, the format, or both. Reviews are posted in the Zulip topic of the proposal. A busy page can have its own reviewers, named in its first line.
[Names.]

21.4 Community

Members of the Zulip, who propose and discuss changes. A new member is endorsed by an editor or by a member at least two years past the PhD. Early-stage researchers (PhD students, and researchers up to two years after the PhD) can choose to be listed as such.
[Names.]

22 Provenance and AI use

Page team: Michele Dolce · Text: [demo text]
Discuss on Zulip: #contributing, topic “Provenance and AI use”.

This page records where each piece of the wiki comes from and who has checked what. The labels are explained on How to contribute. [demo text]

22.1 Rules

- Text drafted by Claude (AI) carries [demo text] until a member of the page team replaces or approves it.
- Arguments produced by AI systems appear only as verbatim quotations, with source and date, or as entries labelled [raw, unchecked] on Raw data.
- The first line of each page records its team, its status and the checks made. A machine check is recorded as such and never counts as a human check.
- Statements attributed to a paper give its theorem number; synthesis proofs say which papers they follow.

- Every accepted proposal is linked, in the edit summary, to the Zulip topic where it was discussed.

22.2 This version

On 8 November 2026 the wiki was reorganized into the parts drawn on the map, and its pages were reduced to a skeleton, to be filled by the editorial board and contributors. The first version (6 November 2026: Couette flow with three proofs, a hypocoercivity page, the toolkit, the history) is kept in the project folder, in `archive/2026-11-06-first-version`. The only computed content is the video on the Overview, with its record on Numerical methods, made by Claude (AI) and not yet checked by a human. The TeX source of each page is generated from the wiki by a script.

22.3 AI-generated documents

Document	Date	Contents	Checks recorded
none	—	—	—

22.4 How a result becomes checked

[To be agreed with the editorial board: how many readers a proof needs before it loses its label, and whether a page records who read which proof.]

23 Literature

Page team: to be named · Status: generated from the reference catalogue (references/catalog.json, checked 2026-11-05)
Discuss on Zulip: #contributing, topic “Literature”.

The papers of the reference catalogue, one line each, grouped by topic. This page is generated by a script from the catalogue; to add or correct an entry, propose the change as a small correction; the catalogue is updated first. Pages cite these entries by their key, for instance Bedrossian and Coti Zelati (2017). Preprints are listed with their arXiv identifier. The list is a working bibliography, not a survey of the field.

23.1 Orientation

1. M. Coti Zelati, G. Crippa, G. Iyer and A. L. Mazzucato, *Mixing in incompressible flows: transport, dissipation, and their interplay*, Notices of the American Mathematical Society **71** (2024), 593–604, arXiv:2308.00358.

2. A. L. Mazzucato, Y. Feng and C. Nobili, *Enhanced dissipation by advection and applications to PDEs*, Applied Mathematical Modelling **150** (2026), Part A, 116310, arXiv:2501.17695.

23.2 General mechanisms and rate limitations

1. P. Constantin, A. Kiselev, L. Ryzhik and A. Zlatoš, *Diffusion and mixing in fluid flow*, Annals of Mathematics **168** (2008), 643–674, arXiv:math/0509663.

2. W. Cooperman, G. Iyer, K. Rowan and S. Son, *Exponentially mixing flows with slow enhanced dissipation*, preprint (2025), arXiv:2507.21305.

3. M. Coti Zelati, M. G. Delgadino and T. M. Elgindi, *On the relation between enhanced dissipation timescales and mixing rates*, Communications on Pure and Applied Mathematics **73** (2020), 1205–1244, arXiv:1806.03258.

4. Y. Feng and G. Iyer, *Dissipation enhancement by mixing*, Nonlinearity **32** (2019), 1810–1851, arXiv:1806.03699.

5. C. Seis, *Bounds on the rate of enhanced dissipation*, Communications in Mathematical Physics **399** (2023), 2071–2081, arXiv:2003.08794.

6. D. Wei, *Diffusion and mixing in fluid flow via the resolvent estimate*, Science China Mathematics **64** (2021), 507–518, arXiv:1811.11904.

23.3 Smooth stationary shears

1. D. Albritton, R. Beekie and M. Novack, *Enhanced dissipation and Hörmander's hypoellipticity*, Journal of Functional Analysis **283** (2022), 109522, arXiv:2105.12308.

2. J. Bedrossian and M. Coti Zelati, *Enhanced dissipation, hypoellipticity, and anomalous small noise inviscid limits in shear flows*, Archive for Rational Mechanics and Analysis **224** (2017), 1161–1204, arXiv:1510.08098.

3. M. Coti Zelati, *Stable mixing estimates in the infinite Péclet number limit*, Journal of Functional Analysis **279** (2020), 108562, arXiv:1909.01310.

4. M. Coti Zelati and T. D. Drivas, *A stochastic approach to enhanced diffusion*, Annali della Scuola Normale Superiore di Pisa, Classe di Scienze **22** (2021), 811–834, arXiv:1911.09995.

5. M. Coti Zelati and T. Gallay, *Enhanced dissipation and Taylor dispersion in higher-dimensional parallel shear flows*, Journal of the London Mathematical Society **108** (2023), 1358–1392, arXiv:2108.11192.

23.4 Circular and Hamiltonian flows and Taylor dispersion

1. M. Beck, O. Chaudhary and C. E. Wayne, *Analysis of enhanced diffusion in Taylor dispersion via a model problem*, Hamiltonian Partial Differential Equations and Applications, Fields Institute Communications **75** (2015), pp. 31–71, arXiv:1501.05877.

2. M. Beck, O. Chaudhary and C. E. Wayne, *Rigorous justification of Taylor dispersion via center manifolds and hypocoercivity*, Archive for Rational Mechanics and Analysis **235** (2020), 1105–1149, arXiv:1804.06916.

3. E. Bruè, M. Coti Zelati and E. Marconi, *Enhanced dissipation for two-dimensional Hamiltonian flows*, Archive for Rational Mechanics and Analysis **248** (2024), Article 84, arXiv:2211.14057.

4. M. Coti Zelati and M. Dolce, *Separation of time-scales in drift-diffusion equations on R^2*, Journal de Mathématiques Pures et Appliquées **142** (2020), 58–75, arXiv:1907.04012.

5. M. Coti Zelati, M. Dolce and C. Lo, *Diffusion enhancement and Taylor dispersion for rotationally symmetric flows in discs and pipes*, Journal of Mathematical Fluid Mechanics **26** (2024), Article 12, arXiv:2305.18162.

6. M. Dolce, C. J. P. Johansson and M. Sorella, *Dissipation enhancing properties for a class of Hamiltonian flows with closed streamlines*, Communications in Partial Differential Equations **50** (2025), 301–352, arXiv:2407.06884.

7. Y. Feng, A. L. Mazzucato and C. Nobili, *Enhanced dissipation by circularly symmetric and parallel pipe flows*, Physica D: Nonlinear Phenomena **445** (2023), 133640, doi:10.1016/j.physd.2022.133640.

8. G. Iyer and H. Zhou, *Quantifying the dissipation enhancement of cellular flows*, SIAM Journal on Mathematical Analysis **55** (2023), 6496–6516, arXiv:2209.11645.

9. T. Li and L. Zhang, *Enhanced dissipation and Taylor dispersion by a parallel shear flow in an infinite cylinder with unbounded cross section*, preprint (2025), arXiv:2510.13097.

23.5 Time dependent shears and localized estimates

1. J. Bedrossian, S. He, S. Iyer and F. Wang, *Pseudo-Gevrey smoothing for the passive scalar equations near Couette*, Journal of Functional Analysis **289** (2025), 110987, arXiv:2405.19233.

2. J. Benthaus, G. M. Coclite and C. Nobili, *Mixing and enhanced dissipation in a time-translating shear flow*, preprint (2026), arXiv:2603.14624.

3. J. Benthaus and C. Nobili, *Enhanced dissipation via time-modulated velocity fields*, Evolution Equations and Control Theory **15** (2026), 21–50, arXiv:2501.16905.

4. D. Coble and S. He, *A note on enhanced dissipation of time-dependent shear flows*, Communications in Mathematical Sciences **22** (2024), 1685–1700, arXiv:2309.15738.

5. S. He, *Localized Enhanced Dissipation: A Hypocoercivity Approach*, preprint (2026), arXiv:2603.14657.

23.6 Deterministic mixing constructions

1. T. M. Elgindi, K. Liss and J. C. Mattingly, *Optimal enhanced dissipation and mixing for a time-periodic, Lipschitz velocity field on T2*, Duke Mathematical Journal **174** (2025), 1209–1260, arXiv:2304.05374.

23.7 Rough velocities and generalized diffusion

1. M. Colombo, M. Coti Zelati and K. Widmayer, *Mixing and diffusion for rough shear flows*, Ars Inveniendi Analytica, Paper No. **2** (2021), 21 pp., arXiv:2009.12268.

2. Y. Feng and A. L. Mazzucato, *Dissipation enhancement by shear flows for generalized diffusion operators in high dimension*, Communications in Mathematical Sciences **23** (2025), 2203–2215, doi:10.4310/CMS.250930031923.

3. L. Galeati and M. Gubinelli, *Mixing for generic rough shear flows*, SIAM Journal on Mathematical Analysis **55** (2023), 7240–7272, arXiv:2107.12115.

4. S. He, *Enhanced dissipation, hypoellipticity for passive scalar equations with fractional dissipation*, Journal of Functional Analysis **282** (2022), 109319, arXiv:2103.07906.

5. M. Romito and L. Roveri, *Instantaneous Total Enhanced Dissipation For Very Rough Shear Flows*, preprint (2025), arXiv:2510.27331.

23.8 Random stirring and transport noise

1. J. Bedrossian, A. Blumenthal and S. Punshon-Smith, *Almost-sure enhanced dissipation and uniform-in-diffusivity exponential mixing for advection-diffusion by stochastic Navier-Stokes*, Probability Theory and Related Fields **179** (2021), 777–834, arXiv:1911.01561.

2. A. Blumenthal, M. Coti Zelati and R. S. Gvalani, *Exponential mixing for random dynamical systems and an example of Pierrehumbert*, Annals of Probability **51** (2023), 1559–1601, arXiv:2204.13651.

3. W. Cooperman, G. Iyer and S. Son, *A Harris theorem for enhanced dissipation, and an example of Pierrehumbert*, Nonlinearity **38** (2025), 045027, arXiv:2403.19858.

4. M. Coti Zelati, M. Hairer and D. Villringer, *A Stochastic RAGE Theorem and Enhanced Dissipation for Transport Noise*, preprint (2025), arXiv:2507.11422.

5. A. Del Zotto and M. Nualart, *Exponential mixing and enhanced dissipation on the unit sphere with Rossby-Haurwitz flows*, preprint (2026), arXiv:2606.11055.

6. F. Flandoli and D. Luo, *Enhanced dissipation and Lyapunov exponents for stochastic transport-diffusion equations with small molecular diffusivity and small noise intensity*, Stochastics and Partial Differential Equations: Analysis and Computations **14** (2026), 1062–1076, doi:10.1007/s40072-026-00426-y.

23.9 Inviscid mixing anomalous dissipation and further lower bounds

1. G. Alberti, G. Crippa and A. L. Mazzucato, *Exponential self-similar mixing by incompressible flows*, Journal of the American Mathematical Society **32** (2019), 445–490, arXiv:1605.02090.

2. G. Crippa, T. Elgindi, G. Iyer and A. L. Mazzucato, *Growth of Sobolev norms and loss of regularity in transport equations*, Philosophical Transactions of the Royal Society A **380** (2022), 20210024, arXiv:2109.14975.

3. T. D. Drivas, T. M. Elgindi, G. Iyer and I. Jeong, *Anomalous dissipation in passive scalar transport*, Archive for Rational Mechanics and Analysis **243** (2022), 1151–1180, arXiv:1911.03271.

4. T. M. Elgindi and A. Zlatoš, *Universal mixers in all dimensions*, Advances in Mathematics **356** (2019), 106807, arXiv:1809.09614.

5. T. M. Elgindi and K. Liss, *Norm growth, non-uniqueness, and anomalous dissipation in passive scalars*, Archive for Rational Mechanics and Analysis **248** (2024), arXiv:2309.08576.

6. G. Iyer, A. Kiselev and X. Xu, *Lower bounds on the mix norm of passive scalars advected by incompressible enstrophy-constrained flows*, Nonlinearity **27** (2014), 973–985, arXiv:1310.2986.

7. C. Nobili and S. Pottel, *Lower bounds on mixing norms for the advection diffusion equation in Rd*, Nonlinear Differential Equations and Applications NoDEA **29** (2022), Article 12, arXiv:2006.04614.

References

A. Blumenthal, M. Coti Zelati and R. S. Gvalani, *Exponential mixing for random dynamical systems and an example of Pierrehumbert*, Annals of Probability **51** (2023), 1559–1601, arXiv:2204.13651.

W. Cooperman, G. Iyer and S. Son, *A Harris theorem for enhanced dissipation, and an example of Pierrehumbert*, Nonlinearity **38** (2025), 045027, arXiv:2403.19858.