

# Impact of model resolution on the representation of deep water formation and its link with the AMOC


**Eneko Martin-Martinez<sup>1,2</sup>, Eduardo Moreno-Chamarro<sup>1</sup>, Pablo Ortega<sup>1</sup>, Amanda Frigola<sup>1</sup>, Bernat Jiménez-Esteve<sup>1</sup>, Daria Kuznetsova<sup>1</sup>, Aude Carréric<sup>1</sup>, Saskia Loosveldt-Tomas<sup>1</sup>, Pierre-Antoine Bretonnière<sup>1</sup>, Margarida Samsó Cabré<sup>1</sup>**

1: Barcelona Supercomputing Center (BSC)

2: Departament de Dinàmica de la Terra i l'Oceà, Facultat de Ciències de la Terra, Universitat de Barcelona (UB)


## Preliminary results

- Mixing region is shifted to Labrador Sea when increasing the model resolution
- Surface wind-stress is the main mixing driver
- *EC-Earth3P-VHR* shows no propagation of the mixing imprint in the *MOC* to lower latitudes




**Figure 1:** Representation of the ORCA grids used in the EC-Earth3P configurations. Each dot represents 10x10 grid points. Each figure shows one of the regions we have used: subpolar gyre box (left, red box), North Atlantic box (middle, blue box), and Atlantic basin (right, green area).


Argo



EN4-v4.2.2



## Mixed layer depth (MLD)



**Figure 2:** MLD time-average.

## Methods

- 1950-control experiments
- HighResMIP protocol
- 76 years (removed the first 30 years with model drift)
- 95 % of confidence (non-significant regions masked).

## Discussion

- First PC of  $\sigma_2$  profile is related to the first *MLD* series, around 0.95 correlation coefficient in all the cases
- Positive/negative NAO induced wind-stress enhances/diminishes the mixing in the *EC-Earth3P-VHR* and *EC-Earth3P-HR* models
- Influence of westward propagated salinity anomalies in *EC-Earth3P-VHR*
- Propagation downward and southward of the density anomalies forced by deep-water mixing
- Different responses on the *AMOC* to the deep-water-mixing: *EC-Earth3P* shows a deeper propagation which arrives to low latitudes, *EC-Earth3P-HR* shows propagation to middle latitudes, *EC-Earth3P-VHR* shows only local changes


## Ongoing and future analyses

- Continue exploring the impacts of deep-water mixing in the three different resolutions
- Study the impact of Greenland land-ice melting in the deep-water mixing and the *MOC*
- Explore other climate impacts related to the mesoscale processes and deep-water mixing


## Contact

eneko.martin@bsc.es


linkedin.com/in/eneko-martin




**Figure 3:** First and second EOFs of density profiles for  $\sigma_2$  average in the subpolar gyre box (see Figure 1).




**Figure 4:** Correlation of the *MLD* average in the subpolar gyre box (see Figure 1) with  $\tau_x$  (top row) and SSS (bottom row).



**Figure 5:** Correlation of the *MLD* average in the subpolar gyre box (see Figure 1) with  $\sigma_2$  lagged one year at 500 m (top row) and 1000 m (bottom row).



**Figure 6:** Correlation of the 3-year low-pass filtered *MLD* average in the subpolar gyre box (see Figure 1) with 3-year low-pass filtered and lagged *MOC* removing the Ekman transport in the Atlantic basin (see Figure 1). No-lag (top row) and lags of two years, four years, six years with the *MOC* lagging (second row, third row, bottom row).



**Figure 7:** Correlation of the yearly *maxMOC* without the Ekman transport at 45° with the yearly *maxMOC* without the Ekman transport from 10°N to 60°N.



**Barcelona Supercomputing Center**  
Centro Nacional de Supercomputación

